

# UC Santa Barbara

## UC Santa Barbara Previously Published Works

### Title

Dynamic social balance and convergent appraisals via homophily and influence mechanisms

### Permalink

<https://escholarship.org/uc/item/6r95h8sb>

### Authors

Mei, Wenjun  
Cisneros-Velarde, Pedro  
Chen, Ge  
[et al.](#)

### Publication Date

2019-12-01

### DOI

10.1016/j.automatica.2019.108580

Peer reviewed



# Dynamic social balance and convergent appraisals via homophily and influence mechanisms<sup>☆</sup>

Wenjun Mei<sup>a,\*</sup>, Pedro Cisneros-Velarde<sup>b</sup>, Ge Chen<sup>d</sup>, Noah E. Friedkin<sup>b,c</sup>, Francesco Bullo<sup>b</sup>

<sup>a</sup> Automatic Control Laboratory, ETH Zurich, Switzerland

<sup>b</sup> Center of Control, Dynamical Systems and Computation, University of California, Santa Barbara, USA

<sup>c</sup> Department of Sociology, University of California, Santa Barbara, USA

<sup>d</sup> Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing, China

## ARTICLE INFO

### Article history:

Received 25 October 2017

Received in revised form 20 August 2019

Accepted 11 September 2019

Available online 24 September 2019

### Keywords:

Structural balance

Multi-agent systems

Homophily/Influence mechanisms

Nonlinear network dynamics

## ABSTRACT

Social balance theory describes allowable and forbidden configurations of the topologies of signed directed social appraisal networks. In this paper, we propose two discrete-time dynamical systems that explain how an appraisal network converges to social balance from an initially unbalanced configuration. These two models are based on two different socio-psychological mechanisms, respectively: the homophily mechanism and the influence mechanism. Our main theoretical contribution is a comprehensive analysis for both models in three steps. First, we establish the well-posedness and bounded evolution of the interpersonal appraisals. Second, we fully characterize the set of equilibrium points; for both models, each equilibrium network is composed of an arbitrary number of complete subgraphs satisfying structural balance. Third, we establish the equivalence among three distinct properties: non-vanishing appraisals, convergence to all-to-all appraisal networks, and finite-time achievement of social balance. In addition to theoretical analysis, Monte Carlo validations illustrate how the non-vanishing appraisal condition holds for generic initial conditions in both models. Moreover, a numerical comparison between the two models indicates that the homophily-based model might be a more universal explanation for the emergence of social balance. Finally, adopting the homophily-based model, we present numerical results on the mediation and globalization of local conflicts, the competition for allies, and the asymptotic formation of a single versus two factions.

© 2019 Elsevier Ltd. All rights reserved.

## 1. Introduction

**Motivation and problem description.** Social systems involving friendly/antagonistic relationships are often modeled as signed networks. *Social balance* (also referred to as *structural balance*) theory, which originated from several seminal works by Heider (1944, 1946), characterizes the stable configurations of signed social networks, summarized as the famous Heider's axioms: "Friends' friends are friends; Friends' enemies are enemies; Enemies' friends are enemies; Enemies' enemies are friends". Empirical studies for both large-scale networks (Facchetti, Iacono, & Altafini, 2011; Leskovec, Huttenlocher, & Kleinberg, 2010) and

small groups (Harary, 1961; King, 1964; Taylor, 1970) indicate that social balance is a type of stable configurations frequently observed in real social networks. Dynamic social balance theory, aiming to explain how an initially unbalanced network evolves to a balanced state, has recently attracted much interest. Despite recent progress, it remains a valuable open problem to propose dynamic models that enjoy desirable boundedness and convergence properties. Such models make it possible to further study meaningful predictions and control strategies for the evolution of social networks to balance.

In this paper, we propose two novel discrete-time dynamic social balance models, in which a group of individuals repeatedly update their interpersonal appraisals via two socio-psychological mechanisms, respectively: the homophily mechanism and the influence mechanism. Loosely speaking, for the homophily mechanism, the interpersonal appraisals of any two individuals in a social group are adjusted based on whether they agree on the appraisals of the group members. For the influence mechanism, each individual assigns influence to others proportionally to her/his appraisal of them. Both mechanisms are well established in the social sciences literature, e.g., see the seminal work by Lazarsfeld and Merton (1954), and the award-winning book

<sup>☆</sup> This material is based upon work supported by, or in part by, the U.S. Army Research Laboratory and the U.S. Army Research Office under grant number W911NF-15-1-0577. The material in this paper was not presented at any conference. This paper was recommended for publication in revised form by Associate Editor Julien M. Hendrickx under the direction of Editor Christos G. Cassandras.

\* Corresponding author.

E-mail addresses: [wmei@ethz.ch](mailto:wmei@ethz.ch) (W. Mei), [pacisneros@ece.ucsb.edu](mailto:pacisneros@ece.ucsb.edu) (P. Cisneros-Velarde), [cheng@amss.ac.cn](mailto:cheng@amss.ac.cn) (G. Chen), [friedkin@soc.ucsb.edu](mailto:friedkin@soc.ucsb.edu) (N.E. Friedkin), [bullo@engineering.ucsb.edu](mailto:bullo@engineering.ucsb.edu) (F. Bullo).

by Friedkin and Johnsen (2011), respectively. For both models, we characterize their sets of equilibrium and their dynamical behavior. Moreover, we compare these two models via both theoretical analysis and numerical comparisons and give a tentative answer that, compared to the influence mechanism, the homophily mechanism is a more universal explanation for the evolution of appraisal networks to social balance.

*Literature review.* Following the early works by Heider (1944, 1946), static social balance theory has been extensively studied in the last seven decades, including the characterization of the balanced configurations for both complete networks (Cartwright & Harary, 1956; Harary, 1953) and arbitrary networks (de Nooy, 1999; Easley & Kleinberg, 2010); the measure of the degree of balance (Cartwright & Gleason, 1966; Henley, Horsfall, & De Soto, 1969); the clustering and its relation to balance (Davis, 1967; Doreian & Mrvar, 1996); as well as the relevant partitioning algorithms (Doreian & Krackhardt, 2001; Kim & Candan, 2012). Numerous empirical studies have been conducted for different social systems, including social systems at the national level (Harary, 1961; McDonald & Rosecrance, 1985), at the group level (King, 1964; Rawling & Friedkin, 2017), and at the individual level (Facchetti et al., 2011; Taylor, 1970). For a comprehensive review we refer to Zheng, Zeng, and Wang (2015).

In the last decade, researchers have started to incorporate dynamical systems into the social balance theory, aiming to explain how a signed network evolves to a structurally balanced state. Early works include the discrete-time *local triad dynamics* (LTD) (Antal, Krapivsky, & Redner, 2005) and *constrained triad dynamics* (Antal, Krapivsky, & Redner, 2006). These models suffer from the existence of unbalanced equilibria, i.e., the *jammed states*. Other works based on network games are proposed by van de Rijt (2011) and Malekzadeh, Fazli, Jalaly Khalidabadi, Rabiee, and Safari. In all the aforementioned models, the link weights in the signed networks only take values from the set  $\{-1, 0, 1\}$ .

Our models are closely related to the continuous-time dynamic social balance models (Kuřakowski, Gawroński, & Gronek, 2005; Marvel, Kleinberg, Kleinberg, & Strogatz, 2011; Traag, Van Dooren, & De Leenheer, 2013), in which the link weights can take arbitrary real values. The model proposed by Kuřakowski et al. (2005) is based on an influence-like mechanism. Theoretical analysis by Marvel et al. (2011) reveals that for symmetric initial conditions, the probability of achieving social balance in finite time tends to 1 as the network size tends to infinity. Traag et al. (2013) extend the set of initial conditions to normal matrices and provide a sufficient condition for finite-time social balance. In Traag et al. (2013), the authors also propose an alternative continuous-time model based on a homophily mechanism, and prove that the homophily-based model leads to finite-time social balance for generic initial conditions. In addition to theoretical analysis, Kuřakowski et al. (2005) investigate numerically the relation between the formation of factions and the initial appraisal distribution, for the influence-like model. The corresponding results for the homophily-based model is unavailable in previous literature. A non-negligible shortcoming of all the models mentioned above is that the interpersonal appraisals diverge to infinity in finite time. To remedy this shortcoming, in Kuřakowski et al. (2005), the authors impose a predetermined upper bound of the interpersonal appraisals. As the consequence, the magnitudes of all the appraisals converge to the predetermined upper bound, see the rigorous analysis in Wongkaew, Caponigro, Kuřakowski, and Borzi (2015). In addition to those continuous-time models, Jia, Friedkin, and Bullo (2016) propose a discrete-time model, with a generalized notion of social balance and a modified influence mechanism, and establish its convergence to the generalized balance.

*Contributions.* The contribution of this paper are manifold. Our paper is the first to propose two well-behaved discrete-time models that explain the evolution of interpersonal appraisal networks towards the classic Heider's social balance, via the homophily and the influence mechanisms, respectively. Both mechanisms are cast in the language of influence systems; indeed the key novelty is the formulation of appropriate influence matrices such that both models are well-behaved and enjoy the desirable properties of bounded evolution and convergent appraisals.

Regarding the theoretical analysis, we first fully characterize the two models' respective equilibrium sets, each of which turn out to include all possible balanced configurations in terms of sign pattern. Second, we establish the equivalence relations among the non-vanishing appraisal condition, the convergence of appraisal networks to all-to-all balanced configurations, and the achievement of social balance in finite time.

Numerical study of our both models leads to various insightful results. First, Monte-Carlo validations indicate that the non-vanishing appraisal condition holds for generic initial conditions, while, for the influence-based model, the non-vanishing appraisal condition holds almost surely if the initial appraisals satisfy some generalized notion of symmetry. Second, further simulation results show that, for the influence-based model with generic initial conditions, the probability that the appraisal network converges to social balance monotonically decays to 0 as the network size tends to infinity. Based on this observation, we conclude that the homophily-based model might be a more universal explanation than the influence-based model for the evolution to social balance. Third, for the homophily-based model, we numerically investigate its behavior under perturbation when the appraisal network is composed of multiple structurally balanced subnetworks. Such numerical study reveals some insightful and realistic interpretations such as the escalation and mediation of local conflicts. Finally, we study by simulation the effect of the initial appraisal distribution on the formation of factions, i.e., whether an appraisal network converges to two antagonistic factions or an all-friendly network.

The main advantage of our models, compared with the previous continuous-time models (Kuřakowski et al., 2005; Traag et al., 2013), is that our models are well-behaved, in the sense that our models enjoy the desirable property of convergent appraisals (as opposed to the undesirable property of finite-time divergence). The convergence property makes it possible to characterize the systems' fixed points and their stability, as well as the transition from one equilibrium to another. In our models, the convergent appraisals are due to the introduction of either homophily or interpersonal influence networks, which also provide a connection between the field of dynamic social balance and the field of opinion dynamics with antagonistic interactions, e.g. Altafini (2013). In addition, our models have the desired property that they are invariant under scaling, i.e., if a solution is scaled by a constant, it remains a solution. This feature is particularly important in the modeling of social systems, in which quantities are usually meaningful only in the relative sense. Compared with the model proposed in Kuřakowski et al. (2005) with bounded evolution, our models do not rely on any predetermined bound to prevent divergence and the asymptotic appraisals in our models are determined by the initial condition rather than the manually determined bound. Some additional advantages of our models are discussed in Section 5.1.

*Organization.* Section 2 introduces some notations and basic concepts. Section 3 and 4 contain the theoretical analyses of our models. Section 5 provides further discussions and numerical results. Section 6 gives the conclusion. An auxiliary lemma is provided in Appendix. Some proofs are provided in the technical report (Mei, Cisneros-Velarde, Friedkin, & Bullo) with full details.

**Table 1**  
Notations frequently used in this paper.

$\mathbb{1}_n$ ( $0_n$ )	The all-ones (all-zeros) $n \times 1$ vector
$\mathbb{R}$ ( $\mathbb{Z}_{\geq 0}$ )	Set of real numbers (non-negative integers)
$>$ ( $<$ )	Entry-wise greater than (less than)
$ X $	Entry-wise absolute value of matrix $X$
$\text{sign}(X)$	Entry-wise sign of $X$ , i.e., $\text{sign}(X)_{ij} = 1$ if $X_{ij} > 0$ , $\text{sign}(X)_{ij} = -1$ if $X_{ij} < 0$ , and $\text{sign}(X)_{ij} = 0$ if $X_{ij} = 0$ .
$ X _{\max}$	The max norm of $X$ , i.e., $\max_{i,j}  X_{ij} $
$X_{i*}$ ( $X_{*i}$ )	the $i$ th row (column) vector of $X$
$G(X)$	Weighted digraph associated with adjacency matrix $X$ . We allow negative link weights. That is, if $X_{ij} < 0$ , then there exists a link in $G(X)$ from $i$ to $j$ with negative weight $X_{ij}$ .

**2. Notations and basic concepts**

*Notations.* Some frequently used notations are defined in Table 1. The following sets will be used throughout this paper:

$$\mathcal{S}_{\text{nz-row}} = \{X \in \mathbb{R}^{n \times n} \mid \text{for every } i, X_{i*} \neq 0_n^T\}, \tag{1}$$

$$\mathcal{S}_{\text{s-symm}}^+ = \{X \in \mathbb{R}^{n \times n} \mid \text{sign}(X) = \text{sign}(X)^T \text{ and } X_{ij} > 0 \text{ for every } i\}, \tag{2}$$

$$\mathcal{S}_{\text{rs-symm}}^+ = \{X \in \mathcal{S}_{\text{s-symm}}^+ \mid \text{there exists } \gamma > 0_n \text{ such that } \text{diag}(\gamma)X = (\text{diag}(\gamma)X)^T\}. \tag{3}$$

By definition,  $\mathcal{S}_{\text{rs-symm}}^+ \subset \mathcal{S}_{\text{s-symm}}^+ \subset \mathcal{S}_{\text{nz-row}}$ . In addition,  $\mathcal{S}_{\text{s-symm}}^+$  and  $\mathcal{S}_{\text{rs-symm}}^+$  are both invariant under permutations. That is, given any  $X \in \mathcal{S}_{\text{s-symm}}^+$  (or  $X \in \mathcal{S}_{\text{rs-symm}}^+$  resp.) and a permutation matrix  $P$ , we have  $PXP^T \in \mathcal{S}_{\text{s-symm}}^+$  (or  $PXP^T \in \mathcal{S}_{\text{rs-symm}}^+$  resp.).

*Appraisal matrices and social balance.* Given a group of  $n$  agents, the interpersonal appraisals are given by the appraisal matrix  $X \in \mathbb{R}^{n \times n}$ . The sign of  $X_{ij}$  determines whether  $i$ 's appraisal of  $j$  is positive, i.e.,  $i$  "likes"  $j$ , or negative, i.e.,  $i$  "dislikes"  $j$ . The magnitude of  $X_{ij}$  represents the intensity of the sentiment. When  $X_{ij} = 0$ , the appraisal is one of indifference. The diagonal entry  $X_{ii}$  represents agent  $i$ 's self-appraisal. The weighted digraph  $G(X)$  associated to  $X$  as the adjacency matrix is referred to as the appraisal network.

**Definition 2.1** (Social Balance (Harary, 1953; Heider, 1946)). An appraisal network  $G(X)$  satisfies social balance, or, equivalently, is structurally balanced, if the appraisal matrix  $X$  satisfies the following properties: (S1)  $X_{ii} > 0$  for any  $i \in \{1, \dots, n\}$ ; (S2)  $\text{sign}(X_{ij}) \text{sign}(X_{jk}) \text{sign}(X_{ki}) = 1$  for any  $i, j, k \in \{1, \dots, n\}$ .

According to Harary (1953), a structurally balanced appraisal network either has only one faction in which the interpersonal appraisals are all positive, or is composed of two antagonistic factions such that individuals in the same faction positively appraise each other while all the inter-faction appraisals are negative.

**Lemma 2.2** (Equivalent Conditions for Social Balance). For any  $X \in \mathbb{R}^{n \times n}$  such that all of its entries are non-zero,  $G(X)$  satisfies social balance if and only if it satisfies (S1) in Definition 2.1 and (S3):  $\text{sign}(X_{i*}) = \pm \text{sign}(X_{j*})$ , for all  $i, j \in \{1, \dots, n\}$ . Moreover, for  $G(X)$  satisfying social balance,  $X$  is sign-symmetric, i.e.,  $\text{sign}(X) = \text{sign}(X)^T$ .

**Proof.** Suppose that (S1) and (S3) hold. For any  $i, j \in \{1, \dots, n\}$ ,  $\text{sign}(X_{i*}) = \delta \text{sign}(X_{j*})$ , where  $\delta$  is either  $-1$  or  $1$ . Therefore,  $\text{sign}(X_{ij}) \text{sign}(X_{ji}) = \delta^2 \text{sign}(X_{ij}) \text{sign}(X_{ii}) = 1$ , i.e.,  $\text{sign}(X_{ij}) = \text{sign}(X_{ji})$ . Moreover, for any  $k$ , since  $\text{sign}(X_{ij}) = \delta \text{sign}(X_{jj})$  and  $\text{sign}(X_{jk}) = \delta \text{sign}(X_{ik})$ , we have

$$\begin{aligned} \text{sign}(X_{ij}) \text{sign}(X_{jk}) \text{sign}(X_{ki}) \\ = \delta^2 \text{sign}(X_{jj}) \text{sign}(X_{ik}) \text{sign}(X_{ki}) = 1. \end{aligned}$$

Therefore, (S1) and (S3) imply (S1) and (S2) in Definition 2.1, as well as the sign symmetry of  $X$ .

Now suppose (S1) and (S2) in Definition 2.1 hold. The sign symmetry of  $X$  is obtained by letting  $k = j$  in (S2). Moreover, due to the sign symmetry and (S2), we obtain  $\text{sign}(X_{ij}) \text{sign}(X_{jk}) \text{sign}(X_{ik}) = 1$ . Therefore,  $\text{sign}(X_{ik}) \text{sign}(X_{jk})$  does not depend on  $k$  and is equal to  $\text{sign}(X_{ij}) \in \{-1, 1\}$ . That is,  $\text{sign}(X_{i*}) = \pm \text{sign}(X_{j*})$  for any  $i$  and  $j$ . This concludes the proof.  $\square$

**3. Homophily-based model**

In this and the next section, we propose and analyze two dynamic social balance models, respectively. These two models are distinct in the microscopic individual interaction mechanisms.

**Definition 3.1** (Homophily-based Model). Given an initial appraisal matrix  $X(0) \in \mathcal{S}_{\text{s-symm}}^+ \subset \mathbb{R}^{n \times n}$ , the homophily-based model is defined by

$$X(t+1) = \text{diag}(|X(t)|\mathbb{1}_n)^{-1}X(t)X^T(t). \tag{4}$$

**Remark 3.2** (Interpretation). Eq. (4) updates the appraisals based on what can be considered as the homophily mechanism. For any  $i, j \in \{1, \dots, n\}$ , agent  $i$ 's appraisal of agent  $j$  at time step  $t+1$  depends on to what extent they are in agreement with each other on the appraisals of all the agents in the group. For any  $k \in \{1, \dots, n\}$ , if  $\text{sign}(X_{ik}(t)) = \text{sign}(X_{jk}(t))$ , then the term  $X_{ik}(t)X_{jk}(t)$  contributes positively to  $X_{ij}(t+1)$ , and vice versa. The matrix  $W(X(t)) = \text{diag}(|X(t)|\mathbb{1}_n)^{-1}X(t)$  can be regarded as the influence matrix constructed from the appraisals through homophily mechanism. Since  $X_{ij}(t+1) = \sum_k W_{ik}(t)X_{jk}(t)$ , each  $|W_{ik}(t)|$  represents how much weight individual  $i$  assigns to the agreement on the appraisal of individual  $k$ . Note that the entry-wise absolute value, i.e.,  $|W(t)|$ , is row-stochastic. Such type of influence matrices has been widely studied in the opinion dynamics with antagonism, see Hendrickx (2014), Proskurnikov and Cao (2017) and Xia, Cao, and Johansson (2016).

The proposition below presents some useful results on the finite-time behavior of the homophily-based model.

**Proposition 3.3** (Invariant Set and Finite-Time Behavior of HbM). Consider the dynamical system (4) and define  $f_{\text{homophily}}(X) = \text{diag}(|X|\mathbb{1}_n)^{-1}XX^T$ . Pick  $X_0 \in \mathcal{S}_{\text{nz-row}}$ . The following statements hold:

- (i) the map  $f_{\text{homophily}}$  is well-defined for any  $X \in \mathcal{S}_{\text{nz-row}}$  and maps  $\mathcal{S}_{\text{nz-row}}$  to  $\mathcal{S}_{\text{s-symm}}^+$ ;
- (ii) the solution  $X(t)$ ,  $t \in \mathbb{Z}_{\geq 0}$ , to Eq. (4) from initial condition  $X(0) = X_0$  exists and is unique;
- (iii) the max norm of any solution  $X(t)$  satisfies

$$|X(t+1)|_{\max} \leq |X(t)|_{\max} \leq |X(0)|_{\max};$$

- (iv) for any  $c > 0$ , the trajectory  $cX(t)$  is the solution to Eq. (4) from initial condition  $X(0) = cX_0$ .

**Proof.** For simplicity, denote  $X^+ = f_{\text{homophily}}(X)$ . For any  $X \in \mathcal{S}_{\text{nz-row}}$ , since, for any  $i$  and  $j$ ,  $X_{ij}^+ = \frac{1}{\|X_{i*}\|_1} \sum_k X_{ik}X_{jk}$  and  $\|X_{i*}\|_1 > 0$ ,  $f_{\text{homophily}}(X)$  is well-defined. Moreover,

$$X_{ii}^+ = \frac{1}{\|X_{i*}\|_1} \sum_k X_{ik}X_{ik} = \frac{\|X_{i*}\|_2^2}{\|X_{i*}\|_1} > 0, \quad \text{and}$$

$$X_{ij}^+ = \frac{\|X_{j*}\|_1}{\|X_{i*}\|_1} X_{ji}^+, \quad \text{for any } i \text{ and } j.$$

Therefore,  $f_{\text{homophily}}$  maps  $\mathcal{S}_{\text{nz-row}}$  to  $\mathcal{S}_{\text{s-symm}}^+$ . This concludes the proof of statement (i). Statement (ii) is a direct consequence of



statement (i), since, for any  $t \in \mathbb{Z}_{\geq 0}$ ,  $X(t) \in \mathcal{S}_{\text{nz-row}}$  defines a unique  $X(t+1) = f_{\text{homophily}}(X(t)) \in \mathcal{S}_{\text{s-symm}}^+$ . In addition,

$$\begin{aligned} |X_{ij}^+| &\leq \frac{1}{\|X_{i*}\|_1} \sum_{k=1}^n |X_{ik}X_{jk}| \leq \frac{1}{\|X_{i*}\|_1} \sum_{k=1}^n |X_{ik}| |X_{jk}| \\ &\leq \max_k |X_{jk}| \leq |X|_{\max} \end{aligned}$$

immediately leads to statement (iii). Finally, statement (iv) is obtained by replacing  $X(t)$  with  $cX(t)$  on the right-hand side of Eq. (4).  $\square$

According to statement (iii) of Proposition 3.3, for any  $a > 0$ , the set  $\mathcal{S}_{\text{nz-row}} \cap [-a, a]^{n \times n}$  is positively invariant under dynamics (4). This desired bounded-evolution property makes our model substantially different from some previous models, in which  $X(t)$  diverges in finite time (Marvel et al., 2011; Traag et al., 2013).

The theorem below characterizes the set of fixed points of system (4), i.e., the steady-state appraisal matrix  $X$  satisfying  $X = f_{\text{homophily}}(X)$ . Fixed points are sociologically interesting because they correspond to the states that can often be observed in the real world.

**Theorem 3.4 (Fixed Points and Balance).** Consider the dynamical system (4) in domain  $\mathcal{S}_{\text{nz-row}}$ . Define

$$\begin{aligned} Q_{\text{homophily}} &= \left\{ PYP^T \in \mathcal{S}_{\text{nz-row}} \mid P \text{ is a permutation matrix,} \right. \\ &\quad \left. Y \text{ is a block diagonal matrix with blocks of} \right. \\ &\quad \left. \text{the form } \alpha bb^T, \alpha > 0, b \in \{-1, +1\}^m, m \leq n \right\}. \end{aligned}$$

Then

- (i)  $Q_{\text{homophily}}$  is the set of all the fixed points of (4).
- (ii) for any  $X \in Q_{\text{homophily}}$ ,  $G(X)$  is composed by isolated complete subgraphs that satisfy social balance.

**Proof.** We first prove that any  $X^* \in Q_{\text{homophily}}$  is a fixed point of system (4). For any  $\alpha > 0$  and  $b \in \{-1, +1\}^n$ , the matrix  $Y = \alpha bb^T$  satisfies

$$f_{\text{homophily}}(Y) = \text{diag}(n\alpha \mathbb{1}_n)^{-1} \alpha^2 bb^T bb^T = \alpha bb^T = Y.$$

This arguments extend to block diagonal matrices  $Y$ . By the definition of  $f_{\text{homophily}}$ , for any block diagonal matrix  $Y = \text{diag}(Y^{(1)}, \dots, Y^{(K)})$ ,  $Y = f_{\text{homophily}}(Y)$  if and only if  $Y^{(i)} = \text{diag}(|Y^{(i)}| \mathbb{1}_n)^{-1} Y^{(i)} Y^{(i)T}$  for any  $i$ . Therefore,  $Y$  is a fixed point of system (4) if each  $Y^{(i)}$  in  $Y = \text{diag}(Y^{(1)}, \dots, Y^{(K)})$  is a  $n_i \times n_i$  matrix of the form  $\alpha_i b^{(i)} b^{(i)T}$ , with  $\alpha_i > 0$ ,  $b^{(i)} \in \{-1, +1\}^{n_i}$ , and  $n_1 + \dots + n_K = n$ . Moreover, given any fixed point  $Y$ , for any permutation matrix  $P \in \mathbb{R}^{n \times n}$ ,

$$\begin{aligned} PYP^T &= P \text{diag}(|Y| \mathbb{1}_n)^{-1} Y Y^T P^T \\ &= \text{diag}(|PYP^T| \mathbb{1}_n)^{-1} (PYP^T) (PYP^T)^T \\ &= f_{\text{homophily}}(PYP^T). \end{aligned}$$

Therefore, any  $X^* \in Q_{\text{homophily}}$  is a fixed point of (4).

Now we prove by induction that  $Q_{\text{homophily}}$  is the set of all the fixed points of system (4). For the trivial case of  $n = 1$ ,  $Q_{\text{homophily}}$  represents the set of all the positive scalars and one can easily check that any positive scalar  $X$  is a fixed point of system (4) with  $n = 1$ . Suppose statement (i) holds for any system with dimension  $\tilde{n} < n$ . For system (4) with dimension  $n$ , suppose  $X$  is a fixed point, i.e.,  $X = f_{\text{homophily}}(X)$ . For any  $i, j \in \{1, \dots, n\}$ , by comparing the  $(i, j)$ -th and the  $(j, i)$ -th equations of  $X = f_{\text{homophily}}(X)$ , we conclude that  $X_{ij}$  and  $X_{ji}$  always have the same sign. In addition, since  $X_{ii} = \sum_{k=1}^n X_{ik}^2 / \|X_{i*}\|_1$ , we have  $X_{ii} > 0$  for

any  $i$ . Since  $X$  is a fixed point of  $f_{\text{homophily}}$ , we have that, for any  $i, j \in \{1, \dots, n\}$ ,

$$\begin{aligned} |X_{ij}| &= \frac{1}{\|X_{i*}\|_1} \left| \sum_k X_{ik} X_{jk} \right| \\ &\leq \frac{1}{\|X_{i*}\|_1} \sum_k |X_{ik}| |X_{jk}| \leq |X|_{\max}. \end{aligned}$$

Moreover, there exists  $(i, j)$  such that  $|X_{ij}| = |X|_{\max}$ . For any such  $(i, j)$ , either of the following two cases hold:

Case 1:  $i = j$  and there does not exist  $k \neq i$  such that  $|X_{ik}| = |X|_{\max}$ . In this case,  $|X_{ii}| = |X|_{\max}$ . Since

$$\begin{aligned} |X_{ii}| &= \frac{1}{\|X_{i*}\|_1} \left| \sum_k X_{ik} X_{ik} \right| \\ &\leq \frac{1}{\|X_{i*}\|_1} \sum_k |X_{ik}| |X_{ik}| \leq |X|_{\max}, \end{aligned}$$

in order for  $|X_{ii}| = |X|_{\max}$  to hold,  $X_{i*}$  must satisfy  $|X_{ik}| = |X|_{\max}$ , for any  $k$  such that  $X_{ik} \neq 0$ . By the definition of Case 1, we conclude that there does not exist  $k \neq i$  such that  $X_{ik} \neq 0$ . Therefore, there exists a permutation matrix  $P$  such that

$$PXP^T = \begin{bmatrix} |X|_{\max} & \mathbb{0}_{n-1}^T \\ \mathbb{0}_{n-1} & \tilde{X}_{(n-1) \times (n-1)} \end{bmatrix}.$$

Since  $PXP^T$  is also a fixed point of system (4), one can check that  $\tilde{X}$  satisfies  $\tilde{X} = \text{diag}(|\tilde{X}| \mathbb{1}_n)^{-1} \tilde{X} \tilde{X}^T$ . Therefore,  $\tilde{X}$  is a fixed point of system (4) with dimension  $n-1$ . Since we have assumed that statement (i) holds for dimension  $\tilde{n} < n$ , there exists an  $(n-1) \times (n-1)$  permutation matrix  $\tilde{P}$  and a block diagonal  $\tilde{Y}$ , with blocks of the form  $\alpha bb^T$ , where  $\alpha > 0$ ,  $b \in \{-1, +1\}^m$ ,  $m < n-1$ , such that  $\tilde{X} = \tilde{P} \tilde{Y} \tilde{P}^T$ . Therefore,

$$X = P^T \begin{bmatrix} 1 & \mathbb{0}_{n-1}^T \\ \mathbb{0}_{n-1} & \tilde{P} \end{bmatrix} \begin{bmatrix} |X|_{\max} & \mathbb{0}_{n-1}^T \\ \mathbb{0}_{n-1} & \tilde{Y} \end{bmatrix} \begin{bmatrix} 1 & \mathbb{0}_{n-1}^T \\ \mathbb{0}_{n-1} & \tilde{P} \end{bmatrix}^T P.$$

The matrix  $P^T \begin{bmatrix} 1 & \mathbb{0}_{n-1}^T \\ \mathbb{0}_{n-1} & \tilde{P} \end{bmatrix}$  is also a permutation matrix. Therefore  $X \in Q_{\text{homophily}}$ .

Case 2:  $j \neq i$  and  $|X_{ij}| = |X|_{\max}$ . We first define some notations used in the following proof: For any  $k$ , let  $\theta_k = \{\ell \mid X_{k\ell} \neq 0\}$  and  $|\theta_k|$  be the cardinality of the set  $\theta_k$ . Note that, since  $X = f_{\text{homophily}}(X) \in \mathcal{S}_{\text{s-symm}}^+$ ,  $k$  is always in  $\theta_k$  and  $X_{kk} > 0$ . Let  $X_{\ell*, \theta_k} \in \mathbb{R}^{1 \times |\theta_k|}$  be the  $\ell$ -th row vector of  $X$  with all the  $X_{\ell p}$  entries such that  $p \notin \theta_k$  is removed.

We point out a general result that, for any  $k$  and  $\ell$ , if

$$|X_{k\ell}| = \frac{1}{\|X_{k*}\|_1} \left| \sum_{p=1}^n X_{kp} X_{\ell p} \right| = |X|_{\max},$$

then, for the second equality to hold,  $X$  must satisfy that: (1)  $\theta_k \subset \theta_\ell$ ; (2)  $|X_{\ell p}| = |X|_{\max}$  for any  $p \in \theta_k$ ; (3)  $\text{sign}(X_{\ell*, \theta_k}) = \pm \text{sign}(X_{k*, \theta_k})$ . Therefore, for the  $i, j$  indexes such that  $|X_{ij}| = |X|_{\max}$  and  $i \neq j$ , we have:  $|X_{jk}| = |X|_{\max}$ , for any  $k \in \theta_i$ ;  $\theta_i \subset \theta_j$ ; and  $\text{sign}(X_{j*, \theta_i}) = \pm \text{sign}(X_{i*, \theta_i})$ . Since  $i \in \theta_i$  and  $X = f_{\text{homophily}}(X)$ , we obtain  $|f_{\text{homophily}}(X)_{ji}| = |X_{ji}| = |X|_{\max}$ . Therefore,  $|f_{\text{homophily}}(X)_{ik}| = |X_{ik}| = |X|_{\max}$ , for any  $k \in \theta_j$ , and  $\theta_j \subset \theta_i$ , which in turn leads to  $\theta_i = \theta_j$  and  $|X_{ik}| = |X|_{\max}$  for any  $k \in \theta_i$ . Therefore, for any  $k \in \theta_i$ ,  $|f_{\text{homophily}}(X)_{ik}| = |X|_{\max}$ , which implies  $|X_{k\ell}| = |X|_{\max}$  for any  $\ell \in \theta_i$ . Since  $|f_{\text{homophily}}(X)_{k\ell}| = |X_{k\ell}|$ , we further obtain that  $\theta_k \subset \theta_\ell$  and  $\text{sign}(X_{k*, \theta_k}) = \pm \text{sign}(X_{\ell*, \theta_k})$ . Moreover, due to the fact that the indexes  $k$  and  $\ell$  are interchangeable, we conclude that, for any  $k, \ell \in \theta_i$ : (a)  $\theta_k = \theta_\ell = \theta_i$ ; (b)  $|X_{k\ell}| = |X|_{\max}$ ; (c)  $\text{sign}(X_{k*}) = \pm \text{sign}(X_{\ell*})$ .

If  $|\theta_i| = n$ , let  $\alpha = X_{11}$  and  $b = \text{sign}(X_{1*})^T$ , then we have  $X = \alpha bb^T$ . If  $|\theta_i| < n$ , there exists a permutation matrix  $P$  such

that

$$PXP^T = \begin{bmatrix} X^{(\theta_i)} & \mathbb{0}_{|\theta_i| \times (n-|\theta_i|)} \\ \mathbb{0}_{(n-|\theta_i|) \times |\theta_i|} & \tilde{X} \end{bmatrix},$$

where  $X^{(\theta_i)}$  is a  $|\theta_i| \times |\theta_i|$  matrix. Moreover,  $X^{(\theta_i)} = |X|_{\max} bb^T$ , where  $b = \text{sign}(X_{i\cdot, \theta_i})^T$ . Following the same line of argument for Case 1, we know that  $\tilde{X}$  is of the form  $\tilde{P}\tilde{Y}\tilde{P}^T$  and thereby  $X \in Q_{\text{homophily}}$ . This concludes the proof for statement (i).

For any  $X^* \in Q_{\text{homophily}}$ , there exists a permutation matrix  $P$  and a block diagonal matrix  $Y = \text{diag}(Y^{(1)}, \dots, Y^{(K)})$  such that  $X^* = PYP^T$ . Note that  $G(Y)$  has exactly the same topology as  $G(X)$ , but with the nodes re-indexed. Therefore, we only need to analyze the structure of  $G(Y)$ . The graph  $G(Y)$  is made up of  $K$  isolated complete subgraphs and  $Y^{(i)} = \alpha_i b^{(i)} b^{(i)T}$  for each such subgraph  $G(Y^{(i)})$ , where  $b^{(i)} = (b_1^{(i)}, \dots, b_{n_i}^{(i)})^T$ . Therefore, according to Lemma 2.2, each subgraph  $G(Y^{(i)})$  satisfies social balance. This concludes the proof for statement (ii). □

**Remark 3.5 (Social Balance with Multiple Isolated Subgraphs).** An appraisal matrix  $X \in Q_{\text{homophily}}$  can be a block-diagonal matrix  $\text{diag}(X_1, \dots, X_k)$  and thus corresponds to an appraisal network  $G(X)$  composed of  $k$  isolated subgraphs, each of which satisfies social balance as in Definition 2.1. With the notion of social balance extended to graphs with multiple isolated subgraphs, in terms of sign pattern, the set of fixed points  $X$  of the homophily-based model (4) corresponds to exactly the set of all the possible structurally balanced configurations of the appraisal network  $G(X)$ . Such characterization of fixed points is impossible in the previous continuous-time models (Marvel et al., 2011; Traag et al., 2013) since those models diverge in finite time. Moreover, for any  $X \in Q_{\text{homophily}}$  such that  $G(X)$  has  $k$  isolated subgraphs,  $X$  is a rank- $k$  matrix.

Before presenting the main results on the convergence of the appraisal matrix  $X(t)$  to social balance, we define a property of  $X(t)$  as the solution to Eq. (4).

**Definition 3.6 (Non-vanishing Appraisal Condition).** A solution  $X(t)$  satisfies the non-vanishing appraisal condition if  $\liminf_{t \rightarrow \infty} \min_{i,j} |X_{ij}(t)| > 0$ .

**Theorem 3.7 (Convergence and Social Balance in HbM).** Consider the homophily-based model given by Eq. (4). The following statements hold:

- (i) Each element in  $Q_{\text{homophily}}$  of rank one is a locally stable fixed point of  $f_{\text{homophily}}$ ;
- (ii) For any  $X(0) \in S_{\text{nz-row}}$ , the following three statements are equivalent:
  - (a) the solution  $X(t)$  satisfies the non-vanishing appraisal condition;
  - (b) there exists  $t_0 > 0$  such that  $G(X(t))$  satisfies social balance for all  $t \geq t_0$ ;
  - (c) there exists  $X^* \in Q_{\text{homophily}}$  of rank one such that  $\lim_{t \rightarrow \infty} X(t) = X^*$ .

**Proof.** For simplicity of notations, let  $|X|_{\min} = \min_{k,l} |X_{k,l}|$ . We start by proving the following two claims. For any given  $t_0 \geq 0$ , if all the entries of  $X(t_0)$  are non-zero and  $G(X(t_0))$  satisfies social balance, then,

(C.1) for any  $t \geq t_0$ ,  $G(X(t))$  satisfies social balance and  $\text{sign}(X(t)) = \text{sign}(X(t_0))$ ;

(C.2) for any  $t \geq t_0$ ,  $|X(t)|_{\max}$  is non-increasing and  $|X(t)|_{\min}$  is non-decreasing.

To prove claim (C.1), it suffices to prove that  $G(X(t_0 + 1))$  satisfies social balance and  $\text{sign}(X(t_0 + 1)) = \text{sign}(X(t_0))$ , as the cases for  $t \geq t_0 + 1$  follow by induction. For any  $i$  and  $j$ , since  $G(X(t_0))$  satisfies social balance, according to Lemma 2.2, we have  $\text{sign}(X_{i\cdot, \theta_i}(t_0)) = \pm \text{sign}(X_{j\cdot, \theta_j}(t_0))$ . In addition, we have  $X_{ij}(t_0) > 0$  for any  $j$ . Therefore,

$$\begin{aligned} \text{sign}(X_{ij}(t_0 + 1)) &= \text{sign}\left(\frac{1}{\|X_{i\cdot}(t_0)\|_1} \sum_{k=1}^n X_{ik}(t_0) X_{jk}(t_0)\right) \\ &= \text{sign}(X_{ij}(t_0) X_{jj}(t_0)) = \text{sign}(X_{ij}(t_0)), \end{aligned}$$

for any  $i$  and  $j$ . This concludes the proof for claim (C.1). For any  $t \geq t_0$ , since  $G(X(t))$  satisfies social balance,

$$|X_{ij}(t + 1)| = \frac{1}{\|X_{i\cdot}(t)\|_1} \sum_{k=1}^n |X_{ik}(t)| |X_{jk}(t)| \text{ for any } i, j,$$

we have  $|X(t + 1)|_{\min} \geq |X(t)|_{\min} \geq |X(t_0)|_{\min}$  and  $|X(t + 1)|_{\max} \leq |X(t)|_{\max} \leq |X(t_0)|_{\max}$ .

Now we prove statement (i), i.e., each  $X^* \in Q_{\text{homophily}}$  with rank 1 is locally stable. Let  $X^* = \alpha bb^T$ , where  $\alpha > 0$  and  $b \in \{-1, +1\}^n$ . For any matrix  $\Delta \in \mathbb{R}^{n \times n}$  such that  $|\Delta|_{\max} = \zeta < \alpha$ , we have  $\text{sign}(X^* + \Delta) = \text{sign}(X^*)$ . Due to claim (C.1) and (C.2), we know that, for  $X(0) = X^* + \Delta$ ,  $X(t)$  satisfies that, for any  $t \geq 0$ : (1)  $\text{sign}(X(t)) = \text{sign}(X(0)) = \text{sign}(X^*)$ ; (2)  $\alpha - \zeta \leq |X(t)|_{\min} \leq |X(t)|_{\max} \leq \alpha + \zeta$ . Therefore, for any  $i$  and  $j$ ,  $X_{ij}(t)$  is of the form  $\alpha_{ij}(t) \text{sign}(X_{ij}^*)$ , where  $0 < \alpha - \zeta \leq \alpha_{ij}(t) \leq \alpha + \zeta$ . We thereby have

$$\begin{aligned} |X(t) - X^*|_{\max} &= \max_{ij} |\alpha_{ij}(t) \text{sign}(X_{ij}^*) - \alpha \text{sign}(X_{ij}^*)| \\ &= \max_{ij} |\alpha_{ij}(t) - \alpha| \leq \zeta. \end{aligned}$$

Therefore, for any  $\epsilon > 0$ , there exists  $\zeta = \min\{\frac{\epsilon}{2}, \frac{\epsilon}{2}\}$  such that, for any  $X(0)$  satisfying  $|X(0) - X^*|_{\max} < \zeta$ ,  $|X(t) - X^*|_{\max} < \epsilon$  for any  $t \geq 0$ , i.e.,  $X^*$  is locally stable.

Now we prove (ii)(a)  $\Rightarrow$  (ii)(b). We first establish the convergence of the solution  $X(t)$  to some set of structurally balanced states via the LaSalle invariance principle. For simplicity, denote  $X^+ = f_{\text{homophily}}(X)$ . The map  $f_{\text{homophily}}(X)$  is continuous for any  $X \in S_{\text{s-symm}}^+$  and, by Proposition 3.3, for any given  $X(0) \in S_{\text{s-symm}}^+$ ,  $|X(t)|_{\max} \leq |X(0)|_{\max}$  for any  $t \in \mathbb{Z}_{\geq 0}$ . In addition, letting  $\delta = \liminf_{t \rightarrow \infty} \min_{i,j} |X_{ij}(t)| > 0$ , we see that there exists  $\tilde{t} \in \mathbb{Z}_{\geq 0}$  such that  $\min_{i,j} |X_{ij}(t)| \geq \delta/2$  for any  $t \geq \tilde{t}$ . Therefore, the set

$$G_c = \left\{ X \in S_{\text{s-symm}}^+ \mid \begin{aligned} &\min_{i,j} |X_{ij}| \geq \delta/2, \\ &|X|_{\max} \leq |X(0)|_{\max} \end{aligned} \right\}$$

is a compact subset of  $S_{\text{s-symm}}^+$  and  $X(t) \in G_c$  for any  $t \geq \tilde{t}$ . Thirdly, define  $V(X) = |X|_{\max}$ . The function  $V$  is continuous on  $S_{\text{s-symm}}^+$  and, by Proposition 3.3, satisfies  $V(X^+) - V(X) \leq 0$  for any  $X \in S_{\text{s-symm}}^+$ . According to the extended LaSalle invariance principle in Theorem 2 of Mei and Bullo,  $X(t)$  converges to the largest invariant set  $M$  of the set  $E = \{X \in G_c \mid V(X^+) - V(X) = 0\}$ .

Now we characterize the largest invariant set  $M$ . For any  $X \in M \subset E$ ,  $V(X^+) = V(X) = |X|_{\max}$ . Suppose  $|X_{ij}^+| = \max_{k,\ell} |X_{k\ell}^+|$ . Since  $X^+ = f_{\text{homophily}}(X)$ , we have

$$|X_{ij}^+| \leq \frac{1}{\|X_{i\cdot}\|_1} \sum_{\ell=1}^n |X_{i\ell}| |X_{j\ell}| \leq |X|_{\max}. \tag{5}$$

In order for all these inequalities to hold with equality and noticing that  $|X_{i\ell}| > 0$  for any  $\ell$  since  $X \in G_c$ ,  $X$  must satisfy that

- (a)  $X_{i*}$  and  $X_{j*}$  have the same or opposite sign pattern, i.e.,  $\text{sign}(X_{i*}) = \pm \text{sign}(X_{j*})$ ,
- (b) All entries of  $X_{j*}$  have the magnitude  $|X|_{\max}$ .

Therefore, for any  $X \in E$ , there exist some  $i$  and  $j$  such that the aforementioned conditions (a) and (b) hold. Moreover, since the set  $M$  is invariant,  $X \in M$  implies  $X^+ \in M \subset E$ . Applying Condition (b) to  $X^+$ , there exists a  $\tilde{j}$  such that, for any  $p$ ,  $|X_{jp}^+| = |X^+|_{\max} = |X|_{\max}$ . In order for  $|X_{jp}^+| = |X|_{\max}$  to hold, following the same argument on the conditions such that the inequalities (5) become equalities, we know that, for any  $p$ ,  $\text{sign}(X_{j*}) = \pm \text{sign}(X_{p*})$  and  $|X_{pk}| = |X|_{\max}$  for any  $k$ . As these relationships hold for any  $p$ , we conclude that for any  $i, j \in \{1, \dots, n\}$ ,  $X_{i*}$  and  $X_{j*}$  must have the same or the opposite sign pattern. Let  $\alpha = |X|_{\max}$  and  $b = \text{sign}(X_{1*}^\top)$ . Each row of  $X$  is thereby equal to either  $\alpha b^\top$  or  $-\alpha b^\top$ . Therefore,  $X$  is of the form  $X = \alpha c b^\top$ , where  $c \in \{-1, 1\}^n$ . Moreover, since all the diagonal entries of  $X$  are positive, the column vector  $c$  satisfies  $c_i b_i = 1$  for any  $i$ , which implies  $c = b$ . In short, we have proved that  $X \in M$  leads to  $X = \alpha b b^\top$ . In addition, by Theorem 3.4, any matrix  $X = \alpha b b^\top$ , with  $\alpha > 0$  and  $b \in \{-1, 1\}^n$ , is a fixed point of  $f_{\text{homophily}}$  and is thus invariant. Therefore, we conclude the compactness of

$$M = \left\{ X = \alpha b b^\top \mid \frac{\delta}{2} \leq \alpha \leq |X(0)|_{\max}, b \in \{-1, 1\}^n \right\}.$$

For any  $\hat{X} \in M$ , since  $\hat{X}$  satisfies social balance (see Theorem 3.4) and  $\min_{i,j} |\hat{X}_{ij}| \geq \delta/2 > 0$ , there exists an open neighbor set defined as  $\mathcal{U}(\hat{X}) = \{X = \hat{X} + \Delta \mid |\Delta|_{\max} < \min_{i,j} |\hat{X}_{ij}|\}$  such that any  $X \in \mathcal{U}(\hat{X})$  satisfies social balance. According to the Heine–Borel theorem, there exists a finite set  $\{\hat{X}_1, \dots, \hat{X}_k\} \subset M$  such that  $M \subset \cup_{k=1}^k \mathcal{U}(\hat{X}_k)$ . Since  $\cup_{k=1}^k \mathcal{U}(\hat{X}_k)$  is an open set, there exists  $\epsilon > 0$  such that the neighbor set of  $M$ , defined as  $\mathcal{U}(M, \epsilon) = \{X \in S_{\text{symm}}^+ \mid |X - M|_{\max} < \epsilon\}$ , satisfies that  $\mathcal{U}(M, \epsilon) \subset \cup_{k=1}^k \mathcal{U}(\hat{X}_k)$  and thereby any  $X \in \mathcal{U}(M, \epsilon)$  satisfies social balance.

Since  $X(t) \rightarrow M$  as  $t \rightarrow \infty$ , there exists  $t_0 \in \mathbb{Z}_{\geq 0}$  such that  $X(t) \in \mathcal{U}(M, \epsilon)$  for any  $t \geq t_0$ . Therefore,  $X(t)$  satisfies social balance for any  $t \geq t_0$ , which concludes the proof for (ii)(a)  $\Rightarrow$  (ii)(b).

Now we prove (ii)(b)  $\Rightarrow$  (ii)(c). Suppose  $G(X(t_0))$  satisfies social balance for some  $t_0 > 0$ . If  $|X(t_0)|_{\max} = |X(t_0)|_{\min}$ , then there exists some  $\alpha > 0$  such that  $X(t_0) = \alpha B$ , where  $B \in \{-1, 1\}^{n \times n}$ . Since  $G(X(t_0))$  satisfies social balance, we have  $B_{ii} > 0$  and  $B_{j*} = \pm B_{1*}$ , which in turn implies that  $B = B_{1*}^\top B_{1*} = \text{sign}(X_{1*}(t_0))^\top \text{sign}(X_{1*}(t_0))$ . Therefore,  $X(t_0)$  is already a rank-one fixed point in the set  $Q_{\text{homophily}}$ .

Suppose  $G(X(t_0))$  satisfies social balance but  $|X(t_0)|_{\max} > |X(t_0)|_{\min}$ . For any  $t \geq t_0$ , let  $|X_{pq}(t)| = |X(t)|_{\min}$ . We have that, for any  $i$  and  $j$ ,

$$\begin{aligned} |X_{jp}(t+1)| &= \frac{1}{\|X_{j*}(t)\|_1} \sum_{k=1}^n |X_{jk}(t)| |X_{pk}(t)| \\ &\leq \frac{|X_{jq}(t)|}{\|X_{j*}(t)\|_1} |X_{pq}(t)| + \left(1 - \frac{|X_{jq}(t)|}{\|X_{j*}(t)\|_1}\right) |X(t)|_{\max} \\ &\leq |X(t)|_{\max} - \frac{|X_{jq}(t)|}{\|X_{j*}(t)\|_1} (|X(t)|_{\max} - |X(t)|_{\min}) \\ &\leq |X(t)|_{\max} - \frac{|X(t)|_{\min}}{n |X(t)|_{\max}} (|X(t)|_{\max} - |X(t)|_{\min}), \end{aligned}$$

and, similarly,

$$\begin{aligned} |X_{ij}(t+2)| &= \frac{1}{\|X_{i*}(t+1)\|_1} \sum_{k=1}^n |X_{ik}(t+1)| |X_{jk}(t+1)| \\ &\leq \frac{|X_{ip}(t+1)|}{\|X_{i*}(t+1)\|_1} |X_{jp}(t+1)| \end{aligned}$$

$$\begin{aligned} &+ \left(1 - \frac{|X_{ip}(t+1)|}{\|X_{i*}(t+1)\|_1}\right) |X(t+1)|_{\max} \\ &\leq \frac{|X_{ip}(t+1)|}{\|X_{i*}(t+1)\|_1} |X_{jp}(t+1)| + \left(1 - \frac{|X_{ip}(t+1)|}{\|X_{i*}(t+1)\|_1}\right) |X(t)|_{\max} \\ &= |X(t)|_{\max} - \frac{|X_{ip}(t+1)|}{\|X_{i*}(t+1)\|_1} (|X(t)|_{\max} - |X_{jp}(t+1)|) \\ &\leq |X(t)|_{\max} \\ &\quad - \frac{|X(t+1)|_{\min}}{n |X(t+1)|_{\max}} \frac{|X(t)|_{\min}}{n |X(t)|_{\max}} (|X(t)|_{\max} - |X(t)|_{\min}) \\ &\leq |X(t)|_{\max} - \frac{|X(t)|_{\min}^2}{n^2 |X(t)|_{\max}^2} (|X(t)|_{\max} - |X(t)|_{\min}). \end{aligned}$$

Therefore,

$$\begin{aligned} |X(t+2)|_{\max} - |X(t+2)|_{\min} &\leq \left(1 - \frac{|X(t)|_{\min}^2}{n^2 |X(t)|_{\max}^2}\right) (|X(t)|_{\max} - |X(t)|_{\min}) \\ &\leq \left(1 - \frac{|X(t_0)|_{\min}^2}{n^2 |X(t_0)|_{\max}^2}\right) (|X(t)|_{\max} - |X(t)|_{\min}). \end{aligned}$$

Now we have established the exponential convergence of  $|X(t)|_{\max} - |X(t)|_{\min}$  to 0. Therefore, there exists  $\alpha > 0$  such that  $\lim_{t \rightarrow \infty} |X_{ij}(t)| = \alpha$  for any  $i, j$ . Moreover, since  $\text{sign}(X(t)) = \text{sign}(X(t_0))$  for any  $t \geq t_0$ , we have  $\lim_{t \rightarrow \infty} X(t) = \alpha b b^\top$ , where  $b = \text{sign}(X_{1*}(t_0))^\top$ . This concludes the proof for (ii)(b)  $\Rightarrow$  (ii)(c).

The proof for (ii)(b)  $\Rightarrow$  (ii)(a) is straightforward. If  $G(X(t_0))$  satisfies social balance, then, according to claim C.2),  $|X(t)|_{\min} \geq |X(t_0)|_{\min}$  for any  $t \geq t_0$ , which means that  $\liminf_{t \rightarrow \infty} \min_{ij} |X_{ij}(t)| \geq |X(t_0)|_{\min} > 0$ .

Now we prove (ii)(c)  $\Rightarrow$  (ii)(b). Suppose  $X(t) \rightarrow X^*$  as  $t \rightarrow \infty$ . For any  $X^* \in Q_{\text{homophily}}$  of rank one, there exists  $\alpha > 0$  and  $b \in \{-1, 1\}^n$  such that  $X^* = \alpha b b^\top$ . Since  $\alpha > 0$ , there exists a neighbor set  $\mathcal{U}(X^*)$  such that for any  $X \in \mathcal{U}(X^*)$ ,  $\text{sign} X = \text{sign} X^*$ , which implies that, for any  $X \in \mathcal{U}(X^*)$ ,  $G(X)$  satisfies social balance. Moreover, since  $X(t) \rightarrow X^*$ , there exists  $t_0 > 0$  such that  $X(t) \in \mathcal{U}(X^*)$  for any  $t \geq t_0$ . Therefore,  $G(X(t))$  achieves social balance at  $t_0$ . This concludes the proof.  $\square$

As Theorem 3.7 points out, the appraisal matrix  $X(t)$  converge to some rank-one matrix  $\alpha b b^\top$  if and only if  $X(t)$  achieves social balance (see Definition 2.1) at some time  $t_0$ . The mathematical intuition behind the convergence to rank-one matrices is that, after achieving social balance, the quantity  $\max_{ij} |X_{ij}(t)| - \min_{k\ell} |X_{k\ell}(t)|$  is monotonically vanishing. In reality, various factors such as noisy disturbances and individual prejudice (see Friedkin & Johnsen, 1990) may prevent the appraisal matrix from converging to rank-one matrices.

Monte-Carlo validation of the non-vanishing appraisal condition indicates that statement (ii)(b) of Theorem 3.7 holds for generic initial conditions. The detailed simulation results are presented in Section 5. In fact, there exist some counter examples of  $X(0)$  with which the non-vanishing condition on the solution  $X(t)$  does not hold. Example 1: if  $X(0)$  is block-diagonal, then the dynamics of the blocks are decoupled. While statement (ii) of Theorem 3.7 still holds block-wisely, the non-vanishing condition on the entire matrix  $X(t)$  does not hold; Example 2: if all the off-diagonal entries of  $X(0) \in \mathbb{R}^{n \times n}$  are equal to some  $-b < 0$  and all the diagonal entries are equal to  $a = (n-2)b/2$ , one can check by computation that  $X(1)$  becomes a diagonal matrix with strictly positive diagonals, i.e.,  $X(1)$  is a rank- $n$  fixed point and therefore the non-vanishing condition does not hold. However, for both Examples 1 and 2, the sets of initial conditions are zero-measure and simulation results indicate that the zero-pattern of  $X(t)$  with those specifically constructed  $X(0)$  are not robust under perturbation: For Example 1, if  $X(0)$  has two diagonal

blocks, any perturbation of any of its zero-entries render the convergence of  $X(t)$  to a rank-one matrix, and therefore the non-vanishing appraisal condition holds again; For [Example 2](#), under any perturbation of any entry of  $X(0)$ ,  $X(t)$  converges to a rank-one matrix and the non-vanishing appraisal condition holds as well. Moreover, even for [Examples 1](#) and [2](#), the systems are still well-behaved and the solutions  $X(t)$  achieve social balance with  $k$  isolated subgraphs, as defined in [Remark 3.5](#).

We end this section with some remarks on the homophily-based model.

**Remark 3.8** (Sufficient Conditions for Non-Vanishing Appraisals). Since the non-vanishing appraisal condition is satisfied if  $X(t)$  achieves social balance at finite time, by writing down the closed-form expressions of  $X(1)$  and  $X(2)$  and applying [Lemma 2.2](#), we obtain the following sufficient conditions on the initial appraisals  $X$  for non-vanishing appraisals: (i) either  $(X_{i*}X_{1*}^\top)(X_{1*}X_{j*}^\top)(X_{i*}X_{j*}^\top) > 0$  for any  $i, j$ , (ii) or  $(X_{i*}X^\top XX_{1*}^\top)(X_{1*}X^\top XX_{j*}^\top)(X_{i*}X^\top XX_{j*}^\top) > 0$  for any  $i, j$ . Here the condition (i) ((ii) resp.) corresponds to the case when  $X(1)$  ( $X(2)$  resp.) is structurally balanced. Both condition (i) and (ii) define sets of initial appraisal matrices  $X$  with non-zero measure.

**Remark 3.9.** Our homophily-based model exhibits the following somehow unrealistic behavior: for any  $X(0) \in \mathcal{S}_{\text{nz-row}}$ , the solution  $X(t)$  immediately becomes sign-symmetric at time step 1. However, if we adopt a simple modification by considering individual memory, i.e., if the dynamics are given by

$$X(t + 1) = \epsilon f_{\text{homophily}}(X(t)) + (1 - \epsilon)X(t), \quad (6)$$

for some  $\epsilon \in (0, 1]$ , then, following the same argument as in the proofs for [Proposition 3.3](#), [Theorems 3.4](#), and [3.7](#), we conclude that

- (i) The set  $\mathcal{S}_{\text{pos-diag}} = \{X \in \mathbb{R}^{n \times n} \mid X_{ii} > 0 \text{ for any } i\}$  is invariant under dynamics (6);
- (ii) [Theorem 3.4](#) still holds, while statements (ii)–(iv) of [Proposition 3.3](#) and [Theorem 3.7](#) still hold for any  $X(0) \in \mathcal{S}_{\text{pos-diag}}$ .

The proof is provided in the technical report ([Mei et al.](#)).

#### 4. Influence-based model

In this section, we propose the *influence-based model* (IbM) and present some important theoretical results parallel to the results on the homophily-based model. Since the proof methods are similar to those in [Section 3](#), we leave out all the proofs in this section and refer the readers to the technical report ([Mei et al.](#)).

**Definition 4.1** (Influence-based Model). Given an initial appraisal matrix  $X(0) \in \mathcal{S}_{\text{rs-symm}}^+ \subset \mathbb{R}^{n \times n}$ , the influence-based model is defined by

$$X(t + 1) = \text{diag}(|X(t)|_{\mathbb{1}_n})^{-1}X(t)X(t). \quad (7)$$

**Remark 4.2** (Interpretation). Compared with the homophily-based model (4), the only difference here is that the term  $X(t)X(t)^\top$  on the right-hand side of (4) is changed to  $X(t)X(t)$ . Eq. (7) now describes an interpersonal influence process: Individuals adjust their appraisals of each other via the opinion dynamics  $X(t + 1) = W(t)X(t)$ . Here the opinion of each individual is how she/he appraise every one in the group, and each  $W_{ij}(t)$  denotes the weight that individual  $i$  assigns to individual  $j$ 's opinions. The construction of the influence matrix  $W(t) = \text{diag}(|X(t)|_{\mathbb{1}_n})^{-1}X(t)$  implies that the interpersonal influences are proportional to the interpersonal appraisals.

Next, we present some results on the invariant set and finite-time behavior of the influence-based model.

**Proposition 4.3** (Finite-time Properties of the IbM). Consider the dynamical system (7) and define  $f_{\text{influence}}(X) = \text{diag}(|X|_{\mathbb{1}_n})^{-1}XX$ . Pick any  $X_0 \in \mathcal{S}_{\text{rs-symm}}^+$ . The following statements hold:

- (i) the map  $f_{\text{influence}}$  is well-defined for any  $X \in \mathcal{S}_{\text{nz-row}}$  and maps  $\mathcal{S}_{\text{rs-symm}}^+$  to  $\mathcal{S}_{\text{rs-symm}}^+$ ;
- (ii) the solution  $X(t)$ ,  $t \in \mathbb{Z}_{\geq 0}$ , to Eq. (7) from initial condition  $X(0) = X_0$  exists and is unique;
- (iii)  $|X(t)|_{\max}$  is non-increasing for any  $t \geq 0$ ;
- (iv) for any  $c > 0$ , the trajectory  $cX(t)$  is the solution to Eq. (7) from initial condition  $X(0) = cX_0$ .

Notice that  $\mathcal{S}_{\text{nz-row}}$  is not an invariant set of the map  $f_{\text{influence}}$ . For example,

$$X(0) = \begin{bmatrix} 1 & 2 \\ -0.5 & -1 \end{bmatrix} \in \mathcal{S}_{\text{nz-row}}$$

leads to  $X(1) \notin \mathcal{S}_{\text{nz-row}}$  and, moreover,  $f_{\text{influence}}(X(1))$  is not defined. For the influence-based model, we consider  $\mathcal{S}_{\text{rs-symm}}^+$  as the domain of system (7) due to its invariance under the map  $f_{\text{influence}}$ . According to [Proposition 4.3](#), for any  $X(0) \in \mathcal{S}_{\text{rs-symm}}^+$  and any  $t \geq 0$ , each entry of  $|X(t)|$  is uniformly upper bounded, which is a desired property the previous models in [Marvel et al. \(2011\)](#) and [Traag et al. \(2013\)](#) do not have.

The following theorem characterizes the set of fixed points of the map  $f_{\text{influence}}$  in  $\mathcal{S}_{\text{rs-symm}}^+$ .

**Theorem 4.4** (Fixed Points and Social Balance). Consider system (7) in domain  $\mathcal{S}_{\text{rs-symm}}^+$ . Define

$$\begin{aligned} Q_{\text{influence}} &= \left\{ PYP^\top \in \mathcal{S}_{\text{rs-symm}}^+ \mid P \text{ is a permutation matrix,} \right. \\ &\quad \left. Y \text{ is a block diagonal matrix with blocks of the} \right. \\ &\quad \left. \text{form } \text{sign}(w)w^\top, w \in \mathbb{R}^m \text{ and } |w| > 0_m, m \leq n \right\}. \end{aligned}$$

Then

- (i)  $Q_{\text{influence}}$  is the set of all the fixed points of system (7) in domain  $\mathcal{S}_{\text{rs-symm}}^+$ ;
- (ii) for any  $X \in Q_{\text{influence}}$ ,  $G(X)$  is composed by isolated complete subgraphs that satisfy social balance.

**Remark 4.5.** The proof of [Theorem 4.4](#) implies that  $Q_{\text{influence}}$  is actually the set of all the fixed points of the map  $f_{\text{influence}}$  in  $\mathcal{S}_{\text{rs-symm}}^+$ . However, the set  $Q_{\text{influence}}$  does not contain all the fixed points in  $\mathcal{S}_{\text{nz-row}}$ . For example, let  $X = \alpha bb^\top$  for some  $\alpha > 0$  and  $b \in \{-1, +1\}^n$ . Then, pick one  $i \in \{1, \dots, n\}$  and set  $X_{*i} = 0_n$ . It can be easily verified that  $X = f_{\text{influence}}(X)$  but  $X \notin Q_{\text{influence}}$ .

Now we present the main results on the convergence of the appraisal network to social balance.

**Theorem 4.6** (Convergence and Social Balance in the IbM). Consider the influence-based model given by Eq. (7). The following statements hold:

- (i) Each element in  $Q_{\text{influence}}$  of rank one is a locally stable fixed point of  $f_{\text{influence}}$ ;
- (ii) For any  $X(0) \in \mathcal{S}_{\text{rs-symm}}^+$ , the following three statements are equivalent:
  - (a) the solution  $X(t)$  satisfies the non-vanishing appraisal condition given by [Definition 3.6](#);



- (b) there exists  $t_0 \geq 0$  such that  $G(X(t))$  satisfies social balance for all  $t \geq t_0$ ;
- (c) there exists  $X^* \in \mathcal{Q}_{\text{influence}}$  of rank one such that  $\lim_{t \rightarrow \infty} X(t) = X^*$ .

## 5. Further discussion and numerical simulations

### 5.1. Numerical validation of the non-vanishing appraisal condition and model comparisons

Monte Carlo validation indicates that, for the homophily-based model, the non-vanishing appraisal condition, given by Definition 3.6, holds for generic initial conditions in  $\mathcal{S}_{\text{nz-row}}$ . By generic initial condition, we mean each of  $X(0)$ 's entries is independently randomly generated from the uniform distribution on some support  $[-a, a]$ . Since the homophily-based model is independent of scaling, we only need to consider the support  $[-1, 1]$ . For any randomly generated  $X(0) \in \mathcal{S}_{\text{nz-row}} \cap [-1, 1]^{n \times n}$ , define the random variable  $Z : \mathcal{S}_{\text{nz-row}} \rightarrow \{0, 1\}$  as

$$Z(X(0)) = \begin{cases} 1, & \text{if } \min_{100 \leq t \leq 1000} \min_{i,j} |X_{ij}(t)| \geq 0.001, \\ 0, & \text{otherwise.} \end{cases}$$


Let  $p = \mathbb{P}[Z(X(0)) = 1]$ . For  $N$  such independent random samples  $Z_1, \dots, Z_N$ , define  $\hat{p}_N = \sum_{i=1}^N Z_i / N$ . For any accuracy  $1 - \varepsilon \in (0, 1)$  and confidence level  $1 - \xi \in (0, 1)$ ,  $|\hat{p}_N - p| < \varepsilon$  with probability greater than  $1 - \xi$  if the Chernoff bound is satisfied:  $N \geq \frac{1}{2\varepsilon^2} \log \frac{2}{\xi}$ . For  $\varepsilon = \xi = 0.01$ , the bound is satisfied by  $N = 27000$ . We ran the 27000 independent simulations of the homophily-based model with  $n = 8$ , and found that  $\hat{p} = 1$ . Therefore, we conclude that, for any generic initial condition  $X(0) \in \mathcal{S}_{\text{nz-row}}$ , with 99% confidence level, there is at least 0.99 probability that every entry of  $|X(t)|$  is lower bounded by a positive scalar (set to be 0.001 in this simulation) for all  $t \in \{100, \dots, 10000\}$ .

We remark that the continuous-time homophily-based model (Traag et al., 2013) has a similar property that the interpersonal appraisals reach social balance in finite time, however they diverge later also in finite time.


The same Monte Carlo validation is also applied to the influence-based model, except that now the generic initial conditions  $X(0) \in \mathcal{S}_{\text{rs-symm}}^+ \subset \mathbb{R}^{n \times n}$  is generated by the following steps: (1) Randomly and independently generate the diagonal and the upper triangular entries of a matrix  $\hat{X} \in \mathbb{R}^{n \times n}$  from the uniform distribution on  $[-1, 1]$ ; (2) Let  $\hat{X}_{ij} = \hat{X}_{ji}$  for any  $i > j$ ; (3) Randomly and independently generate the entries of a  $n \times 1$  vector  $\gamma$  from the uniform distribution on  $[0, 1]$ ; (4) Let  $X(0) = \text{diag}(\gamma)\hat{X}$ . We obtained that, for any initial condition  $X(0) \in \mathcal{S}_{\text{rs-symm}}^+$ , with 99% confidence level, there is at least 0.99 probability that every entry of  $|X(t)|$  is uniformly strictly lower bounded from 0 for all  $t \in \{100, \dots, 10000\}$ .

In the continuous-time influence-like model (Marvel et al., 2011; Traag et al., 2013), when the initial appraisal matrix  $X(0)$  is a normal matrix, i.e., when  $X(0)X(0)^T = X(0)^T X(0)$ , the appraisal network  $G(X(t))$  almost surely reaches social balance only in the limit case when the network size  $n$  tends to infinity. Compared with these models, besides the desired convergence property, our influenced-based model has the following advantages: (1) Unlike the set of normal matrices, of which the sociological meaning is not explicit, the almost-sure convergence to social balance in our influence-based model holds for any  $X(0) = \text{diag}(\gamma)\hat{X}$ , where  $\hat{X}$  is symmetric and  $\text{diag}(\gamma)$  has positive diagonals. With the term  $\text{diag}(\gamma)$ , our model allows for individuals' heterogeneous scaling of appraisals, which is sociologically more reasonable; (2) In our influence-based model, the almost-sure finite-time achievement of social balance holds for any finite network size  $n$ .

For both homophily-based and influence-based models, Monte Carlo validations with uniform but asymmetric initial appraisal



**Fig. 1.** Error-bar plot of the estimated probability of converging to social balance for both the homophily-based model and the influence model. For each network size, we run 1000 realizations, each with an initial condition  $X(0)$  randomly generated from  $\mathcal{S}_{\text{nz-row}} \cap [-1, 1]^{n \times n}$  in the same way as in the first paragraph of Section 5.1. Numerical convergence is determined by whether the non-vanishing appraisal condition holds. The error bars are taken as the estimated standard deviations of the probability estimation and turn out to be very small (0 for the homophily-based model).



**Fig. 2.** Visualization of the evolution of the appraisal matrix under perturbations in the homophily-based model. For each entry, the dark gray color indicates a negative appraisal, while the light gray indicates a positive one. The white color indicates no appraisal. The appraisal network has 11 nodes and is initially in a social balance state with 3 isolated subgraphs. With 6 links (4 positive and 2 negative links) added to the network, the appraisal network evolves to a single-clique structurally balanced state after 5 iterations.


distributions lead to the same results, but are not presented here due to the limit of space.

We further numerically estimate, for our influence-based model, the probability that the non-vanishing appraisal condition holds for generic initial conditions  $X(0) \in \mathcal{S}_{\text{nz-row}} \cap [-1, 1]^{n \times n}$ . According to Theorem 4.6, this probability is also the probability that the appraisal network converges to social balance. As shown in Fig. 1, for the influence-based model, the probability of converging to social balance is quite low and decays to zero as the network size increases. Such feature indicates that, if system (4) and (7) correctly characterize the homophily and influence mechanisms, respectively, then the homophily mechanism is a more universal explanation for the convergence of appraisal networks to social balance. That is, it is more probable that the empirically observed structurally balanced social networks are formed via the homophily mechanism rather than the influence mechanism.

### 5.2. Social balance under perturbation

For the homophily model, extensive simulation observations indicate that social balance with  $k > 1$  isolated subgraphs is unstable under perturbations. With some links added to the network, the subnetworks connected by the added links merge into larger subnetworks and the perturbed network converges to another balanced state with fewer isolated subgraphs, see Fig. 2 as an example and the following two insightful scenarios.

**Example 1 (Globalization of Local Conflicts).** Consider the appraisal network with two isolated subgraphs. Each subgraph is structurally balanced and made up of two antagonistic factions. The two factions in subgraph 1 are node sets  $V_1$  and  $V_2$ , respectively, while the two factions in subgraph 2 are  $V_3$  and  $V_4$ , respectively. Suppose one link with weight  $\eta$  is added from one node in  $V_1$  to one node in  $V_3$ . By computing the closed form expression of  $X(2)$ , we obtain that the perturbed appraisal network always recovers



**Fig. 3.** Visual illustration of the behavior of the social balance with 2 isolated subgraphs with the addition of one inter-subgraph link.

to a complete and structurally balanced network composed of two antagonistic factions. Moreover, if  $\eta > 0$ , then the two factions are  $V_1 \cup V_3$  and  $V_2 \cup V_4$ ; if  $\eta < 0$ , then the two factions are  $V_1 \cup V_4$  and  $V_2 \cup V_3$ . Fig. 3 visualizes the behavior described above. In reality, such behavior could be interpreted as the escalation of local conflicts. One real example of such phenomena is the formation of the globalized conflicts between the Axis and the Ally in World War II, after the Nazi German allied with the Imperial Japan.

**Example 2 (Competition for Ally and Mediation of Conflicts).** Consider an appraisal network with two isolated subgraphs: subgraph 1 with two antagonistic factions  $V_1 = \{1, \dots, n_1\}$  and  $V_2 = \{n_1 + 1, \dots, n_1 + n_2\}$ , and subgraph 2 with only one faction  $V_3 = \{n_1 + n_2 + 1, \dots, n_1 + n_2 + n_3\}$ . Suppose the appraisal matrix associated with subgraph 1 is given by  $\alpha bb^T$ , where  $b = (\mathbb{1}_{n_1}^T, -\mathbb{1}_{n_2}^T)^T$ , and  $\alpha > 0$  represents the sentiment strength inside subgraph 1. Similarly, the appraisal matrix associated with subgraph 2 is given by  $\hat{\alpha} \hat{b} \hat{b}^T$ , where  $\hat{b} = \mathbb{1}_{n_3}$  and  $\hat{\alpha} > 0$  represents the sentiment strength inside subgraph 2. Imagine then that both  $V_1$  and  $V_2$  aim to ally with  $V_3$ . Accordingly, suppose that, in order to ally with  $V_3$ , each node in  $V_1$  builds a bilateral link with each node in  $V_3$ , with link weight  $\epsilon_1 > 0$ , while each node in  $V_2$  builds a bilateral link with each node in  $V_3$  with weight  $\epsilon_2 > 0$ . With all these links added, the associated appraisal matrix takes the following form:


$$X(0) = \begin{bmatrix} \alpha \mathbb{1}_{n_1} \mathbb{1}_{n_1}^T & -\alpha \mathbb{1}_{n_1} \mathbb{1}_{n_2}^T & \epsilon_1 \mathbb{1}_{n_1} \mathbb{1}_{n_3}^T \\ -\alpha \mathbb{1}_{n_2} \mathbb{1}_{n_1}^T & \alpha \mathbb{1}_{n_2} \mathbb{1}_{n_2}^T & \epsilon_2 \mathbb{1}_{n_2} \mathbb{1}_{n_3}^T \\ \epsilon_1 \mathbb{1}_{n_3} \mathbb{1}_{n_1} & \epsilon_2 \mathbb{1}_{n_3} \mathbb{1}_{n_2} & \hat{\alpha} \mathbb{1}_{n_3} \mathbb{1}_{n_3}^T \end{bmatrix}.$$

Along the evolution of  $X(t)$  determined by  $X(0)$ , we obtain the following numerical results.

(i) If  $\epsilon_1 n_1 > \epsilon_2 n_2$ , i.e., faction  $V_1$  takes greater effort than  $V_2$  in allying with  $V_3$ , then  $V_1$  gains at least one ally, either  $V_2$  or  $V_3$ . Moreover, the following conditions  $\epsilon_1 n_1 - \epsilon_2 n_2 \geq \hat{\alpha} \epsilon_2 n_3 / \alpha$  and  $\epsilon_1 \epsilon_2 n_3 \leq \alpha^2 (n_1 + n_2)$  guarantee that  $V_1$  ally with  $V_3$ ; This statement also holds when all the subscripts 1 and 2 are switched;

(ii) If  $\epsilon_1 \epsilon_2 n_3 \leq \alpha^2 (n_1 + n_2)$ , then  $V_3$  eventually gains at least one ally. That is,  $V_3$  avoids the situation in which  $V_1$  and  $V_2$  end up allying with each other against  $V_3$ ;

(iii) Any of the following conditions guarantees that no negative link exists in the asymptotic appraisal network: (1)  $\epsilon_1 \epsilon_2 n_3 \geq \alpha^2 (n_1 + n_2)$  and  $\epsilon_1 n_1 - \epsilon_2 n_2 = 0$ ; (2)  $\epsilon_1 \epsilon_2 n_3 \geq \alpha^2 (n_1 + n_2)$  and  $0 < \epsilon_1 n_1 - \epsilon_2 n_2 \leq \epsilon_2 \hat{\alpha} n_3$ ; (3)  $\epsilon_1 \epsilon_2 n_3 \geq \alpha^2 (n_1 + n_2)$  and  $0 < \epsilon_2 n_2 - \epsilon_1 n_1 \leq \epsilon_1 \hat{\alpha} n_3$ . Notice that the inequality  $\epsilon_1 \epsilon_2 n_3 \geq \alpha^2 (n_1 + n_2)$  is required for all the three sufficient conditions. The right-hand side of this inequality above reflects the “scale” of the conflicts between factions  $V_1$  and  $V_2$ , while the left-hand side is  $V_1$  and  $V_2$ ’s average efforts in allying with  $V_3$ , multiplied by the size of  $V_3$ . From the three sufficient conditions, we learn that the larger the size of  $V_3$ , the more capable it is of mediating the conflicts between  $V_1$  and  $V_2$ . In addition,  $V_1$  and  $V_2$ ’s strong willingness to ally with  $V_3$ , as well as the sentiment strength inside  $V_3$ , i.e.,  $\hat{\alpha}$ , also helps mediate the conflicts.



**Fig. 4.** Formation of factions under different initial condition distributions for the homophily-based model. The white color indicates the presence of two factions in all the 30 random samples, while the dark gray color indicates the presence of one faction in all of the samples. The light gray color indicates any other case.

### 5.3. Distribution of initial conditions and formation of factions in the homophily-based model

We investigate numerically, for the homophily-based model, how initial appraisal distribution determines whether the appraisal network evolves to only one faction or two antagonistic factions. We randomly and independently sample the entries of  $X(0)$  from the uniform distribution on  $[x_{\min}, x_{\max}]$ , for which  $\text{ave}(x_{\min}, x_{\max}) = (x_{\max} + x_{\min})/2$  indicates how the initial appraisals are biased towards being positive. We set  $x_{\max} - x_{\min} = 2$  and change the values of  $\text{ave}(x_{\min}, x_{\max})$  and the number of agents. Given  $[x_{\min}, x_{\max}]$ , 30 samples of the initial condition  $X(0)$  are independently randomly generated and for each  $X(0)$  we count how many factions appear at  $X(500)$ . Since any  $X(0)$  and  $-X(0)$  lead to the same  $X(1)$  and  $X(t)$  thereafter, we only consider different values of  $\text{ave}(x_{\min}, x_{\max}) \geq 0$ . Fig. 4 shows that, for fixed network size, the smaller the value of  $\text{ave}(x_{\min}, x_{\max})$ , the more likely it is to find two antagonistic factions; for fixed value of  $\text{ave}(x_{\min}, x_{\max})$ , the larger the network size, the more likely that only one faction emerges.

Note that similar numerical study in Marvel et al. (2011) for the continuous-time influence-like model indicates that the appraisal network evolves to two antagonistic factions if the initial mean appraisal is non-positive. The appraisal network evolves to all-friendly state if the initial mean is positive. However, such results in Marvel et al. (2011) only hold for the limit case of infinitely large network size  $n$ .

## 6. Conclusion

This paper proposes both homophily-based and influence-based discrete-time models for the bounded evolution of interpersonal appraisal networks towards social balance. For either model, the set of fixed points includes all the possible balanced configurations, in the sense of sign pattern, of the appraisal network. Under the non-vanishing appraisal condition, we prove that both models exhibit asymptotic convergence to structurally balanced networks, while the convergence property holds for larger initial conditions set in the homophily-based model than in the influence-based model. Moreover, our models admit the existence of multiple isolated subgraphs in the final structure of the evolved appraisal network. Numerical study indicates how the final emergence of factions in the social network is sensitive to the initial appraisal distribution, and how the system transits from one fixed point to another under perturbations.

We remark that our models and the previous continuous-time models (Jia et al., 2016; Kuřakowski et al., 2005; Marvel et al., 2011; Traag et al., 2013) all adopt the definition of social balance for complete graphs, or isolated complete subgraphs in our paper, which implies that individuals interact with everyone

in the group/subgroup. This assumption limits the scope of the application of our models to (groups of) small-size groups, which are usually assumed to be complete graphs.

Possible future research directions include a better understanding of the influence-based model for arbitrary initial conditions, a validation of the proposed models with laboratory and/or field data, the study of asynchronous models with pairwise updates, and further study of conditions and cases in which one socio-psychological mechanism dominates the other.

## References

- Altafani, C. (2013). Consensus problems on networks with antagonistic interactions. *IEEE Transactions on Automatic Control*, 58(4), 935–946. <http://dx.doi.org/10.1109/TAC.2012.2224251>.
- Antal, T., Krapivsky, P. L., & Redner, S. (2005). Dynamics of social balance on networks. *Physical Review E*, 72(3), 036121. <http://dx.doi.org/10.1103/PhysRevE.72.036121>.
- Antal, T., Krapivsky, P. L., & Redner, S. (2006). Social balance on networks: The dynamics of friendship and enmity. *Physica D: Nonlinear Phenomena*, 224(1), 130–136. <http://dx.doi.org/10.1016/j.physd.2006.09.028>.
- Cartwright, D., & Gleason, T. C. (1966). The number of paths and cycles in a digraph. *Psychometrika*, 31, 179–199. <http://dx.doi.org/10.1007/BF02289506>.
- Cartwright, D., & Harary, F. (1956). Structural balance: A generalization of Heider's theory. *Psychological Review*, 63(5), 277. <http://dx.doi.org/10.1037/h0046049>.
- Davis, J. A. (1967). Clustering and structural balance in graphs. *Human Relations*, 20, 181–187. <http://dx.doi.org/10.1177/001872676702000206>.
- de Nooy, W. (1999). The sign of affection: Balance-theoretic models and incomplete signed digraphs. *Social Networks*, 21(3), 269–286. [http://dx.doi.org/10.1016/S0378-8733\(99\)00012-X](http://dx.doi.org/10.1016/S0378-8733(99)00012-X).
- Doreian, P., & Krackhardt, D. (2001). Pre-transitive balance mechanisms for signed networks. *Journal of Mathematical Sociology*, 25(1), 43–67. <http://dx.doi.org/10.1080/0022250X.2001.9990244>.
- Doreian, P., & Mrvar, A. (1996). A partitioning approach to structural balance. *Social Networks*, 18, 149–168. [http://dx.doi.org/10.1016/0378-8733\(95\)00259-6](http://dx.doi.org/10.1016/0378-8733(95)00259-6).
- Easley, D., & Kleinberg, J. (2010). *Networks, crowds, and markets: reasoning about a highly connected world*. Cambridge University Press.
- Facchetti, G., Iacono, G., & Altafani, C. (2011). Computing global structural balance in large-scale signed social networks. *Proceedings of the National Academy of Sciences*, 108(52), 20953–20958. <http://dx.doi.org/10.1073/pnas.1109521108>.
- Friedkin, N. E., & Johnsen, E. C. (1990). Social influence and opinions. *Journal of Mathematical Sociology*, 15(3–4), 193–206. <http://dx.doi.org/10.1080/0022250X.1990.9990069>.
- Friedkin, N. E., & Johnsen, E. C. (2011). *Social influence network theory: A sociological examination of small group dynamics*. Cambridge University Press.
- Harary, F. (1953). On the notion of balance of a signed graph. *Michigan Mathematical Journal*, 2(2), 143–146. <http://dx.doi.org/10.1307/mmj/1028989917>.
- Harary, F. (1961). A structural analysis of the situation in the Middle East in 1956. *Journal of Conflict Resolution*, 5, 167–178. <http://dx.doi.org/10.1177/002200276100500204>.
- Heider, F. (1944). Social perception and phenomenal causality. *Psychological Review*, 51(6), 358–374. <http://dx.doi.org/10.1037/h0055425>.
- Heider, F. (1946). Attitudes and cognitive organization. *Journal of Psychology*, 21(1), 107–112. <http://dx.doi.org/10.1080/00223980.1946.9917275>.
- Hendrickx, J. M. (2014). A lifting approach to models of opinion dynamics with antagonisms. In *IEEE conf. on decision and control* (pp. 2118–2123). <http://dx.doi.org/10.1109/CDC.2014.7039711>.
- Henley, N. M., Horsfall, R. B., & De Soto, C. B. (1969). Goodness of figure and social structure. *Psychological Review*, 76, 194–204. <http://dx.doi.org/10.1037/h0027358>.
- Jia, P., Friedkin, N. E., & Bullo, F. (2016). The coevolution of appraisal and influence networks leads to structural balance. *IEEE Transactions on Network Science and Engineering*, 3(4), 286–298. <http://dx.doi.org/10.1109/TNSE.2016.2600058>.
- Kim, M., & Candan, K. S. (2012). SBV-Cut: Vertex-cut based graph partitioning using structural balance vertices. *Data & Knowledge Engineering*, 72, 285–303. <http://dx.doi.org/10.1016/j.datak.2011.11.004>.
- King, M. G. (1964). Structural balance, tension, and segregation in a university group. *Human Relations*, 17, 221–225. <http://dx.doi.org/10.1177/001872676401700303>.
- Kuřakowski, K., Gawroński, P., & Gronek, P. (2005). The Heider balance: A continuous approach. *International Journal of Modern Physics C*, 16(05), 707–716. <http://dx.doi.org/10.1142/S012918310500742X>.
- Lazarsfeld, P. F., & Merton, R. K. (1954). Friendship as a social process: A substantive and methodological analysis. In M. Berger, & T. Abel (Eds.), *Freedom and control in modern society: Vol. 18*, (pp. 18–66). Van Nostrand.
- Leskovec, J., Huttenlocher, D., & Kleinberg, J. (2010). Signed networks in social media. In *28th international conference on human factors in computing systems* (pp. 1361–1370). <http://dx.doi.org/10.1145/1753326.1753532>, Atlanta, USA.
- Malekzadeh, M., Fazli, M., Jalaly Khalidabadi, P., Rabiee, H. R., & Safari, M. A. (2011). Social balance and signed network formation games. In *Proceedings of 5th KDD workshop on social network analysis*. San Diego, USA.
- Marvel, S. A., Kleinberg, J., Kleinberg, R. D., & Strogatz, S. H. (2011). Continuous-time model of structural balance. *Proceedings of the National Academy of Sciences*, 108(5), 1771–1776. <http://dx.doi.org/10.1073/pnas.1013213108>.
- McDonald, H. B., & Rosecrance, R. (1985). Alliance and structural balance in the international system: A reinterpretation. *Journal of Conflict Resolution*, 29(1), 57–82. <http://dx.doi.org/10.1177/0022002785029001004>.
- Mei, W., & Bullo, F. (2017). LaSalle invariance principle for discrete-time dynamical systems: A concise and self-contained tutorial. arXiv preprint. URL: <https://arxiv.org/abs/1710.03710>.
- Mei, W., Cisneros-Velarde, P., Friedkin, N. E., & Bullo, F. (2017). Dynamic social balance and convergent appraisals via homophily and influence mechanisms. arXiv preprint arXiv:1710.09498. URL: <https://arxiv.org/abs/1710.09498>.
- Proskurnikov, A. V., & Cao, M. (2017). Modulus consensus in discrete-time signed networks and properties of special recurrent inequalities. In *IEEE conf. on decision and control* (pp. 2003–2008). <http://dx.doi.org/10.1109/CDC.2017.8263942>, Melbourne, Australia.
- Rawling, C. M., & Friedkin, N. E. (2017). The structural balance theory of sentiment networks: Elaboration and test. *American Journal of Sociology*, 123(2), 510–548. <http://dx.doi.org/10.1086/692757>.
- Taylor, H. F. (1970). Balance in small groups. *sociological concepts*. In *Methods and data series*. Van Nostrand Reinhold.
- Traag, V. A., Van Dooren, P., & De Leenheer, P. (2013). Dynamical models explaining social balance and evolution of cooperation. *PLoS One*, 8(4), e60063. <http://dx.doi.org/10.1371/journal.pone.0060063>.
- van de Rijt, A. (2011). The micro-macro link for the theory of structural balance. *Journal of Mathematical Sociology*, 35(1–3), 94–113. <http://dx.doi.org/10.1080/0022250X.2010.532262>.
- Wongkaew, S., Caponigro, M., Kuřakowski, K., & Borzi, A. (2015). On the control of the Heider balance model. *The European Physical Journal Special Topics*, 224(17), 3325–3342. <http://dx.doi.org/10.1140/epjst/e2015-50087-9>.
- Xia, W., Cao, M., & Johansson, K. H. (2016). Structural balance and opinion separation in trust-mistrust social networks. *IEEE Transactions on Control of Network Systems*, 3(1), 46–56. <http://dx.doi.org/10.1109/TCNS.2015.2437528>.
- Zheng, X., Zeng, D., & Wang, F.-Y. (2015). Social balance in signed networks. *Information Systems Frontiers*, 17(5), 1077–1095. <http://dx.doi.org/10.1007/s10796-014-9483-8>.



**Wenjun Mei** received the Ph.D. in Mechanical Engineering from University of California, Santa Barbara, and the Bachelor of Science degree in Theoretical and Applied Mechanics from Peking University. He is currently a postdoc researcher at the Automatic Control Laboratory, ETH Zurich. His current research interests include dynamics of social systems, control of network systems, network games, and complex network analysis.




**Pedro Cisneros-Velarde** received the Bachelor of Science degree in Electrical Engineering from Pontifical Catholic University of Peru. He is currently a Ph.D. student in Mechanical Engineering at University of California, Santa Barbara. His current research interests include the modeling and analysis of social network dynamics.



**Ge Chen** received the B.Sc. degree in mathematics from the University of Science and Technology of China Hefei, China, in 2004, and the Ph.D. degree in mathematics from the University of Chinese Academy of Sciences, China, Beijing, China, in 2009. He joined the National Center for Mathematics and Interdisciplinary Sciences, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, in 2011, and is currently an Associate Professor. His current research interests include the collective behavior of multiagent systems.



Dr. Chen was the recipient of the First Prize of the Application Award from the Operations Research Society of China, in 2010. One of his papers was selected as a SIGEST paper by the SIAM Review, in 2014. He was also a Finalist of the IFORS ( International Federation of Operations Research Societies) Prize for OR in Development (2011).



**Noah E. Friedkin** received the B.A. and Ph.D. degrees from the University of Chicago, Chicago, IL, USA, in 1969 and 1978. He is currently with the University of California, Santa Barbara, CA, USA, where he is a Professor of sociology and a Faculty Member of the Center for Control, Dynamical-Systems, and Computation. His journal publications concentrate on the investigation of social networks, and in particular, attend to the social psychological foundations of social influence systems. His research has contributed new mathematical models, findings from experiments on groups of human subjects, and findings from statistical analyses of field setting observations. His current research interests include understanding how natural dynamical systems of interpersonal influence operate and the implications of such systems.

Prof. Friedkin has served as the Chair of the Mathematical Sociology Section of the American Sociological Association twice, the Chair of the Sociology Department, and is currently an Editor-in-Chief of the Journal of Mathematical Sociology. He was the recipient of two Harrison White Outstanding Book

Awards from the Mathematical Sociology Section of the American Sociological Association and election to the Sociological Research Association, an association of leading sociologists.



**Francesco Bullo** is a Professor with the Mechanical Engineering Department and the Center for Control, Dynamical Systems and Computation at the University of California, Santa Barbara. He was previously associated with the University of Padova (Laurea degree in Electrical Engineering, 1994), the California Institute of Technology (Ph.D. degree in Control and Dynamical Systems, 1999), and the University of Illinois. His research interests focus on network systems and distributed control with application to robotic coordination, power grids and social networks. He is the coauthor of "Geometric Control of Mechanical Systems" (Springer, 2004) and "Distributed Control of Robotic Networks" (Princeton, 2009); his forthcoming "Lectures on Network Systems" is available on his website.

He received best paper awards for his work in IEEE Control Systems, Automatica, SIAM Journal on Control and Optimization, IEEE Transactions on Circuits and Systems, and IEEE Transactions on Control of Network Systems. He is a Fellow of IEEE and IFAC. He has served on the editorial boards of IEEE, SIAM, and ESAIM journals, and serves as 2018 IEEE CSS President.