

## UC Davis

### UC Davis Previously Published Works

#### Title

Proactive classical biological control of *Lycorma delicatula* (Hemiptera: Fulgoridae) in California (U.S.): Host range testing of *Anastatus orientalis* (Hymenoptera: Eupelmidae).

#### Permalink

<https://escholarship.org/uc/item/6s1131w2>

#### Authors

Gómez Marco, Francesc

Yanega, Douglas

Ruiz, Marta

et al.

#### Publication Date

2023

#### DOI

10.3389/finsc.2023.1134889

Peer reviewed



## OPEN ACCESS

## EDITED BY

Miriam Cooperband,  
USDA APHIS PPQ, United States

## REVIEWED BY

Jana Collatz,  
Agroscope (Switzerland), Switzerland  
Teodora Borislavova Toshova,  
Bulgarian Academy of Sciences, Bulgaria

## \*CORRESPONDENCE

Francesc Gómez Marco  
✉ francegm@ucr.edu

## SPECIALTY SECTION

This article was submitted to  
Invasive Insect Species,  
a section of the journal  
Frontiers in Insect Science

RECEIVED 31 December 2022

ACCEPTED 08 February 2023

PUBLISHED 20 February 2023

## CITATION

Gómez Marco F, Yanega D, Ruiz M and  
Hoddle MS (2023) Proactive classical  
biological control of *Lycorma delicatula*  
(Hemiptera: Fulgoridae) in California (U.S.):  
Host range testing of *Anastatus orientalis*  
(Hymenoptera: Eupelmidae).  
*Front. Insect Sci.* 3:1134889.  
doi: 10.3389/finsc.2023.1134889

## COPYRIGHT

© 2023 Gómez Marco, Yanega, Ruiz and  
Hoddle. This is an open-access article  
distributed under the terms of the [Creative  
Commons Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The  
use, distribution or reproduction in other  
forums is permitted, provided the original  
author(s) and the copyright owner(s) are  
credited and that the original publication in  
this journal is cited, in accordance with  
accepted academic practice. No use,  
distribution or reproduction is permitted  
which does not comply with these terms.

# Proactive classical biological control of *Lycorma delicatula* (Hemiptera: Fulgoridae) in California (U.S.): Host range testing of *Anastatus orientalis* (Hymenoptera: Eupelmidae)

Francesc Gómez Marco<sup>1\*</sup>, Douglas Yanega<sup>1</sup>, Marta Ruiz<sup>2</sup>  
and Mark S. Hoddle<sup>1,3</sup>

<sup>1</sup>Department of Entomology, University of California, Riverside, CA, United States, <sup>2</sup>Department of Botany & Plant Sciences, University of California, Riverside, CA, United States, <sup>3</sup>Center for Invasive Species Research, University of California, Riverside, CA, United States

*Lycorma delicatula* (Hemiptera: Fulgoridae), the spotted lanternfly, native to China, invaded and established in the northeast U.S. in 2014. Since this time, populations have grown and spread rapidly, and invasion bridgeheads have been detected in mid-western states (i.e., Indiana in 2021). This invasive pest presents a significant threat to Californian agriculture. Therefore, a proactive classical biological control program using *Anastatus orientalis* (Hymenoptera: Eupelmidae), a *L. delicatula* egg parasitoid native to China, was initiated in anticipation of eventual establishment of *L. delicatula* in California. In support of this proactive approach, the potential host range of *A. orientalis* was investigated. Eggs of 34 insect species either native or non-native to the southwestern U.S. were assessed for suitability for parasitism and development of *A. orientalis*. Of the native species tested, 10, 13, and one were Hemiptera, Lepidoptera, and Mantodea, respectively. Of the non-native species, eight Hemiptera and two Lepidoptera were evaluated. Host range tests conducted in a quarantine facility, exposed individually mated *A. orientalis* females (Haplotype C) to non-target and target (i.e., *L. delicatula*) eggs in sequential no-choice and static choice experiments to determine suitability for parasitization and development. Additionally, the sex ratio, fertility, and size of offspring obtained from non-target and target eggs were evaluated. Results of host range testing indicated that *A. orientalis* is likely polyphagous and can successfully parasitize and develop in host species belonging to at least two different orders (i.e., Hemiptera, Lepidoptera) and seven families (Coreidae, Erebidae, Fulgoridae, Lasiocampidae, Pentatomidae, Saturniidae and Sphingidae). Prospects for use of *A. orientalis* as a classical biological control agent of *L. delicatula* in the southwestern U.S. are discussed.

## KEYWORDS

Coreidae, Lasiocampidae, Spotted lanternfly, Saturniidae, parasitoid, polyphagy, southwestern U.S

## 1 Introduction

Classical or introduction biological control is the intentional importation, release, and establishment of natural enemies for suppressing damaging populations of invasive non-native organisms to densities that no longer cause economic or ecological harm. This approach aims to reduce pest population densities by re-associating safe (i.e., host-specific) and efficacious natural enemies with the target pest (1). Host range and host specificity testing are important primary steps in identifying natural enemy species that may have deleterious impacts on pest populations while presenting minimal risk to non-target species (1). Host use evaluation studies are mandatory in the United States of America (U.S.) and provide safety data for review by Federal agencies (i.e., United States Department of Agriculture, Animal and Plant Health Inspection Service [USDA APHIS]), that regulate the importation and release of natural enemies for use in classical biological control programs (2). Host range and host specificity testing evaluations are time consuming, often taking years to complete (3). During this time, newly established invasive pest populations tend to increase in density and spread as management plans are slowly developed and implemented. Proactive biological control attempts to reduce or eliminate this window of opportunity for an invasive pest by evaluating candidate natural enemies for potential use in a classical biological control program in advance of the anticipated incursion and establishment of the target pest in the area of concern (3).

Spotted lanternfly, *Lycorma delicatula* (White) (Hemiptera: Fulgoridae), native to China (4), was detected for the first time in the U.S. in Berks County Pennsylvania, in September 2014 (5). By November 2022, *L. delicatula* infestations were confirmed in an additional 13 states in eastern and mid-western areas of the U.S. (Connecticut, Delaware, Indiana, Maryland, Massachusetts, Michigan, New Jersey, New York, North Carolina, Ohio, Rhode Island, Virginia, and West Virginia) (6). *Lycorma delicatula* is a phloem-feeding fulgorid that has a broad host range (7). Direct feeding damage can cause mortality to highly preferred hosts like *Ailanthus altissima* (Miller) (Sapindales: Simaroubaceae) and grapevines (*Vitis vinifera* L. [Vitales: Vitaceae]). Indirect damage results from the excretion of high quantities of honeydew that promote sooty mold growth (7, 8). *Lycorma delicatula* has been recorded infesting forest and ornamental shade tree species in natural and urban areas, respectively (7, 9), and presents a significant risk to economically important perennial agricultural crops (e.g., grapes and nuts) (10, 11). Long-distance dispersal by *L. delicatula* is almost entirely human-assisted. This occurs primarily through the accidental translocation of egg masses that are often laid indiscriminately on inert substrates (e.g., wooden pallets and railcars) that undergo subsequent transportation into uninfested areas (12, 13). This type of inadvertent relocation in the U.S. likely resulted in the establishment of invasion bridgeheads in Indiana (2021) and Michigan and North Carolina (2022) (6). Ecological niche models indicate that *L. delicatula* has a potential distribution that includes large areas of the west coast of the U.S., and other parts of the world (e.g., Europe) (14). For California, a western U.S. state, with an agricultural economy worth ~\$50 billion per year (15), *L.*

*delicatula* is viewed as a significant invasion threat that could cause major problems for producers of specialty crops, like grapes and nuts, which are multi-billion-dollar industries (15).

The egg parasitoid, *Anastatus orientalis* Yang and Choi (Hymenoptera: Eupelmidae), was found parasitizing *L. delicatula* eggs in northern China in 2011 during foreign exploration for natural enemies for use in South Korea, where *L. delicatula* is also invasive (16, 17). Following the invasion and spread of *L. delicatula* on the east coast of the U.S. there was renewed interest in the potential use of *A. orientalis* as a classical biological control agent (18, 19). Molecular analyses identified six different haplotypes of *A. orientalis* collected from the native range. Importations of *A. orientalis* into the U.S. were initially comprised of Haplotype C, which was first evaluated as a classical biological control agent against *L. delicatula* (20). The majority of *Anastatus* spp. Motschulsky are primary endoparasitoids attacking eggs of Diptera, Dictyoptera, Coleoptera, Hemiptera, Lepidoptera, Orthoptera, and Mantodea (21–27). Numerous *Anastatus* species have been considered or used as classical biological control agents against various pests around the world. For example, *A. japonicus* Ashmead was released in the eastern U.S. for control of *Lymantria dispar* L. (Lepidoptera: Erebidae) (27, 28) and against *Tessaratomia papillosa* Drury (Hemiptera: Pentatomidae) in China (29, 30). *Anastatus* sp. was released in Nepal to control *Rhynchocoris humeralis* (Thunberg) (Hemiptera: Pentatomidae) (31). *Anastatus bifasciatus* (Geoffroy) was evaluated to study levels of biotic resistance of central European natural enemies against invasive *Halyomorpha halys* Stål (Hemiptera: Pentatomidae) populations (32, 33). In the eastern U.S., *A. reduvii*, a native species, has been detected as one of the most common parasitoids emerging from eggs of invasive *H. halys* (34). Some *Anastatus* spp. are commercially-available and used for augmentative biological control of *Amblyopelta nitida* Stål and *A. lutescens lutescens* Distant (Hemiptera: Coreidae) in Australia (35, 36).

Given the obvious threat posed to California agriculture by the westward migration of *L. delicatula* in the U.S., a proactive biological control program was initiated. Proactive research efforts focused on the suitability of *A. orientalis* (Haplotype C) as a potential classical biological control agent of *L. delicatula* in advance of its expected establishment in California (3). Consequently, the objective of this study was to investigate the physiological host range of *A. orientalis* on native and non-native non-target species from the southwestern U.S. (i.e., California and Arizona) to determine whether or not this natural enemy would be a suitable candidate to release for classical biological control of *L. delicatula* should it eventually establish in California. The results of these studies are presented here.

## 2 Materials and methods

### 2.1 Source and collection of test insects

*Anastatus orientalis* Haplotype C specimens were obtained from colonies established at USDA APHIS PPQ (Plant Protection and Quarantine) Forest Pest Methods Laboratory in Massachusetts,

U.S. Initial *A. orientalis* populations were shipped to the University of California Riverside Insectary and Quarantine Facility (UCR-IQF) as parasitized *L. delicatula* egg masses under USDA-APHIS permit P526P-22-03022 and P526P-22-04208 and California Department of Food and Agriculture (CDFA) permit 3888. Colonies of *A. orientalis* were established in UCR-IQF in October 2019 and reared continuously on cold stored *L. delicatula* egg masses.

*Lycorma delicatula* egg masses were field collected in winter (December to March) of 2019 to 2022. Collections were made in Pennsylvania, U.S. (Berks, Dauphin, Huntingdon, Lancaster and Lebanon Counties) predominantly from *A. altissima* (>90%). Entire egg masses attached to underlying bark were removed using chisels and shipped or hand carried to the UCR-IQF under USDA-APHIS permit P526P-19-02058 and CDFA Permit 3458. In quarantine, all field collected egg masses were stored at 5°C and 60-75% R.H. Egg masses were randomly selected and used for experiments reported here.

Selection of non-target species for host range testing was made based on phylogenetic relationships amongst species within the family Fulgoridae and their representation in the southwestern U.S. Five native fulgorid genera in California and Arizona, *Amcyle* spp., *Cyrpoptus* spp., *Poblicia* spp., *Scaralina* spp. (described incorrectly as genus *Alphina* spp. Stål, Yanega et al. unpublished), and *Scolopsella* spp. (37). were targeted for field collections and use in host range tests. Previous host range studies on *Anastatus* suggested that species may potentially have broad host ranges and are capable of utilizing hosts from different orders (32). Consequently, to determine if *A. orientalis* potentially exhibits oligophagy or polyphagy, additional non-target species belonging to Hemiptera (Cicadellidae, Coreidae, Liviidae, Pentatomidae, Reduviidae, and Rhopalidae), Lepidoptera (Erebidae, Lasiocampidae, Saturniidae, and Sphingidae), and Mantodea (Mantidae) were included in host range testing for *A. orientalis* (Table 1). These families were also selected to compliment simultaneous testing conducted by collaborators at the USDA APHIS PPQ Forest Pest Methods Laboratory of potential native and non-native non-target species found in the eastern U.S. All non-target insect colonies used for host range testing, unless otherwise stated, were maintained on each test species preferred host plant species held in cages (BugDorm-2120 61×61×61 cm, MegaView Science Co. Ltd., Taiwan) at the UCR-IQF at 25°C, 65%RH, L:D 16:8. Colonies were checked daily for egg masses which were harvested and used immediately or held at 10°C until used for experiments.

## 2.2.1 Hemiptera

### 2.2.1.1 Fulgoridae collections

Fulgorids collected in the Chiricahua mountains near Portal, Santa Cruz County, in southeastern Arizona included *Scaralina* spp. (comprised of three undescribed species and incorrectly placed as *Alphina* genera, Yanega et al. unpublished) and *Cyrpoptus vanduzeei* Ball (Table 1). Adult *Scaralina* spp. were hand collected as they were attracted to mercury vapor and UV lights. Immediately after capture, adult males and females were caged (i.e., sleeve cages made of mesh with fiber spacing of 160 µm (Figure 1)

on trunks of oak trees, *Quercus arizonica* Sarg. (Fagales: Fagaceae), at the American Museum of Natural History's Southwestern Research Station, Portal Arizona. Cages were inspected daily for oviposited egg masses which were collected and maintained at ~10°C until use in experiments with *A. orientalis*. Three species of *Scaralina* were collected and relatively low numbers of egg masses per species were obtained. Therefore, all egg masses (n = 9) used for experiments were pooled and referred to as "*Scaralina* spp.". *Poblicia fuliginosa* (Olivier) and *C. vanduzeei* adults were collected during the day from *Baccharis sarothroides* Gray (Asterales: Asteraceae) from different locations in Arizona (Table 1). Adult *P. fuliginosa* and *C. vanduzeei* were maintained on potted *B. sarothroides* plants held in cages. Live insects used for experiments were moved to UCR-IQF under USDA-APHIS Permit number P526P-19-00766 and CDFA Permit number 3457.

### 2.2.1.2 Cicadellidae collections

*Homalodisca vitripennis* Germar (Hemiptera: Cicadellidae), a pest of grapes, were collected with sweep nets in citrus orchards in Riverside, California and maintained on potted basil, *Ocimum basilicum* (Lamiales: Lamiaceae), a host plant that supports adult feeding and oviposition.

### 2.2.1.3 Coreidae collections

*Acanthocephala thomasi* Uhler, *Chelinidea vittiger* Uhler, *Leptoglossus zonatus* (Dallas) and *Thasus neocalifornicus* Brailovskiy and Barrera were included in host range tests. *Acanthocephala thomasi* specimens were hand collected in Portal Arizona (Table 1) and maintained on potted *B. sarothroides* plants. *Chelinidea vittiger* specimens were collected in Riverside, California and maintained on *Opuntia* sp. (Caryophyllales: Cactaceae). *Leptoglossus zonatus* specimens were obtained from research colonies maintained in the Department of Entomology at UC Riverside. *Thasus neocalifornicus* were collected in Sonoita, Arizona and maintained on *Prosopis velutina* Wootton (Fabales: Fabaceae) trees (Table 1).

### 2.2.1.4 Liviidae collections

*Diaphorina citri* Kuwayama (Hemiptera: Liviidae) eggs were obtained from colonies maintained in the UCR-IQF building of the Department of Entomology at UCR. *Diaphorina citri*, collected in southern California in 2011 and certified free of the citrus killing bacterium *Candidatus Liberibacter asiaticus* (CLAs), were reared on *Murraya koenigii* (Sapindales: Rutaceae), a non-propagative host for CLAs.

### 2.2.1.5 Pentatomidae collections

*Banasa dimidiata* (Say) (native), *Bragada hilaris* (Burmeister) (invasive), *Chinavia hilaris* Say (native), *Nezara viridula* L. (invasive) and *H. halys* (invasive) were used in host range tests. *Banasa dimidiata* specimens were collected in Riverside, California and maintained on *Hirschfeldia incana* (Brassicales: Brassicaceae) (Table 1). *Bragada hilaris* and *Chinavia hilaris* were obtained from

TABLE 1 Non-target species tested, selection criteria for use in evaluations, and collection information.

| Order     | Family       | Genera                | Species                  | Native, non-native or invasive | Egg deposition type <sup>1</sup> | Selection criteria (Ref)                               | GPS coordinates of collection sites <sup>2</sup> / Commercially obtained | Collection date     |
|-----------|--------------|-----------------------|--------------------------|--------------------------------|----------------------------------|--|--|---------------------|
| Hemiptera | Cicadellidae | <i>Homalodisca</i>    | <i>vitripennis</i>       | Invasive                       | M                                | Egg masses laid under leaf epidermis                   | 33° 58' 20.26"N - 117° 19' 3.33"W (CA)                                   | April 2020          |
|           | Coreidae     | <i>Acanthocephala</i> | <i>thomasi</i>           | Native                         | I                                | Recorded host family for <i>Anastatus</i> sp (29, 32). | 31° 54' 54.38"N - 109° 8' 9.17"W (AZ)                                    | August 2021         |
|           |              | <i>Chelinidea</i>     | <i>vittiger</i>          | Native                         | G                                | Recorded host family for <i>Anastatus</i> sp (29, 32). | 33° 58' 29.32"N - 117° 18' 59.98"W (CA)                                  | July 2020           |
|           |              | <i>Leptoglossus</i>   | <i>zonatus</i>           | Invasive                       | G                                | Recorded host family for <i>Anastatus</i> sp (29, 32). | Colonies maintained at UC Riverside                                      | -                   |
|           |              | <i>Thasus</i>         | <i>neocalifornicus</i>   | Native                         | G                                | Recorded host family for <i>Anastatus</i> sp (29, 32). | 31° 40' 42.93"N - 110° 39' 37.16"W (AZ)                                  | August 2021         |
|           | Fulgoridae   | <i>Cyrpoptus</i>      | <i>vanduzeei</i>         | Native                         | M                                | Family-level relatedness to <i>L. delicatula</i>       | 31° 54' 48.28"N - 109° 8' 22.75"W (AZ)                                   | August 2020         |
|           |              | <i>Lycorma</i>        | <i>delicatula</i>        | Invasive                       | M                                | Target   | Pennsylvania, U.S.   | 2019, 2020 and 2021 |
|           |              | <i>Poblicia</i>       | <i>fuliginosa</i>        | Native                         | M                                | Family-level relatedness to <i>L. delicatula</i>       | Southeastern AZ  | August 2021         |
|           |              | <i>Scaralina</i>      | <i>unidentified</i> spp. | Native                         | M                                | Family-level relatedness to <i>L. delicatula</i>       | 31° 53' 12.77"N - 109° 12' 40.37"W (AZ)                                  | August 2019 to 2021 |
|           | Liviidae     | <i>Diaphorina</i>     | <i>citri</i>             | Invasive                       | G                                | Eggs readily available from research colonies.         | Colonies maintained at UC Riverside                                      | -                   |
|           | Pentatomidae | <i>Banasa</i>         | <i>dimidiata</i>         | Native                         | M                                | Recorded host family for <i>Anastatus</i> sp (29).     | 33° 58' 20.26"N - 117° 19' 3.34"W (CA)                                   | April 2020          |
|           |              | <i>Bragada</i>        | <i>hilaris</i>           | Invasive                       | I                                | Recorded host family for <i>Anastatus</i> sp (29).     | Colonies maintained at UC Riverside                                      | -                   |

(Continued)

TABLE 1 Continued

| Order       | Family        | Genera                  | Species                 | Native, non-native or invasive | Egg deposition type <sup>1</sup> | Selection criteria (Ref)                                | GPS coordinates of collection sites <sup>2</sup> / Commercially obtained | Collection date     |
|-------------|---------------|-------------------------|-------------------------|--------------------------------|----------------------------------|---|--|---------------------|
|             |               | <i>Chinavia</i>         | <i>hilaris</i>          | Native                         | M                                | Recorded host family for <i>Anastatus</i> sp (29).      | Colonies maintained at UC Riverside                                      | –                   |
|             |               | <i>Nezara</i>           | <i>viridula</i>         | Invasive                       | M                                | Recorded host species for <i>Anastatus</i> sp (21, 29). | 34° 3' 46.03"N - 118° 21' 16.12"W (CA)                                   | April 2020          |
|             |               | <i>Halyomorpha</i>      | <i>halys</i>            | Invasive                       | M                                | Recorded host species for <i>Anastatus</i> sp (29, 31). | 34° 3' 46.03"N - 118° 21' 16.12"W (CA)                                   | April 2020          |
|             | Reduviidae    | <i>Zelus</i>            | <i>renardii</i>         | Non-native                     | M                                | Beneficial insect. Readily available                    | Commercially available   | –                   |
|             | Rhopalidae    | <i>Jadera</i>           | <i>haematoloma</i>      | Invasive                       | I                                | Easily collected from field sites                       | 33° 58' 26.67"N - 117° 19' 1.55"W (CA)                                   | April 2020          |
| Mantodea    | Mantidae      | <i>Stagmomantis</i>     | <i>californica</i>      | Native                         | M                                | Recorded host family for <i>Anastatus</i> sp (22, 23).  | 33° 40' 12.00"N - 116° 24' 44.43"W (CA)                                  | August 2020         |
| Lepidoptera | Erebidae      | <i>Apantesis</i>        | <i>unidentified</i> sp. | Native                         | I                                | Recorded host family for <i>Anastatus</i> sp (29).      | 33° 27' 57.48"N - 117° 2' 29.93"W (CA)                                   | June 2020           |
|             |               | <i>Pseudohemihyalea</i> | <i>edwardsii</i>        | Native                         | G                                | Recorded host family for <i>Anastatus</i> sp (29).      | 31° 53' 12.77"N-109° 12' 40.37"W (AZ)                                    | August 2019 to 2021 |
|             | Lasiocampidae | <i>Gloveria</i>         | <i>arizonensis</i>      | Native                         | G                                | Recorded host family for <i>Anastatus</i> sp (29).      | 31° 53' 12.77"N-109° 12' 40.37"W (AZ)                                    | August 2019 to 2021 |
|             | Saturniidae   | <i>Actias</i>           | <i>luna</i>             | Non-native                     | I                                | Recorded host family for <i>Anastatus</i> sp (29).      | Commercially available   | –                   |
|             |               | <i>Agapema</i>          | <i>anona</i>            | Native                         | I                                | Recorded host family for <i>Anastatus</i> sp (29).      | 31° 43' 5.15"N - 110° 52' 56.22"W (AZ)                                   | September 2021      |

(Continued)

TABLE 1 Continued

| Order | Family     | Genera              | Species               | Native, non-native or invasive | Egg deposition type <sup>1</sup> | Selection criteria (Ref)                            | GPS coordinates of collection sites <sup>2</sup> / Commercially obtained | Collection date |
|-------|------------|---------------------|-----------------------|--------------------------------|----------------------------------|---|--|-----------------|
|       |            | <i>Anisota</i>      | <i>oslari</i>         | Native                         | G                                | Recorded host family for <i>Anastatus</i> sp (29).  | 31° 53' 12.77"N-109° 12' 40.37"W (AZ)                                    | August 2021     |
|       |            | <i>Antheraea</i>    | <i>oculea</i>         | Native                         | G                                | Recorded host genera for <i>A. orientalis</i> (38). | 31° 53' 12.77"N-109° 12' 40.37"W (AZ)                                    | August 2021     |
|       |            | <i>Automeris</i>    | <i>cecrops pamina</i> | Native                         | G                                | Recorded host family for <i>Anastatus</i> sp (29).  | 31° 53' 12.77"N-109° 12' 40.37"W (AZ)                                    | August 2021     |
|       |            |                     | <i>metzli</i>         | Non-native                     | G                                | Recorded host family for <i>Anastatus</i> sp (29).  | Commercially available   | -               |
|       |            | <i>Eupackardia</i>  | <i>calleta</i>        | Native                         | I                                | Recorded host family for <i>Anastatus</i> sp (29).  | 31° 43' 5.15"N - 110° 52' 56.22"W (AZ)                                   | August 2021     |
|       |            | <i>Hyalophora</i>   | <i>euryalus</i>       | Native                         | I                                | Recorded host family for <i>Anastatus</i> sp (29).  | 32° 54' 55.26"N - 116° 53' 50.67"W (CA)                                  | May 2021        |
|       |            | <i>Rothschildia</i> | <i>cineta</i>         | Native                         | I                                | Recorded host family for <i>Anastatus</i> sp (29).  | 31° 43' 5.15"N - 110° 52' 56.22"W (AZ)                                   | August 2021     |
|       |            | <i>Saturnia</i>     | <i>walterorum</i>     | Native                         | I                                | Recorded host family for <i>Anastatus</i> sp (29).  | 32° 54' 55.26"N - 116° 53' 50.67"W (CA)                                  | May 2021        |
|       | Sphingidae | <i>Pachysphinx</i>  | <i>occidentalis</i>   | Native                         | I                                | Recorded host family for <i>Anastatus</i> sp (29).  | 31° 53' 12.77"N-109° 12' 40.37"W (AZ)                                    | August 2025     |

(1) Egg type: M = Egg mass [eggs laid very close together with some protective material covering the eggs]; I = Individual eggs; G = Group of individual eggs laid in patches of irregular number of eggs.  
 (2) CA, California; AZ, Arizona. California and Arizona are southwestern states in the U.S. (Ref), Reference.

research colonies maintained in the Department of Entomology at UCR. *Halyomorpha halys* and *N. viridula* were established from adults collected from Hancock Park, in Los Angeles, California. All live insects were transported to UCR-IQF under USDA-APHIS Permit number P526P-22-03011 and CDFA Permit number 3887 (Table 1). *Halyomorpha halys* colonies were maintained on a mixed diet of avocados, carrots, apples, green beans, table grapes and *A.*

*altissima*. *Nezara viridula* colonies were maintained on green bean plants and raw peanuts.

#### 2.2.1.6 Reduviidae collections

Egg masses of *Zelus renardii* Kolenati were purchased from Arbico Organics (Oro Valley, Arizona). Purchased eggs were




FIGURE 1

(A) Sleeve cages set up on branches of *Quercus arizonica* in Portal, AZ were used to confine adult native fulgorids on putative host plants for mating and oviposition. (B) A *Scaralina* sp. specimen captured at night by black lighting is seen resting on the bark of *Q. arizonica* branch enclosed by a sleeve cage (the red arrow indicates position of *Scaralina* sp.).

approximately 2 days of age upon receipt. Egg masses were exposed to *A. orientalis* immediately.

#### 2.2.1.7 Rhopalidae collections

*Jadera haematoloma* Herrich-Schäffer adults were collected in the Botanic Gardens at the University of California Riverside campus, Riverside, California. Specimens were not feed, adults were kept on ventilated plastic containers, provided with a water-saturated cotton wick, and eggs were collected daily and exposed immediately to *A. orientalis*.

#### 2.2.2 Lepidoptera

All Lepidoptera (Erebidae, Lasiocampidae, Saturniidae, and Sphingidae) species, used in host range testing (except for *Automeris metzli* Sallé which were purchased from an online vendor as pupae) were field collected as adults (Table 1). *Automeris metzli* pupae were held at 10°C for two months to simulate exposure to winter temperatures. After this chilling period, pupae were moved to a temperature cabinet set at 25 ± 2° C and 60% R.H. until adults emerged. Field captured adult moths were kept in bug-dorms (BugDorm-2120 61×61×61 cm, MegaView



TABLE 2 Non-target species tested for host range suitability of *A. orientalis*.

| Order       | Family        | Genera                  | Species                | 7 days exposure to <i>A. orientalis</i> |                 |                                      |                       | 24 hours exposure to <i>A. orientalis</i> |     |                                      |                       |
|-------------|---------------|-------------------------|------------------------|---|-----------------|--------------------------------------|-----------------------|---|-----|--------------------------------------|-----------------------|
|             |               |                         |                        | Percent parasitism ( $\pm$ SE)          | (n)             | Average number of eggs/n ( $\pm$ SE) | Sex ratio ( $\pm$ SE) | Percent parasitism ( $\pm$ SE)            | (n) | Average number of eggs/n ( $\pm$ SE) | Sex ratio ( $\pm$ SE) |
| Hemiptera   | Cicadellidae  | <i>Homalodisca</i>      | <i>vitripennis</i>     | 0                                       | 5               | 10 $\pm$ 0.32                        | –                     | –   | –   | –                                    | –                     |
|             | Coreidae      | <i>Acanthocephala</i>   | <i>thomasi</i>         | 100                                     | 1               | 1                                    | 1                     | –   | –   | –                                    | –                     |
|             |               | <i>Chelinidea</i>       | <i>vittiger</i> *      | 0                                       | 1               | 37                                   | –                     | –   | –   | –                                    | –                     |
|             |               | <i>Leptoglossus</i>     | <i>zonatus</i>         | 0                                       | 11              | 38.27 $\pm$ 2.67                     | –                     | –   | –   | –                                    | –                     |
|             |               | <i>Thasus</i>           | <i>neocalifornicus</i> | 0                                       | 7               | 4.71 $\pm$ 0.75                      | –                     | –   | –   | –                                    | –                     |
|             | Fulgoridae    | <i>Cyrpoptus</i>        | <i>vanduzeei</i> *     | 0                                       | 2               | 54.5                                 | –                     | –   | –   | –                                    | –                     |
|             |               | <i>Lycorma</i>          | <i>delicatula</i> **   | 58.24 $\pm$ 10.09                       | 13              | 39.08 $\pm$ 3.84                     | 0.77 $\pm$ 0.07       | 28.15 $\pm$ 5.04                          | 38  | 40.66 $\pm$ 3.79                     | 0.76 $\pm$ 0.04       |
|             |               | <i>Poblicia</i>         | <i>fuliginosa</i>      | 12.37 $\pm$ 8.42                        | 7               | 29.86 $\pm$ 2.69                     | 0                     | –   | –   | –                                    | –                     |
|             |               | <i>Scaralina</i>        | <i>spp.</i>            | 0                                       | 7               | 26.57 $\pm$ 2.57                     | –                     | –   | –   | –                                    | –                     |
|             | Liviidae      | <i>Diaphorina</i>       | <i>citri</i>           | 0                                       | 3               | 115.67 $\pm$ 40.71                   | –                     | –   | –   | –                                    | –                     |
|             | Pentatomidae  | <i>Banasa</i>           | <i>dimidiata</i>       | 0                                       | 1               | 14                                   | –                     | –   | –   | –                                    | –                     |
|             |               | <i>Bragada</i>          | <i>hilaris</i>         | 0                                       | 3               | 6 $\pm$ 0.58                         | –                     | –   | –   | –                                    | –                     |
|             |               | <i>Chinavia</i>         | <i>hilaris</i>         | 68.79 $\pm$ 12.41                       | 11              | 17.73 $\pm$ 2.51                     | 0.22 $\pm$ 0.1        | 7.41                                      | 2   | 30.5                                 | 0                     |
|             |               | <i>Nezara</i>           | <i>viridula</i>        | 18.34 $\pm$ 6.87                        | 17 <sup>a</sup> | 77.35 $\pm$ 5.79                     | 0.07 $\pm$ 0.05       | 4.79 $\pm$ 1.91                           | 17  | 59.58 $\pm$ 3.78                     | 0.02 $\pm$ 0.01       |
|             |               | <i>Halyomorpha</i>      | <i>halys</i>           | 48.05 $\pm$ 3.5                         | 92 <sup>b</sup> | 26.85 $\pm$ 0.57                     | 0.05 $\pm$ 0.02       | 37.27 $\pm$ 13.96                         | 8   | 23 $\pm$ 2.46                        | 0                     |
|             | Reduviidae    | <i>Zelus</i>            | <i>renardii</i>        | 0                                       | 4               | 24.25 $\pm$ 2.02                     | –                     | –   | –   | –                                    | –                     |
|             | Rhopalidae    | <i>Jadera</i>           | <i>haematoloma</i>     | 0                                       | 3               | 43 $\pm$ 3.21                        | –                     | –   | –   | –                                    | –                     |
| Mantodea    | Mantidae      | <i>Stagmomantis</i>     | <i>californica</i>     | 0                                       | 2               | 150                                  | –                     | –   | –   | –                                    | –                     |
| Lepidoptera | Erebidae      | <i>Apantesis</i>        | <i>sp.</i>             | 0                                       | 2               | 130 $\pm$ 8                          | –                     | –   | –   | –                                    | –                     |
|             |               | <i>Pseudohemihyalea</i> | <i>edwardsii</i>       | 16.7 $\pm$ 6                            | 6               | 22.17 $\pm$ 3.12                     | 0                     | –   | –   | –                                    | –                     |
|             | Lasiocampidae | <i>Gloveria</i>         | <i>arizonensis</i>     | 38.74 $\pm$ 9.62                        | 13 <sup>a</sup> | 18.92 $\pm$ 1.72                     | 0.11 $\pm$ 0.06       | 23.33 $\pm$ 9.55                          | 5   | 16 $\pm$ 1                           | 0.72 $\pm$ 0.21       |
|             | Saturniidae   | <i>Actias</i>           | <i>luna</i>            | 25.78 $\pm$ 6.89                        | 9               | 19.11 $\pm$ 0.79                     | 0.04 $\pm$ 0.04       | –   | –   | –                                    | –                     |
|             |               | <i>Agapema</i>          | <i>anona</i>           | 76.48 $\pm$ 14.10                       | 5               | 42.2 $\pm$ 7.43                      | 0                     | –   | –   | –                                    | –                     |
|             |               | <i>Anisota</i>          | <i>oslari</i> *        | 0                                       | 16              | 14.63 $\pm$ 1.22                     | –                     | –   | –   | –                                    | –                     |

(Continued)

TABLE 2 Continued

| Order | Family     | Genera              | Species               | 7 days exposure to <i>A. orientalis</i> |     |                                 |                  | 24 hours exposure to <i>A. orientalis</i> |     |                                 |                  |   |
|-------|------------|---------------------|-----------------------|---|-----|---------------------------------|------------------|---|-----|---------------------------------|------------------|---|
|       |            |                     |                       | Percent parasitism (± SE)               | (n) | Average number of eggs/n (± SE) | Sex ratio (± SE) | Percent parasitism (± SE)                 | (n) | Average number of eggs/n (± SE) | Sex ratio (± SE) |   |
|       |            | <i>Antheraea</i>    | <i>oculea</i>         | 0                                       | 1   | 6                               | -                | -   | -   | -                               | -                | - |
|       |            | <i>Automeris</i>    | <i>cecrops pamina</i> | 19.12 ± 14.41                           | 3   | 13 ± 3                          | 0                | -   | -   | -                               | -                | - |
|       |            |                     | <i>metzli</i>         | 0                                       | 1   | 10                              | -                | -   | -   | -                               | -                | - |
|       |            | <i>Eupackardia</i>  | <i>callata</i>        | 100                                     | 1   | 12                              | 0                | -   | -   | -                               | -                | - |
|       |            | <i>Hyalophora</i>   | <i>euryalus</i>       | 100                                     | 3   | 5.5                             | 0.16             | -   | -   | -                               | -                | - |
|       |            | <i>Rothschildia</i> | <i>cincta</i>         | 45                                      | 2   | 10                              | 0                | -   | -   | -                               | -                | - |
|       |            | <i>Saturnia</i>     | <i>walterorum</i>     | 19.60 ± 17.41                           | 5   | 11.2 ± 1.02                     | 0.5 ± 0.32       | 42.96 ± 25.7                              | 3   | 9.67 ± 0.33                     | 0.5 ± 0.2        | - |
|       | Sphingidae | <i>Pachysphinx</i>  | <i>occidentalis</i>   | 1.56 ± 1.56                             | 4   | 17.5 ± 2.72                     | 0                | -   | -   | -                               | -                | - |

Male and female parasitoid offspring produced (female sex ratio) on each non-target species, number of repetitions per species (n), average number of eggs used per species per repetition and average parasitism rate of *A. orientalis* on eggs of each species.

(\*) 100% of the *A. orientalis* females did not parasitize *L. delicatula* in the sequential exposure.

(\*\*) target host.

<sup>a</sup>non-target host eggs of different ages. See section 1 and Figure 2.

Science Co. Ltd., Taiwan) and maintained outdoors near collection sites or in a temperature chamber (25 ± 2°C; 60% of R.H.) when moved into UCR-IQF for mating and oviposition. Adults were not provided with food or water as test species do not feed in adult stage (except *Apanteles* sp. which was provided 50% honey water solution). Eggs oviposited onto walls of cages or on cardboard oviposition strips were collected daily and either used immediately or maintained at 10°C until used for experiments.

### 2.2.3 Mantodea

#### 2.2.3.1 Mantidae collections

Adult female *Stagmomantis californica* Rehn & Hebard were collected in Riverside County, California (Table 1) and fed with *H. halys* nymphs and adults. Ootheca, ~48 h of age, were collected and presented to *A. orientalis*.

### 2.3 No-choice sequential host-testing

Five female *A. orientalis*, less than 24 hours of age, were placed in a test unit with one male and a thin smear of pure honey on the mesh of the unit's lid as a carbohydrate source. Each experimental egg mass-parasitoid test arena was comprised of a clear plastic container 3 cm × 4 cm × 5 cm (180 mL clear RPTE hinged lid deli containers, AD16 GenPak, Charlotte, NC) with a modified lid that had a ventilated mesh window (1.5 cm x 2.5 cm) to facilitate air exchange. One *L. delicatula* egg mass was placed in the test unit and exposed to the five females and the male of *A. orientalis* for seven days. This seven day period is a pre-oviposition period during which host feeding occurs (pers. obs. F. Gomez Marco) at temperatures that simulate the fall (i.e., September when parasitoid oviposition in the field occurs) in Beijing (average daily high 25°C, average daily low 14°C, lights on 6:00 AM, lights off 6:30 PM [i.e., L:D 12.5:11.5], 75% R.H.; referred to as Beijing-fall regimen), the general area where *A. orientalis* was collected for colony establishment in the U.S (16–19). All experiments were conducted under the Beijing-fall regimen. After this one-week pre-oviposition period, females were moved individually and placed singly without males in new separate test units for a total of 274 *A. orientalis* females. One non-target host egg mass [number of eggs in the egg mass and the physical size of the egg mass varied on species being tested (Table 2)] was placed into each test unit containing a single mated female for seven days. After the seven-day exposure period, non-target eggs were removed, and replaced with *L. delicatula* egg masses, and females were left to host feed and oviposit for an additional seven days. Thus, the sequential non-choice tests were performed in this order; target host (pre-oviposition period, seven days) – non-target host (seven days) – target host (seven days). Parasitism of *L. delicatula* egg masses in the exposure trial following exposure to non-target eggs confirmed female competency if no parasitism was observed from non-target exposures. The total time taken to complete each no choice sequential host test cycle for each female was 21 days.

A variation of this experiment that reduced female exposure time to non-target egg masses from five species [*C. hilaris*, *Gloveria*

*arizonensis* Packard, *H. halys*, *N. viridula* and *Saturnia walterorum* Hogue & Johnson (Table 1)] was conducted. In this set of experiments, five *A. orientalis* females had a seven day pre-oviposition period with a male and access to a *L. delicatula* egg mass. Following this seven-day exposure period, individual mated females without males were exposed to non-target eggs (Tables 1, 2) for 24 hours then moved to a target *L. delicatula* egg mass for 24 hours to confirm competency.

For both experimental designs, female *A. orientalis*, that did not produce offspring either on the non-target eggs or on the second exposure to the *L. delicatula* egg mass, were classified as “incompetent” and discarded from analyses. This rule was not followed for *Anisota oslari* Rothschild due to the high mortality (100%) of *A. orientalis* females following exposure to eggs of this species, and for *C. vittiger* and *C. vanduzeei*, due the low number of repetitions due to low egg availability (Table 2), which resulted from difficulty in acquiring sufficient test eggs of non-target species for experiments. Each *L. delicatula* egg mass (from pre-oviposition and post-non-target exposure) and non-target species eggs were held under the Beijing-fall regimen for one month, for development of parasitoid larvae. After this four-week period, eggs were transferred to 25°C, 16:8 L:D, and 75% R.H. for emergence of parasitoid offspring following previously published protocols (19). The host species from which *A. orientalis* emerged were recorded. Target eggs that did not produce parasitoids were dissected to detect failed parasitism (i.e., presence of dead parasitoid larvae or pupae were recorded). Percentage of parasitism was calculated as:

% Parasitism

$$= \frac{\text{Total number of parasitoids (i.e., emerged adults + failed larvae + failed pupae)}}{\text{Total number of host eggs exposed}} \times 100 \quad (1)$$

Mortality of non-target host species due to exposure to *A. orientalis* was calculated using the Henderson–Tilton formula (Equation 2) (38), which calculates percent mortality based on the initial and final insect counts in the control relative to treatments with parasitoid exposure. Rates of naturally occurring mortality for non-target species host eggs were calculated with control eggs that were held under similar ambient conditions to test eggs but were not exposed to *A. orientalis*. The percentage of mortality by parasitoids was calculated as:

% Mortality by parasitoids

$$= \left( 1 - \frac{(\text{Average number of eggs in controls}) \times (\text{Number of juveniles after exposure to } A. orientalis)}{(\text{Number of eggs exposed}) \times (\text{Average number of juveniles in controls})} \right) \times 100 \quad (2)$$

Using percent mortality caused by parasitoids and percent parasitism, percent mortality of non-target hosts resulting from parasitoid activity, but that did not result in *A. orientalis* offspring (i.e., excess mortality [due to host feeding and/or oviposition attempts]) was calculated with the equation:

% Excess mortality

$$= \% \text{ Mortality by parasitoid} - \% \text{ Parasitism} \quad (3)$$

Percent excess mortality caused by parasitoids to non-target hosts was compared with percent mortality in controls not exposed to parasitoids when the number of successfully completed repetitions for each treatment exceeded a minimum of three.

## 2.4 Choice host-testing

To assess host preference on parasitization by *A. orientalis* when given a simultaneous choice between eggs of non-target species and *L. delicatula*, choice host-tests were performed with two non-target species, *G. arizonensis* and *H. halys*. Pairs of egg masses of target and non-target species were presented simultaneously to female parasitoids ~48 hours of age in exposure arenas which were constructed using two stacked transparent U-shaped acrylic risers 15cm×15cm×15cm (SW Plastics F2191, Riverside, CA), that formed a rectangular cage 15cm×15cm×30cm with two open sides. One open face was covered with white semi-opaque no-see-um netting (Skeeta, Bradenton, FL) and the other was fitted with a 30cm-long sleeve sewn from no-see-um netting). Choice tests were run either for 24 hours or seven days. Inside arenas, egg masses were separated by 26 cm and randomly placed on the floor of arenas for each repetition. After exposure time, each group of eggs (target, non-target, and control eggs not exposed to parasitoids) were isolated in ventilated clear plastic test arenas (3 cm × 4 cm × 5 cm, see section 2) and held under the Beijing-fall regimen for four weeks before being moved to 25°C until parasitoids or immature non-target species emerged from eggs, or eggs were classified as dead and dissected for evidence of parasitism when possible.

## 2.5 *Anastatus orientalis* offspring sex ratio, fertility and size when reared from target and non-target hosts

*Anastatus orientalis* offspring that emerged from non-target host species and target host (i.e., *L. delicatula*) eggs were evaluated for offspring sex ratio, fertility of males and females, and adult size. Parasitoid sex ratio was calculated as the number of female parasitoids divided by the total number of female and male parasitoids combined that emerged from each experimental egg mass. Three different offspring fertility evaluations were performed on five non-target host species: *Actias luna* L., *Agapema anona* Ottolengui, *G. arizonensis*, *H. halys* and *P. fuliginosa*. First, males and females emerging from the same non-target host species (< 48 hours of age) were set up in test arenas (see section 2 for details). Second, males that emerged from non-target host species were coupled with unmated *A. orientalis* females that emerged from *L. delicatula* eggs. Finally, females that emerged from non-target host species were coupled with *A. orientalis* males that emerged from *L. delicatula* egg masses. All mating couples were exposed to *L. delicatula* egg masses for seven days and a thin smear of pure honey on the mesh of the ventilated lid of the test arena as a carbohydrate source. After seven days, male-female pairs were removed, and each egg mass was held under the Beijing-fall




FIGURE 2

Percentage of parasitism ( $\pm$  SE) by *A. orientalis* on non-target host eggs of (A) *H. halys* (<24 hours; n = 27, 1 day; n = 23; 2 days; n = 27, 3 days; n = 17), (B) *N. viridula* (<24 hours; n = 11, 1 day; n = 6) and (C) *G. arizonensis*. Different letters indicate significant differences for percent parasitism of eggs of different ages.

regimen for four weeks then moved to 25°C for the emergence of parasitoids and non-target hosts.

To evaluate the effect of host species on the size of *A. orientalis* male and female parasitoids that successfully emerged from non-target and target host eggs, measurements of right hind tibia lengths were used as a proxy for parasitoid size and subsequent fitness (i.e., parasitoids with larger hind tibia are assumed to be bigger and more fit than parasitoids with smaller tibia) (39, 40). Excised right hind tibiae were placed onto glass slides and covered with a second glass slide. Hind tibia length was measured from its point of attachment

on the femur to the attachment point with the tarsi using a Leica S8AP0 microscope. Slide mounted hind tibiae were photographed at a magnification of 25 $\times$  with an attached Leica DMC2900 camera and length was measured using the Leica Application Suite version 4.6.2. A total of ~10 randomly selected *A. orientalis* males and females from each non-target host were measured and compared to 10 randomly selected males and females reared from *L. delicatula*.

To evaluate the effects of non-target host egg age on parasitism rates/host acceptance of *A. orientalis*, the age of eggs from all non-target host species exposed to *A. orientalis* was recorded. Data from

eggs exposed to *A. orientalis* for seven days in the non-choice experiment (see section 2) were used. Three species, *G. arizonensis*, *N. viridula*, and *H. halys* resulted in sufficient repetitions and/or age variability to be analyzed. *Gloveria arizonensis* Packard eggs age ranged from 5 to 35 days old. *Nezara viridula* egg age used in this study were  $\leq 24$  hours of age. Finally, *H. halys* egg age ranged from  $< 24$  hours to three days of age. Percent parasitism was compared between egg age for each of these three species.

## 2.6 Statistical analysis

All statistical analyses were conducted in R version 4.1.3 (41) using the development environment RStudio (42). Percent excess mortality that resulted from non-reproductive behavior of *A. orientalis* was compared with natural-occurring mortality rates in the paired controls using a generalized linear model (GLM) with a quasi-binomial distribution to account for high variance in data sets. The relation between the hind tibia size of the male parasitoids offspring and the sex ratio of parasitoid offspring emerging from the same host was analyzed using linear regression. Differences in mean hind tibiae lengths between male and female parasitoids emerging from different non-target species eggs and target eggs were analyzed using ANOVA followed by a Tukey posthoc test at the 0.05 level of significance using the package ‘multcomp’. All means are presented  $\pm$  standard error (SE).

## 7 Results

### 7.1 No-choice sequential host-testing experiments

In addition to the target host *L. delicatula*, eggs from a total of 34 non-target host species, however, eggs of three *Scaralina* spp. were pooled as *Scaralina* sp. giving a functional total of 32 species that were exposed to *A. orientalis* females in no-choice sequential host testing experiments. From the total of 244 female *A. orientalis* that completed the entire sequential exposure series (non-target and target), 23 (9.4%) females failed to parasitize the non-target host and the target host. They were considered incompetent and were excluded from data analyses. There were three exceptions for non-target hosts; *C. vittiger* and *C. vanduzeei*, due to the low number of repetitions because of the low numbers of non-target eggs available for testing, and *A. oslari* due to the high mortality of parasitoids ( $n = 16$ ; 100% of females tested died) after exposure to non-target eggs (Table 2). Additionally, 41 (16.8%) parasitoids were able to parasitize non-target host eggs and then failed to parasitize *L. delicatula* eggs. The results from these trials were included in data analyses.

*Anastatus orientalis* parasitized five species in the order Hemiptera: *A. thomasi* (Coreidae), *C. hilaris*, *H. halys* and *N. viridula* (all Pentatomidae) and *P. fuliginosa* (Fulgoridae) (Table 2). *Anastatus orientalis* parasitized 10 species in the order

Lepidoptera: *A. luna*, *A. anona*, *Automeris cecrops pamina* Neumoegen, *Eupackardia calleta* Westwood, *Hyalophora euryalus* Boisduval, *Rothschildia cincta* Tepper, *S. walterorum* (Saturniidae), *G. arizonensis* (Lasiocampidae), *Pseudohemihyalea edwardsii* Packard (Erebidae) and *Pachysphinx occidentalis* Edwards (Sphingidae) (Table 2). The maximum percent parasitism of eggs for non-target hosts in Hemiptera and Lepidoptera were obtained on the native pentatomid, *C. hilaris*, and the native saturniid, *A. anona*, at  $68.79\% \pm 12.41$  and  $76.48\% \pm 14.10$ , respectively.

Percent non-target host mortality of eggs exposed to *A. orientalis* was corrected with their paired controls (Equation 2) and excess mortality for non-target hosts following exposure to parasitoids was calculated (Equation 3). Excess mortality was compared across nine non-target hosts species (four suitable for parasitism and five unsuitable for parasitism, Figure 3) with their paired controls. Percent excess mortality resulting from death other than parasitism that resulted in the emergence of an adult parasitoid (i.e., mortality from host feeding and/or failed parasitism) for non-target host species were not affected by exposure to *A. orientalis* and no significant differences were found for nine species (*A. oslari*;  $F_{1,18} = 0.184$ ,  $P = 0.67$ , *C. hilaris*;  $F_{1,19} = 0.11$ ,  $P = 0.75$ , *D. citri*;  $F_{1,4} = 0.82$ ,  $P = 0.41$ , *H. halys*;  $F_{1,116} = 0.097$ ,  $P = 0.75$ , *J. haematoloma*;  $F_{1,18} = 3.5$ ,  $P = 0.13$ , *L. zonatus*;  $F_{1,13} = 0.69$ ,  $P = 0.42$ , *N. viridula*;  $F_{1,30} = 1.98$ ,  $P = 0.16$ , *S. walterorum*;  $F_{1,6} = 0.71$ ,  $P = 0.43$ , *T. neocalifornicus*;  $F_{1,8} = 4.89$ ,  $P = 0.058$ ) (Figure 3).

Of the five non-target host species exposed to *A. orientalis* females for 24 hours, three pentatomids, *C. hilaris*, *H. halys* and *N. viridula*, and two Lepidoptera species, *G. arizonensis* (Lasiocampidae) and *S. walterorum* (Saturniidae), were all parasitized by *A. orientalis*. The maximum average percent parasitism was observed for *S. walterorum* ( $42.96 \pm 25.7$ ) and the minimum average percent parasitism was recorded for *N. viridula* ( $4.79 \pm 1.91$ ) (Table 2).

The effect of non-target host egg age on parasitism by *A. orientalis* was evaluated for three species, *G. arizonensis*, *H. halys* and *N. viridula*. Rates of parasitism decreased as non-target host egg age increased for *H. halys* ( $F_{1,92} = 1.54$ ,  $P < 0.001$ ), and no effect of egg age on parasitism was observed for *N. viridula* ( $F_{1,17} = 1.54$ ,  $P = 0.23$ ). Similarly, increasing age of *G. arizonensis* eggs did not affect parasitism rates ( $F_{1,13} = 0.07$ ,  $P = 0.79$ ) (Figure 2).

### 7.2 Choice host-testing experiments

The two non-target species used in choice experiments, *G. arizonensis* and *H. halys*, were parasitized in both exposure periods, 24 hours and 7 days, when exposed to *A. orientalis* in the presence of *L. delicatula* egg masses (Figure 4). For all the *G. arizonensis* vs *L. delicatula* choice trials ( $n = 12$ ), four parasitoids failed to parasitize one of the two hosts species exposed; *G. arizonensis* and *L. delicatula* eggs were not parasitized three times and one time, respectively. For *H. halys* vs *L. delicatula* experiments ( $n = 19$ ), three *A. orientalis* females did not parasitize either the




FIGURE 3

Percent non-target host mortality after exposure to *A. orientalis* from which parasitoids emerged (suitable hosts) and failed to emerge (unsuitable hosts). Species with no excess/natural mortality following parasitoid exposure were excluded from figure (Correction of mortality for exposed non-target host was null [Equation 2]). Total percent mortality of non-target hosts is the sum of the percentage of parasitism (in black) and the excess mortality (in grey) ( $\pm$  SE). (ns) Indicates no significant differences between the percent excess mortality after exposure to *A. orientalis* and the percentage of eggs alive without exposure to *A. orientalis* (control or naturally-occurring mortality).

target or the non-target host. Eight parasitoids failed to parasitize one of the two host species exposed; *H. halys* eggs were not parasitized in three trials and *L. delicatula* eggs were not parasitized in five trials.

### 7.3 *Anastatus orientalis* offspring from non-target hosts; sex ratio, male and female fertility, and size

#### 7.3.1 Sex ratio

A total of 15 non-target host species were suitable hosts for *A. orientalis* and a total of 1,870 parasitoids, 1,780 males and 90

females, emerged from susceptible non-target species following seven-day exposure time. The highest female sex ratio was obtained from *S. walterorum* eggs ( $0.5 \pm 0.32$ ) (there was one exception, *A. thomasi*, which only one egg was exposed and parasitized and resulted in a female parasitoid). Seven non-target species did not produce females after being parasitized by *A. orientalis* (i.e., parasitism of *A. anona*, *A. cecrops pamina*, *E. calleta*, *P. edwardsii*, *P. fuliginosa*, *P. occidentalis* and *R. cincta* eggs produced only male offspring) (Table 2).

The five non-target species exposed to *A. orientalis* for 24 hours resulted in a total of 137 parasitoids, 110 males, 24 females and three larvae. The highest sex ratio between these five species was obtained from *G. arizonensis* eggs ( $0.72 \pm 0.21$ ).




FIGURE 4

Average percent parasitism of non-target host eggs (A) *G. arizonensis* and (B) *H. halys*, when presented simultaneously with the target host, *L. delicatula* for two different exposure times, 24 hours and 7 days (n = number of repetitions).

Only *C. hilaris* and *H. halys* did not produce females (i.e., male offspring only produced) after being parasitized by *A. orientalis* (Table 2).

### 7.3.2 Fertility

Male parasitoids were observed mating with females reared from the same non-target host species and female offspring were subsequently produced confirming mating was successful (*A.*

*orientalis* is arrhenotokous and female offspring are produced from fertilized eggs). Consequently, male-female pairs of *A. orientalis* that emerged from the same non-target host species produced male and female offspring. (Table 3). Males that emerged from non-target host species successfully inseminated females emerging from *L. delicatula* egg masses as female offspring were produced from these male-female pairings (Table 3).

TABLE 3 Parasitism rates and offspring sex ratio (i.e., proportion of female offspring) produced by i) *A. orientalis* couples that emerged from five different non-target host species or ii) males that emerged from non-target host species mated with unmated females that emerged from *L. delicatula* eggs or iii) females that emerged from non-target host species mated with males that emerged from *L. delicatula* eggs (n = number of repetitions).

| Non-target host species     | Host source of <i>A. orientalis</i> mating pairs and resulting parasitism rates and offspring sex ratio |     |             |     |   |     |             |     |   |     |           |     |
|-----------------------------|---|-----|-------------|-----|---|-----|-------------|-----|---|-----|-----------|-----|
|                             | Male-female pair from same non-target host species  |     |             |     | Male from non-target host, Female from <i>L. delicatula</i> |     |             |     | Female from non-target host, male from <i>L. delicatula</i> |     |           |     |
|                             | Percentage of parasitism  | (n) | sex ratio   | (n) | Percentage of parasitism                                    | (n) | sex ratio   | (n) | Percentage of parasitism                                    | (n) | sex ratio | (n) |
| <i>Actias luna</i>          | 10.39 ± 0.81  | 3   | 0.33 ± 0.19 | 3   | 58.19 ± 8.02  | 3   | 0.53 ± 0.26 | 3   | –   | –   | –         | –   |
| <i>Agapema anona</i>        | –   | –   | –           | –   | 11.64 ± 8.02  | 3   | 0.57        | 2   | –   | –   | –         | –   |
| <i>Gloveria arizonensis</i> | 20.89 ± 13.27   | 7   | 0.88 ± 0.07 | 3   | 0   | 3   | –           | –   | 13.04*  | 1   | –         | –   |
| <i>Halyomorpha halys</i>    | 30.96 ± 7.93  | 13  | 0.36 ± 0.1  | 11  | –   | –   | –           | –   | –   | –   | –         | –   |
| <i>Poblicia fuliginosa</i>  | –   | –   | –           | –   | 0   | 1   | –           | –   | –   | –   | –         | –   |

\*Parasitoids failed to emerge and parasitism was calculated by counting numbers of parasitoid larvae (alive) in dissected eggs.

### 7.3.3 Size

A wide range of offspring sizes as well as a pronounced sexual dimorphism with larger females and smaller males was observed for *A. orientalis* when reared from non-target host species (Table 4). Average male hind tibia lengths ranged from  $0.309 \pm 0.005$  mm (host: *P. edwardsii*) to  $0.587 \pm 0.022$  mm (host: *G. arizonensis*). Average hind tibia lengths for females ranged from  $0.613 \pm 0.007$  mm (host: *C. hilaris*) to 1.04 mm (host: *A. thomasi*) (Table 4). The largest *A. orientalis* males emerged from the target host, *L. delicatula* ( $F_{6,62} = 100.9$ ,  $P < 0.001$ ). The largest female emerged from the non-target host, *A. thomasi*. All other non-target hosts from which females emerged were smaller when compared with females that emerged from *L. delicatula* eggs ( $F_{3,26} = 285.4$ ,  $P < 0.001$ ) (Table 4). The average size of male offspring was significantly correlated with the sex ratio of the parasitoids emerging from the same host species ( $F_{1,6} = 7.668$ ,  $P = 0.032$ ;  $R^2 = 0.561$ ) (Figure 5). Non-target hosts that produced *A. orientalis* males were smaller and

the sex ratio of emerged parasitoids was lower (i.e., male biased) and larger parasitoids typically emerged from host eggs that had female-biased sex ratios (Figure 5).

## 8 Discussion


This is the first study to assess the physiological host range and specificity of *A. orientalis* (Haplotype C) with respect to potential non-target species that occur in the southwestern U.S. In the quarantine laboratory, no-choice tests assessing the developmental suitability of non-target host species for *A. orientalis* demonstrated that 15 out of 34 non-target host species were suitable hosts for *A. orientalis*. This work indicated *H. halys* eggs could successfully support development of *A. orientalis* (offspring sex ratio was strongly male biased), a finding contrary to previous reports (43). *Antheraea* sp. (Saturniidae) eggs (i.e., *A.*

TABLE 4 Average hind tibia length (mm) of *A. orientalis* females and males that emerged from eggs of non-target species.

| Host species                      | Average of hind tibia length (mm) |    |                 |    |
|-----------------------------------|-----------------------------------|----|-----------------|----|
|                                   | Males                             | n  | Females         | n  |
| <i>Acanthocephala thomasi</i>     | –                                 | –  | 1.04            | 1  |
| <i>Chinavia hilaris</i>           | 0.485 ± 0.005 cd                  | 9  | 0.613 ± 0.007 c | 4  |
| <i>Gloveria arizonensis</i>       | 0.587 ± 0.022 b                   | 10 | 0.881 ± 0.018 b | 6  |
| <i>Halyomorpha halys</i>          | 0.492 ± 0.013 c                   | 10 | 0.629 ± 0.008 c | 10 |
| <i>Lycorma delicatula</i>         | 0.689 ± 0.018 a                   | 10 | 0.959 ± 0.007 a | 10 |
| <i>Nezara viridula</i>            | 0.367 ± 0.004 e                   | 10 | –               | –  |
| <i>Pachysphinx occidentalis</i>   | 0.501                             | 1  | –               | –  |
| <i>Poblicia fuliginosa</i>        | 0.433 ± 0.011 d                   | 10 | –               | –  |
| <i>Pseudohemihyalea edwardsii</i> | 0.309 ± 0.005 f                   | 10 | –               | –  |

Different letters indicate significant differences between hind tibia size of parasitoids that emerged from different host species.





*pernyi*) have been reported to be suitable reproductive hosts for *A. orientalis* (43). However, parasitization of *Antheraea ocella* Neumoegen eggs by *A. orientalis* was not recorded in this study. Five new hemipteran hosts in three families (Coreidae [*A. thomasi* {native}], Fulgoridae [*P. fuliginosa* {native}], and Pentatomidae [*C. hilaris* {native}, *H. halys* {invasive}, *N. viridula* {invasive}]) and ten lepidopteran hosts in four families (Erebidae [*P. edwardsii* {native}], Lasiocampidae [*G. arizonensis* {native}], Saturniidae [*A. luna* {non-native}, *A. anona* {native}, *A. cecrops pamina* {native}, *E. calleta* {native}, *H. euryalus* {native}, *R. cincta* {native}, *S. walterorum* {native}] and Sphingidae [*P. occidentalis* {native}]) were identified as suitable hosts and are added to an increasing list of identified species that *A. orientalis* can successfully use as reproductive hosts in the laboratory (Table 2).

In terms of percent parasitism, eggs of two non-target species, *C. hilaris* (Pentatomidae [native]) and *A. anona* (Saturniidae [native]), were similar to or better hosts, than the target, *L. delicatula*. In the first case, *C. hilaris*, percent parasitism ( $68.79\% \pm 12.41$ ; sex ratio  $0.22 \pm 0.1$ ) was similar to the target, *L. delicatula* ( $58.24 \pm 10.09$ ; sex ratio  $0.77 \pm 0.07$ ). This result may need to be interpreted cautiously, because *C. hilaris* eggs masses had on average a smaller number of eggs ( $17.73 \pm 2.51$ ) when compared to the average number of eggs per egg mass ( $39.08 \pm 3.84$ ) from the target, *L. delicatula*. In the second case, *A. anona*, percent parasitism was higher ( $76.48 \pm 14.10$ ; sex ratio 0) than the target *L. delicatula* ( $58.24 \pm 10.09$ ), and both egg masses were approximately equal in size (i.e.,  $42.2 \pm 7.43$  eggs and  $39.08 \pm 3.84$  eggs for *A. anona* and *L. delicatula*, respectively). Collectively, data reported here and findings from companion studies (i.e., Broadley et al. [USDA], Submitted) suggest that *A. orientalis*, is at a minimum oligophagous, but more likely to be a polyphagous species.

Similar results from host specificity tests from other *Anastatus* spp. have been reported further supporting findings that *Anastatus* spp. potentially have broad host ranges. Stahl et al. (32, 33) studied the physiological host range of *A. bifasciatus* (this species is native

to Europe) for use as a biological control agent against *H. halys*, an invasive pest in Europe. In this study, *A. bifasciatus* successfully parasitized eggs of eight pentatomid species (including *N. viridula* [tested in this study]) and 14 lepidopteran species belonging to seven different families (Endromidae, Erebididae, Lasiocampidae, Notodontidae, Papilionidae, Saturniidae, and Sphingidae). Results reported here indicate that *A. orientalis* can also parasitize hosts eggs from Erebididae, Lasiocampidae, Saturniidae and Sphingidae. Host range tests that expose *A. orientalis* females to eggs of Notodontidae and Papilionidae, two lepidopteran families with species representation in the southwest U.S., may be warranted to further understanding of potential non-target host use (the family Endromidae is not present in North America). For *Anastatus* spp., results reported here and those of Stahl et al. (32, 33), indicate strongly that selecting non-target species which are restricted to close taxonomic relatedness to the target pest (e.g., families in the Fulgoroidea [Hemiptera: Auchenorrhyncha]) may be inadequate as *Anastatus* spp. may tend to be generalists capable of parasitizing species across different orders.

No additional significant levels of excess mortality (i.e., mortality from causes other than parasitism) to non-target species exposed to *A. orientalis* was observed (Figure 3). Failure to detect excess mortality of non-target eggs exposed to *A. orientalis* when compared to levels of naturally-occurring mortality in control eggs not exposed to *A. orientalis* may have at least two explanations: i) parasitized non-target eggs develop successfully when parasitoid larvae and/or envenomation failed to kill the host egg (host eggs are incapable of encapsulating parasitoid eggs and larvae), or ii) *A. orientalis* females only host fed on eggs that were parasitized and it is possible that *A. orientalis* is a concurrent parasitoid (i.e., host feeds on parasitized eggs). Thus, egg mortality from host feeding alone was not observed and egg mortality was attributed to solely to parasitism. Additional studies confirming the lack of excess mortality due to unsuccessful parasitization or host feeding following exposure of non-target host species to *A. orientalis* are needed. Excess mortality of non-target host eggs following exposure to *A. orientalis* should be considered an important deleterious non-target impact if it occurs (44).

For three non-target host species tested, *A. oslari* (Saturniidae [native]), *C. vanduzeei* (Fulgoridae [native]) and *C. vittiger* (Coreidae [native]), all parasitoid females exposed to egg masses of these species failed to parasitize *L. delicatula* egg masses in sequential exposure tests. For *C. vanduzeei* and *C. vittiger*, too few non-target eggs were available for experiments and replication was low for each test species and therefore were not excluded from the results. Interestingly, for female *A. orientalis* exposed to *A. oslari* eggs excess egg mortality increased slightly but not significantly, and all females ( $n=16$ ) died following exposure to *A. oslari* eggs prior to sequential exposure to *L. delicatula* egg masses. These results, a slight but non-significant increase in excess egg mortality and premature mortality of females, suggest that *A. orientalis* females may have host fed on *A. oslari* eggs and egg contents were possibly toxic to parasitoids. This could be explained by chemical protection of eggs by secondary plant compounds, like tannins, which are used as chemical defenses by host plants (i.e., *Quercus* spp.) of *A. oslari* (45). Sequestration of protective

compounds, like tannins, in *A. oslari* eggs could reduce survivorship rates of third trophic level organisms like *A. orientalis* (46).

To evaluate a more realistic exposure time of *A. orientalis* to non-target species, 24 hour exposure tests (as opposed to a 7 day exposure period) were performed with five non-target species (*C. hilaris*, *G. arizonensis*, *H. halys*, *N. viridula* and *S. walterorum*). *Anastatus orientalis* was able to parasitize the five non-target species in this shorter exposure period following the seven-day pre-oviposition exposure period on *L. delicatula* eggs. These findings suggest that prior host exposure, especially to the target, *L. delicatula*, does not deter use of subsequent non-target host eggs. In addition, when non-target and target host eggs were exposed to *A. orientalis* females at the same time (i.e., choice experiments), that non-target species were attacked under both exposure time scenarios (i.e., 24 hours and 7 days). These results further suggest that *A. orientalis* is probably a generalist parasitoid capable of using any suitable non-target host species upon encounter. In many instances, female parasitoids engaged in parasitism within minutes of introduction into test arenas and contact with non-target eggs (pers. obs. F. Gomez Marco).

The age of non-target host eggs can affect the acceptance behavior of *A. orientalis* females and rates of successful parasitism. For example, percent parasitism by *A. orientalis* was higher on young eggs (<24 hours of age) vs. older eggs (>24 hours to 3 days) of *H. halys*. However, for two other non-target species, *G. arizonensis* and *N. viridula*, tested in egg age acceptance studies, no significant effect of egg age on parasitism was found. Therefore, age of non-target eggs and the non-target species may be an important covariables to consider when host tests are being designed and executed. Additionally, defensive chemical compounds (see above) may also affect the acceptance behavior of parasitoids (and survivorship rates) and this may also warrant consideration in design, analysis, and interpretation of host range tests (46).

The sex ratio of *A. orientalis* offspring was strongly dependent on host egg size. All non-target host eggs which were parasitized by *A. orientalis* with an average egg size visibly (not measured in this study) smaller than *L. delicatula* produced a male biased sex ratio (< 0.5) and smaller adult males and females. However, the sex ratio from hosts which produced smaller *A. orientalis* males decreased, indicating that there was a strong correlation between offspring size and offspring sex ratio. Only one non-target host with eggs larger than *L. delicatula* eggs (pers. obs. F. Gomez Marco) was used in our tests (*A. thomasi*) and this resulted in the largest female parasitoid (hind tibia size of 1.04 mm) obtained from host-range studies. These findings tentatively support conditional sex allocation theory, where, with decreasing host quality (i.e., host egg size in this study), parasitoid offspring sex ratio becomes more male biased and males are generally smaller, both of which correlate with decreased fitness (47). In support of results presented here, two previous studies, Hou et al. (48) and Stahl et al. (32), reported more male-biased sex ratios when *Anastatus* spp. were reared on host eggs that were smaller than the target host. Interestingly, differences in *A. orientalis* offspring sex ratio from two non-target host species, *C. hilaris* and *H. halys*, existed when egg masses were exposed for 24 hours (*C. hilaris* and *H. halys*: 0, no females) or seven days (*C. hilaris*: 0.22 ± 0.1, *H. halys*: 0.05 ± 0.02). This finding might indicate a preference

of *A. orientalis* females to first oviposit (within at least the first 24 hours) non-fertilized eggs (i.e., produce male offspring) and then oviposit fertilized eggs (i.e., produce female offspring) when exposure times are longer and there is more time to repeatedly assess host quality. Sex ratio of offspring has important demographic implications as it affects rates of population growth. Male-biased progeny production on non-target hosts of marginal quality may limit or negate adverse population-level impacts on non-target species (32).

*Anastatus orientalis* offspring reared from non-target host species successfully parasitized *L. delicatula* egg masses. Percent parasitism and sex ratios resulting from mated couples that emerged from the same non-target host species were similar to values recorded for *A. orientalis* reared continuously on *L. delicatula*. Interestingly, males that emerged from *G. arizonensis* and *P. fuliginosa* when paired with unmated *A. orientalis* females that emerged from *L. delicatula* failed to produce any offspring. Due to the low number of repetitions for these experiments this finding should be viewed with caution. However, no obvious biological explanation (i.e., size of the males or inability to copulate with females) was observed to explain these results as unmated *A. orientalis* females should be able to oviposit unfertilized eggs that produce male offspring without mating with males that emerged from *G. arizonensis* and *P. fuliginosa*.

In laboratory studies in a quarantine facility, sequential no-choice and choice exposure studies that exposed female *A. orientalis* to non-target eggs of 34 native and non-native species in three orders and 12 families and target eggs (*L. delicatula*) for either 24 hours or 7 days, indicate that this egg parasitoid potentially has a wide host range as it successfully parasitized 15 species in 6 families in two orders (Hemiptera and Lepidoptera). Results presented here are for *A. orientalis* (Haplotype C). Other haplotypes of *A. orientalis* have been identified and are being assessed to determine if differences (i.e., greater specificity) in host preferences exist (20). A well-founded criticism of host-range tests is that parasitoids are constrained in small ventilated containers spaces with easily accessible hosts for long periods of time (i.e., 24 hours to 7 days) and are unable to engage in behaviors (e.g., rapid abandonment of patches with sub-optimal hosts) that could reduce or eliminate non-target use (49–52). Consequently, in the absence of comprehensive field data on host use and non-target species - target species - parasitoid phenology in the native range (i.e., China) from where *A. orientalis* was sourced, it is difficult to determine if high levels of non-target host use observed in host range tests reported here occurs in the field. Until detailed field data from the native range are available and based on results of host range tests reported here that suggest *A. orientalis* [and possibly most species of *Anastatus* (23–27, 32, 33)] has a very broad host range, use of this natural enemy in classical biological control of *L. delicatula* in the western U.S. should be assessed critically and with extreme caution.

## Data availability statement

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

## Author contributions

FG-M conceived and designed the experiments, performed collections and experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft. DY performed collections, reviewed drafts of the paper, and approved the final draft. MR performed collections, reviewed drafts of the paper, and approved the final draft. MH obtained the funding, conceived and designed the experiments, performed collections authored or reviewed drafts of the paper, and approved the final draft. All authors contributed to the article and approved the submitted version.

## Funding

Funding for this work was provided by the California Department of Agriculture Office of Environmental Farming and Innovation's Proactive Integrated Pest Management Solutions Grant Program, award number 18-0632-000-SG "Proactive Biological Control of Spotted Lantern Fly, *Lycorma delicatula* (Hemiptera: Fulgoridae)".

## Acknowledgments

We thank Dr. Juli Gould, Dr. Hannah Broadley (USDA APHIS PPQ, Forest Pest Methods Laboratory), and Dr. Kim Hoelmer (USDA ARS, Beneficial Insects Introduction Research Unit) for their assistance and contributions in starting and maintaining this project and for providing *A. orientalis* to start colonies at UC Riverside. We are indebted to USDA APHIS field crews for their enthusiastic assistance with identification of infested field sites and *L. delicatula* egg mass collections at field sites in Pennsylvania over winter. We acknowledge Prof. Jocelyn Millar, Prof. Kent Deena,

Prof. Richard Stouthamer and Prof. Thomas Perring and their respective lab members for providing non-target eggs. Ruth Amrich and Ishika Sameth provided assistance with colony maintenance and experiments in the Insectary & Quarantine facility at UC Riverside. We thank Bryan David, Dr. Serguei Triapitsyn (Principal Museum Scientist at UC Riverside) Michael Wilson, and Dr. Steve McElfresh for their help to collect non-target native fauna. Dr. Jorge Torres, Departamento de Agronomia, Entomologia, Universidade Federal Rural de Pernambuco, Recife, Brazil provided extraordinary assistance with establishment and maintenance of pentatomid colonies during his 12-month sabbatical program with the Hoddle lab. We thank the American Museum of Natural History's Southwestern Research Station (Portal, Arizona) team for generous assistance and access to field sites. Dr. A. James Downer, Barbara Roth, and Reed Peters (Cave Creek Ranch) for hosting us during our summer field trips to Portal, Arizona.

## Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

## Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

## References

- Hoddle MS, Lake EC, Minter CR, Daane KM. Importation biological control. In: Mason PG (ed). *Biological control: Global impacts, challenges and future directions of pest management*. Clayton South: CSIRO PUBLISHING. (2021). 67–89.
- United States Department of Agriculture (USDA). *Technical advisory group for biological control agents of weeds manual* Washington DC. (2017) (Accessed October 15, 2022).
- Hoddle MS, Mace K, Steggall J. Proactive biological control: a cost-effective management option for invasive pests. *California Agric* (2018) 72:148–50. doi: 10.3733/ca.2018a0027
- Kim H, Kim M, Kwon DH, Park S, Lee Y, Huang J, et al. Molecular comparison of *Lycorma delicatula* (Hemiptera: Fulgoridae) isolates in Korea, China, and Japan. *J Asia-Pac Entomol* (2013) 16:503–6. doi: 10.1016/j.aspen.2013.07.003
- Barringer LE, Donovall LR, Spichiger S-E, Lynch D, Henry D. The first new world record of *Lycorma delicatula* (Insecta: Hemiptera: Fulgoridae). *Entomol News* (2015) 125:20–4. doi: 10.3157/021.125.0105
- Management NYSSIP. *Spotted lanternfly range in the U.S* (2022) (Accessed 2022 November).
- Dara SK, Barringer L, Arthurs SP. *Lycorma delicatula* (Hemiptera: Fulgoridae): a new invasive pest in the united states. *J Int Pest Manag* (2015) 6:1–6. doi: 10.1093/jipm/pmv021
- Urban JM, Calvin D, Hills-Stevenson J. Early response (2018–2020) to the threat of spotted lanternfly, *Lycorma delicatula* (Hemiptera: Fulgoridae) in Pennsylvania. *Ann Entomol Soc Am* (2021) 114:709–18. doi: 10.1093/aesa/saab030
- Liu H, Hartlieb RJ. Spatial distribution of *Lycorma delicatula* (Hemiptera: Fulgoridae) egg masses on tree-of-heaven, black walnut, and Siberian elm in north America. *J Econ Entomol* (2020) 113:1028–32. doi: 10.1093/jeet/toz350
- Nixon LJ, Jones SK, Tang L, Urban J, Felton K, Leskey TC. Survivorship and development of the invasive *Lycorma delicatula* (Hemiptera: Fulgoridae) on wild and cultivated temperate host plants. *Environ Entomol* (2022) 51:222–8. doi: 10.1093/ee/nvab137
- Lee D-H, Park Y-L, Leskey TC. A review of biology and management of *Lycorma delicatula* (Hemiptera: Fulgoridae), an emerging global invasive species. *J Asia-Pac Entomol* (2019) 22:589–96. doi: 10.1016/j.aspen.2019.03.004
- Baker TC, Smyers EC, Urban JM, Meng Z, Pagadala Damadaram KJ, Myrick AJ, et al. Progression of seasonal activities of adults of the spotted lanternfly, *Lycorma delicatula*, during the 2017 season of mass flight dispersal behavior in eastern Pennsylvania. *J Asia-Pac Entomol* (2019) 22:705–13. doi: 10.1016/j.aspen.2019.05.006
- Urban JM, Leach H. Biology and management of the spotted lanternfly, *Lycorma delicatula* (Hemiptera: Fulgoridae), in the united states. *Ann Rev Entomol* (2023) 68:151–67. doi: 10.1146/annurev-ento-120220-111140

14. Wakie TT, Neven LG, Yee WL, Lu Z. The establishment risk of *Lycorma delicatula* (Hemiptera: Fulgoridae) in the united states and globally. *J Econ Entomol* (2019) 113:306–14. doi: 10.1093/jeet/toz259
15. California Department of Food and Agriculture. *California Agricultural production statistics* (2019) (Accessed 2022 February).
16. Choi M-Y, Yang Z-Q, Wang X-Y, Tang Y-L, Hou Z-R, Kim JH, et al. Parasitism rate of egg parasitoid *Anastatus orientalis* (Hymenoptera: Eupelmidae) on *Lycorma delicatula* (Hemiptera: Fulgoridae) in China. *K J App Entomol* (2014) 53:135–9. doi: 10.5656/KSAE.2014.01.1.075
17. Yang Z-Q, Choi W-Y, Cao L-M, Wang X-Y, Hou Z-R. A new species of *Anastatus* (Hymenoptera: Eupelmidae) from China, parasitizing eggs of *Lycorma delicatula* (Homoptera: Fulgoridae). *Zool Syst* (2015) 40:290–302. doi: 10.11865/zs.20150305
18. Xin B, Zhang Y-l, Wang X-y, Cao L-m, Hoelmer KA, Broadley HJ, et al. Exploratory survey of spotted lanternfly (Hemiptera: Fulgoridae) and its natural enemies in China. *Environ Entomol* (2020) 50:36–45. doi: 10.1093/ee/nvaa137
19. Broadley HJ, Gould JR, Sullivan LT, Wang X-y, Hoelmer KA, Hickin ML, et al. Life history and rearing of *Anastatus orientalis* (Hymenoptera: Eupelmidae), an egg parasitoid of the spotted lanternfly (Hemiptera: Fulgoridae). *Environ Entomol* (2021) 50:28–35. doi: 10.1093/ee/nvaa124
20. Wu Y, Broadley HJ, Vieira KA, McCormack JJ, Losch C, McGraw AR, et al. Cryptic genetic diversity and associated ecological differences of *Anastatus orientalis*, an egg parasitoid from China of the spotted lanternfly. *Front Insect Sci – Invasive Insect Species*. (Submitted).
21. Askew A. A new European species of *Anastatus motschulsky* (Hym., eupelmidae). *Entomol Monthly Mag* (2005) 141:1697–9.
22. Jones WA. World review of the parasitoids of the southern green stink bug, *Nezara viridula* (L.) (Heteroptera: Pentatomidae). *Ann Entomol Soc Am* (1988) 81:262–73. doi: 10.1093/aesa/81.2.262
23. Burks BD. The north American species of *Anastatus motschulsky* (Hymenoptera, eupelmidae). *Trans Am Entomological Soc* (1967) . 93:423–32.
24. Pieltain BC. *Estudio monográfico de las especies españolas del género anastatus Motsch. (Hym. chalc.)*. Revista Española de Entomología 10. (1935). 273–92.
25. Chen YM, Gibson GAP, Peng LF, Iqbal A, Zang LS. *Anastatus motschulsky* (Hymenoptera, eupelmidae): egg parasitoids of *Caligula japonica* Moore (Lepidoptera, saturniidae) in China. *Zookeys* (2019) 881:109–34. doi: 10.3897/zookeys.881.34646
26. Noyes JS. *Universal chalcidoidea database*. World Wide Web electronic publication (2019). (Accessed October 15, 2022).
27. Hayat M. Some Indian species of *Anastatus* (Hymenoptera: Chalcidoidea, eupelmidae). *Oriental Insects* (1975) 9:261–71. doi: 10.1080/00305316.1975.10434497
28. Crossman S. Two imported egg parasites of the gipsy moth, *Anastatus bifasciatus* fonsc and *Schedius kuvanae* Howard. *J Agric Res* (1925) 30:643–75.
29. Chen Y, Lin L, Chen J. Preliminary report of study on the release of *Anastatus* for the control of the stink bug (*Tessaratomia papillosa*). *Fujian Agric Sci Technol* (1990) 2:15–6.
30. Huang M, Mai S, Wu W, Poo C. The bionomics of *Anastatus* sp. and its utilization for the control of lichee stink bug, *Tessaratomia papillosa* Drury. *Act Entomol Sin* (1974) 17:362–75.
31. Shrestha R. Productivity improvement of citrus fruits through effective fruit drop management technique in the mid and far Western development region of Nepal. *Tech Rep* (2011).
32. Stahl JM, Babendreier D, Haye T. Using the egg parasitoid *Anastatus bifasciatus* against the invasive brown marmorated stink bug in Europe: Can non-target effects be ruled out? *J Pest Sci* (2018) 91:1005–17. doi: 10.1007/s10340-018-0969-x
33. Stahl JM, Babendreier D, Marazzi C, Caruso S, Costi E, Maistrello L, et al. Can *Anastatus bifasciatus* be used for augmentative biological control of the brown marmorated stink bug in fruit orchards? *Insects* (2019) 10(4):108. doi: 10.3390/insects10040108
34. Ogburn EC, Bessin R, Dieckhoff C, Dobson R, Grieshop M, Hoelmer KA, et al. Natural enemy impact on eggs of the invasive brown marmorated stink bug, *Halyomorpha halys* (Stål) (Hemiptera: Pentatomidae), in organic agroecosystems: a regional assessment. *Biol Control* (2016) 101:39–51. doi: 10.1016/j.biocontrol.2016.06.002
35. Govender AW. *Australian Fruitspotting bugs, amblypelta nitida stål and a. lutescens lutescens distant* (Hemiptera: Coreidae), and the potential for their biologically based management in macadamia orchards. Queensland, Australia: University of Queensland (2015).
36. Huwer R, Maddox C, Fay H, Llewellyn R, Danne A, Hickey M, et al. A multi target approach to fruitspotting bug management. In: *Proceedings of the VII world avocado congress*. Cairns, Australia (2011).
37. Bartlett CR. *Planthoppers of north America* (2020) (Accessed February 2019).
38. Henderson CF, Tilton EW. Tests with acaricides against the brown wheat mite. *J Econom Entomol* (1955) 48:157–61. doi: 10.1093/jeet/48.2.157
39. Gebiola M, Gomez-Marco F, Simmons GS, Stouthamer R. Effect of host feeding on life history traits of *Tamarixia radiata*, parasitoid of the Asian citrus psyllid, diaphorina citri. *BioControl* (2018) 63:763–71. doi: 10.1007/s10526-018-9903-7
40. Ameri M, Rasekh A, Michaud JP, Allahyari H. Morphometric indicators for quality assessment in the aphid parasitoid, *Lysiphlebus fabarum* (Braconidae: Aphidiinae). *Eur J Entomol* (2013) 110:519–25. doi: 10.14411/eje.2013.069
41. R Core Team. *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing (2022).
42. RStudio Team. *RStudio: Integrated development for r*. Boston, MA: RStudio, PBC (2020).
43. Seo M, Kim JH, Seo BY, Park C, Choi BR, Kim KH, et al. Mass-rearing techniques of *Anastatus orientalis* (Hymenoptera: Eupelmidae), as the egg-parasitoid of *Lycorma delicatula* (Hemiptera: Fulgoridae): an using method of *Antheraea pernyi* (Lepidoptera: Saturniidae) and *I. delicatula* eggs in laboratory. *K J App Entomol* (2018) 57:243–51. doi: 10.5656/KSAE.2018.08.0.035
44. Abram PK, Brodeur J, Urbaneja A, Tena A. Nonreproductive effects of insect parasitoids on their hosts. *Annu Rev Entomol* (2019) 64:259–76. doi: 10.1146/annurev-ento-011118-111753
45. Forkner RE, Marquis RJ, Lill JT. Feeny revisited: condensed tannins as anti-herbivore defences in leaf-chewing herbivore communities of quercus. *Ecol Entomol* (2004) 29:174–87. doi: 10.1111/j.1365-2311.2004.0590.x
46. Fatouros NE, Cusumano A, Bin F, Polaszek A, van Lenteren JC. How to escape from insect egg parasitoids: A review of potential factors explaining parasitoid absence across the insecta. *Proc R Soc B: Biol Sci* (2020) 287:20200344. doi: 10.1098/rspb.2020.0344
47. Charnov EL, Los-den Hartogh RL, Jones WT, van den Assem J. Sex ratio evolution in a variable environment. *Nature* (1981) 289:27–33. doi: 10.1038/289027a0
48. Hou Z, Liang H, Chen Q, Hu Y, Tian H. Application of *Anastatus* sp. against *Halyomorpha halys*. *For Pest Dis* (2009) 4:39.
49. Haye T, Goulet H, Mason PG, Kuhlmann U. Does fundamental host range match ecological host range? a retrospective case study of a lygus plant bug parasitoid. *Biol Control* (2005) 35:55–67. doi: 10.1016/j.biocontrol.2005.06.008
50. Haye T, Mason PG, Gillespie DR, Miall JH, Gibson GAP, Diaconu A, et al. Determining the host specificity of the biological control agent *Trichomalus perfectus* (Hymenoptera: Pteromalidae): The importance of ecological host range. *Biocontrol Sci Techn* (2015) 25:21–47. doi: 10.1080/09583157.2014.945900
51. van Lenteren JC, Loomans AJ. Environmental risk assessment: Methods for comprehensive evaluation and quick scan. In: Bigler F, Babendreier D, Kuhlmann U, editors. *Environmental impact of invertebrates for biological control of arthropods: methods and risk assessment*, vol. 10. Wallingford: CABI Publishing (2006). p. 254–72.
52. Whitters T, Mansfield S. Choice or no-choice tests? effects of experimental design on the expression of host range. In: Hoddle M (ed), *Second international symposium on biological control of arthropods*. Davos, Switzerland: USDA Forest Service Bulletin, Morgantown, West Virginia, USA (2005). 620–32.