

**UCLA**

**Proceedings of UCLA Health**

**Title**

A Case of Amyloid Cardiomyopathy

**Permalink**

<https://escholarship.org/uc/item/6s30w4z6>

**Journal**

Proceedings of UCLA Health, 24(1)

**Author**

Khafaf, Mana

**Publication Date**

2020-11-09

## CLINICAL VIGNETTE

---

# A Case of Amyloid Cardiomyopathy

---

Mana Khafaf, MD

A 65-year-old male presented with episodes of chest pain and shortness of breath. He had a long history of medical non-compliance with smoking, obesity and hypertension. Myocardial perfusion test showed nonspecific apical changes more apparent at rest, suggestive of prior injury versus artifact. His left ventricular chamber size appeared enlarged with hypertrophied muscle mass. Initial prior echocardiogram showed left atrial enlargement with moderately increased ventricular wall thickness. This progressed to severely increased left ventricular wall thickness and severe bi-atrial enlargement on repeat echocardiogram a few years later. He still had preserved ejection fraction of 65 to 70%, with no evidence of pulmonary hypertension.

As his shortness of breath worsened, chest CT scan demonstrated persistent enlargement of the main pulmonary artery with pleural effusion. Abdominal ultrasound showed hepatosplenomegaly.

With suspicion of infiltrative heart disease, cardiology ordered studies for restrictive cardiomyopathy. Kappa/Lambda ratio was abnormal, and Hematology recommended obtaining tissue for possible amyloid infiltration. Abdominal fat pad biopsy showed only benign fibrofatty tissue. Endomyocardial biopsy and cardiac catheterization confirmed left ventricular hypertrophy, normal right heart and pulmonary capillary wedge pressures. Biopsies of septal wall supported the diagnosis of amyloidosis and indicated AL (lambda)-type amyloid deposition. Bone marrow biopsy was compatible with plasma cell myeloma. Flow cytometry demonstrated monotypic lambda restricted plasma cells, in addition to evidence of amyloid deposit by Congo red stain. He was started on chemotherapy with CyBORd Regimen. Disease progression coupled with medication non-compliance and delayed treatments led to worsened cardiomyopathy, heart failure and atrial flutter with left atrial thrombus and he was started on anticoagulation. Renal involvement progressed to end stage renal disease dependent on dialysis.

He developed hepatomegaly with elevated liver enzymes, which worsened to acute liver failure likely due to ischemic hepatic insult with occasional low blood pressures.

With further deterioration treatment was directed at comfort and he died less than two years after his diagnosis of AL cardiac Amyloidosis.

### **Introduction**

Amyloidosis is caused by the deposition of one of over 30 different precursor proteins which fold abnormally into a beta pleated sheet configuration. The unstable tertiary structure of these proteins cause them to aggregate and deposit as insoluble amyloid fibrils in the extracellular space of different organs including the heart.<sup>1</sup>

Amyloidosis is classified based on the precursor protein type. Amyloid deposits stained with Congo red dye show an apple-green birefringence, while they display a cross-beta-pleated sheet configuration under electron microscopy.

More than 95% of all clinical cardiac amyloidosis cases are attributed to transthyretin amyloidosis (ATTR) or light chain amyloidosis (AL).<sup>2-4</sup> ATTR is caused by deposition of dissociated monomers and oligomers of TTR as amyloid fibrils. Transthyretin (TTR), previously known as prealbumin is synthesized in the liver, circulates as a stable tetramer which transports thyroid hormone and vitamin A. "Wild type ATTR" is an under-diagnosed cause of heart failure, found in about 13% of the patients who present with heart failure with preserved ejection fraction.<sup>5</sup>

Primary systemic amyloidosis (AL) is due to immunoglobulin light chain deposits related to a plasma cell dyscrasia. This type of amyloidosis is rare, with incidence of about 1 per 100,000 representing about 2500-5000 new cases per year in the United States.<sup>6</sup>

Other rare causes of amyloid cardiomyopathy are serum amyloid A amyloidosis (AA) and apolipoprotein A-1 (ApoA-1) amyloidosis which are not discussed.

### **Clinical Manifestation**

Age of onset of amyloidosis symptoms varies based on the type of amyloid deposit.<sup>6</sup> AL (light chain) cardiac amyloidosis usually has an age of onset of above 40, compared to ATTR cardiac amyloidosis which usually presents at above 60-year-old patients. Systemic AL amyloidosis can involve the liver, kidneys, spleen, autonomic and peripheral nervous system, lungs and heart. Fatigue, carpal tunnel syndrome, GI bleed, macroglossia, periorbital purpura and bleeding diathesis are some manifestations.

Majority of AL amyloidosis patients (50-70%) have cardiac infiltration which determines their prognosis.<sup>7,8</sup> Cardiac

amyloidosis usually presents with restrictive cardiomyopathy with right ventricular failure symptoms, including lower extremity edema, elevated JVP, hepatic congestion, ascites and dyspnea. As the disease progresses, low cardiac output symptoms develop.

The worse outcome of amyloidosis caused by AL (light chains) compared to ATTR patients may be attributed to the toxicity of light chains to the myocardial cells.<sup>9,10</sup> Bradyarrhythmia and advanced atrioventricular block are most common causes of syncope in amyloidosis patients. Progressive conduction system disease needing pacemaker implantation is more common in ATTR cases. High degree atrioventricular block or symptomatic sinus node dysfunction is not as prevalent in AL amyloidosis.<sup>11</sup>

The risk of cardiac thromboembolism is increased in AL amyloidosis with atrial fibrillation. amyloid deposition in atrial and ventricular walls contributes to electromechanical dissociation during sinus rhythm which increases the risk of atrial thrombus formation.<sup>12,13</sup>

Restrictive cardiomyopathy associated with ATTR can lead to low gradient, low flow aortic stenosis.<sup>14</sup> Some non-cardiac manifestations of wild type ATTR are spinal stenosis, biceps tendon rupture, and bilateral carpal tunnel syndrome with amyloid deposits found with decompression procedures.<sup>15</sup>

### **Diagnosis**

Initial tests can reveal proteinuria and renal failure, elevated liver enzymes with congestive hepatopathy, elevated natriuretic peptide and troponin levels. Cardiac enzymes are usually mildly elevated in ATTR compared to higher levels in AL amyloidosis which is likely secondary to light chains cardiotoxicity. Common EKG findings are increased left ventricular wall thickness and reduced QRS voltage, the latter seen most in AL amyloidosis compared to ATTR cases.<sup>16,17</sup>

About 30 percent of patients with wild type ATTR, have left ventricular hypertrophy or left bundle branch block with pseudo-infarction patterns seen in 70 percent of the cases in addition to conduction abnormalities. About 15 percent of patients also have atrial fibrillation. Cardiac amyloidosis should be suspected when unexplained LVH is noted on echocardiogram, low gradient/low flow aortic stenosis and heart failure.<sup>14,18</sup>

Early impaired ventricular relaxation leads to diastolic dysfunction with progression to restrictive physiology.<sup>19</sup> Marked elevation in LV diastolic pressure can cause pulmonary hypertension. In AL amyloidosis, small pericardial and pleural effusions may also be present.

After physical exam, echocardiograms are initially ordered in suspected cases with reduced global longitudinal strain as an early marker with high sensitivity and specificity.<sup>20</sup> Left ventricular hypertrophy is usually non-dilated with small

ventricles. Thickening of the valves and interatrial septum, speckled myocardium and bi-atrial dilatation are common. Symmetric LVH is seen more in AL amyloidosis versus asymmetry with predominant septal hypertrophy being more common in ATTR. Cardiovascular magnetic resonance imaging (CMR) can detect cardiac abnormalities prior to LVH development. It can assess cardiac structure, function and tissue characteristics, but cannot distinguish cardiac AL from ATTR.<sup>21,22</sup>

Inconsistent results usually make diagnosis of cardiac amyloidosis unlikely, other cause of LVH including hypertrophic cardiomyopathy, hypertension or other etiologies should be considered if the above tests are not conclusive.<sup>23</sup>

If the CMR results are consistent with possible diagnosis, serum kappa/lambda free light chain ratio analysis, serum protein immunofixation and urine protein immunofixation should be obtained to identify monoclonal proteins and bone marrow biopsy and evaluation for treatment options. Tissue biopsy from fat pad or other organs may be indicated for both diagnostic and typing purposes. In majority of the case, noncardiac tissue biopsy in addition to CMR findings is sufficient.<sup>24</sup> Typical apple-green birefringence with Congo red dye under polarized light microscopy and cross-beta-pleated sheets under electron microscopy are the diagnostic characteristics of extracellular amyloid deposits. Although immunohistochemistry, immunofluorescence and immunoelectron microscopy are used to determine the type of amyloid fibril, laser microdissection with mass spectrometry is considered the gold standard.<sup>25</sup>

If monoclonal proteins are not detected in above tests, a bone tracer cardiac scintigraphy scan can be performed for cardiac uptake evaluation and grading. Grade zero suggests that cardiac amyloidosis is unlikely and other causes of LVH should be considered. Grade 1 suggests additional tests including endomyocardial biopsy for confirmation. Grade 2 or 3, in addition to lack of evidence of plasma cell dyscrasia, is highly specific for ATTR cardiac disease and in these cases tissue biopsy is not needed.<sup>26</sup>

### **Treatment**

The two approaches in treating patients with amyloidosis are management of heart failure and treatment of the underlying disease. Heart failure treatment is more responsive in wild type ATTR compared to AL amyloidosis in which control of the plasma cell dyscrasia leads to rapid decrease in serum biomarkers of heart failure.<sup>27</sup>

In amyloidosis heart failure, treatment with loop diuretics is commonly used in addition to aldosterone therapy. The treatment options in these patients differs from other types of heart failure due to potential hypotension side effects of beta blockers, ACE inhibitors and calcium channel blockers. Amyloid fibrils can also bind to digoxin which can increase its toxicity. However, if atrial fibrillation is present, low dose beta

blockade and digoxin can be used cautiously to assist with rate control and symptom management. Amiodarone seems to be well tolerated in these patients as well.<sup>28</sup> Anticoagulation is used to decrease the increased risk of intraatrial thrombus, regardless of their CHADS2 or CHADS2VA2SC score. It is even suggested in patients with normal sinus rhythm due to associated atrial thrombus in AL type group.<sup>29,30</sup>

Since sudden cardiac death is common in cardiac AL amyloidosis patients and prophylactic implantable cardioverter-defibrillators (ICDs) have been suggested despite its unclear role in electromechanical dissociation.<sup>31</sup>

Heart transplant is not usually recommended for these patients as most of them have significant extracardiac amyloid associated disease. Heart transplant would not treat the underlying plasma cell dyscrasia which leads to its progression and involvement the transplanted heart.<sup>32-34</sup>

Prognosis of AL amyloidosis patients depends on the degree of their multiple organ involvement. Routine therapy for underlying plasma cell clones include chemotherapy and/or autologous stem cell transplant, with the goal of 90 percent reduction in serum free light chains. NYHA functional class III or IV Heart failure is considered a contraindication for autologous stem cell transplant.<sup>35</sup>

## REFERENCES

1. **Fontana M, Ćorović A, Scully P, Moon JC.** Myocardial Amyloidosis: The Exemplar Interstitial Disease. *JACC Cardiovasc Imaging.* 2019 Nov;12(11 Pt 2):2345-2356. doi: 10.1016/j.jcmg.2019.06.023. Epub 2019 Aug 14. PMID: 31422120.
2. **Falk RH, Alexander KM, Liao R, Dorbala S.** AL (Light-Chain) Cardiac Amyloidosis: A Review of Diagnosis and Therapy. *J Am Coll Cardiol.* 2016 Sep 20;68(12):1323-41. doi: 10.1016/j.jacc.2016.06.053. PMID: 27634125.
3. **Gertz MA, Benson MD, Dyck PJ, Grogan M, Coelho T, Cruz M, Berk JL, Plante-Bordeneuve V, Schmidt HHJ, Merlini G.** Diagnosis, Prognosis, and Therapy of Transthyretin Amyloidosis. *J Am Coll Cardiol.* 2015 Dec 1;66(21):2451-2466. doi: 10.1016/j.jacc.2015.09.075. PMID: 26610878.
4. **Sipe JD, Benson MD, Buxbaum JN, Ikeda SI, Merlini G, Saraiva MJ, Westermark P.** Amyloid fibril proteins and amyloidosis: chemical identification and clinical classification International Society of Amyloidosis 2016 Nomenclature Guidelines. *Amyloid.* 2016 Dec;23(4):209-213. doi: 10.1080/13506129.2016.1257986. Epub 2016 Nov 24. PMID: 27884064.
5. **González-López E, Gallego-Delgado M, Guzzo-Merello G, de Haro-Del Moral FJ, Cobo-Marcos M, Robles C, Bornstein B, Salas C, Lara-Pezzi E, Alonso-Pulpon L, Garcia-Pavia P.** Wild-type transthyretin amyloidosis as a cause of heart failure with preserved ejection fraction. *Eur Heart J.* 2015 Oct 7;36(38):2585-94. doi: 10.1093/eurheartj/ehv338. Epub 2015 Jul 28. PMID: 26224076.
6. **Wechalekar AD, Gillmore JD, Hawkins PN.** Systemic amyloidosis. *Lancet.* 2016 Jun 25;387(10038):2641-2654. doi: 10.1016/S0140-6736(15)01274-X. Epub 2015 Dec 21. PMID: 26719234.
7. **Merlini G, Bellotti V.** Molecular mechanisms of amyloidosis. *N Engl J Med.* 2003 Aug 7;349(6):583-96. doi: 10.1056/NEJMra023144. PMID: 12904524.
8. **Kyle RA, Linos A, Beard CM, Linke RP, Gertz MA, O'Fallon WM, Kurland LT.** Incidence and natural history of primary systemic amyloidosis in Olmsted County, Minnesota, 1950 through 1989. *Blood.* 1992 Apr 1;79(7):1817-22. PMID: 1558973.
9. **Liao R, Jain M, Teller P, Connors LH, Ngoy S, Skinner M, Falk RH, Apstein CS.** Infusion of light chains from patients with cardiac amyloidosis causes diastolic dysfunction in isolated mouse hearts. *Circulation.* 2001 Oct 2;104(14):1594-7. Erratum in: *Circulation.* 2013 Sep 10;128(11):e174. PMID: 11581134.
10. **Shi J, Guan J, Jiang B, Brenner DA, Del Monte F, Ward JE, Connors LH, Sawyer DB, Semigran MJ, Macgillivray TE, Seldin DC, Falk R, Liao R.** Amyloidogenic light chains induce cardiomyocyte contractile dysfunction and apoptosis via a non-canonical p38alpha MAPK pathway. *Proc Natl Acad Sci U S A.* 2010 Mar 2;107(9):4188-93. doi: 10.1073/pnas.0912263107. Epub 2010 Feb 11. PMID: 20150510; PMCID: PMC2840082.
11. **Mathew V, Olson LJ, Gertz MA, Hayes DL.** Symptomatic conduction system disease in cardiac amyloidosis. *Am J Cardiol.* 1997 Dec 1;80(11):1491-2. doi: 10.1016/s0002-9149(97)82785-3. PMID: 9399732.
12. **Martinez-Naharro A, Gonzalez-Lopez E, Corovic A, Mirelis JG, Baksi AJ, Moon JC, Garcia-Pavia P, Gillmore JD, Hawkins PN, Fontana M.** High Prevalence of Intracardiac Thrombi in Cardiac Amyloidosis. *J Am Coll Cardiol.* 2019 Apr 9;73(13):1733-1734. doi: 10.1016/j.jacc.2019.01.035. PMID: 30947929.
13. **El-Am EA, Dispenzieri A, Melduni RM, Ammash NM, White RD, Hodge DO, Noseworthy PA, Lin G, Pislaru SV, Egbe AC, Grogan M, Nkomo VT.** Direct Current Cardioversion of Atrial Arrhythmias in Adults With Cardiac Amyloidosis. *J Am Coll Cardiol.* 2019 Feb 12;73(5):589-597. doi: 10.1016/j.jacc.2018.10.079. PMID: 30732713; PMCID: PMC6378685.
14. **Castaño A, Narotsky DL, Hamid N, Khalique OK, Morgenstern R, DeLuca A, Rubin J, Chiuhan C, Nazif T, Vahl T, George I, Kodali S, Leon MB, Hahn R, Bokhari S, Maurer MS.** Unveiling transthyretin cardiac amyloidosis and its predictors among elderly patients with severe aortic stenosis undergoing transcatheter aortic valve replacement. *Eur Heart J.* 2017 Oct 7;38(38):2879-2887. doi: 10.1093/eurheartj/ehx350. PMID: 29019612; PMCID: PMC5837725.
15. **Sekijima Y, Uchiyama S, Tojo K, Sano K, Shimizu Y, Imaeda T, Hoshii Y, Kato H, Ikeda S.** High prevalence of wild-type transthyretin deposition in patients with idiopathic carpal tunnel syndrome: a common cause of carpal tunnel syndrome in the elderly. *Hum Pathol.* 2011

- Nov;42(11):1785-91. doi: 10.1016/j.humpath.2011.03.004. PMID: 21733562.
16. **Cyrille NB, Goldsmith J, Alvarez J, Maurer MS.** Prevalence and prognostic significance of low QRS voltage among the three main types of cardiac amyloidosis. *Am J Cardiol.* 2014 Oct 1;114(7):1089-93. doi: 10.1016/j.amjcard.2014.07.026. Epub 2014 Jul 18. PMID: 25212550.
  17. **Mussinelli R, Salinaro F, Alogna A, Boldrini M, Raimondi A, Musca F, Palladini G, Merlini G, Perlini S.** Diagnostic and prognostic value of low QRS voltages in cardiac AL amyloidosis. *Ann Noninvasive Electrocardiol.* 2013 May;18(3):271-80. doi: 10.1111/anec.12036. Epub 2013 Jan 20. PMID: 23714086; PMCID: PMC6932192.
  18. **Cavalcante JL, Rijal S, Abdelkarim I, Althouse AD, Sharbaugh MS, Fridman Y, Soman P, Forman DE, Schindler JT, Gleason TG, Lee JS, Schelbert EB.** Cardiac amyloidosis is prevalent in older patients with aortic stenosis and carries worse prognosis. *J Cardiovasc Magn Reson.* 2017 Dec 7;19(1):98. doi: 10.1186/s12968-017-0415-x. PMID: 29212513; PMCID: PMC5719789.
  19. **Knight DS, Zumbo G, Barcella W, Steeden JA, Muthurangu V, Martinez-Naharro A, Treibel TA, Abdel-Gadir A, Bulluck H, Kotecha T, Francis R, Rezk T, Quarta CC, Whelan CJ, Lachmann HJ, Wechalekar AD, Gillmore JD, Moon JC, Hawkins PN, Fontana M.** Cardiac Structural and Functional Consequences of Amyloid Deposition by Cardiac Magnetic Resonance and Echocardiography and Their Prognostic Roles. *JACC Cardiovasc Imaging.* 2019 May;12(5):823-833. doi: 10.1016/j.jcmg.2018.02.016. Epub 2018 Apr 18. PMID: 29680336.
  20. **Phelan D, Collier P, Thavendiranathan P, Popović ZB, Hanna M, Plana JC, Marwick TH, Thomas JD.** Relative apical sparing of longitudinal strain using two-dimensional speckle-tracking echocardiography is both sensitive and specific for the diagnosis of cardiac amyloidosis. *Heart.* 2012 Oct;98(19):1442-8. doi: 10.1136/heartjnl-2012-302353. Epub 2012 Aug 3. PMID: 22865865.
  21. **Brownrigg J, Lorenzini M, Lumley M, Elliott P.** Diagnostic performance of imaging investigations in detecting and differentiating cardiac amyloidosis: a systematic review and meta-analysis. *ESC Heart Fail.* 2019 Oct;6(5):1041-1051. doi: 10.1002/ehf2.12511. Epub 2019 Sep 5. PMID: 31487121; PMCID: PMC6816075.
  22. **Dorbala S, Cuddy S, Falk RH.** How to Image Cardiac Amyloidosis: A Practical Approach. *JACC Cardiovasc Imaging.* 2020 Jun;13(6):1368-1383. doi: 10.1016/j.jcmg.2019.07.015. Epub 2019 Oct 11. PMID: 31607664; PMCID: PMC7148180.
  23. **Martinez-Naharro A, Treibel TA, Abdel-Gadir A, Bulluck H, Zumbo G, Knight DS, Kotecha T, Francis R, Hutt DF, Rezk T, Rosmini S, Quarta CC, Whelan CJ, Kellman P, Gillmore JD, Moon JC, Hawkins PN, Fontana M.** Magnetic Resonance in Transthyretin Cardiac Amyloidosis. *J Am Coll Cardiol.* 2017 Jul 25;70(4):466-477. doi: 10.1016/j.jacc.2017.05.053. PMID: 28728692.
  24. **Ruberg FL, Grogan M, Hanna M, Kelly JW, Maurer MS.** Transthyretin Amyloid Cardiomyopathy: JACC State-of-the-Art Review. *J Am Coll Cardiol.* 2019 Jun 11;73(22):2872-2891. doi: 10.1016/j.jacc.2019.04.003. PMID: 31171094; PMCID: PMC6724183.
  25. **Nativi-Nicolau J, Maurer MS.** Amyloidosis cardiomyopathy: update in the diagnosis and treatment of the most common types. *Curr Opin Cardiol.* 2018 Sep;33(5):571-579. doi: 10.1097/HCO.0000000000000547. PMID: 30015648.
  26. **Gillmore JD, Maurer MS, Falk RH, Merlini G, Damy T, Dispenzieri A, Wechalekar AD, Berk JL, Quarta CC, Grogan M, Lachmann HJ, Bokhari S, Castano A, Dorbala S, Johnson GB, Glaudemans AW, Rezk T, Fontana M, Palladini G, Milani P, Guidalotti PL, Flatman K, Lane T, Vonberg FW, Whelan CJ, Moon JC, Ruberg FL, Miller EJ, Hutt DF, Hazenberg BP, Rapezzi C, Hawkins PN.** Nonbiopsy Diagnosis of Cardiac Transthyretin Amyloidosis. *Circulation.* 2016 Jun 14;133(24):2404-12. doi: 10.1161/CIRCULATIONAHA.116.021612. Epub 2016 Apr 22. PMID: 27143678.
  27. **Maurer MS, Elliott P, Comenzo R, Semigran M, Rapezzi C.** Addressing Common Questions Encountered in the Diagnosis and Management of Cardiac Amyloidosis. *Circulation.* 2017 Apr 4;135(14):1357-1377. doi: 10.1161/CIRCULATIONAHA.116.024438. PMID: 28373528; PMCID: PMC5392416.
  28. **Tan NY, Mohsin Y, Hodge DO, Lacy MQ, Packer DL, Dispenzieri A, Grogan M, Asirvatham SJ, Madhavan M, McLeod CJ.** Catheter Ablation for Atrial Arrhythmias in Patients With Cardiac Amyloidosis. *J Cardiovasc Electrophysiol.* 2016 Oct;27(10):1167-1173. doi: 10.1111/jce.13046. Epub 2016 Aug 26. PMID: 27422772.
  29. **Feng D, Edwards WD, Oh JK, Chandrasekaran K, Grogan M, Martinez MW, Syed IS, Hughes DA, Lust JA, Jaffe AS, Gertz MA, Klarich KW.** Intracardiac thrombosis and embolism in patients with cardiac amyloidosis. *Circulation.* 2007 Nov 20;116(21):2420-6. doi: 10.1161/CIRCULATIONAHA.107.697763. Epub 2007 Nov 5. Erratum in: *Circulation.* 2008 Aug 19;118(8):e131. Syed, Imran I [corrected to Syed, Imran S]. PMID: 17984380.
  30. **Feng D, Syed IS, Martinez M, Oh JK, Jaffe AS, Grogan M, Edwards WD, Gertz MA, Klarich KW.** Intracardiac thrombosis and anticoagulation therapy in cardiac amyloidosis. *Circulation.* 2009 May 12;119(18):2490-7. doi: 10.1161/CIRCULATIONAHA.108.785014. Epub 2009 May 4. PMID: 19414641.
  31. **Kristen AV, Dengler TJ, Hegenbart U, Schonland SO, Goldschmidt H, Sack FU, Voss F, Becker R, Katus HA, Bauer A.** Prophylactic implantation of cardioverter-defibrillator in patients with severe cardiac amyloidosis and high risk for sudden cardiac death. *Heart Rhythm.* 2008 Feb;5(2):235-40. doi: 10.1016/j.hrthm.2007.10.016. Epub 2007 Oct 9. PMID: 18242546.
  32. **Dubrey SW, Cha K, Anderson J, Chamarthi B, Reisinger J, Skinner M, Falk RH.** The clinical features of immunoglobulin light-chain (AL) amyloidosis with

heart involvement. *QJM*. 1998 Feb;91(2):141-57. doi: 10.1093/qjmed/91.2.141. PMID: 9578896.

33. **Dubrey S, Simms RW, Skinner M, Falk RH.** Recurrence of primary (AL) amyloidosis in a transplanted heart with four-year survival. *Am J Cardiol*. 1995 Oct 1;76(10):739-41. doi: 10.1016/s0002-9149(99)80214-8. PMID: 7572642.
34. **Hosenpud JD, DeMarco T, Frazier OH, Griffith BP, Uretsky BF, Menkis AH, O'Connell JB, Olivari MT, Valantine HA.** Progression of systemic disease and reduced long-term survival in patients with cardiac amyloidosis undergoing heart transplantation. Follow-up results of a multicenter survey. *Circulation*. 1991 Nov;84(5 Suppl):III338-43. PMID: 1934428.
35. **Wechalekar AD, Schonland SO, Kastritis E, Gillmore JD, Dimopoulos MA, Lane T, Foli A, Foard D, Milani P, Rannigan L, Hegenbart U, Hawkins PN, Merlini G, Palladini G.** A European collaborative study of treatment outcomes in 346 patients with cardiac stage III AL amyloidosis. *Blood*. 2013 Apr 25;121(17):3420-7. doi: 10.1182/blood-2012-12-473066. Epub 2013 Mar 11. PMID: 23479568.