

# UCLA

## UCLA Previously Published Works

### Title

Age at start of using tobacco on the risk of head and neck cancer: Pooled analysis in the International Head and Neck Cancer Epidemiology Consortium (INHANCE)

### Permalink

<https://escholarship.org/uc/item/6sc8v1nd>

### Authors

Chang, Chun-Pin  
Chang, Shen-Chih  
Chuang, Shu-Chun  
et al.

### Publication Date

2019-12-01

### DOI

10.1016/j.canep.2019.101615

Peer reviewed



# HHS Public Access

Author manuscript

*Cancer Epidemiol.* Author manuscript; available in PMC 2023 April 04.

Published in final edited form as:

*Cancer Epidemiol.* 2019 December ; 63: 101615. doi:10.1016/j.canep.2019.101615.

## Age at start of using tobacco on the risk of head and neck cancer: Pooled analysis in the International Head and Neck Cancer Epidemiology Consortium (INHANCE)

*A full list of authors and affiliations appears at the end of the article.*

### Abstract

**Background:** Tobacco use is a well-established risk factor for head and neck cancer (HNC). However, less is known about the potential impact of exposure to tobacco at an early age on HNC risk.

**Methods:** We analyzed individual-level data on ever tobacco smokers from 27 case-control studies (17,146 HNC cases and 17,449 controls) in the International Head and Neck Cancer Epidemiology (INHANCE) consortium. Adjusted odds ratios (ORs) and 95% confidence intervals (CIs) were estimated using random-effects logistic regression models.

**Results:** Without adjusting for tobacco packyears, we observed that younger age at starting tobacco use was associated with an increased HNC risk for ever smokers (OR<sub><10 years vs. 30 years</sub>: 1.64, 95% CI: 1.35, 1.97). However, the observed association between age at starting tobacco use and HNC risk became null after adjusting for tobacco packyears (OR<sub><10 years vs. 30 years</sub>: 0.97, 95% CI: 0.80, 1.19). In the stratified analyses on HNC subsites by tobacco packyears or years since quitting, no difference in the association between age at start and HNC risk was observed.

**Conclusions:** Results from this pooled analysis suggest that increased HNC risks observed with earlier age at starting tobacco smoking are largely due to longer duration and higher cumulative tobacco exposures.

---

\*Corresponding author at: Division of Public Health, Department of Family and Preventive Medicine, University of Utah School of Medicine, Huntsman Cancer Institute, 375 Chipeta Way, Suite A, Salt Lake City, UT, 84108, United States. amy.lee@utah.edu (Y.-C.A. Lee).

<sup>†</sup>These authors contributed equally to this work.

Authorship contribution statement

C. Chang, S.C., M.H., and Y.A.L contributed to write, review, and/or revision of the manuscript; S.C. contributed to the analysis, and C. Chang performed statistical support; J.B., G.F., K.M., V.W., T.N.T., M.B.C., C.L.V., A.F.O., J.P.Z., D.S., J.M., E.M.S., G.L., H.M., F.L., L.D.M., E.S., K.K., M.M., T.L.V., P.L., H.R., C. Chen, S.M.S., D.M.W., C.B., V.E., W.G., E.N., R.B.H., M.P.P., S.B., G.C., O.S., R.K., M.P.C., M.V., B.S., R.H., S.F., S.B., L.F., A.M.B.M., A.W.D., D.M., S.S., G.Y., J.L., H.B., E.F., P.R., P. Brennan, and Z.Z. conducted research and provided single-study databases; P. Boffetta, Y.A.L., and M.H. are the INHANCE study coordinators; all authors approved the final version of the manuscript.

Conflict of interest statement

None declared.

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## Keywords

age at start of tobacco use; head and neck cancer; oral cancer; oropharyngeal cancer; hypopharyngeal cancer; laryngeal cancer

---

## 1. Introduction

Head and neck cancer refers to a group of neoplasms originating from the upper aerodigestive tract, including the oral cavity, oropharynx, hypopharynx and larynx. According to the GLOBOCAN 2018 estimates, there were around 705,781 new head and neck cancer cases diagnosed and 358,144 deaths from head and neck cancer worldwide in 2018, consisting of 3.9% and 3.7% of all cancer sites for incidence and mortality, respectively [1]. The IARC Monographs on the evaluation of carcinogenic risks to humans have concluded that there is sufficient evidence that tobacco smoking is causally related to cancers of the oral cavity, pharynx, and larynx in humans [2]. In previous INHANCE analyses, we reported that the population attributable risk for head and neck cancer was 33% due to tobacco alone [3].

The risk of head and neck cancer increases with the duration and intensity of smoking tobacco [2, 4]. Smokeless tobacco, such as chewing tobacco and snuff, also increases the risk of head and neck cancer [5–8]. However, whether teenagers are more vulnerable to the exposure of tobacco in terms of head and neck cancer risk, independent of the amount exposed, is less conclusive and is of special interest. Previous studies on the association between younger age at start of smoking and head and neck cancer reported inconsistent results. While several studies indicated elevated risks of head and neck cancer with earlier age at starting to smoke [9–16], others showed little evidence [17–21]. Even fewer studies have investigated the association of age at start of using smokeless tobacco products with head and neck cancer risk. Lewin et al. reported no association between age at start of using Swedish snuff and squamous cell carcinoma of the head and neck without adjustments for tobacco smoking duration or frequency [14]. In a cohort study in India, Jayalekshmi et al. reported no clear trend between age at start of chewing tobacco and oral cavity cancer without adjustments for tobacco smoking [22].

There are several limitations in the previous studies on the association of age at start of using tobacco with head and neck cancer risk. Some studies included people who have never used tobacco as the reference group [9–12, 14, 16, 18, 21, 22] while others did not [13, 15, 17, 19, 20]. Second, individual studies might not have sufficient statistical power for the comparisons using fine age categories. The pooled data in the International Head and Neck Cancer Epidemiology (INHANCE) consortium [23] provides an opportunity to investigate the effect of age at start of using tobacco products on head and neck cancer with a very large sample size.

## 2. Material and methods

The current analysis included 27 individual case-control studies [12, 18, 24–47] comprising 19,416 head and neck cancer cases and 27,934 controls from the INHANCE pooled data

version 1.4. After excluding subjects with missing data on age, sex, or race/ethnicity, and cases with missing information on the site of origin of their cancer (147 cases and 134 controls), the final data for analyses included 19,269 head and neck cancer cases and 27,800 controls. We further restricted the analyses to ever-tobacco users, which included 27 studies with 17,146 head and neck cancer cases and 17,449 controls (Table 1).

Characteristics of individual studies included in the pooled data are provided in Supplementary Table 1. Among these 27 studies, 19 were hospital-based case-control studies, and in most studies, the control subjects were frequency matched to the case subjects on at least age, sex and additional factors (such as study center, hospital, and race/ethnicity). The Los Angeles study individually matched the controls to the cases on age decade, sex, and neighbourhood, although the matching was broken in the study analysis.

Questionnaire interviews in all studies were conducted face-to-face, except for the Iowa, Boston, and North Carolina (2002–2006) studies, in which study participants completed self-administered questionnaires. Written informed consent was obtained from every study subject, and the individual studies were approved by Institutional Review Boards at each of the institutes involved. Questionnaires were collected from all the individual studies, to assess the comparability of the data and wording of interview questions. Each data item was checked for illogical or missing values. Queries were sent to investigators and inconsistencies were resolved.

Cases included patients with invasive tumors of the oral cavity, oropharynx, hypopharynx, oral cavity/oropharynx/hypopharynx not otherwise specified (NOS), larynx, or head and neck cancer unspecified. Cancers of the salivary gland (ICD-O-2 C07-C08) were excluded from our analysis due to the different etiologic pattern from head and neck cancers [48]. The pooled data excluded lymphomas, minor salivary gland cancers, and sarcomas of the upper aerodigestive tract cases. Studies provided tumor site data using either the International Classification of Diseases-Oncology, Version 2 (ICD-O-2) or ICD 9 or 10. Among the 19,269 head and neck cancer cases, there were a total of 5,452 oral cavity cancer cases, 4,880 oropharyngeal cancer cases, 1,417 hypopharyngeal cancer cases, 1,852 oral cavity/pharynx not specified cases, 5,347 laryngeal cancer cases and 321 unspecified head and neck cancer cases. Of the 14,878 head and neck cancer cases for which histological information was available, 14,355 were squamous cell carcinomas (96.5%).

In the tobacco section of the questionnaires, subjects were asked if they were users of each specific type of tobacco including cigarettes, cigars, pipes, chewing tobacco, snuff, cigarillos, and straw-cigarettes. Type of tobacco use varied across studies as follows: 1) cigarettes, cigars, pipes, chewing tobacco, or snuff in 13 studies including Houston, Iowa, Los Angeles, Memorial Sloan Kettering Cancer Center (MSKCC), New York Multicenter, North Carolina (1994–1997), North Carolina (2002–2006), Seattle (1985–1995), Seattle-Leo, Tampa, US Multicenter, Puerto Rico, and International Multicenter studies (Tampa study combined information on chewing tobacco and snuff); 2) cigarettes, cigars, pipes or cigarillos in France (1987–1992) and Germany-Heidelberg study (France study combined information on cigar and cigarillo); 3) cigarettes, cigars, pipes, or straw-cigarettes in Sao Paulo study; 4) cigarettes, cigars, pipes, or chewing tobacco in Boston study; 5) cigarettes,

cigars, or pipes in 8 studies including Aviano, Central Europe, Italy Multicenter, Milan (1984–1989), Milan (2006–2009), Rome, Switzerland, and Latin America studies; and 6) cigarettes in Germany-Saarland and Japan (2001–2005) studies. We assumed that studies which did not collect information on a specific tobacco product had negligible prevalence of use in their population. Among ever tobacco users, age at start of using each specific tobacco product was asked. Houston and MSKCC studies collected information on the use of cigarettes, cigars, pipes, chewing tobacco, and snuff, but information on age at start of smoking was only available for cigarette smokers.

## 2.1. Statistical Methods

We categorized age at start of using tobacco products into 5-year categories as <10, 10–14, 15–19, 20–24, 25–29 and 30 years. We estimated odds ratios (OR) and 95% confidence intervals (95% CI) using random effect regression models (PROC GLIMMIX procedure) with study center as the random effect intercept. In one model, covariates included age (categories shown in Table 1), sex, education (categories shown), race/ethnicity (categories shown), alcohol drinks per day (continuous). In another model, covariates additionally included duration of using chewing tobacco (continuous), duration of using snuff (continuous), and years since quitting tobacco use (continuous). Finally, covariates in the final model additionally included packyears of tobacco smoking (continuous) and the product term for interaction between alcohol drinks per day and packyears of tobacco smoking. Packyears of tobacco smoking was defined as the sum of 1) the number of packs of cigarettes smoked per day multiplied by the years of smoking cigarettes, 2) cigar-, 3) pipe-, 4) cigarillo-, and 5) straw-cigarettes-equivalent cigarette packyears. Tobacco contents were taken to be the equivalent of four cigarettes for one roll of cigar, 3.5 cigarettes for one roll of pipe, two cigarettes for one roll of cigarillo, and five cigarettes for one roll of straw-cigarette.[49] One tobacco packyear was equal to smoking 20 cigarettes (1 pack) per day for a year.

Age at the start, duration, time since quitting tobacco use, and age at diagnosis are four time-related variables when modeling tobacco history. To reduce multicollinearity, we used tobacco packyears to replace the use of intensity and duration as separate variables and used categorized ages instead of continuous variables [50].

For subjects missing education level (1,658 cases and 3,999 controls), we applied multiple imputation with the PROC MI procedure in the SAS statistical software. We assumed that the education data are missing at random (MAR), i.e. whether education is missing or not does not depend on any other unobserved or missing values [51]. We used the logistic regression model [52] to predict education level with age, sex, race/ethnicity, study, and case/control status as the covariates, for each of the geographic regions (Europe, South America, and North America) separately. The logistic regression results to assess summary estimates for the five imputations were combined by the PROC MIANALYZE procedure in SAS. We summarized estimates of associations in a random effect logistic regression model, which allowed for unexplained sources of heterogeneity between studies [53].

Stratified analyses were conducted by cancer subsites (oral, oropharynx, hypopharynx, and larynx), years since quitting tobacco use (current, < 20 years, and 20 years), tobacco

packyears (< 20 tobacco packyears and > 20 tobacco packyears), and duration of using tobacco (1–19 years, 20–39 years, and ≥ 40 years).

### 3. Results

The distributions of age, sex, race/ethnicity, and education level were different between the case and control groups (Table 1). The controls had a higher proportion of younger subjects, men, and more educated participants than the cases. Among the individuals who started using tobacco products before 10 year-old, more than half were from South or Central America, where straw cigarettes are used (data not shown).

Without adjusting for tobacco-related covariates, we observed that age at start of using tobacco was positively associated with head and neck cancer risk with a linear trend ( $P$  for trend < 0.01, OR of <10 years vs. ≥ 30 years: 1.49, 95% CI: 1.25, 1.79, in the basic model; Table 2). However, after the adjustment on tobacco exposures, we observed no association and no dose-response relationship ( $P$  for trend = 0.21) between age at start of using tobacco and head and neck cancer risk. When we changed the reference group to ≥ 25 years for all the models we tested, the inference remained the same (Supplementary Table 2).

When stratified by head and neck cancer subsites, younger age at start of using tobacco products was associated with increased risks of oropharyngeal and laryngeal cancer in the model without adjusting for tobacco packyears. In the model adjusting for tobacco packyears, however, age at start of using tobacco products was not associated with oropharyngeal and laryngeal cancer risk (Table 3). When we changed the reference group to starting tobacco habits at ≥ 25 years old, the inference remained the same (Supplementary Table 3). For oral cavity cancer, younger age at start of using tobacco suggested an inverse dose-response with younger ages at starting tobacco being associated with lower ORs, although the CIs were wide ( $P$  for trend < 0.01).

In stratified analyses by time since quitting tobacco use (Table 4), younger age at start of using tobacco was associated with head and neck cancer risk ( $P$  for trend < 0.01) among current tobacco users. However, after adjustment for tobacco packyears, no associations were detected. Stratified analysis by tobacco packyears was performed to further examine how tobacco packyears modified the trend. No association was observed regardless of tobacco packyears strata (Table 5).

We did not adjust for duration of using tobacco because it was highly correlated with age at start of using tobacco products. The distribution of cases and controls showed that more people with younger age at start of using tobacco products were in the group with ≥ 40 years duration of using tobacco products (Table 6). Within the group with similar duration of using tobacco products, an association was observed between age at start of using tobacco products and head and neck cancer risk within those with duration of 20–39 years. However, such observations disappeared with adjustment for tobacco packyears. The results remained similar when we changed the reference group from ≥ 30 years to ≥ 25 years (Supplementary Table 4 to Table 6).

## 4. Discussion

In this large data pooling project, we observed an association between younger age at start of using tobacco and the risk of head and neck cancer among people who ever used tobacco products. However, the association was no longer observed after adjusting for tobacco related factors. Stratified analyses by head and neck cancer subsites, years since quitting tobacco use, tobacco packyears, and duration of using tobacco products showed little evidence of the association between earlier age at start of using tobacco and head and neck cancer risk after adjustment by tobacco related factors.

Previous studies [9–16] reported the potential association between younger age at start of using tobacco and increased risks of the head and neck cancer, which implied that early tobacco exposure is associated with a higher head and neck cancer risk later in life. In a study of former smokers with lung cancer, Wiencke et al. [54] reported higher DNA damage levels in lungs of patients who started to smoke at an earlier age, indicating that either early-age smoking may impair normal DNA repair function or young smokers may be more susceptible to DNA adduct formation. However, the reported association could simply reflect longer duration, which is a key aspect in the process of tobacco-related carcinogenesis. In our study, to elucidate the role of age at start of using tobacco, we carefully adjusted for other smoking-related factors in the indirect pathways and confounding factors, such as tobacco packyears, interaction between alcohol drinks per day and tobacco packyears, duration of chewing tobacco, duration of using snuff, and years since quitting tobacco use. We estimated the cancer risk in tobacco-users by including people who started using tobacco at 30-year-old as the reference group. The association between earlier age at starting tobacco and head and neck cancer risk was no longer present after adjusting for smoking-related factors, implying that the previously observed association of age at start of using tobacco with the risk of head and neck cancer may not be a direct effect but was mediated by smoking duration and frequency.

In this study, adjusting for human papillomavirus (HPV) infection was not feasible since the information was not available across all of the individual studies. HPV infection is primarily linked with oropharyngeal cancer [55]. Moreover, HPV-positive oropharyngeal cancer patients tend to be younger, male, and less likely to smoke [56]. This study restricted to ever-tobacco users with adjustment for age and sex in the analyses. If people with HPV infection tended to start smoking earlier in their life, then ORs of head and neck cancer may be biased away from the null without adjustment for HPV exposure. However, since the association between age at start and the risk of head and neck cancer was null after adjustment for tobacco packyears, HPV infection status is unlikely to influence our conclusion.

It is logically impossible to compare age at start while adjusting for duration of using tobacco products in the case-control study because age at start of using tobacco has less variation among those with similar age at diagnosis and duration, and it is difficult to assess their effects separately. We used tobacco packyears, which combined smoking duration and frequency to adjust for the effect of smoking and to minimize the multicollinearity issue. We conducted stratified analyses to alleviate the effect of years of using tobacco products



on head and neck cancer risk. Null associations were observed after adjustment for tobacco related variables.

One possible limitation in our study is recall bias. Being aware of disease status and understanding that smoking is related to their disease, cases might tend to recall an earlier age at start of using these products. This could have resulted in an overestimation of the risk associated with younger age at start. On the other hand, since the general legal age worldwide of buying tobacco was between 16 and 18 years old, individuals who started to smoke earlier than that age might report their age at starting smoking later. Thus, we might have underestimated the risk association of starting using tobacco products before the age of 16 to 18. However, we did not observe heterogeneity of the results between European studies and studies from other regions of the world, and we believe that reporting bias would be minimal. Another limitation of this study is that we were not able to adjust for the intensity of using chewing tobacco and snuff because the units were not comparable across studies (such as grams of tobacco or pinch of tobacco). Therefore, only the duration of chewing tobacco and snuff were adjusted for, and the results might be susceptible to residual confounding. However, the predominant tobacco habit was cigarette smoking, and we were able to fully adjust for frequency and duration of cigarette smoking.

The major strength of this study is the large sample size, which provides adequate statistical power to examine potential associations between age at start of using tobacco and head and neck cancer risk among ever tobacco users. We were also able to compare the associations across different head and neck cancer subsites. Furthermore, the detailed information on tobacco habits enabled us to adjust for potential confounding factors while accounting for the multicollinearity issue of various time-related factors.

In summary, the results from this large pooling project from the INHANCE consortium support an association between the risk of head and neck cancer and the younger age at start of using tobacco products, which is mediated by cumulative tobacco smoking. Even though a null association was observed between age at start of using tobacco products and the risk of head and neck cancer after adjusting for tobacco related factors, younger age at start of using tobacco products was closely related to longer duration or higher frequency of tobacco smoking, which are important risk factors of head and neck cancer.

## Funding

This pooled analysis was supported by grants National Cancer Institute (NCI) R03CA113157, National Institute of Dental and Craniofacial Research (NIDCR) R03DE016611 from the National Institutes of Health (NIH). Shen-Chih Chang was supported by a Special Training Award from the International Agency for Research on Cancer. The Aviano and Italy Multicenter studies were supported by Italian Association for Research on Cancer (AIRC), Italian League Against Cancer, and Italian Ministry of Research. The Boston study was supported by grants R01CA078609 and R01CA100679 from the National Institutes of Health. The Central Europe study was supported by grant IC15- CT98-0332 from World Cancer Research Fund and the European Commission INCO-COPERNICUS Program. The France (1987-1992) study was supported by grant



KFS1069–09–2000 from Swiss League against Cancer, grant FOR381.88 from Fribourg League against Cancer, grant AKT 617 from Swiss Cancer Research and grant 88D28 from Gustave-Roussy Institute. The Germany-Heidelberg study was supported by grant 01GB9702/3 from German Ministry of Education and Research. The Germany-Saarland study was supported by Ministry of Science, Research and Arts Baden-Wuerttemberg. The Houston study was supported by grants R01ES11740 and R01CA100264 from the National Institutes of Health. The International Multicenter study was supported by grants FIS 97/0024 and FIS 97/0662 from Fondo de Investigaciones Sanitarias (FIS) of the Spanish Government, BAE 01/5013 from International Union Against Cancer (UICC), and Yamagiwa-Yoshida Memorial International Cancer Study Grant. The Iowa study was supported by grants NIDCR R01DE011979, NIDCR R01DE013110 and FIRCA TW001500 from National Institutes of Health and was supported by Veterans Affairs Merit Review Funds. The Japan (2001–2005) study was supported by grant 17015052 from Scientific Research grant from the Ministry of Education, Science, Sports, Culture and Technology of Japan, and grant H20–002 from the Ministry of Health, Labor and Welfare of Japan. The Latin America study was supported from grant 01/01768–2 from Fondo para la Investigacion Cientifica y Tecnologica (FONCYT) Argentina, IMIM (Barcelona), Fundaco de Amparo à Pesquisa no Estado de Sao Paulo (FAPESP) , and grant IC18-CT97–0222 from European Commission. The Los Angeles study was supported by grants P50CA090388, R01DA011386, R03CA077954, T32CA009142, U01CA096134 and R21ES011667 from the National Institutes of Health, and was supported by the Alper Research Program for Environmental Genomics of the UCLA Jonsson Comprehensive Cancer Center. The Milan study was supported by the Italian Association for Cancer Research. The MSKCC was supported by grant R01CA051845 from the National Institutes of Health. The New York Multicenter study was supported by grants P01CA068384 and K07CA104231 from the National Institutes of Health. The North Carolina (1994–1997) and North Carolina (2002–2006) studies were supported by grant R01CA061188 from the National Institutes of Health, and grant P30ES010126 from the National Institute of Environmental Health Sciences. The Puerto Rico study was jointly funded by National Cancer Institute (NCI) and National Institute of Dental and Craniofacial Research (NIDCR) intramural programs, National Institutes of Health. The Rome study was supported by the Italian Association for Cancer Research (AIRC). The Sao Paulo study was supported by grant 10/51168–0 from the State of São Paulo Research Foundation (FAPESP). The Seattle (1985–1995) study was supported by R01CA048896 and R01DE012609 from the National Institutes of Health. The Seattle-LEO study was supported by R01CA030022 from the National Institutes of Health. The Switzerland study was supported by grants KFS-700 and OCS-1633 from Swiss League against Cancer and the Swiss Research against Cancer/Oncosuisse. The Tampa study was supported by grants P01CA068384, K07CA104231 and R01DE013158 from the National Institutes of Health. The US Multicenter study was supported by the Intramural Program of the National Cancer Institute (NCI), National Institutes of Health.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Authors

Chun-Pin Chang<sup>a,1</sup>, Shen-Chih Chang<sup>b,c,1</sup>, Shu-Chun Chuang<sup>d</sup>, Julien Berthiller<sup>b,e</sup>, Gilles Ferro<sup>b</sup>, Keitaro Matsuo<sup>f,g</sup>, Victor Wünsch-Filho<sup>h</sup>, Tatiana N. Toporcov<sup>h</sup>, Marcos Brasilino de Carvalho<sup>i</sup>, Carlo La Vecchia<sup>j</sup>, Andrew F. Olshan<sup>k</sup>, Jose P. Zevallos<sup>l</sup>, Diego Serraino<sup>m</sup>, Joshua Muscat<sup>n</sup>, Erich M. Sturgis<sup>o</sup>, Guojun Li<sup>o</sup>, Hal Morgenstern<sup>p</sup>, Fabio Levi<sup>q</sup>, Luigino Dal Maso<sup>m</sup>, Elaine Smith<sup>r</sup>, Karl Kelsey<sup>s</sup>, Michael McClean<sup>s</sup>, Thomas L. Vaughan<sup>t</sup>, Philip Lazarus<sup>u</sup>, Heribert Ramroth<sup>v</sup>, Chu Chen<sup>t</sup>, Stephen M. Schwartz<sup>t</sup>, Deborah M. Winn<sup>w</sup>, Cristina Bosetti<sup>x</sup>, Valeria Edefontij<sup>y</sup>, Werner Garavello<sup>y</sup>, Eva Negri<sup>z</sup>, Richard B. Hayes<sup>A</sup>, Mark P. Purdue<sup>w</sup>, Stefania Boccia<sup>B,C</sup>, Gabriella Cadoni<sup>D,E</sup>, Oxana Shangina<sup>F</sup>, Rosalina Koifman<sup>G</sup>, Maria Paula Curado<sup>H</sup>, Marta Vilensky<sup>I</sup>, Beata Swiatkowska<sup>J</sup>, Rolando Herrero<sup>b</sup>, Silvia Franceschi<sup>K</sup>, Simone Benhamou<sup>L</sup>, Leticia Fernandez<sup>M</sup>, Ana M.B. Menezes<sup>N</sup>, Alexander W. Daudt<sup>O</sup>, Dana Mates<sup>P</sup>, Stimson Schantz<sup>Q</sup>, Guo-Pei Yu<sup>R</sup>, Jolanta Lissowska<sup>S</sup>, Hermann Brenner<sup>T,U,V</sup>, Eleonora Fabianova<sup>W</sup>, Peter Rudnai<sup>X</sup>, Paul Brennan<sup>b</sup>, Paolo Boffetta<sup>Y</sup>, Zuo-Feng Zhang<sup>c</sup>, Mia Hashibe<sup>a</sup>, Yuan-Chin Amy Lee<sup>a,\*</sup>

## Affiliations

<sup>a</sup>Division of Public Health, Department of Family & Preventive Medicine, University of Utah School of Medicine, and Huntsman Cancer Institute, Salt Lake City, UT, United States

<sup>b</sup>International Agency for Research on Cancer, Lyon, France

<sup>c</sup>Department of Epidemiology, UCLA Fielding School of Public Health, Los Angeles, CA, United States

<sup>d</sup>Institute of Population Health Sciences, National Health Research Institutes, Taiwan

<sup>e</sup>Pôle Information Médicale Evaluation Recherche, Hospices Civils de Lyon, France

<sup>f</sup>Division of Cancer Epidemiology and Prevention, Aichi Cancer Center Research Institute, Nagoya, Japan

<sup>g</sup>Department of Cancer Epidemiology, Nagoya University Graduate School of Medicine, Nagoya, Japan

<sup>h</sup>Department of Epidemiology, School of Public Health, University of São Paulo, São Paulo, Brazil

<sup>i</sup>Department of Head and Neck, Heliopolis Hospital, São Paulo, Brazil

<sup>j</sup>Department of Clinical Sciences and Community Health, University of Milan, Milan, Italy

<sup>k</sup>Department of Epidemiology, Gillings School of Global Public Health, Chapel Hill, NC, United States

<sup>l</sup>Division of Head and Neck Surgical Oncology, Department of Otolaryngology/Head and Neck Surgery, Washington University School of Medicine, Saint Louis, MO, United States

<sup>m</sup>Cancer Epidemiology Unit, Centro di Riferimento Oncologico di Aviano (CRO) IRCCS, Italy

<sup>n</sup>Penn State College of Medicine, Hershey, PA, United States

<sup>o</sup>University of Texas M. D. Anderson Cancer Center, Houston, TX, United States

<sup>p</sup>Departments of Epidemiology and Environmental Health Sciences, School of Public Health, and Department of Urology, Medical School, University of Michigan, Ann Arbor, MI, United States

<sup>q</sup>Institute of Social and Preventive Medicine (IUMSP), Lausanne University Hospital (CHUV), Lausanne, Switzerland

<sup>r</sup>College of Public Health, University of Iowa, Iowa City, IA, United States <sup>s</sup> Brown University, Providence, RI, United States

<sup>t</sup>Fred Hutchinson Cancer Research Center, Seattle, WA, United States

<sup>u</sup>Washington State University College of Pharmacy, Spokane, WA, United States <sup>v</sup> University of Heidelberg, Heidelberg, Germany

<sup>w</sup>Division of Cancer Epidemiology and Genetics, National Cancer Institute, Rockville, MD, United States

<sup>x</sup>Department of Oncology, Istituto di Ricerche Farmacologiche Mario Negri IRCCS, Milan, Italy

<sup>y</sup>Department of Otorhinolaryngology, School of Medicine and Surgery, University of Milano-Bicocca, Milan, Italy

<sup>z</sup>Department of Biomedical and Clinical Sciences, University of Milan, Italy

<sup>A</sup>Division of Epidemiology, New York University School of Medicine, New York, United States

<sup>B</sup>Institute of Public Health, Section of Hygiene, Università Cattolica del Sacro Cuore, Rome, Italy

<sup>C</sup>Fondazione Policlinico Universitario A.Gemelli IRCCS, Rome, Italy

<sup>D</sup>Department of Head and Neck Surgery, Institute of Clinical Otorhinolaryngology, Università Cattolica del Sacro Cuore, Rome, Italy

<sup>E</sup>Istituto di Clinica Otorinolaringoiatrica, Fondazione Policlinico Universitario A.Gemelli IRCCS, Rome, Italy

<sup>F</sup>Cancer Research Centre, Moscow, Russia

<sup>G</sup>Escola Nacional de Saude Publica, Fundacao Oswaldo Cruz, Rio de Janeiro, Brazil

<sup>H</sup>ACCAMARGO Cancer Center, São Paulo, Brazil

<sup>I</sup>Institute of Oncology Angel H. Roffo, University of Buenos Aires, Argentina

<sup>J</sup>Nofer Institute of Occupational Medicine, Lodz, Poland

<sup>K</sup>Scientific Directorate, Centro di Riferimento Oncologico di Aviano (CRO) IRCCS, Italy

<sup>L</sup>National Institute of Health and Medical Research, Inserm U1018, Villejuif, France

<sup>M</sup>Institute of Oncology and Radiobiology, Havana, Cuba

<sup>N</sup>Universidade Federal de Pelotas, Pelotas, Brazil

<sup>O</sup>Hospital de Clinicas de Porto Alegre, Porto Alegre, Brazil

<sup>P</sup>National Institute of Public Health, Bucharest, Romania

<sup>Q</sup>New York Eye and Ear Infirmary, New York, NY, United States

<sup>R</sup>Department of Otolaryngology, School of Medicine, New York Medical College, Valhalla, NY, United States

<sup>S</sup>The M. Skasodowska-Curie Memorial Cancer Center and Institute of Oncology, Dept. of Cancer Epidemiology and Prevention, Warsaw, Poland

<sup>T</sup>Division of Clinical Epidemiology and Aging Research, German Cancer Research Center (DKFZ), Heidelberg, Germany

<sup>U</sup>Division of Preventive Oncology, German Cancer Research Center (DKFZ) and National Center for Tumor Diseases (NCT), Heidelberg, Germany

<sup>V</sup>German Cancer Consortium (DKTK), German Cancer Research Center (DKFZ), Heidelberg, Germany

<sup>W</sup>Regional Authority of Public Health in Banska Bystrica, Slovakia

<sup>X</sup>National Institute of Environmental Health, Budapest, Hungary

<sup>Y</sup>The Tisch Cancer Institute, Icahn School of Medicine at Mount Sinai, New York, NY, United States

## Acknowledgments

We would like to thank Dr. Xavier Castellsague for his expertise and assistance throughout all aspects of our study.

## Abbreviations:

**HNC**                      Head and Neck Cancer

## References

- [1]. Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A, Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries, *CA Cancer J Clin* (2018).
- [2]. IARC, List of Classifications by cancer sites with sufficient or limited evidence in humans, Volumes 1–122, Lyon, 2018.
- [3]. Hashibe M, Brennan P, Chuang SC, Boccia S, Castellsague X, Chen C, Curado MP, Dal Maso L, Daudt AW, Fabianova E, Fernandez L, Wunsch-Filho V, Franceschi S, Hayes RB, Herrero R, Kelsey K, Koifman S, La Vecchia C, Lazarus P, Levi F, Lence JJ, Mates D, Matos E, Menezes A, McClean MD, Muscat J, Eluf-Neto J, Olshan AF, Purdue M, Rudnai P, Schwartz SM, Smith

E, Sturgis EM, Szeszenia-Dabrowska N, Talamini R, Wei Q, Winn DM, Shangina O, Pilarska A, Zhang ZF, Ferro G, Berthiller J, Boffetta P, Interaction between tobacco and alcohol use and the risk of head and neck cancer: pooled analysis in the International Head and Neck Cancer Epidemiology Consortium, *Cancer epidemiology, biomarkers & prevention : a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology* 18(2) (2009) 541–50.

- [4]. Di Credico G, Edefonti V, Polesel J, Pauli F, Torelli N, Serraino D, Negri E, Luce D, Stucker I, Matsuo K, Brennan P, Vilensky M, Fernandez L, Curado MP, Menezes A, Daudt AW, Koifman R, Wunsch-Filho V, Holcatova I, Ahrens W, Lagiou P, Simonato L, Richiardi L, Healy C, Kjaerheim K, Conway DI, Macfarlane TV, Thomson P, Agudo A, Znaor A, Boaventura Rios LF, Toporcov TN, Franceschi S, Herrero R, Muscat J, Olshan AF, Zevallos JP, La Vecchia C, Winn DM, Sturgis EM, Li G, Fabianova E, Lissowska J, Mates D, Rudnai P, Shangina O, Swiatkowska B, Moysich K, Zhang ZF, Morgenstern H, Levi F, Smith E, Lazarus P, Bosetti C, Garavello W, Kelsey K, McClean M, Ramroth H, Chen C, Schwartz SM, Vaughan TL, Zheng T, Menvielle G, Boccia S, Cadoni G, Hayes RB, Purdue M, Gillison M, Schantz S, Yu GP, Brenner H, D'Souza G, Gross ND, Chuang SC, Boffetta P, Hashibe M, Lee YA, Dal Maso L, Joint effects of intensity and duration of cigarette smoking on the risk of head and neck cancer: A bivariate spline model approach, *Oral Oncol* 94 (2019) 47–57. [PubMed: 31178212]
- [5]. Boffetta P, Hecht S, Gray N, Gupta P, Straif K, Smokeless tobacco and cancer, *Lancet Oncol* 9(7) (2008) 667–75. [PubMed: 18598931]
- [6]. Lee PN, Hamling J, Systematic review of the relation between smokeless tobacco and cancer in Europe and North America, *BMC Med* 7 (2009) 36. [PubMed: 19638245]
- [7]. Zhou J, Michaud DS, Langevin SM, McClean MD, Eliot M, Kelsey KT, Smokeless tobacco and risk of head and neck cancer: evidence from a case-control study in New England, *International journal of cancer. Journal international du cancer* 132(8) (2013) 1911–7. [PubMed: 22987222]
- [8]. Wyss AB, Hashibe M, Lee YA, Chuang SC, Muscat J, Chen C, Schwartz SM, Smith E, Zhang ZF, Morgenstern H, Wei Q, Li G, Kelsey KT, McClean M, Winn DM, Schantz S, Yu GP, Gillison ML, Zevallos JP, Boffetta P, Olshan AF, Smokeless Tobacco Use and the Risk of Head and Neck Cancer: Pooled Analysis of US Studies in the INHANCE Consortium, *Am J Epidemiol* 184(10) (2016) 703–716. [PubMed: 27744388]
- [9]. Barra S, Baron AE, Franceschi S, Talamini R, La Vecchia C, Cancer and non-cancer controls in studies on the effect of tobacco and alcohol consumption, *Int J Epidemiol* 20(4) (1991) 845–51. [PubMed: 1800421]
- [10]. Choi SY, Kahyo H, Effect of cigarette smoking and alcohol consumption in the aetiology of cancer of the oral cavity, pharynx and larynx, *Int J Epidemiol* 20(4) (1991) 878–85. [PubMed: 1800426]
- [11]. Day GL, Blot WJ, Austin DF, Bernstein L, Greenberg RS, Preston-Martin S, Schoenberg JB, Winn DM, McLaughlin JK, Fraumeni JF Jr., Racial differences in risk of oral and pharyngeal cancer: alcohol, tobacco, and other determinants, *J Natl Cancer Inst* 85(6) (1993) 465–73. [PubMed: 8445674]
- [12]. Franceschi S, Talamini R, Barra S, Baron AE, Negri E, Bidoli E, Serraino D, La Vecchia C, Smoking and drinking in relation to cancers of the oral cavity, pharynx, larynx, and esophagus in northern Italy, *Cancer Res* 50(20) (1990) 6502–7. [PubMed: 2208109]
- [13]. Garrote LF, Herrero R, Reyes RM, Vaccarella S, Anta JL, Ferbey L, Munoz N, Franceschi S, Risk factors for cancer of the oral cavity and oro-pharynx in Cuba, *Br J Cancer* 85(1) (2001) 46–54. [PubMed: 11437401]
- [14]. Lewin F, Norell SE, Johansson H, Gustavsson P, Wennerberg J, Biorklund A, Rutqvist LE, Smoking tobacco, oral snuff, and alcohol in the etiology of squamous cell carcinoma of the head and neck: a population-based case-referent study in Sweden, *Cancer* 82(7) (1998) 1367–75. [PubMed: 9529030]
- [15]. Llewellyn CD, Johnson NW, Warnakulasuriya KA, Risk factors for oral cancer in newly diagnosed patients aged 45 years and younger: a case-control study in Southern England, *J Oral Pathol Med* 33(9) (2004) 525–32. [PubMed: 15357672]

- [16]. Merletti F, Boffetta P, Ciccone G, Mashberg A, Terracini B, Role of tobacco and alcoholic beverages in the etiology of cancer of the oral cavity/oropharynx in Torino, Italy, *Cancer Res* 49(17) (1989) 4919–24. [PubMed: 2758421]
- [17]. Balaram P, Sridhar H, Rajkumar T, Vaccarella S, Herrero R, Nandakumar A, Ravichandran K, Ramdas K, Sankaranarayanan R, Gajalakshmi V, Munoz N, Franceschi S, Oral cancer in southern India: the influence of smoking, drinking, paan-chewing and oral hygiene, *Int J Cancer* 98(3) (2002) 440–5. [PubMed: 11920597]
- [18]. Blot WJ, McLaughlin JK, Winn DM, Austin DF, Greenberg RS, Preston-Martin S, Bernstein L, Schoenberg JB, Stemhagen A, Fraumeni JF Jr., Smoking and drinking in relation to oral and pharyngeal cancer, *Cancer Res* 48(11) (1988) 3282–7. [PubMed: 3365707]
- [19]. Lissowska J, Pilarska A, Pilarski P, Samolczyk-Wanyura D, Piekarczyk J, Bardin-Mikollajczak A, Zatonski W, Herrero R, Munoz N, Franceschi S, Smoking, alcohol, diet, dentition and sexual practices in the epidemiology of oral cancer in Poland, *Eur J Cancer Prev* 12(1) (2003) 25–33. [PubMed: 12548107]
- [20]. Menvielle G, Luce D, Goldberg P, Bugel I, Leclerc A, Smoking, alcohol drinking and cancer risk for various sites of the larynx and hypopharynx. A case-control study in France, *Eur J Cancer Prev* 13(3) (2004) 165–72. [PubMed: 15167214]
- [21]. Talamini R, Bosetti C, La Vecchia C, Dal Maso L, Levi F, Bidoli E, Negri E, Pasche C, Vaccarella S, Barzan L, Franceschi S, Combined effect of tobacco and alcohol on laryngeal cancer risk: a case-control study, *Cancer Causes Control* 13(10) (2002) 957–64. [PubMed: 12588092]
- [22]. Jayalekshmi PA, Gangadharan P, Akiba S, Koriyama C, Nair RR, Oral cavity cancer risk in relation to tobacco chewing and bidi smoking among men in Karunagappally, Kerala, India: Karunagappally cohort study, *Cancer Sci* 102(2) (2011) 460–7. [PubMed: 21129124]
- [23]. Winn DM, Lee YC, Hashibe M, Boffetta P, consortium I, The INHANCE consortium: toward a better understanding of the causes and mechanisms of head and neck cancer, *Oral Dis* 21(6) (2015) 685–93. [PubMed: 25809224]
- [24]. Benhamou S, Tuimala J, Bouchardy C, Dayer P, Sarasin A, Hirvonen A, DNA repair gene XRCC2 and XRCC3 polymorphisms and susceptibility to cancers of the upper aerodigestive tract, *Int J Cancer* 112(5) (2004) 901–4. [PubMed: 15386379]
- [25]. Boing AF, Antunes JL, de Carvalho MB, de Gois Filho JF, Kowalski LP, Michaluart P Jr., Eluf-Neto J, Boffetta P, Wunsch-Filho V, How much do smoking and alcohol consumption explain socioeconomic inequalities in head and neck cancer risk?, *J Epidemiol Community Health* 65(8) (2011) 709–14. [PubMed: 20724282]
- [26]. Bosetti C, Gallus S, Trichopoulou A, Talamini R, Franceschi S, Negri E, La Vecchia C, Influence of the Mediterranean diet on the risk of cancers of the upper aerodigestive tract, *Cancer epidemiology, biomarkers & prevention : a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology* 12(10) (2003) 1091–4.
- [27]. Cui Y, Morgenstern H, Greenland S, Tashkin DP, Mao J, Cao W, Cozen W, Mack TM, Zhang ZF, Polymorphism of Xeroderma Pigmentosum group G and the risk of lung cancer and squamous cell carcinomas of the oropharynx, larynx and esophagus, *Int J Cancer* 118(3) (2006) 714–20. [PubMed: 16094634]
- [28]. Dietz A, Ramroth H, Urban T, Ahrens W, Becher H, Exposure to cement dust, related occupational groups and laryngeal cancer risk: results of a population based case-control study, *Int J Cancer* 108(6) (2004) 907–11. [PubMed: 14712496]
- [29]. Divaris K, Olshan AF, Smith J, Bell ME, Weissler MC, Funkhouser WK, Bradshaw PT, Oral health and risk for head and neck squamous cell carcinoma: the Carolina Head and Neck Cancer Study, *Cancer Causes Control* 21(4) (2010) 567–75. [PubMed: 20049634]
- [30]. Elahi A, Zheng Z, Park J, Eyring K, McCaffrey T, Lazarus P, The human OGG1 DNA repair enzyme and its association with orolaryngeal cancer risk, *Carcinogenesis* 23(7) (2002) 1229–34. [PubMed: 12117782]
- [31]. Galli P, Cadoni G, Volante M, De Feo E, Amore R, Giorgio A, Arzani D, Paludetti G, Ricciardi G, Boccia S, A case-control study on the combined effects of p53 and p73 polymorphisms on head and neck cancer risk in an Italian population, *BMC Cancer* 9 (2009) 137. [PubMed: 19426493]



- [32]. Hashibe M, Boffetta P, Zaridze D, Shangina O, Szeszenia-Dabrowska N, Mates D, Janout V, Fabianova E, Bencko V, Moullan N, Chabrier A, Hung R, Hall J, Canzian F, Brennan P, Evidence for an important role of alcohol- and aldehyde-metabolizing genes in cancers of the upper aerodigestive tract, *Cancer Epidemiol Biomarkers Prev* 15(4) (2006) 696–703. [PubMed: 16614111]
- [33]. Hayes RB, Bravo-Otero E, Kleinman DV, Brown LM, Fraumeni JF Jr., Harty LC, Winn DM, Tobacco and alcohol use and oral cancer in Puerto Rico, *Cancer Causes Control* 10(1) (1999) 27–33. [PubMed: 10334639]
- [34]. Herrero R, Castellsague X, Pawlita M, Lissowska J, Kee F, Balamam P, Rajkumar T, Sridhar H, Rose B, Pintos J, Fernandez L, Idris A, Sanchez MJ, Nieto A, Talamini R, Tavani A, Bosch FX, Reidel U, Snijders PJ, Meijer CJ, Viscidi R, Munoz N, Franceschi S, Human papillomavirus and oral cancer: the International Agency for Research on Cancer multicenter study, *J Natl Cancer Inst* 95(23) (2003) 1772–83. [PubMed: 14652239]
- [35]. Levi F, Pasche C, La Vecchia C, Lucchini F, Franceschi S, Monnier P, Food groups and risk of oral and pharyngeal cancer, *International journal of cancer* 77(5) (1998) 705–9. [PubMed: 9688303]
- [36]. Muscat JE, Richie JP Jr., Thompson S, Wynder EL, Gender differences in smoking and risk for oral cancer, *Cancer Res* 56(22) (1996) 5192–7. [PubMed: 8912856]
- [37]. Negri E, La Vecchia C, Franceschi S, Tavani A, Attributable risk for oral cancer in northern Italy, *Cancer Epidemiol Biomarkers Prev* 2(3) (1993) 189–93. [PubMed: 8318870]
- [38]. Olshan AF, Weissler MC, Watson MA, Bell DA, GSTM1, GSTT1, GSTP1, CYP1A1, and NAT1 polymorphisms, tobacco use, and the risk of head and neck cancer, *Cancer Epidemiol Biomarkers Prev* 9(2) (2000) 185–91. [PubMed: 10698480]
- [39]. Peters ES, McClean MD, Liu M, Eisen EA, Mueller N, Kelsey KT, The ADH1C polymorphism modifies the risk of squamous cell carcinoma of the head and neck associated with alcohol and tobacco use, *Cancer epidemiology, biomarkers & prevention : a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology* 14(2) (2005) 476–82.
- [40]. Rogers MA, Thomas DB, Davis S, Vaughan TL, Nevissi AE, A case-control study of element levels and cancer of the upper aerodigestive tract, *Cancer Epidemiol Biomarkers Prev* 2(4) (1993) 305–12. [PubMed: 8348053]
- [41]. Rosenblatt KA, Daling JR, Chen C, Sherman KJ, Schwartz SM, Marijuana use and risk of oral squamous cell carcinoma, *Cancer Res* 64(11) (2004) 4049–54. [PubMed: 15173020]
- [42]. Schantz SP, Zhang ZF, Spitz MS, Sun M, Hsu TC, Genetic susceptibility to head and neck cancer: Interaction between nutrition and mutagen sensitivity, *Laryngoscope* 107(6) (1997) 765–781. [PubMed: 9185733]
- [43]. Smith EM, Ritchie JM, Summersgill KF, Klussmann JP, Lee JH, Wang D, Haugen TH, Turek LP, Age, sexual behavior and human papillomavirus infection in oral cavity and oropharyngeal cancers, *Int J Cancer* 108(5) (2004) 766–72. [PubMed: 14696105]
- [44]. Suzuki T, Wakai K, Matsuo K, Hirose K, Ito H, Kuriki K, Sato S, Ueda R, Hasegawa Y, Tajima K, Effect of dietary antioxidants and risk of oral, pharyngeal and laryngeal squamous cell carcinoma according to smoking and drinking habits, *Cancer Sci* 97(8) (2006) 760–7. [PubMed: 16800818]
- [45]. Szymanska K, Hung RJ, Wunsch-Filho V, Eluf-Neto J, Curado MP, Koifman S, Matos E, Menezes A, Fernandez L, Daudt AW, Boffetta P, Brennan P, Alcohol and tobacco, and the risk of cancers of the upper aerodigestive tract in Latin America: a case-control study, *Cancer Causes Control* 22(7) (2011) 1037–46. [PubMed: 21607590]
- [46]. Twardella D, Loew M, Rothenbacher D, Stegmaier C, Ziegler H, Brenner H, The diagnosis of a smoking-related disease is a prominent trigger for smoking cessation in a retrospective cohort study, *J Clin Epidemiol* 59(1) (2006) 82–9. [PubMed: 16360565]
- [47]. Zhang Z, Shi Q, Liu Z, Sturgis EM, Spitz MR, Wei Q, Polymorphisms of methionine synthase and methionine synthase reductase and risk of squamous cell carcinoma of the head and neck: a case-control analysis, *Cancer Epidemiol Biomarkers Prev* 14(5) (2005) 1188–93. [PubMed: 15894670]



- [48]. Sun EC, Curtis R, Melbye M, Goedert JJ, Salivary gland cancer in the United States, *Cancer Epidemiol Biomarkers Prev* 8(12) (1999) 1095–100. [PubMed: 10613342]
- [49]. Franco EL, Kowalski LP, Oliveira BV, Curado MP, Pereira RN, Silva ME, Fava AS, Torloni H, Risk factors for oral cancer in Brazil: a case-control study, *Int J Cancer* 43(6) (1989) 992–1000. [PubMed: 2732011]
- [50]. Leffondre K, Abrahamowicz M, Siemiatycki J, Rachet B, Modeling smoking history: a comparison of different approaches, *Am J Epidemiol* 156(9) (2002) 813–23. [PubMed: 12396999]
- [51]. Greenland S, Finkle WD, A critical look at methods for handling missing covariates in epidemiologic regression analyses, *Am J Epidemiol* 142(12) (1995) 1255–64. [PubMed: 7503045]
- [52]. Rubin DB, *Multiple imputation for nonresponse in surveys*, John Wiley and Sons, Inc, New York, 1987.
- [53]. DerSimonian R, Laird N, Meta-analysis in clinical trials, *Control Clin Trials* 7(3) (1986) 177–88. [PubMed: 3802833]
- [54]. Wiencke JK, Thurston SW, Kelsey KT, Varkonyi A, Wain JC, Mark EJ, Christiani DC, Early age at smoking initiation and tobacco carcinogen DNA damage in the lung, *J Natl Cancer Inst* 91(7) (1999) 614–9. [PubMed: 10203280]
- [55]. Maxwell JH, Grandis JR, Ferris RL, HPV-Associated Head and Neck Cancer: Unique Features of Epidemiology and Clinical Management, *Annu Rev Med* 67 (2016) 91–101. [PubMed: 26332002]
- [56]. Young D, Xiao CC, Murphy B, Moore M, Fakhry C, Day TA, Increase in head and neck cancer in younger patients due to human papillomavirus (HPV), *Oral Oncol* 51(8) (2015) 727–30. [PubMed: 26066977]

### Highlights

- Younger age at starting tobacco use was associated with higher HNC risk.
- Age of starting tobacco was not associated with HNC risk after adjusting for tobacco use.
- Increased HNC risk with younger start age appears to be mediated by cumulative tobacco use.

**Table 1.**

Distribution of demographic characteristics among head and neck cancer cases and controls who have ever used tobacco products, in the INHANCE Consortium

	Cases (N = 17,146)		Controls (N=17,449)	
	n	%	n	%
<b>Age, years</b>				
< 40	467	2.7	936	5.4
40–44	868	5.1	1051	6.0
45–49	1787	10.4	1662	9.5
50–54	2658	15.5	2558	14.7
55–59	3207	18.7	3012	17.3
60–64	3032	17.7	2929	16.8
65–69	2513	14.7	2466	14.1
70–74	1710	10.0	1886	10.8
75	904	5.3	949	5.4
$P^a$		<0.001		
<b>Sex</b>				
Female	3123	18.2	3062	17.6
Male	14023	81.8	14387	82.5
$P^a$		0.1061		
<b>Ethnicity</b>				
White	11071	64.6	12037	69.0
Black	1011	5.9	689	4.0
Hispanic	166	1.0	231	1.3
Asian/Pacific Islanders	953	5.6	2228	12.8
Brazilian	3786	22.1	2144	12.3
Others	159	0.9	120	0.7
$P^a$		<0.001		
<b>Education level</b>				
No formal educaion	574	3.7	218	1.5
Junior high school	6056	38.6	5030	33.7
Some high school	2711	17.3	2151	14.4
High school graduate	2505	16.0	2176	14.6
Technical s'hool, some college	2473	15.7	3029	20.3
College graduate or higher	1390	8.9	2307	15.5
Missing	1437		2538	
$P^a$		<0.001		
<b>Alcohol drinks per day</b>				

	Cases (N = 17,146)		Controls (N=17,449)	
	n	%	n	%
Never drinkers	1711	10.5	2958	17.6
>0-1	2806	17.3	4881	29.1
1-2	3305	20.3	4101	24.4
3-4	2213	13.6	2017	12.0
5	6211	38.2	2838	16.9
Missing	900		654	
$P^a$		<0.001		
<b>Tobacco packyears</b>				
>0-10	1134	7.0	3633	21.7
11-20	1535	9.4	3021	18.0
21-30	2216	13.6	2682	16.0
31-40	2645	16.3	2352	14.0
41-50	2348	14.5	1639	9.8
>50	6376	39.2	3438	20.5
Missing	892		684	
$P^a$		<0.001		
<b>Years since quitting tobacco use</b>				
Current users or quit 1 year	12717	74.7	8423	48.7
1-9	1967	11.6	2402	13.9
10-19	1173	6.9	2699	15.5
20	1163	6.8	3767	21.8
Missing	126		158	
$P^a$		<0.001		
<b>Cancer subsite</b>				
Oral cavity	4668	27.2		
Oropharynx	4275	24.9		
Hypopharynx	1340	7.8		
Oral/pharynx NOS	1507	8.8		
Larynx	5079	29.6		
Unspecified	277	1.6		

<sup>a</sup>Chi-square test (two-sided)

Age at start of using tobacco products and the risk of head and neck cancer, in the INHANCE Consortium

Table 2.

	Cases (%)	Controls (%)	Basic Model		Model without tobacco packyears		Model adjusted for tobacco packyears	
			OR <sup>a</sup>	95% CI	OR <sup>b</sup>	95% CI	OR <sup>c</sup>	95% CI
30	713 (4.3)	1079 (6.3)	1.00	Referent	1.00	Referent	1.00	Referent
25-29	841 (5.1)	1164 (6.8)	1.14	0.99, 1.32	1.28	1.10, 1.49	1.05	0.89, 1.23
20-24	3297 (19.9)	4512 (26.4)	1.31	1.16, 1.48	1.50	1.33, 1.71	1.13	0.98, 1.29
15-19	7432 (44.8)	7386 (43.2)	1.35	1.20, 1.51	1.59	1.41, 1.79	1.08	0.94, 1.23
10-14	3594 (21.7)	2573 (15.1)	1.43	1.26, 1.62	1.60	1.40, 1.82	0.99	0.85, 1.14
< 10	707 (4.3)	380 (2.2)	1.49	1.25, 1.79	1.64	1.35, 1.97	0.97	0.80, 1.19
Missing	562	355						
<b>P trend</b>			<0.01		<0.01			0.21

Abbreviations: CI, confidence interval; INHANCE, International Head and Neck Cancer Epidemiology; OR, odds ratio.

<sup>a</sup>Model was adjusted for age, sex, race/ethnicity, education levels, and alcohol drinks per day.

<sup>b</sup>Model was adjusted for age, sex, race/ethnicity, education levels, alcohol drinks per day, duration of using chewing tobacco, duration of using snuff, and years since quitting tobacco use

<sup>c</sup>Model was adjusted for age, sex, race/ethnicity, education levels, alcohol drinks per day, tobacco packyears, interaction between alcohol drinks per day and tobacco packyears, duration of using chewing tobacco, duration of using snuff, and years since quitting tobacco use.

**Table 3**

Age at start of using tobacco products and the risk of head and neck cancer, by subsite, in the INHANCE Consortium

		Cases/Controls	Model without tobacco packyears		Model adjusted for tobacco packyears	
			OR <sup>a</sup>	95% CI	OR <sup>b</sup>	95% CI
<b>Oral cavity</b>						
	30	250/1043	1.00	Referent	1.00	Referent
	25–29	260/1124	1.20	0.95, 1.52	1.00	0.78, 1.29
	20–24	881/4402	1.29	1.06, 1.56	0.99	0.80, 1.23
	15–19	1968/7049	1.33	1.10, 1.60	0.94	0.76, 1.16
	10–14	890/2538	1.20	0.98, 1.47	0.77	0.62, 0.96
	< 10	193/377	1.37	1.04, 1.81	0.84	0.62, 1.13
	Missing	226/355				
	<i>P</i> <sub>trend</sub>		0.09		<0.01	
<b>Oropharynx</b>						
	30	157/1043	1.00	Referent	1.00	Referent
	25–29	210/1124	1.31	1.02, 1.70	1.10	0.84, 1.44
	20–24	797/4402	1.45	1.17, 1.80	1.13	0.91, 1.42
	15–19	1972/7049	1.51	1.23, 1.86	1.08	0.87, 1.35
	10–14	889/2538	1.46	1.17, 1.82	0.98	0.77, 1.24
	< 10	159/377	1.51	1.12, 2.03	0.92	0.67, 1.27
	Missing	105/355				
	<i>P</i> <sub>trend</sub>		<0.01		0.16	
<b>Hypopharynx</b>						
	30	54/941	1.00	Referent	1.00	Referent
	25–29	65/1065	1.14	0.75, 1.73	0.88	0.57, 1.37
	20–24	308/4137	1.63	1.16, 2.30	1.16	0.80, 1.68
	15–19	547/6520	1.48	1.06, 2.07	0.92	0.64, 1.33
	10–14	277/2313	1.44	1.00, 2.06	0.81	0.55, 1.20
	< 10	64/349	1.51	0.95, 2.39	0.81	0.50, 1.33
	Missing	25/260				
	<i>P</i> <sub>trend</sub>		0.12		0.04	
<b>Larynx</b>						
	30	160/855	1.00	Referent	1.00	Referent
	25–29	217/969	1.58	1.23, 2.04	1.13	0.87, 1.47
	20–24	2218/5954	1.96	1.59, 2.42	1.25	1.00, 1.55
	15–19	1142/2052	2.23	1.82, 2.73	1.23	0.99, 1.53
	10–14	219/313	2.46	1.98, 3.06	1.20	0.95, 1.51
	< 10	2218/5954	2.38	1.79, 3.16	1.16	0.86, 1.57

	Cases/Controls	Model without tobacco packyears		Model adjusted for tobacco packyears	
		OR <sup>a</sup>	95% CI	OR <sup>b</sup>	95% CI
<b>Oral cavity</b>					
	Missing	173/246			
	<i>P trend</i>		<0.01	0.40	

Abbreviations: CI, confidence interval; INHANCE, International Head and Neck Cancer Epidemiology; OR, odds ratio.

<sup>a</sup>Model was adjusted for age, sex, race/ethnicity, education levels, alcohol drinks per day, duration of using chewing tobacco, duration of using snuff, and years since quitting tobacco use.

<sup>b</sup>Model was additionally adjusted for tobacco packyears and interaction between alcohol drinks per day and tobacco packyears.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript



**Table 4**

Age at start of using tobacco products and the risk of head and neck cancer, by years since quitting tobacco use, in the INHANCE Consortium,

	Cases/Controls		Model without tobacco packyears		Model adjusted for tobacco packyears	
			OR <sup>a</sup>	95% CI	OR <sup>b</sup>	95% CI
<b>Current tobacco users</b>						
	30	529/659	1.00	Referent	1.00	Referent
	25–29	616/575	1.49	1.23, 1.79	1.11	0.91, 1.36
	20–24	2389/2147	1.79	1.53, 2.08	1.15	0.97, 1.37
	15–19	5399/3282	1.83	1.58, 2.13	1.01	0.85, 1.19
	10–14	2767/1307	1.86	1.58, 2.18	0.90	0.75, 1.08
	< 10	543/203	1.88	1.49, 2.37	0.86	0.66, 1.10
	Missing	474/250				
	<i>P</i> trend		<0.01		<0.01	
<b>Quit tobacco use &lt; 20 years</b>						
	30	132/268	1.00	Referent	1.00	Referent
	25–29	158/345	0.93	0.68, 1.28	0.89	0.64, 1.22
	20–24	651/1346	1.09	0.84, 1.42	0.99	0.75, 1.30
	15–19	1414/2250	1.18	0.91, 1.51	1.04	0.80, 1.36
	10–14	633/763	1.18	0.90, 1.55	1.02	0.76, 1.37
	< 10	130/108	1.28	0.87, 1.89	1.10	0.73, 1.64
	Missing	22/21				
	<i>P</i> trend		0.03		0.33	
<b>Quit tobacco use ≥ 20 years</b>						
	30	41/124	1.00	Referent	1.00	Referent
	25–29	54/230	0.78	0.47, 1.29	0.74	0.44, 1.23
	20–24	237/997	0.83	0.55, 1.26	0.77	0.50, 1.19
	15–19	600/1837	0.96	0.64, 1.44	0.86	0.57, 1.31
	10–14	189/498	0.93	0.60, 1.43	0.78	0.49, 1.23
	< 10	32/69	0.87	0.47, 1.62	0.76	0.40, 1.44
	Missing	10/12				
	<i>P</i> trend		0.48		0.85	

Abbreviations: CI, confidence interval; INHANCE, International Head and Neck Cancer Epidemiology; OR, odds ratio.

<sup>a</sup> Model was adjusted for age, sex, race/ethnicity, education levels, alcohol drinks per day, duration of using chewing tobacco, duration of using snuff, and years since quitting tobacco use.

<sup>b</sup> Model was additionally adjusted for tobacco packyears and interaction between alcohol drinks per day and tobacco packyears.

**Table 5**

Age at start of using tobacco products and the risk of head and neck cancer, by tobacco packyears, in the INHANCE Consortium

		Cases/Controls	Model without tobacco packyears		Model adjusted for tobacco packyears	
			OR <sup>a</sup>	95% CI	OR <sup>b</sup>	95% CI
<b>20 tobacco packyears</b>						
	30	306/705	1.00	Referent	1.00	Referent
	25–29	223/609	0.94	0.74, 1.19	0.89	0.70, 1.13
	20–24	632/1823	1.01	0.84, 1.23	0.94	0.78, 1.15
	15–19	1056/2725	1.01	0.83, 1.21	0.92	0.76, 1.11
	10–14	324/615	1.00	0.80, 1.27	0.90	0.71, 1.14
	< 10	58/74	1.02	0.67, 1.56	0.94	0.61, 1.44
	Missing	70/103				
	<i>P trend</i>		0.78		0.51	
<b>&gt; 20 tobacco packyears</b>						
	30	304/280	1.00	Referent	1.00	Referent
	25–29	570/507	1.14	0.90, 1.44	1.06	0.83, 1.34
	20–24	2542/2591	1.23	1.00, 1.51	1.10	0.89, 1.35
	15–19	6181/4499	1.22	1.00, 1.48	1.03	0.84, 1.26
	10–14	3162/1866	1.18	0.96, 1.45	0.94	0.76, 1.17
	< 10	610/286	1.20	0.93, 1.55	0.94	0.73, 1.22
	Missing	216/82				
	<i>P trend</i>		0.49		0.03	

Abbreviations: CI, confidence interval; INHANCE, International Head and Neck Cancer Epidemiology; OR, odds ratio.

<sup>a</sup>Model was adjusted for age, sex, race/ethnicity, education levels, alcohol drinks per day, duration of using chewing tobacco, duration of using snuff, and years since quitting tobacco use.

<sup>b</sup>Model was additionally adjusted for tobacco packyears and interaction between alcohol drinks per day and tobacco packyears.

**Table 6.**

Age at start of using tobacco products and the risk of head and neck cancer, by duration of using tobacco products, in the INHANCE Consortium

Smoking Duration	Cases/Controls	Model without tobacco packyears		Model adjusted for tobacco packyears		
		OR <sup>a</sup>	95% CI	OR <sup>b</sup>	95% CI	
<b>1–19 years</b>						
30	222/529	1.00	Referent	1.00	Referent	
25–29	94/364	0.81	0.58, 1.12	0.76	0.54, 1.06	
20–24	268/1092	0.85	0.65, 1.13	0.78	0.59, 1.04	
15–19	526/1814	0.97	0.74, 1.26	0.88	0.66, 1.16	
10–14	128/374	0.95	0.68, 1.34	0.86	0.61, 1.23	
< 10	10/14	1.04	0.44, 2.46	1.00	0.41, 2.42	
Missing	0/0					
<i>P</i> <sub>trend</sub>		0.63		0.96		
<b>20–39 years</b>						
30	410/474	1.00	Referent	1.00	Referent	
25–29	525/625	1.26	1.02, 1.56	1.06	0.85, 1.32	
20–24	1790/2332	1.55	1.28, 1.86	1.21	0.99, 1.47	
15–19	3531/3540	1.60	1.32, 1.93	1.14	0.93, 1.40	
10–14	1231/1091	1.58	1.27, 1.96	1.03	0.82, 1.30	
< 10	101/114	1.45	0.99, 2.12	0.90	0.61, 1.34	
Missing	0/0					
<i>P</i> <sub>trend</sub>		<0.01		0.63		
<b>40 years</b>						
30	68/45	1.00	Referent	1.00	Referent	
25–29	208/156	0.93	0.57, 1.51	0.86	0.51, 1.45	
20–24	1217/1060	0.96	0.62, 1.50	0.79	0.49, 1.27	
15–19	3355/2002	0.98	0.63, 1.52	0.72	0.44, 1.15	
10–14	2230/1098	0.98	0.63, 1.54	0.67	0.41, 1.08	
< 10	594/242	1.02	0.64, 1.64	0.68	0.41, 1.13	
Missing	0/0					
<i>P</i> <sub>trend</sub>		0.59		0.01		

Abbreviations: CI, confidence interval; INHANCE, International Head and Neck Cancer Epidemiology; OR, odds ratio.

<sup>a</sup>Model was adjusted for age, sex, race/ethnicity, education levels, alcohol drinks per day, duration of using chewing tobacco, duration of using snuff, and years since quitting tobacco use.

<sup>b</sup>Model was additionally adjusted for tobacco packyears and interaction between alcohol drinks per day and tobacco packyears.