

UCSF

UC San Francisco Previously Published Works

Title

The Demographic and Medical Correlates of Plasma A β 40 and A β 42

Permalink

<https://escholarship.org/uc/item/6vg493w5>

Journal

Alzheimer Disease & Associated Disorders, 27(3)

ISSN

0893-0341

Authors

Metti, Andrea L
Cauley, Jane A
Ayonayon, Hilsa N
et al.

Publication Date

2013-07-01

DOI

10.1097/wad.0b013e318260a8cb

Peer reviewed



Published in final edited form as:

Alzheimer Dis Assoc Disord. 2013 ; 27(3): 244–249. doi:10.1097/WAD.0b013e318260a8cb.

The Demographic and Medical Correlates of Plasma A β 40 and A β 42

Andrea L. Metti, MPH, Jane A. Cauley, DrPH, Hilsa N. Ayonayon, PhD, Tamara B. Harris, MD, MS, Caterina Rosano, MD, Jeff D. Williamson, MD, and Kristine Yaffe, MD for the Health, Aging and Body Composition Study

Department of Epidemiology, Graduate School of Public Health, University of Pittsburgh, Pittsburgh, PA (AM, JC, CR); Departments of Psychiatry (KY), Neurology (KY), and Epidemiology and Biostatistics (HA, KY), University of California San Francisco, San Francisco, CA; Laboratory of Epidemiology, Demography and Biometry, National Institute on Aging, Bethesda, MD (TH); Department of Geriatrics and Gerontology, Wake Forest University Medical Center, Winston-Salem, NC (JW); San Francisco Veteran's Affairs Medical Center, San Francisco, CA (KY)

Abstract

Plasma amyloid beta-42 (A β 42) and A β 42/A β 40 are increasingly recognized as biomarkers for dementia, with low levels indicating increased risk. Little is known about the demographic and medical correlates of plasma A β 40 or A β 42. In 997 community-dwelling, non-demented older adults from the Health, Aging and Body Composition Study, we determined the cross-sectional association between a wide range of demographic and medical variables with A β 40 and A β 42. In multivariate stepwise linear regression models, A β 40 was significantly associated with race (β = -14.70, F = 22.01, p < 0.0001), age (β = 1.34, F = 6.39, p = 0.01), creatinine (β = 52.91, F = 151.77, p < 0.0001), and serum brain-derived neurotrophic factor (BDNF) (β = -0.0004, F = 7.34, p = 0.007); A β 42 was significantly associated with race (β = -3.72, F = 30.83, p < 0.0001), sex (β = 1.39, F = 4.32, p = 0.04), education (β = 1.50, F = 4.78, p = 0.03), Apolipoprotein E (APOE) e4 allele status (β = -2.82, F = 16.57, p < 0.0001), and creatinine (β = 9.32, F = 120.09, p < 0.0001). These correlates should be considered as potential confounders in future studies investigating plasma A β as a biomarker of dementia. Understanding fully how these correlates mediate or modify the association between plasma A β and dementia will be a fundamental step in determining the biological pathways through which plasma A β 40 and A β 42 are associated with dementia, and in determining their full potential as biomarkers.

Keywords

Plasma amyloid beta; dementia; cognitive decline; biomarker; epidemiology

Introduction

Recent studies have indicated that plasma amyloid beta-40 (A β 40), A β 42 and the ratio (A β 42/A β 40) may be promising biological markers for cognitive impairment or dementia in older adults. For example, decreased plasma A β 42 was shown to be associated with increased risk for developing Alzheimer's disease (AD) and increased cognitive decline over

Corresponding Author: Andrea L. Metti, MPH, Bellefield Professional Building, Center for Aging and Population Health, University of Pittsburgh Department of Epidemiology, 130 N. Bellefield, Fourth Floor, 456, Pittsburgh, PA 15213, Phone: 412-624-2957, Fax: 412-624-7805, Alw111@pitt.edu.

Conflicts of Interest

For the remaining authors no conflicts of interest were reported.

time.¹⁻⁴ Similarly, a prospective longitudinal study found that low plasma A β 40 was associated with incident AD in older men.⁵ Low A β 42/A β 40 has also been associated with increased cognitive decline over time among non-demented elders, and with increased risk for AD.^{1, 6} These two proteins are of interest because they accumulate in the brain, forming the plaques found in AD patients. Furthermore, these proteins have been well established as biomarkers for AD when measured from cerebrospinal fluid (CSF) samples. For example, CSF A β 42 declines in patients with incident dementia, and has been documented as one of the most sensitive biomarkers for detecting prevalent AD.^{7, 8} Similarly, some studies have shown that low CSF A β 40 is associated with mild cognitive impairment (MCI) or the early stages of dementia.⁹ The possibility of identifying A β as a useful biomarker from plasma is clinically important because a simple blood draw is much less invasive and less expensive than the lumbar puncture required to obtain CSF.

In spite of this ongoing research, little is known about the general demographic and medical correlates of plasma A β 40 and A β 42, and how correlates may impact the relationship between plasma A β and cognitive decline. This information is important so these proteins can be fully understood as biomarkers, and so future studies investigating the potential use of plasma A β 40 and A β 42 as biomarkers for cognitive decline can measure and account for all potential confounders. Furthermore, understanding what factors are related to plasma A β 40 and A β 42 may shed some light on the mechanisms underlying an association between plasma A β and cognitive function. Importantly, one previous study has investigated correlates of plasma A β , but the study population consisted of only 205 cognitively normal controls, while the remaining sample was comprised of patients with MCI (n=348) or AD (n=162).¹⁰ As previous studies have shown that plasma A β levels change at different stages of disease progression^{11, 12}, more studies are needed to investigate correlates of these markers in community-dwelling, non-demented older adults. Thus, the objective of this study is to describe the demographic and medical correlates of plasma A β 40 and A β 42 in a population of community-dwelling, non-demented, black and white older adults, age 70–79 at baseline.

Methods

Study Population

Community-dwelling white and black older adults were enrolled in the ongoing Health Aging and Body Composition (Health ABC) study. This prospective cohort study began in 1997, and adults ranged in age from 70 to 79 years at enrollment, and lived in Memphis, TN or Pittsburgh, PA. Participants were recruited from a random sample of Medicare eligible adults living within the designated zip codes, and were eligible if they reported no difficulties performing activities of daily living, walking a quarter mile, or climbing 10 steps without resting. They also had to be free of life-threatening cancers, and plan to remain within the study area for at least three years.¹³ Plasma A β 42 and A β 40 were measured in a random sample of 1000 sex and race stratified participants. Of those, 3 had incomplete plasma amyloid data due to insufficient amounts of stored blood or measurement error; thus our analytic cohort was comprised of 997 participants. All participants included in this analytic cohort were free of cognitive impairment at baseline; consistent with previous literature, cognitive impairment was defined as a Modified Mini-Mental Status Exam (3MS) score <80.¹⁴ Compared to Health ABC participants who did not have plasma A β measured, those in the subgroup were more likely to be black, female, and to have lower mean education, but did not differ on other characteristics. This study was approved by the institutional review boards of the University of Pittsburgh and the University of Tennessee, Memphis, and that of the Coordinating Center, the University of California, San Francisco. All participants signed a written informed consent.

Beta Amyloid

Stored plasma obtained at the first Health ABC follow-up visit was used to measure A β 40 and A β 42. Plasma was stored at -70° C at Fisher BioServices, Inc. Laboratories and shipped directly to the analytical laboratory. Plasma A β was measured at the laboratory of Dr. Steven Younkin at the Mayo Clinic using Innogenetics INNO-BIA assays. The detection limit for this assay is 12 pg/ml for A β 40 and 5 pg/ml for A β 42. Mean inter-assay coefficient of variation was 9.9% for A β 40 and 9.3% for A β 42 and mean intra-assay coefficient of variation was 3.5% for A β 40 and 2.3% for A β 42. Consistent with previous literature, tertile cutoffs were used to categorize A β 42 and A β 42/A β 40 into “low” “medium” and “high” groups.

Potential Correlates

At baseline, demographic data including self-reported participant age, race, sex and education were recorded. Literacy was measured at baseline with the Rapid Estimate of Adult Literacy in Medicine (REALM).¹⁵ REALM scores were categorized as 8th grade and 8th grade, as it has been reported that this cutoff differentiates those who are functionally illiterate versus literate in a healthcare setting.^{16, 17} Cognitive function at baseline was measured with the Modified Mini-Mental Status Exam (3MS). The 3MS is an assessment of global cognitive function with components for orientation, concentration, language, praxis, and immediate and delayed memory.¹⁸ Social support scores were based on the average frequency of visits per week with friends, neighbors, and relatives. Scores were dichotomized into two groups: < the median or \geq the median. Self-reported information on smoking history was recorded and dichotomized into never versus ever having smoked.

Prevalent disease algorithms based on both self-report and physician diagnoses, recorded medications and laboratory data were used to create comorbidity variables indicating presence of diabetes mellitus, hypertension, stroke or transient ischemic attack (TIA), and myocardial infarction (MI). The Center for Epidemiologic Studies Depression Scale (CES-D) was used to assess depressive symptoms with a score ≥ 16 consistent with possible depression.¹⁹ Body mass index (BMI) (kg/m²) was calculated from direct height and weight measurements at baseline. Physical activity level was determined using a standardized questionnaire designed specifically for the Health ABC study, which has been previously described.²⁰ Total physical activity including household chores, paid work, volunteer work, care giving, stair climbing, non-exercise walking, walking for exercise and other moderate and vigorous exercise activities, were measured in kilocalories per week (kcal/wk).²⁰

Apolipoprotein E (APOE) e4 allele status was determined using standard Single Nucleotide Polymorphism (SNP) genotyping techniques and dichotomized into having one or more APOE e4 allele versus no allele.²¹ Brain-derived neurotrophic factor (BDNF) was measured from frozen serum collected at the first follow-up visit. Assays were performed by R&D Systems' Analytical Testing Service, using their own commercial kit employing an enzyme-linked immunosorbent assay method. The detection limit for this assay is 1250 pg/mL. The mean inter-assay coefficient of variation is 9.0%, and the mean coefficient of variation within assay is 5.0%.

At baseline, low-density lipoprotein cholesterol (LDL) (mg/dl), high-density lipoprotein cholesterol (HDL) (mg/dl), total cholesterol (mg/dl) and triglycerides (mg/dl) were measured from fasting blood serum. Measures of high sensitivity C-reactive protein (CRP), tumor necrosis factor- α (TNF α), interleukin-6 (IL-6) and creatinine were obtained from frozen serum or plasma collected at baseline after an overnight fast. Samples were frozen at -70° C and were shipped to the Core Laboratory at the University of Vermont.^{22, 23} Serum CRP was measured in duplicate by enzyme-linked immunosorbent assay on the basis of

purified protein and polyclonal anti-CRP antibodies, and assays were standardized according to the World Health Organization First International Reference Standard with a sensitivity of 0.08 $\mu\text{g/ml}$.²² Plasma IL-6 and TNF α were measured in duplicate by ELISA kits from the R&D Systems (Minneapolis, MN).²⁴ The detectable limit for IL-6 was 0.10 pg/ml, and for TNF α was 0.18 pg/ml.²⁴ Creatinine was measured with the colorimetric technique on a Johnson & Johnson VITROS 950 Chemistry Analyzer (Johnson & Johnson, New Brunswick, N.J., USA) using the enzymatic method.²³

Statistical Analyses

To describe the distribution of plasma A β 40 and A β 42, a variety of descriptive statistics were calculated including assessment of the mean, median, range and standard deviation. The relationship between A β 40 and A β 42 was examined using Spearman's correlation coefficient. Fisher's exact, Pearson's chi-square, and t-tests were used, as appropriate, to determine the association between plasma A β 40 and A β 42 tertile and baseline participant characteristics. Finally, stepwise multivariate linear regression models were examined for both A β 40 and A β 42 to determine which characteristics independently predicted plasma level. Variables that were associated in bivariate analyses ($p < 0.10$) with A β 40 or A β 42, respectively, were considered for these models. We used a forward stepwise selection process with variables significant at the $p = 0.15$ entry and exit criteria to determine the final multivariate models. The relative strength of the associations was expressed as an absolute difference in units of change chosen to approximate 1 SD in the distribution for each continuous variable or null category for dichotomous variables, similar to previous studies investigating correlates of markers in older adults.^{25, 26} The formula used to calculate the absolute difference in rate of change in plasma A β per unit change (SD) of the independent variable was $\beta^* = \text{unstandardized } \beta \times \text{unit change in independent variable}$. The corresponding 95% confidence intervals (CIs) were calculated using the following formula: $\beta \pm (1.96 \times \text{SE}) \times \text{unit change}$. All statistical analyses were performed using SAS version 9.1 (SAS Institute Inc., Cary, NC).

Results

Among the 997 community-dwelling older adults, the mean (standard deviation) of A β 40 and A β 42 were 191.6 pg/ml (50.3) and 33.9 (9.6), respectively. The median of A β 40 was 192.5 pg/ml, and of A β 42 was 32.8 pg/ml. The Spearman correlation coefficient was 0.51 ($p < 0.0001$) for both markers as continuous variables, and 0.42 ($p < 0.0001$) for both markers with tertile cutoffs.

Those with low plasma A β 40 were significantly younger (73.6 ± 3.0 years for low tertile vs. 74.5 ± 2.9 for high tertile, p -value = 0.0001), more likely to have less than a high school education (N(%), 223 (67.2%) vs. 197 (59.5%), p -value = 0.03) to be black (201 (60.4%) vs. 147 (44.1%), p -value < 0.0001), to have lower serum creatinine (mg/dl, 1.0 ± 0.2 vs. 1.2 ± 0.5 , p -value < 0.0001), to have higher HDL cholesterol (mg/dl, 56.6 ± 17.7 vs. 52.9 ± 16.5 , p -value = 0.02), to have higher BDNF (pg/ml, 23361.6 ± 10648.3 vs. 20944.4 ± 10022.9 , p -value = 0.006), and to have lower TNF α (pg/ml, 3.3 ± 2.1 vs. 3.7 ± 1.7 , p -value = 0.0007) (Table 1). There were no differences of A β 40 by sex, literacy, baseline 3MS scores, social support, physical activity, history of stroke/TIA, diabetes history, myocardial infarction, BMI, depressive symptoms, APOE e4 allele status, C-reactive protein, IL-6, total cholesterol, LDL cholesterol, triglycerides, or smoking history.

Those with low plasma A β 42 were more likely to be female (196 (59.8%) vs. 183 (55.5%), p -value = 0.05), to have less than a high school education (231 (70.6%) vs. 193 (58.8%), p -value = 0.006), to be black (214 (65.2%) vs. 151 (45.8%), p -value < 0.0001), to have a history of diabetes (94 (29.8%) vs. 68 (21.3%), p -value = 0.04) to have at least 1 APOE e4

allele (118 (38.3%) vs. 73 (23.1%), p -value=0.0002), to have lower creatinine (mg/dl, 1.0 ± 0.2 vs. 1.1 ± 0.6 , p -value<0.0001), to have greater social support (179 (54.6%) vs. 154 (46.7%), p -value=0.05), and to have higher HDL cholesterol (mg/dl, 57.3 ± 17.9 vs. 53.5 ± 16.9 , p =0.007) (Table 2). There were no differences by A β 42 and age, literacy, baseline 3MS scores, physical activity, history of stroke/TIA, myocardial infarction, BMI, depressive symptoms, C-reactive protein, TNF α , IL-6, BDNF, total cholesterol, LDL cholesterol, triglycerides, or smoking history.

In a stepwise multivariate linear regression model, A β 40 was significantly associated with age (β =1.34, F =6.39, p =0.01), race (β = -14.70, F =22.01, p <0.0001), creatinine (β = 52.91, F =151.77, p <0.0001), and BDNF (β =-0.0004, F =7.34, p =0.007) (Table 3). Similarly, A β 42 was significantly associated with race (β =-3.72, F =30.83, p <0.0001), sex (β =1.39, F =4.32, p =0.04), education (β =1.50, F =4.78, p =0.03), APOE e4 allele status (β =-2.82, F =16.57, p <0.0001), and creatinine (β =9.32, F =120.09, p <0.0001) (Table 3).

The strength of the associations expressed as an absolute difference in units of change (1 SD for continuous variables or null category for dichotomous variables) are shown in Table 4. Age, race, creatinine, and BDNF were all significant correlates of A β 40; race, sex, education, APOE e4, and creatinine were all significant correlates of A β 42, although sex and education were borderline (Table 4).

Discussion

Our results indicated that plasma A β 40 and A β 42 are fairly normally distributed among older adults without dementia. Furthermore, our results suggested that in a community-dwelling sample of white and black older adults, plasma A β 40 significantly differed by age, race, education, serum creatinine, serum BDNF, HDL cholesterol and the inflammatory marker TNF α . Similarly, plasma A β 42 was significantly associated with race, sex, education, social support, a history of diabetes, HDL cholesterol, serum creatinine and APOE e4 allele status.

Our results are supported by prior indirect associations between demographic and medical risk factors and AD. For example, we found black race was associated with significantly lower levels of plasma A β 40 and A β 42, and being female was significantly associated with a lower level of A β 42. As it has been shown that African Americans and females have an increased risk of AD, and other studies have shown low A β 42 is associated with increased risk of dementia, these results seem consistent.^{4, 27}

Our results showing an association between diabetes and plasma A β 42 are in support of earlier studies suggesting that consistently increased insulin levels in the brain may lead to increased A β 42 deposition and plaque formation, and subsequent decreases in peripheral plasma A β 42.^{28, 29} Our results differ from another longitudinal study that recently found higher total cholesterol and higher LDL cholesterol predicted low plasma A β 42.³⁰ Furthermore, this previous study found no association between HDL cholesterol and A β 42 at baseline or over time, as we did with plasma A β 40 and A β 42.³⁰ Similarly, we found no association between A β 40 or A β 42 with LDL, total cholesterol, triglycerides, or history of myocardial infarction or stroke – both of which have high cholesterol as a common risk factor. Finally, the association we found between both plasma A β 40 and A β 42 and serum creatinine – a measure of renal function – is supported by another study which found that in older patients with chronic renal failure, dialysis significantly lowered plasma A β 42 level; these results indicate a close relationship may exist between renal function and plasma A β 42.³¹

It is widely known that APOE e4 allele is associated with increased risk for AD, and in this study we found older adults with at least one APOE e4 allele were more likely to have low A β 42; APOE has also been shown to modify the association between A β 42 and A β 42/A β 40 and cognitive decline.^{6, 32} An interesting finding was the association between higher serum BDNF and low plasma A β 40. BDNF is thought to have neuroprotective effects and to promote brain plasticity.³³ BDNF has also been shown to regulate the Amyloid Precursor Protein (APP) which is thought to reduce the production of amyloid peptides (i.e. A β 40).³³ Thus, it is possible that higher levels of serum BDNF may, at least in part, regulate plasma A β 40, resulting in the lower plasma A β 40 level. There was no relationship found between plasma A β 42 and serum BDNF. The inflammatory marker TNF α has been shown to be associated with increased risk of cognitive decline over time.³⁴ While we found an association between low TNF α and low plasma A β 40, which is contrary to what we expected, we believe this could be due to the less specific association between A β 40 and AD or cognitive decline (versus the relationship between A β 42 and cognitive function).³⁴

Importantly, our results are also supported in part by the previous study investigating correlates and predictors of plasma A β in a population of cognitive normal, MCI and AD patients.¹⁰ Both studies reported relatively normal distributions of both plasma A β 40 and A β 42, indicating no need to log-transform plasma A β 40 or A β 42 when analyzing as continuous variables, although it was reported that the ratio needed log-transformation.¹⁰ Furthermore, both studies found significant associations between higher A β 40 and A β 42 and poorer kidney function in t-tests, and found that creatinine significantly predicted both markers.¹⁰ Other similarities were significant associations between age and A β 42, APOE and A β 42 and cholesterol and A β 40.¹⁰ This study did not support our findings in that it found no association between education or race with either plasma marker; in models the most significant predictors of A β 40 and A β 42 were creatinine, total proteins and cholesterol, while our results showed different significant predictors of A β 42 and A β 40.¹⁰ Interestingly in the other study, age, education nor APOE e4 significantly predicted either marker as we found, and these are known strong risk factors for AD.¹⁰ Finally, the prior study found a much stronger correlation between plasma A β 40 and A β 42 ($r=0.83$ vs. $r=0.51$).¹⁰ We believe differences are most likely attributable to different populations, as our study was made up of community-dwelling older adults who were all dementia free, compared to a population of primarily MCI and AD patients with only a small proportion of cognitively normal controls.¹⁰ Given these characteristics, the contrasting results could reflect that disease progression largely affects plasma A β measurements and correlates. Ultimately, more studies are needed to better understand the fluctuations of plasma A β 40 and A β 42 in older adults who age without cognitive decline, with MCI and with AD to better understand these relationships.

If plasma A β 40 and A β 42 indeed are useful biomarkers of cognitive decline and dementia, then it is important to fully understand how these biomarkers are influenced by demographic and medical conditions for several reasons. First of all, at a very basic level, it will be important so that future studies can measure and adjust for all potential confounders when investigating the association of plasma A β to ensure the most accurate results are presented. Understanding demographic and medical correlates will also be crucial in considering what factors may regulate plasma A β . This will be critical for gaining a better understanding of the mechanisms underlying cognitive decline and dementia. It will also help in identifying possible targets for interventions in reducing risk of or preventing dementia.

This study had several strengths, including the large sample size providing ample analytical power for this study. Measurement of numerous potential confounders allowed us to investigate a large number of demographic and medical characteristics in one sample. Finally, Innogenetics INNO-BIA assays were used for our measurements of A β 42 and

A β 40, and this method may provide more accurate measurements of A β 42 and A β 40 due to its high sensitivity, low variability, and high reproducibility.³⁵ There are also several weaknesses that should be taken into consideration when interpreting these results. These adults were all well functioning and community-dwelling at baseline, so results may not be generalizable to all older adults – for example, nursing home populations. Another potential weakness is that we did not have CSF measurements of A β and thus, we could not correlate plasma A β to measurements of CSF A β . As CSF A β has been such a well-established biomarker for dementia, it would have been useful.

This study identified the demographic and medical correlates of plasma A β 40 and A β 42. The correlates identified in this paper should be considered in future studies as potential confounders of the association between plasma A β and cognitive decline or dementia. Understanding fully how these correlates mediate or modify the association between plasma A β and dementia will be a fundamental step in determining the biological pathways through which plasma A β contributes to dementia and cognitive decline. Future studies should continue to investigate exactly how these demographic and medical correlates may influence the relationship between plasma A β 40 and A β 42, as it may help us target interventions for preventing and reducing the risk of AD and dementia.

Acknowledgments

Study Funding:

The Health, Aging and Body Composition Study is Supported by National Institute on Aging (NIA) Contracts N01-AG-6-2101; N01-AG-6-2103; N01-AG-6-2106; NIA grant R01-AG028050, and NINR grant R01-NR012459. This research was supported in part by the Intramural Research Program of the NIH, National Institute on Aging. Andrea Metti is supported by the National Institutes of Health Training Grant (2T32AG000181). Kristine Yaffe is supported in part by NIA Grant K24 AG031155.

References

1. Graff-Radford NR, Crook JE, Lucas J, et al. Association of low plasma A β 42/A β 40 ratios with increased imminent risk for mild cognitive impairment and Alzheimer disease. *Arch Neurol.* 2007; 64:354–62. [PubMed: 17353377]
2. Lewczuk P, Kornhuber J, Vanmechelen E, et al. Amyloid β peptides in plasma in early diagnosis of Alzheimer's disease: A multicenter study with multiplexing. *Experimental Neurology.* 2009 Aug 5. [Epub Ahead of Print].
3. Pesaresi M, Lovati C, Bertora P, et al. Plasma levels of beta-amyloid (1–42) in Alzheimer's disease and mild cognitive impairment. *Neurobiol Aging.* 2006; 27:904–05. [PubMed: 16638622]
4. Yaffe K, Weston AL, Graff-Radford NR, et al. Plasma beta-amyloid level and cognitive decline: Modulation by cognitive reserve. Unpublished Results. 2010
5. Sundelof J, Giedraitis V, Irizarry MC, et al. Plasma Beta Amyloid and the Risk of Alzheimer Disease and Dementia in Elderly Men. A Propective, Population-Based Cohort Study. *Arch Neurol.* 2008; 65(2):256–63. [PubMed: 18268197]
6. Yaffe K, Weston A, Graff-Radford NR, et al. Association of plasma β -amyloid level and cognitive reserve with subsequent cognitive decline. *JAMA.* 2011; 205(3):261–66. [PubMed: 21245181]
7. Shaw LM, Vanderstichele H, Knapik-Czajka M, et al. Cerebrospinal fluid biomarker signature in Alzheimer's disease neuroimaging initiative subjects. *Ann Neurol.* 2009; 65:403–13. [PubMed: 19296504]
8. Sunderland T, Linker G, Mirza N, et al. Decreased beta-amyloid 1–42 and increased tau levels in cerebrospinal fluid of patients with Alzheimer disease. *JAMA.* 2003; 289(16):2094–2103. [PubMed: 12709467]
9. Jensen M, Schroder J, Blomberg M, et al. Cerebrospinal fluid A β 42 is increased early in sporadic Alzheimer's disease and declines with disease progression. *Ann Neurol.* 1999; 45:504–11. [PubMed: 10211475]

10. Toledo JB, Vanderstichele H, Figurski M, et al. Factors affecting A β plasma levels and their utility as biomarkers in ADNI. *Acta Neuropathol.* 2011; 122:401–413. [PubMed: 21805181]
11. Schupf N, Tang MX, Fukuyama H, et al. Peripheral A β subspecies as risk biomarkers of Alzheimer's disease. *Proc Natl Acad Sci U S A.* 2008; 105(37):14052–7. [PubMed: 18779561]
12. Seppala TT, Herukka SK, Hanninen T, et al. Plasma A β 42 and A β 40 as markers of cognitive change in follow-up: A prospective, longitudinal, population-based cohort study. *J Neurol Neurosurg Psychiatry.* 2010
13. Yaffe K, Barnes D, Lindquist K, et al. Endogenous sex hormone levels and risk of cognitive decline in an older biracial cohort. *Neurobiology of Aging.* 2007; 28:171–78. [PubMed: 17097195]
14. Slinin Y, Paudel ML, Taylor BC, et al. 25-Hydroxyvitamin D levels and cognitive performance and decline in elderly men. *Neurology.* 2010; 74(1):33–41. [PubMed: 19940271]
15. Davis TC, Long SW, Jackson RH, et al. Rapid Estimate of Adult Literacy in Medicine: A shortened screening instrument. *Fam Med.* 1993; 25:391–5. [PubMed: 8349060]
16. Osborn C, Weiss B, Davis T, et al. Measuring adult literacy in health care: performance of the newest vital sign. *Am J Health Behav.* 2007; 31:S36–46. [PubMed: 17931135]
17. Rothman R, Malone R, Bryant B, Horlen C, DeWalt D, Pignone M. The relationship between literacy and glycemic control in a diabetes disease-management program. *The Diabetes Educator.* 2004; 30(2):263–73. [PubMed: 15095516]
18. Teng EL, Chui HC. The Modified Mini-Mental State (3MS) examination. *J Clin Psychiatry.* 1987; 48(8):314–18. [PubMed: 3611032]
19. Radloff L. The CES-D scale: A self-report depression scale for research in the general population. *Appl Psychol Meas.* 1977; 1:385–401.
20. Brach JS, Simonsick EM, Kritchevsky S, Yaffe K, Newman AB. Health Aging and Body Composition Study Research Group. The association between physical function and lifestyle activity and exercise in the health, aging and body composition study. *J Am Geriatr Soc.* 2004; 52(4):502–9. [PubMed: 15066063]
21. Hixson J, Vernier D. Restriction isotyping of human apolipoprotein E by gen amplification and cleavage with HhaI. *J Lipid Res.* 1990; 31(3):545–8. [PubMed: 2341813]
22. Kalogeropoulos A, Georgiopoulou V, Psaty BM, et al. Inflammatory markers and incident heart failure risk in older adults. The Health ABC (Health, Aging and Body Composition) Study. *J Am Coll Cardiol.* 2010; 55(19):2129–37. [PubMed: 20447537]
23. Minev E, Unruh M, Shlipak MG, et al. Association of cystatin C and depression in healthy elders: the health, aging and body composition study. *Nephron Clin Pract.* 2010; 116(3):c241–46. [PubMed: 20606485]
24. Yaffe K, Lindquist K, Penninx B, et al. Inflammatory markers and cognition in well-functioning African-American and white elders. *Neurology.* 2003; 61:76–80. [PubMed: 12847160]
25. Cauley JA, Blackwell T, Zmuda JM, et al. Correlates of trabecular and cortical volumetric bone mineral density at the femoral neck and lumbar spine: The Osteoporotic Fractures in Men Study (MrOS). *J Bone Miner Res.* 2010; 25(9):1958–71. [PubMed: 20572023]
26. Sheu Y, Cauley JA, Wheeler VW, et al. Natural history and correlates of hip BMD loss with aging in men of African ancestry: the Tobago Bone Health Study. *J Bone Miner Res.* 2009; 24(7):1290–98. [PubMed: 19257828]
27. Green RC, Cupples LA, Go R, et al. Risk of dementia among white and African American relatives of patients with Alzheimer disease. *JAMA.* 2002; 287(3):329–36. [PubMed: 11790212]
28. Duckworth WC, Bennett RG, Hamel FG. Insulin acts intracellularly on proteasomes through insulin-degrading enzyme. *Biochem Biophys Res Commun.* 1998; 244:390–94. [PubMed: 9514933]
29. Qiu WQ, Walsh DM, Ye Z, et al. Insulin-degrading enzyme regulates extracellular levels of amyloid beta-protein by degradation. *J Biol Chem.* 1998; 273(49):32730–8. [PubMed: 9830016]
30. Blasko I, Kemmler G, Jungwirth S, et al. Prospective study on association between plasma amyloid beta-42 and atherosclerotic risk factors. *J Neural Transm.* 2011; 118:663–72. [PubMed: 21344239]
31. Rubio I, Caramelo C, Gil A, Lopez MD, Garcia de Yébenes J. Plasma amyloid- β , A β 1–42, load is reduced by haemodialysis. *Journal of Alzheimer's Disease.* 2006; 10:439–43.

32. Jarvik G, Larson EB, Goddard K, Wijsman EM. Influence of apolipoprotein E genotype on the transmission of Alzheimer disease in a community-based sample. *Am J Hum Genet.* 1996; 58:191–200. [PubMed: 8554056]
33. Diniz BS, Teixeira AL. Brain-derived neurotrophic factor and Alzheimer's disease: Physiopathology and beyond. *Neuromolecular Med.* 2011 Sep 7. [Epub ahead of print].
34. Gorelick PB. Role of inflammation in cognitive impairment: results of observational epidemiological studies and clinical trials. *Ann NY Acad Sci.* 2010; 1207:155–62. [PubMed: 20955439]
35. Innogenetics. New prospects for research into Alzheimer's disease with INNO-BIA plasma A β forms. A standardized research test for measuring the concentrations of beta-amyloid isoforms in blood. *A Literature Review.* 2007:1–10.

Table 1Demographic and medical correlates of plasma A β 40 among 997 older adults

Baseline Characteristics N (%) or Mean (SD)	Tertile 1 n=333	Tertile 2 n=331	Tertile 3 N=333	p-value
Age, years	73.6 (3.0)	74.0 (2.8)	74.5 (2.9)	.0001
Black Race	201 (60.4)	190 (57.4)	147 (44.1)	<.0001
Female Sex	184 (55.3)	187 (56.5)	179 (53.8)	.78
Education (HS)	223 (67.2)	226 (68.5)	197 (59.5)	.03
Literacy (REALM, 8 th grade)	94 (29.9)	85 (27.8)	87 (27.8)	.79
3MS	91.0 (5.4)	91.2 (5.5)	91.1 (5.2)	0.90
Social Support *	171 (51.4)	165 (49.9)	152 (45.7)	0.26
Body Mass Index	27.7 (5.2)	28.0 (4.8)	27.5 (4.8)	.35
Physical Activity (kcal per week)	1055.5 (2373.2)	1133.3 (2455.5)	982.4 (1465.0)	0.66
Depressive Symptoms (CES-D 16)	13 (3.9)	22 (6.7)	11 (3.3)	.09
Stroke/TIA History	25 (7.5)	33 (10.1)	37 (11.3)	.25
Diabetes History	74 (23.2)	83 (25.6)	83 (25.9)	.69
Myocardial Infarction History	25 (7.6)	38 (11.6)	42 (12.8)	.07
Cholesterol (mg/dl)	202.7 (37.6)	208.4 (41.1)	202.2 (39.3)	0.08
Triglycerides (mg/dl)	126.8 (80.6)	131.1 (100.5)	138.5 (68.0)	0.19
HDL (mg/dl)	56.6 (17.7)	55.1 (16.7)	52.9 (16.5)	0.02
LDL (mg/dl)	121.4 (34.5)	127.2 (35.5)	121.6 (35.4)	0.06
APOE e4 allele (1 e4 allele)	96 (30.9)	101 (32.2)	93 (29.5)	.77
BDNF (pg/ml)	23361.6 (10648.3)	21206.2 (11064.9)	20944.4 (10022.9)	0.006
Creatinine (mg/dl)	1.0 (0.2)	1.0 (0.3)	1.2 (0.5)	<.0001
C-reactive Protein (μ g/mL)	3.0 (3.9)	2.9 (4.0)	3.4 (7.3)	.42
Tumor Necrosis Factor (pg/ml)	3.3 (2.1)	3.2 (1.3)	3.7 (1.7)	0.0007
IL-6 (pg/ml)	2.3 (1.9)	2.2 (1.7)	2.4 (1.8)	0.28
Smoker (Ever vs. Never)	183 (55.0)	180 (54.4)	180 (54.4)	.99

* N (%) below the median; variable defined as a combination of friends, neighbors and relatives who visit per week.

A β 40 indicates amyloid beta-40; SD standard deviation; HS high school; REALM Rapid Estimate of Adult Literacy in Medicine; 3MS Modified Mini-Mental Status Exam; CES-D Center for Epidemiologic Studies Depression Scale; TIA transient ischemic attack; HDL high-density lipoprotein cholesterol; LDL low-density lipoprotein cholesterol; APOE Apolipoprotein E; BDNF brain-derived neurotrophic factor; IL-6 interleukin-6.

Table 2Demographic and medical correlates of plasma A β 42 among 997 older adults

Baseline Characteristics N (%) or Mean (SD)	Tertile 1 n=328	Tertile 2 n=339	Tertile 3 n=330	p-value
Age, years	74.0 (3.0)	73.8 (2.9)	74.2 (2.9)	.17
Black Race	214 (65.2)	173 (51.0)	151 (45.8)	<.0001
Female Sex	196 (59.8)	171 (50.4)	183 (55.5)	.05
Education (HS)	231 (70.6)	222 (65.7)	193 (58.8)	.006
Literacy (REALM, 8 th grade)	95 (30.8)	97 (30.9)	74 (23.8)	.08
3MS Score	90.7 (5.5)	91.1 (5.3)	91.4 (5.3)	0.30
Social Support *	179 (54.6)	155 (45.7)	154 (46.7)	0.05
Body Mass Index	27.6 (5.2)	27.5 (4.3)	28.1 (5.3)	.26
Physical Activity (kcal per week)	929.8 (1971.7)	1042.1 (1733.5)	1198.4 (2669.9)	0.27
Depressive Symptoms (CES-D 16)	13 (4.0)	12 (3.5)	21 (6.7)	.17
Stroke/TIA History	28 (8.6)	31 (9.3)	36 (11.0)	.56
Diabetes History	94 (29.8)	78 (23.7)	68 (21.3)	.04
Myocardial Infarction History	39 (8.9)	34 (10.2)	42 (12.9)	.24
Cholesterol (mg/dl)	203.2 (38.0)	205.4 (39.2)	204.6 (41.2)	0.76
Triglycerides (mg/dl)	128.5 (80.9)	127.4 (65.6)	140.8 (101.9)	0.08
HDL (mg/dl)	57.3 (17.9)	53.9 (15.9)	53.5 (16.9)	0.007
LDL (mg/dl)	120.8 (34.7)	125.8 (35.4)	123.5 (35.4)	0.20
APOE e4 allele (1 e4 allele)	118 (38.3)	99 (31.3)	73 (23.1)	.0002
BDNF (pg/ml)	22798.0 (10910.9)	21328.8 (10138.0)	21408.9 (10810.7)	0.14
Creatinine (mg/dl)	1.0 (0.2)	1.0 (0.2)	1.1 (0.6)	<.0001
C-reactive Protein (μ g/mL)	3.2 (6.4)	2.7 (3.4)	3.4 (5.8)	.16
Tumor Necrosis Factor (pg/ml)	3.3 (2.1)	3.3 (1.4)	3.6 (1.7)	0.07
IL-6 (pg/ml)	2.2 (1.5)	2.4 (2.0)	2.4 (1.9)	0.58
Smoker (Ever vs. Never)	192 (58.5)	184 (54.3)	167 (50.9)	.15

* N (%) below the median; variable defined as a combination of friends, neighbors and relatives who visit per week.

A β 42 indicates amyloid beta-42; SD standard deviation; HS high school; REALM Rapid Estimate of Adult Literacy in Medicine; 3MS Modified Mini-Mental Status Exam; CES-D Center for Epidemiologic Studies Depression Scale; TIA transient ischemic attack; HDL high-density lipoprotein cholesterol; LDL low-density lipoprotein cholesterol; APOE Apolipoprotein E; BDNF brain-derived neurotrophic factor; IL-6 interleukin-6.

Table 3Stepwise multivariate linear regression associations with plasma A β 40 and A β 42

Variable	Parameter (β)	Standard Error	F-value	p-value
Aβ40				
Age	1.34	0.53	6.39	0.01
Black race	-14.70	3.13	22.01	<0.0001
Creatinine	52.91	4.29	151.77	<0.0001
Serum BDNF	-0.0004	0.0001	7.34	0.007
Aβ42				
Black race	-3.72	0.67	30.83	<0.0001
Femal sex	1.39	0.67	4.32	0.04
Education	1.50	0.69	4.78	0.03
APOE e4	-2.82	0.69	16.57	<0.0001
Creatinine	9.32	0.85	120.09	<0.0001

A β 40 indicates amyloid beta-40; A β 42 amyloid beta-42; BDNF brain-derived neurotrophic factor; APOE Apolipoprotein E.

Table 4

Multivariate correlates of the rate of change in plasma A β 40 and A β 42.

Variable	Unit	Rate of change in plasma A β (95% CI) per unit	
		A β 40	A β 42
Age	2.91	3.90 (0.88, 6.92)	NA
Race	1	-14.70 (-20.83, -8.57)	-3.72 (-5.03, -2.41)
Sex	1	NA	1.39 (0.08, 2.70)
Education	1	NA	1.50 (0.15, 2.85)
Creatinine	0.85 mg/dl	19.58 (16.47, 22.69)	-2.82 (-4.17, -1.47)
APOE e4	1	NA	3.45 (2.83, 4.07)
BDNF	10631.18 pg/ml	-4.25 (-6.33, -2.17)	NA

A β 40 indicates amyloid beta-40; A β 42 amyloid beta-42; CI confidence interval; NA not applicable; APOE Apolipoprotein E; BDNF brain-derived neurotrophic factor.