

# UC Santa Cruz

## UC Santa Cruz Previously Published Works

### Title

The complete mitochondrial genome of *Dascyllus trimaculatus* (Rüppell, 1829)

### Permalink

<https://escholarship.org/uc/item/6xr1r1w0>

### Journal

Mitochondrial DNA Part B, 8(1)

### ISSN

2380-2359

### Authors

Limón, Juliana  
Roberts, May B  
Schultz, Darrin T  
et al.

### Publication Date

2023-01-02

### DOI




10.1080/23802359.2022.2161838

### Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

## The complete mitochondrial genome of *Dascyllus trimaculatus* (Rüppell, 1829)

Juliana Limón<sup>a\*</sup>, May B. Roberts<sup>a\*</sup> , Darrin T. Schultz<sup>b,c,d</sup>  and Giacomo Bernardi<sup>a</sup> 

<sup>a</sup>Department of Ecology and Evolutionary Biology, University of California Santa Cruz, Santa Cruz, CA, USA; <sup>b</sup>Department of Molecular Evolution and Development, University of Vienna, Vienna, Austria; <sup>c</sup>Monterey Bay Aquarium Research Institute, Moss Landing, CA, USA; <sup>d</sup>Department of Biomolecular Engineering and Bioinformatics, University of California, Santa Cruz, CA, USA

### ABSTRACT

Damselfishes (family Pomacentridae) comprise approximately 400 species that play an important ecological role in temperate and coral reefs. Here, for the first time, we assemble and annotate the mitochondrial genome of *Dascyllus trimaculatus*, the three-spot dascyllus, a planktivorous damselfish that primarily recruits in anemones. The circular genome of *D. trimaculatus* is 16,967 bp in length and contains 13 protein-coding genes, 22 transfer RNA genes, two ribosomal RNA genes, and a control region. Gene arrangement and codon usage is similar to reported mitochondrial genomes of other damselfish genera, and a phylogenetic analysis of a set of damselfish representatives is consistent with known evolutionary analyses.

### ARTICLE HISTORY

Received 20 September 2022  
Accepted 20 December 2022

### KEYWORDS

*Dascyllus trimaculatus*  
mitochondrial genome;  
Pomacentridae



The family Pomacentridae, commonly known as damselfishes, contains 29 genera and over 400 species (Allen 1991; Tang et al. 2021), with most of these species occurring in the Indo-west and central Pacific region (Allen 1991; Tang et al. 2021). The three-spot damselfish, *Dascyllus trimaculatus* has a wide distribution that extends from the Red Sea to the Central Pacific and belongs to a species complex with three other recognized species (Bernardi et al. 2001; Leray et al. 2010). Here, we present, for the first time, the fully assembled mitochondrial genome of *D. trimaculatus* (Rüppell, 1829) to aid in future studies of this species complex. We placed this genome in its evolutionary context within the Pomacentridae family, and confirmed the geographic origin of our specimen using a tree based on the control region (d-loop) of several other *D. trimaculatus*.

A juvenile (38 mm TL) *Dascyllus trimaculatus* was acquired through the aquarium trade (LiveAquaria.com), and euthanized following an approved IACUC protocol animal use (Berg 1701.r2). The specimen was deposited at the University of California Museum of Natural History under the voucher number DTR\_Kuro\_0920G (mabrober@ucsc.edu). Total genomic DNA was extracted using muscle tissue from the individual (Qiagen DNeasy Blood & Tissue Kit, Hilden, Germany). A shotgun/whole genome library was prepared and sequenced using a MiSeq Illumina sequencing platform in the Paleogenomics Laboratory at the University of California Santa Cruz resulting in 2,356,971 contigs of 150 bp length each. Data were aligned with the closely related *Abudefduf vaigiensis* (GenBank: NC\_009064) in Geneious R11 and trimmed to create a consensus sequence (Geneious,

Auckland, New Zealand, <https://www.geneious.com>). The final genome was then annotated using MitoAnnotator (Iwasaki et al. 2013). The mitochondrial genome of *D. trimaculatus* was found to be 16,967 bp in length and contains 13 protein-coding genes, 22 transfer RNA genes, two ribosomal RNA genes, and a control region (SRA: SRR20942050, Run: SRR20942050, BioProject: PRJNA828170, BioSample: SAMN27642109-DTR\_Kuro\_0920G; GenBank sequence accession number ON556619) (Supplementary Figure S1). The genome has an overall base composition of 28.54% A, 29.41% C, 15.64% G, and 26.42% T. This composition is similar to the mitogenomes of other vertebrates, with a higher AT (54.95%) than GC (45.05%) content. A comparison of mutations (after removing the variable control region) between *D. trimaculatus* and its close relative *Chromis viridis* (see below) showed, as expected, a higher rate of transitions (10.31%;  $R = 3.17\%$ ,  $Y = 7.14\%$ ) than transversions (4.62%;  $M = 2.13\%$ ,  $W = 1.42\%$ ,  $S = 0.63\%$ ,  $K = 0.43\%$ ) resulting in a tr/tv ratio of 2.3.

A Neighbor-Joining tree based on mitochondrial genomes (all regions except the variable control region, i.e. an alignment of 16,178bp) of seven damselfish species (*Dascyllus trimaculatus*, *Chromis notata*, *Chromis viridis*, *Abudefduf vaigiensis*, *Amblyglyphidodon curacao*, *Chrysiptera biocellata*, and *Amphiprion ocellaris*) and one surfperch, Embiotocidae (*Embiotoca jacksoni*) to serve as an outgroup (following Tang et al. 2021) was established using PAUP (Swofford 2003). Node confidence was obtained by running 1000 bootstrap replicates (Felsenstein 1985; Figure 1).


In addition, a Neighbor-Joining phylogenetic tree based on 375 bp of the variable control region (D-loop) of our

**CONTACT** Giacomo Bernardi  [bernardi@ucsc.edu](mailto:bernardi@ucsc.edu)  Department of Ecology and Evolutionary Biology, University of California Santa Cruz, 115 McAllister Way, Santa Cruz, CA 95060, USA

\*Both authors contributed equally to this work.

© 2023 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



**Figure 1.** Phylogenetic relationships based on complete mitochondrial genomes of representative damselfish species and one surfperch outgroup (Black Surfperch, *Embiotoca jacksoni*). Numbers at nodes represent bootstrap values. The focal species of this study, *Dascyllus trimaculatus*, is labeled in red font. GenBank accession numbers are indicated between parentheses under the species names: *Embiotoca jacksoni* (NC\_029362), *Chrysiptera biocellata* (NC\_040304), *Amphiprion ocellaris* (NC\_009065), *Amblyglyphidodon curacao* (NC\_043918), *Dascyllus trimaculatus* (ON556619), *Chromis viridis* (MT199208), *Chromis notata* (NC\_052722), and *Abudegduf vaigiensis* (NC\_009064). Original drawings by MBR.

sample and an additional 636 different *D. trimaculatus* individuals (obtained from the literature, Leray et al. 2010) was established to locate the geographic origin of our sample (Supplementary Figure S2). Results suggests that the organism we sequenced is from the West Pacific Rim clade, with closest individuals from the Visayas (Philippines) and Manado (Indonesia).

### Author contributions

Conception and design: JL, MBR, DTS, and GB; analysis and interpretation of the data: JL, MBR, DTS, and GB; drafting of the paper: GB; revision and final approval of the version to be published: JL, MBR, DTS, and GB. All authors are accountable for all aspects of the work.

### Disclosure statement

No potential conflict of interest was reported by the authors.

### Funding

This work was supported by Marilyn C Davis Scholarship and American Association of University Women (through STARS Reentry program at UCSC), Burnand-Partridge Foundation and Dearmin Trust Honorary Fund Scholarships, tuition support by a Dissertation Quarter Fellowship (UCSC) to MBR. D.T.S. was supported by the US National Science Foundation [GRFP DGE 1339067], the US National Science Foundation [DEB-1542679] to Steven Haddock, and the European Research Council's Horizon 2020: European Union Research and Innovation Programme, Grant No. [945026] to Oleg Simakov.

### ORCID

May B. Roberts  <http://orcid.org/0000-0001-6882-4266>  
 Darrin T. Schultz  <http://orcid.org/0000-0003-1190-1122>  
 Giacomo Bernardi  <http://orcid.org/0000-0002-8249-4678>

### Data availability statement

The genome sequence data that support the findings of this study are openly available in GenBank of NCBI at <https://www.ncbi.nlm.nih.gov> under the accession no. ON556619. The associated BioProject, SRA, and Bio-Sample numbers are PRJNA828170, SRR20942050, and SAMN27642109-DTR\_Kuro\_0920G, respectively.

### References

- Allen GR. 1991. Damselfishes of the world. Mentor (OH): Aquarium Systems.
- Bernardi G, Holbrook SJ, Schmitt RJ. 2001. Gene flow at three spatial scales in a coral reef fish, the three-spot dascyllus, *Dascyllus trimaculatus*. *Mar Biol.* 138(3):457–465.
- Felsenstein J. 1985. Confidence limits on phylogenies: an approach using the bootstrap. *Evolution.* 39(4):783–791.
- Iwasaki W, Fukunaga T, Isagozawa R, Yamada K, Maeda Y, Satoh TP, Sado T, Mabuchi K, Takeshima H, Miya M, et al. 2013. Mitofish and mitoannotator: a mitochondrial genome database of fish with an accurate and automatic annotation pipeline. *Mol Biol Evol.* 30(11):2531–2540.
- Leray M, Beldade R, Holbrook SJ, Schmitt RJ, Planes S, Bernardi G. 2010. Allopatric divergence and speciation in coral reef fish: the three-spot dascyllus, *Dascyllus trimaculatus*, species complex. *Evolution.* 64(5): 1218–1230.
- Swofford DL. 2003. PAUP\*: phylogenetic analysis using parsimony (\*and other methods) version 4. Sunderland (MA): Sinauer Associates.
- Tang KL, Stiassny MLJ, Mayden RL, Desalle R. 2021. Systematics of damselfishes. *Ichthyol Herpetol.* 109(1):258–318.