

# Lawrence Berkeley National Laboratory

## Recent Work

### Title

MUONIC-DECAY BRANCHING RATIO OF THE LAMBDA HYPERON

### Permalink

<https://escholarship.org/uc/item/6zt6x65j>

### Authors

Kernan, Anne  
Powell, Wilson M.  
Sandier, Carl L.  
et al.

### Publication Date

1963-09-30

University of California  
Ernest O. Lawrence  
Radiation Laboratory

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy  
which may be borrowed for two weeks.  
For a personal retention copy, call  
Tech. Info. Division, Ext. 5545*

MUONIC-DECAY BRANCHING RATIO  
OF THE LAMBDA HYPERON

Berkeley, California

## **DISCLAIMER**

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UNIVERSITY OF CALIFORNIA  
Lawrence Radiation Laboratory  
Berkeley, California  
Contract No. W-7405-eng-48

MUONIC-DECAY BRANCHING RATIO OF THE LAMBDA HYPERON

Anne Kernan, Wilson M. Powell, Carl L. Sandler,  
William L. Knight and F. Russell Stannard

September 30, 1963

The Muonic-Decay Branching Ratio of the  
Lambda Hyperon\*

Anne Kernan, Wilson M. Powell, and Carl L. Sandler

Lawrence Radiation Laboratory  
University of California  
Berkeley, California

and

William L. Knight and F. Russell Stannard

University College London  
London, England

September 30, 1963

A total of 192 000 lambda decays in the charged mode has been examined for evidence of the decay  $\Lambda \rightarrow p\mu^- \bar{\nu}$ . Two possible examples of the process have been found in an effective sample of 19 700 lambdas. The corresponding upper limit for the branching ratio

$$R_{\mu} = \frac{\Gamma(\Lambda \rightarrow p\mu^- \bar{\nu})}{\Gamma(\Lambda \rightarrow p\pi^-) + \Gamma(\Lambda \rightarrow n\pi^0)}$$

is  $R_{\mu} \leq 1 \times 10^{-4}$ .

The experiment was performed with the Berkeley 30-in. heavy-liquid bubble chamber in a 13-kG magnetic field. The lambdas were produced by interactions in the chamber liquid of 0- to 440-MeV/c  $K^-$  mesons. The chamber was filled with a 76-24% mixture by weight of  $CF_3Br-C_3H_8$ ; a detailed account of the properties of this liquid is contained in reference 1.

The fact that 67% of  $\mu^-$ -mesons decay after coming to rest in the chamber liquid was utilized in identifying muonic lambda decays. An initial selection was made of lambda decays conforming to the following criteria:

- (a) The negative track terminated in the chamber with a  $\mu^- \rightarrow e^- \bar{\nu} \nu$  decay, easily recognized in this chamber liquid.
- (b) The proton ended in the chamber.
- (c) The lambda hyperon was produced in a  $K^-$  interaction in the chamber.
- (d) The lambda, proton, and meson tracks were at least 3 mm long.

A typical event satisfying the selection criteria is shown in Fig. 1.

Criterion (a) excludes all  $\Lambda \rightarrow \pi^-$  decays except those in which the pion decayed in flight to a muon, which in turn came to rest and decayed. From the measured pion momentum spectrum of 1000 randomly selected pionic lambda decays, the fraction of pions decaying in flight was calculated to be 0.0189. Thus, criterion (a) separated out a group of lambda decays in which the proportion of  $\Lambda_{\mu}$  events was enhanced by a factor of 53 over that in the original sample.

The remaining criteria ensure that the directions of the lambda, proton, and meson tracks can be measured, and that the proton momentum can be obtained from its range with a mean error of 3%. Multiple scattering in the chamber liquid precluded a reliable determination of the meson momentum by magnetic bending. Neither could the lambda momentum be inferred from the dynamics of the  $K^-$  interactions, since these took place predominantly in complex nuclei.

The 992 lambda decays that satisfied the selection criteria divided between  $\Lambda_{\pi\mu}$  and  $\Lambda_{\mu}$  decays as follows:

$$N(\Lambda_{\pi\mu}) = N_{\Lambda} F_{\pi\mu} S_{\pi\mu} R_{\pi} \quad (1)$$

$$N(\Lambda_{\mu}) = N_{\Lambda} F_{\mu} S_{\mu} R_{\mu} \quad (2)$$

Here  $N_{\Lambda}$  is the total number of charged and neutral pionic lambda decays

in the chamber liquid,  $F_{\pi\mu}$  and  $F_{\mu}$  are the fractions of  $\Lambda \rightarrow p\pi^-$  and  $\Lambda \rightarrow p\mu^-\bar{\nu}$  decays that satisfy conditions (a) through (d), and  $S_{\pi\mu}$  and  $S_{\mu}$  are the respective scanning efficiencies for these events. The branching ratios for charged, pionic lambda decay and for muonic lambda decay are  $R_{\pi}$  and  $R_{\mu}$ , respectively. We take 0.66 for the value of  $R_{\pi}$ .<sup>2</sup> Rearranging and combining Eqs. (1) and (2), we have

$$R_{\mu} = \frac{N(\Lambda_{\mu}) F_{\pi\mu} S_{\pi\mu} R_{\pi}}{N(\Lambda_{\pi\mu}) F_{\mu} S_{\mu}}$$

Two Monte Carlo calculations, outlined below, gave  $F_{\pi\mu}/F_{\mu} = 0.0182$ ; this estimate is independent of the ratio of muon capture to muon decay in the chamber liquid. The ratio  $S_{\pi\mu}/S_{\mu}$  was set equal to unity because the scanning criterion for both types of events was the presence of an electron at the end of the negative track from the lambda decay, and therefore the scanning efficiencies for  $\Lambda_{\pi\mu}$  and  $\Lambda_{\mu}$  decays were expected to be identical. Then we have

$$R_{\mu} = [N(\Lambda_{\mu})/N(\Lambda_{\pi\mu})] (0.0182 \times 0.66). \quad (3)$$

The quantity  $N(\Lambda_{\mu})$  was determined by looking for events in which the length of the negative track was inconsistent with the decay sequence  $\Lambda \rightarrow p\pi^-$ ,  $\pi^- \rightarrow \mu^-\bar{\nu}$  in flight. In this decay sequence the length of the negative-particle track,  $L_{\pi} + L_{\mu}$ , depends upon  $P_{\pi}$ , the magnitude of the pion momentum in lambda decay, and upon the dynamics of the pion decay. For each  $\Lambda_{\pi\mu}$  event,  $L_{\pi} + L_{\mu}$  is restricted by kinematics within an interval  $L_{\min}$  to  $L_{\max}$ , where  $L_{\min}$  ( $L_{\max}$ ) is the range of the muon when the pion decays at the lambda decay vertex, and the muon is emitted in the pion rest frame antiparallel (parallel) to the pion direction.

The kinematics of  $\Lambda \rightarrow p\pi^-$  decay are completely specified by two independent variables, the magnitude of the proton momentum and the angle between the proton and the pion. For all events it was assumed that  $\Lambda \rightarrow p\pi^-$  decay had occurred, and  $P_\pi \pm \Delta P_\pi$  was calculated from the measured values of the proton momentum and  $\psi$ , the angle between the charged decay products. The uncertainty in  $P_\pi$  arises predominantly from the experimental error on  $\psi$ . The limiting lengths of the negative-particle track,  $L_{\max} \pm \Delta L_{\max}$  and  $L_{\min} \pm \Delta L_{\min}$ , were computed from  $P_\pi \pm \Delta P_\pi$ . Whenever  $L$ , the measured length of the negative particle track (see the track connecting origins (b) and (c) in Fig. 1), lay outside the range  $L_{\min}$  to  $L_{\max}$ , the event was remeasured twice. Also, when  $L$ , although within the range  $L_{\min}$  to  $L_{\max}$ , came within two standard deviations of a limiting value, the event was remeasured.

Two events were found with  $L < L_{\min}$ , and therefore, the  $\Lambda_{\pi\mu}$  interpretation was ruled out for these decays. The muon momentum was calculated from the range of the negative track,  $L$ , for the  $\Lambda_\mu$  interpretation, and in both cases the momenta of the proton, muon and neutrino in the plane transverse to the lambda trajectory were consistent with the reaction  $\Lambda \rightarrow p\mu^- \bar{\nu}$ .

A Monte Carlo calculation shows that these two events constitute 24% of the  $\Lambda_\mu$  decays which satisfied the selection criteria. In the calculation, 3000  $\Lambda \rightarrow p\mu^- \bar{\nu}$  decays were generated in a random manner. The momenta of the lambdas and their decay points in the chamber were assigned according to the experimentally determined values of these quantities, obtained from the measurement of 1000 randomly selected pionic lambda decays. Since the form of the lambda decay amplitude is not known, the calculation was performed for both a Lorentz-invariant and a non-Lorentz-invariant



three-body phase space; the same result was obtained in both cases. In 61.7% of the simulated decays, the tracks of both the proton and the muon were at least 3 mm in length and ended in the bubble chamber. For each of these  $\Lambda_\mu$  events a corresponding  $P_\pi$  was calculated for a  $\Lambda \rightarrow p\pi$  decay, with the same proton momentum and the same opening angle between the charged decay products. The length of the muon track,  $L_\mu$ , was compared with the range of values  $L_{\min}$  to  $L_{\max}$ , calculated from  $P_\pi$ . In 24% of the events the length of the muon was outside the interval  $L_{\min}$  to  $L_{\max}$ , ( $L_\mu < L_{\min}$ , 20% :  $L_\mu > L_{\max}$ , 4%).

In a second Monte Carlo calculation, 3700 decay sequences of the type  $\Lambda \rightarrow p\pi^-$ ,  $\pi^- \rightarrow \mu^- \bar{\nu}$  in flight were generated in a random manner. In 59.4% of these decays the proton track and the combined length of the pion and muon tracks were at least 3 mm long and ended in the bubble chamber. The value of  $F_{\pi\mu}/F_\mu$  is therefore  $(0.594 \times 0.0189)/0.617 = 0.0182$ , where 0.0189 is the fraction of pions decaying in flight.

The distribution of  $\theta_{\pi\mu}$  (the projection in the film plane of the  $\pi-\mu$  decay angle) was also examined to determine if  $\Lambda_{\pi\mu}$  decays could be identified by observing the  $\pi-\mu$  decay angle. More than 30% of the  $\theta_{\pi\mu}$  values were less than 10 deg. Detection and measurement of angles of this magnitude was rendered difficult by multiple scattering in the heavy chamber liquid; consequently, no attempt was made to discriminate between  $\Lambda_\mu$  and  $\Lambda_{\pi\mu}$  decays by this means.

The two examples of  $\Lambda_\mu$  decay give an estimate of  $N(\Lambda_\mu) \leq 2/0.24 = 8$ , and  $N(\Lambda_{\pi\mu}) = (992 - 8) = 984$ . For the present, at least, the estimate of  $N(\Lambda_\mu)$  is considered an upper limit because the  $\Lambda_\mu$  events could have been simulated by processes other than  $\Lambda_\mu$  decay. Possible sources of background include:

A.  $\Lambda_{\pi\mu}$  decays in which the pion decayed close to the  $\Lambda$  decay vertex. In this case  $L_{\min}$  and  $L_{\max}$  are computed incorrectly because

the measured angle  $\psi$  is not the angle between the proton and  $\pi^-$  meson.

B.  $\Lambda_{\pi\mu}$  decays in which the pion or proton suffered an unobserved scatter close to the lambda vertex. As in A, the measured  $\psi$  is not the pion-proton angle.

C.  $\Lambda_{\pi\mu}$  decays in which the proton underwent a charge-exchange interaction. This causes the proton momentum, and hence  $L_{\min}$  and  $L_{\max}$  to be incorrectly estimated.

D. Neutron interactions,  $nn \rightarrow n\pi^-$ , with the subsequent decay in flight of the  $\pi^-$  meson.

E. Radiative lambda decays,  $\Lambda \rightarrow p\pi^- \gamma$ , also with subsequent decay in flight of the  $\pi^-$  meson.

The kinematics of the two  $\Lambda_{\mu}$  decay candidates were such as to rule out the possibility that A or C had occurred.

On the basis of two possible cases of  $\Lambda_{\mu}$  decay,  $R_{\mu}$  calculated according to Eq. (3) is

$$R_{\mu} = \left[ \left( \frac{2}{0.24} \right) / (984) \right] (0.0182 \times 0.66) = \frac{2}{19\,700} = 1 \times 10^{-4}.$$

The effective sample size of lambda decays in this experiment is seen to be 19 700.

This experiment gives an upper limit of  $R_{\mu} \leq 4.5 \times 10^{-4}$  at the 5% significance level. A lower limit can be set to the branching ratio from the observation of Good and Lind of one unambiguous case of  $\Lambda_{\mu}$  decay in a total of 2500 lambdas,<sup>3</sup> giving  $R_{\mu} \geq 0.2 \times 10^{-4}$  at the 5% significance level.

The combined results define a 90% confidence interval for  $R_{\mu}$  :

$$0.2 \times 10^{-4} \leq R_{\mu} \leq 4.5 \times 10^{-4} .$$

This estimate is consistent with  $R_{\mu} = (1.3 \pm 0.2) \times 10^{-4}$  deduced from the measured  $\beta$ -decay rate of the lambda hyperon,<sup>1</sup> if we assume that the decay processes  $\Lambda \rightarrow pe^{-}\bar{\nu}$  and  $\Lambda \rightarrow p\mu^{-}\bar{\nu}$  are identical except for the  $\mu^{-} - e^{-}$  mass difference. A theoretical value<sup>4</sup> of  $R_{\mu} = 2.4 \times 10^{-3}$  has been predicted on the basis of the universal V-A weak-interaction theory<sup>5,6</sup> and the hypothesis that the renormalization of the coupling constant due to strong interactions can be ignored in leptonic lambda decays. The measured value of  $R_{\mu}$  clearly disagrees with the calculated one; it has already been shown that the  $\Lambda_{\beta}$  decay rate is similarly depressed relative to the predicted value.<sup>1</sup>

One of us, William L. Knight, gratefully acknowledges a grant from the Department of Scientific and Industrial Research, England.

FOOTNOTES AND REFERENCES

\* Work done under the auspices of the U. S. Atomic Energy Commission.

1. R. P. Ely, G. Gidal, G. E. Kalmus, L. O. Oswald, W. M. Powell, W. J. Singleton, F. W. Bullock, C. Henderson, D. J. Miller, and F. R. Stannard, *Phys. Rev.* 131, 868 (1963).
2. F. S. Crawford, in Proceedings of the 1962 International Conference on High Energy Physics (CERN, Geneva, 1962), p. 832.
3. M. L. Good and V. G. Lind, *Phys. Rev. Letters* 9, 518 (1962).
4. L. Okun, *Ann. Rev. Nucl. Sci.* 9, 82 (1959).
5. R. P. Feynman and M. Gell-Mann, *Phys. Rev.* 109, 193 (1958).
6. E. C. G. Sudarshan and R. E. Marshak, *Phys. Rev.* 109, 1860 (1958).

## FIGURE LEGEND

Fig. 1. A typical event satisfying the selection criteria.

(a)  $K^- + \text{nucleus} \rightarrow \Lambda + \text{fragments}$ ,

(b)  $\Lambda \rightarrow p\pi^-$  or  $\Lambda \rightarrow p\mu^-\bar{\nu}$ ,

(c)  $\mu^- \rightarrow e^-\nu\bar{\nu}$ .




Fig. 1.

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

Faint, illegible text covering the entire page, likely bleed-through from the reverse side of the document.

