

# UC Santa Barbara

## UC Santa Barbara Previously Published Works

### Title

Headwater Stream Microbial Diversity and Function across Agricultural and Urban Land Use Gradients.

### Permalink

<https://escholarship.org/uc/item/7129s59w>

### Journal

Applied and Environmental Microbiology, 86(11)

### ISSN

0099-2240

### Authors

Laperriere, Sarah M  
Hilderbrand, Robert H  
Keller, Stephen R  
[et al.](#)

### Publication Date

2020-05-19

### DOI

10.1128/aem.00018-20

Peer reviewed



# Headwater Stream Microbial Diversity and Function across Agricultural and Urban Land Use Gradients

Sarah M. Laperriere,<sup>a,b\*</sup> Robert H. Hilderbrand,<sup>c</sup> Stephen R. Keller,<sup>c,d</sup> Regina Trott,<sup>c</sup>  Alyson E. Santoro<sup>b</sup>

<sup>a</sup>Horn Point Laboratory, University of Maryland Center for Environmental Science, Cambridge, Maryland, USA

<sup>b</sup>Department of Ecology, Evolution, and Marine Biology, University of California, Santa Barbara, California, USA

<sup>c</sup>Appalachian Laboratory, University of Maryland Center for Environmental Science, Frostburg, Maryland, USA

<sup>d</sup>Department of Plant Biology, University of Vermont, Burlington, Vermont, USA

**ABSTRACT** Anthropogenic activity impacts stream ecosystems, resulting in a loss of diversity and ecosystem function; however, little is known about the response of aquatic microbial communities to changes in land use. Here, microbial communities were characterized in 82 headwater streams across a gradient of urban and agricultural land uses using 16S rRNA gene amplicon sequencing and compared to a rich data set of physicochemical variables and traditional benthic invertebrate indicators. Microbial diversity and community structures differed among watersheds with high agricultural, urban, and forested land uses, and community structure differed in streams classified as being in good, fair, poor, and very poor condition using benthic invertebrate indicators. Microbial community similarity decayed with geodesic distance across the study region but not with environmental distance. Stream community respiration rates ranged from 21.7 to 1,570 mg O<sub>2</sub> m<sup>-2</sup> day<sup>-1</sup> and 31.9 to 3,670 mg O<sub>2</sub> m<sup>-2</sup> day<sup>-1</sup> for water column and sediments, respectively, and correlated with nutrients associated with anthropogenic influence and microbial community structure. Nitrous oxide (N<sub>2</sub>O) concentrations ranged from 0.22 to 4.41 μg N<sub>2</sub>O liter<sup>-1</sup>; N<sub>2</sub>O concentration was negatively correlated with forested land use and was positively correlated with dissolved inorganic nitrogen concentrations. Our findings suggest that stream microbial communities are impacted by watershed land use and can potentially be used to assess ecosystem health.

**IMPORTANCE** Stream ecosystems are frequently impacted by changes in watershed land use, resulting in altered hydrology, increased pollutant and nutrient loads, and habitat degradation. Macroinvertebrates and fish are strongly affected by changes in stream conditions and are commonly used in biotic indices to assess ecosystem health. Similarly, microbes respond to environmental stressors, and changes in community composition alter key ecosystem processes. The response of microbes to habitat degradation and their role in global biogeochemical cycles provide an opportunity to use microbes as a monitoring tool. Here, we identify stream microbes that respond to watershed urbanization and agricultural development and demonstrate that microbial diversity and community structure can be used to assess stream conditions and ecosystem functioning.

**KEYWORDS** Benthic Index of Biotic Integrity, species-area curves, nitrous oxide, respiration, aquatic ecosystems, Chesapeake Bay, species-area relationships

**B**iodiversity is critical to ecosystem functioning and is threatened by anthropogenic activity (1). Microbes perform key ecosystem functions (2–4), necessitating a better understanding of how microbes and microbial diversity respond to environmental stressors (5–7). Streams are examples of threatened ecosystems, where watershed modification decreases stream integrity and water quality (8–11), altering macroinver-

**Citation** Laperriere SM, Hilderbrand RH, Keller SR, Trott R, Santoro AE. 2020. Headwater stream microbial diversity and function across agricultural and urban land use gradients. *Appl Environ Microbiol* 86:e00018-20. <https://doi.org/10.1128/AEM.00018-20>.

**Editor** Shuang-Jiang Liu, Chinese Academy of Sciences

**Copyright** © 2020 American Society for Microbiology. All Rights Reserved.

Address correspondence to Alyson E. Santoro, [asantoro@ucsb.edu](mailto:asantoro@ucsb.edu).

\* Present address: Sarah M. Laperriere, Department of Biological Sciences, University of Southern California, Los Angeles, CA, USA.

**Received** 13 January 2020

**Accepted** 26 March 2020

**Accepted manuscript posted online** 3 April 2020

**Published** 19 May 2020

tebrate, fish, and microbial diversity (11–15). The use of macroinvertebrate and fish indices to assess stream conditions is fundamental to stream ecology (16) and depends on known relationships between stream integrity and community structure (17, 18). The Benthic Index of Biotic Integrity (B-IBI) is one such index, using the abundance and diversity of stream benthic macroinvertebrates to accurately distinguish degraded streams (B-IBI < 2.9) classified based on stream chemical and physical criteria (18). Biotic indices are calibrated to specific regions, as the distribution of stream macroinvertebrates is controlled by a combination of dispersal limitation and local environmental conditions (19). Despite their abundance and ecological importance, natural microbial communities, unlike macroinvertebrates, are not used in stream monitoring programs to assess stream conditions.

As with macroinvertebrates, dispersion and environmental selection control the spatial distribution of microbes along stream continuums (20). Dispersion, or the advection of microbes from the surrounding landscape, impacts headwater stream community composition, and with increasing stream order, environmental sorting becomes more important as stream residence times increase (20). Several studies have demonstrated the influence of the surrounding landscape on stream microbes, showing that watershed urbanization leads to shifts in bacterial communities (21–24). While alpha diversity generally remains constant (22–24), the abundances of taxa associated with anthropogenic activity and high-nutrient conditions increase in urbanized streams (22, 24). Similar to larger organisms, microbes respond to environmental disturbance and are strongly influenced by watershed land use (25, 26); therefore, their distribution may be used to further characterize stream conditions.

Microbes mediate important stream ecosystem functions, controlling the movement of carbon and nitrogen through freshwater ecosystems (2–4). Previous studies demonstrate the effects of urbanization on stream nutrient transformations, such as nitrogen uptake (27), nitrogen retention (28), and carbon processing (29, 30). Community respiration determines the fate of terrestrial carbon in headwater streams, where carbon is either lost as carbon dioxide during respiration or transported farther downstream (31). Community respiration is often used to assess ecosystem function (32), as rates are influenced by watershed land use (33–35), correlated with stream chemistry (30, 36), and sensitive to pollutants (37, 38). The effects of urbanization on stream dissolved organic matter quality (39) and respiration (33–35) have previously been demonstrated, and stream microbial community structure can potentially be used to monitor these ecosystem functions.

In addition to respiration, dissolved organic matter fuels stream denitrification and the microbial reduction of nitrate ( $\text{NO}_3^-$ ) to nitrous oxide ( $\text{N}_2\text{O}$ ) and dinitrogen ( $\text{N}_2$ ) gases (40). Denitrification removes nitrogen from streams and is credited as the major source of the greenhouse gas  $\text{N}_2\text{O}$  (41, 42). Watershed land use and anthropogenic nitrogen loading alter rates of stream denitrification (43), increasing the amount of nitrogen transported downstream (44) and emissions of  $\text{N}_2\text{O}$  to the atmosphere (41). Urbanization has been linked to changes in denitrifier community composition (21, 22, 45, 46), and a previous study linked changes in denitrifier composition to changes in denitrification potential, and therefore nitrogen loss, in urban streams (21). However, it is less clear how changes in microbial community composition in response to land use modification alter  $\text{N}_2\text{O}$  production.

The goal of this study was to identify stream microbes that respond to watershed urbanization and agricultural development. These anthropogenic factors alter microbial diversity and community structure, which can be used to assess stream conditions and ecosystem functioning. We measured microbial diversity using 16S rRNA gene amplicon sequencing across 82 headwater streams within the Chesapeake Bay watershed in the state of Maryland in the spring and summer for 2 years. Measurements were collected in conjunction with stream physicochemical parameters and a macroinvertebrate indicator of stream health. Additionally, at a subset of streams, water column and sediment community respiration were measured using oxygen consumption methods, and  $\text{N}_2\text{O}$  concentrations were measured using gas chromatography. We deter-

mined how stream bacteria and archaea are distributed across gradients of watershed land use and stream conditions, assessed how changes in microbial community composition relate to benthic macroinvertebrate diversity and traditional indices of stream conditions, and determined how these changes influence stream function by relating microbial community composition to rates of microbial respiration and concentrations of  $N_2O$ .


## RESULTS

**Higher microbial alpha diversity in spring.** Operational taxonomic unit (OTU) richness differed according to substrate (water and sediment) and ranged from 115 to 996 and 459 to 895 for water and sediment samples, respectively, with sediment richness being significantly higher than that of water (paired Wilcoxon,  $P < 0.001$ ). Sediment Shannon diversity, at  $6.2 \pm 0.2$  (mean  $\pm$  standard deviation), was greater than water diversity, at  $5.4 \pm 1.1$  (paired Wilcoxon,  $P < 0.001$ ), and sediment communities were more even, at  $0.94 \pm 0.02$ , than were water communities, at  $0.84 \pm 0.11$  (paired Wilcoxon,  $P < 0.001$ ). Similarly, beta diversity (Bray-Curtis dissimilarity) differed by substrate (permutational multivariate analysis of variance [PERMANOVA],  $R^2 = 0.11$ ,  $P < 0.001$ ). Water communities had more *Actinobacteria*, *Bacteroidetes*, and *Proteobacteria*, while *Acidobacteria*, *Planctomycetes*, *Chloroflexi*, *Nitrospirae*, and *Verrucomicrobia* were more abundant in sediments (see Fig. S1 in the supplemental material).

Interannually, there was no difference in alpha diversity between samples collected in 2014 and 2015, and collection year only explained a small fraction of the variance in beta diversity between water and sediment communities (PERMANOVA,  $R^2 = 0.01$ ,  $P < 0.001$ , and  $R^2 = 0.01$ ,  $P < 0.001$ , respectively). Seasonally, water Shannon diversity was greater in spring than in summer (paired Wilcoxon,  $P < 0.001$ ), a result of higher richness (paired Wilcoxon,  $P < 0.001$ ) and evenness (paired Wilcoxon,  $P < 0.001$ ) in spring. There was no seasonal difference in sediment Shannon diversity (paired Wilcoxon,  $P = 0.49$ ), though sediment richness was greater in spring than in summer (paired Wilcoxon,  $P = 0.01$ ), with no significant difference in evenness (paired Wilcoxon,  $P = 0.13$ ). Additionally, there were small seasonal changes in water and sediment community structures (PERMANOVA,  $R^2 = 0.05$ ,  $P < 0.001$ , and  $R^2 = 0.02$ ,  $P < 0.001$ , respectively), with *Actinobacteria* being more abundant in summer water samples than in spring samples (Fig. S2).

**Distance-decay relationships partially drive microbial diversity.** Water sample alpha diversity metrics differed across the three geographic regions, while sediment diversity remained constant (Fig. 1). Streams on the Coastal Plain had lower Shannon diversity than did streams in the Piedmont and Highlands regions (Dunn's Kruskal-Wallis,  $P < 0.001$  and  $P < 0.001$ , respectively). This was driven by both lower evenness (Dunn's Kruskal-Wallis,  $P < 0.001$  and  $P < 0.001$  in Piedmont and Highlands, respectively) and richness (Dunn's Kruskal-Wallis,  $P < 0.001$  and  $P < 0.001$  in Piedmont and Highlands, respectively) in Coastal Plain streams than that in the other regions. Regional differences in Bray-Curtis dissimilarity and taxon abundances were observed in water (PERMANOVA,  $R^2 = 0.07$ ,  $P < 0.001$ ; Fig. 2a and S3a) and sediment (PERMANOVA,  $R^2 = 0.08$ ,  $P < 0.001$ ; Fig. 2b and S3b) communities.


Partial Mantel tests detected correlations between water and sediment Bray-Curtis dissimilarity and geographic distance (the Euclidean distance between sampling locations) ( $\rho = 0.26$ ,  $P = 0.001$ ; and  $\rho = 0.26$ ,  $P = 0.001$ , respectively), and no significant relationship was detected between Bray-Curtis dissimilarities and environmental distance and the Euclidean distance between streams based on the continuous environmental variables ( $\rho = -0.01$ ,  $P = 0.62$ ; and  $\rho = 0.07$ ,  $P = 0.07$ , respectively). Positive distance-decay relationships were observed between the natural-logarithm-transformed least-squares linear regressions of water and sediment community similarity (Sørensen index) and geodesic distance. The absolute value of the regression coefficients (species-area Z-values) for water and sediment communities were 0.12 ( $R^2 = 0.082$ ,  $P < 0.001$ ) and 0.11 ( $R^2 = 0.066$ ,  $P < 0.001$ ) (Fig. 3) for water and sediment communities, respectively.



**FIG 1** (a to f) Alpha-diversity metrics for water column (a, c, and e) and sediment (b, d, and f) samples in spring (white) and summer (gray) when grouped by geographic region. \*,  $P < 0.05$ ; \*\*,  $P < 0.01$ ; \*\*\*,  $P < 0.001$ , Dunn's test, with seasonal differences shown below and regional differences shown above.

The effect of land use on spatial scaling was examined by comparing the distance-decay relationships for streams in high-use urban, agricultural, and forested watersheds. Communities in streams with high urban and agricultural land use had significant distance-decay relationships in both water (urban,  $R^2 = 0.102$ ,  $P < 0.001$ ; agricultural,  $R^2 = 0.100$ ,  $P < 0.001$ ) and sediment (urban,  $R^2 = 0.129$ ,  $P < 0.001$ ; agricultural,  $R^2 = 0.075$ ,  $P < 0.001$ ) samples. No significant distance-decay relationship was observed for samples in highly forested watersheds. The species-area Z-values for highly urban (water, 0.08; sediment, 0.10) and agricultural (water, 0.07; sediment, 0.06) streams were lower than the Z-values when all streams were considered.


**Microbial diversity relates to stream physicochemistry.** Stream physicochemistry varied by geographic region (Table S2), land use (Table S3), and stream conditions (Table S4).  $N_2O$  concentrations ranged from  $0.22 \pm 0.00$  to  $4.41 \pm 0.7 \mu\text{g } N_2O \text{ liter}^{-1}$  (58 to 1,217% saturated; Table 1), with no difference in  $N_2O$  concentrations or saturation between individual streams sampled in 2014 and 2015 (paired  $t$  test,  $P = 0.2$ ).  $N_2O$  concentration negatively correlated with percent forest cover and positively correlated with total nitrogen (TN),  $NO_2^-$ , and Br,  $NO_3^-$ , and  $NH_4^+$  concentrations, and agricultural cover (Table S5).  $N_2O$  concentration did not correlate with water or sediment microbial diversity.



**FIG 2** (a and b) Nonmetric multidimensional scaling (NMDS) plots of Bray-Curtis distances for all water column (a) (stress, 0.18; PERMANOVA, region,  $R^2 = 0.07$ ,  $P < 0.001$ ; season,  $R^2 = 0.06$ ,  $P < 0.001$ ) and sediment (b) (stress, 0.13; PERMANOVA, region,  $R^2 = 0.08$ ,  $P < 0.001$ ; season,  $R^2 = 0.09$ ,  $P < 0.001$ ) samples; symbol color indicates the geographic region, and symbol shape denotes the season.

Respiration rates in Coastal Plain streams ranged from 21.7 to 1,573 mg O<sub>2</sub> m<sup>-2</sup> day<sup>-1</sup> and 31.9 and 3,668 mg O<sub>2</sub> m<sup>-2</sup> day<sup>-1</sup> for water and sediments, respectively (Table 2). There was no significant difference in water rates measured in streams in both 2014 and 2015 (paired *t* test,  $P = 0.9$ ), while sediment rates were higher in 2014 than those in 2015 (paired *t* test,  $P = 0.04$ ). Water respiration rates most strongly negatively correlated with specific conductance, Cl<sup>-</sup>, Ca, and urban cover and positively correlated with forest cover, embeddedness, and Shannon diversity (Table S6). Sediment respiration rates negatively correlated with Zn, total phosphorus (TP), and PO<sub>4</sub><sup>3-</sup> concentrations (Table S7). Both water and sediment microbial community structures correlated with respiration rates (Mantel,  $r = 0.51$ ,  $P = 0.001$ ; and  $r = 0.24$ ,  $P = 0.04$ , respectively).

Water Shannon diversity negatively correlated with several environmental variables, including embeddedness, Cu, thalweg depth, dissolved organic carbon (DOC), and TP and positively correlated with forest cover and stream velocity (Table S8). The best-fit stepwise multiple linear regression model explained 41% of the variance in water Shannon diversity and included pH, DOC, SO<sub>4</sub><sup>2-</sup>, TP, forest cover, Mg, Cu, and thalweg depth as predictor variables. Sediment diversity negatively correlated with DOC, embeddedness, and forest cover, and positively correlated with pH, B-IBI, NO<sub>3</sub><sup>-</sup>, and TN (Table S9). A stepwise linear regression model explained 22% of the variation in sediment Shannon diversity, with pH, DOC concentration, forest cover, Mg concentration, and thalweg depth being the most significant predictors. Similarly, Bray-Curtis



**FIG 3** Relationship between microbial community composition (Sørensen index) and geodesic distance for water column (a) and sediment (b) samples. The model fitting is linear least-squares regression.

**TABLE 1** Nitrous oxide concentrations and percent saturation relative to equilibrium at *in situ* temperature


Stream site	Concn and saturation data by yr			
	2014		2015	
	N <sub>2</sub> O concn (mean ± SD) (μg liter <sup>-1</sup> )	% saturation	N <sub>2</sub> O concn (mean ± SD) (μg liter <sup>-1</sup> )	% saturation
CORS102	1.16	290	1.06 ± 0.03	302
LMON302	0.87 ± 0.01	224		
LOCR102	1.68 ± 0.32	537	1.16 ± 0.04	353
MATT104	0.55 ± 0.01	140		
MATT115	1.17 ± 0.03	286		
MATT320	0.55 ± 0.00	152		
NASS108	0.22 ± 0.00	58	0.48 ± 0.04	120
NASS302	1.33 ± 0.01	394	0.74 ± 0.01	197
PAXL294	0.55 ± 0.01	142	0.65 ± 0.03	168
SEAS109	1.1 ± 0.03	263		
SEAS111	1.41 ± 0.01	385		
UMON134	0.55 ± 0.04	139		
UMON299	2.88 ± 0.10	711		
UPCK102	1.65 ± 0.01	398	1.11 ± 0.05	287
UPCK113	1.28 ± 0.00	316		
UPCR208S			0.76 ± 0.03	179
WIRH215	4.26 ± 0.15	1,217		
WIRH220	4.41 ± 0.07	1,116	4.13 ± 0.12	1,040

dissimilarity correlated with several environmental variables (Tables S10 and S11), including pH, DOC concentration, embeddedness, and Zn concentration. The measured physicochemical variables explained 7% of the variation in community structure according to constrained correspondence analysis in both stream water and sediment communities.

Strong associations between taxon abundance and stream physicochemistry were observed in water and sediment samples (Fig. 4). *Pedospaerales* positively correlated with forest cover and negatively correlated with acid-neutralizing capacity (ANC) and urban cover (Fig. 4a). The order iii1-15 (*Acidobacteria*) strongly negatively correlated with TP and Zn concentrations. *Cenarchaeales* (*Thaumarchaeota*) negatively correlated with DOC and TP concentrations. The strongest sediment associations were positive correlations between *Pirellulales* (*Planctomycetes*) and ANC, Ca concentration, and conductivity, and the strongest negative associations were between *Acidobacteria-*

**TABLE 2** Rates of water column and sediment respiration from streams on the Coastal Plain

Stream site	Respiration rates by yr (mean ± SD) (mg O <sub>2</sub> m <sup>-2</sup> day <sup>-1</sup> )			
	2014		2015	
	Water	Sediment	Water	Sediment
CORS102	148.8	1,563.7	181.9 ± 51.7	204.7 ± 68.8
LMON302	37.5	788.2		
LOCR102	246.0	3,220.3	267.3 ± 79.6	31.9 ± 9.9
LOWI104	221.7	2,441.1		
MATT104	95.0	128.7		
MATT115	52.6	394.2		
MATT320	23.8	1,897.1		
NASS108	348.5	716.5		
NASS302	1,573.2	3,667.9	248.2 ± 31.8	96.3 ± 19.6
PAXL294	45.3	235.7	264.2 ± 49.9	498.2 ± 123.4
SEAS109	21.7	264.2		
SEAS111	35.9			
UMON134	452.6	1,741.3		
UPCK102	314.3			
UPCK113	112.3		77.2 ± 8.0	98.2 ± 18.8
UPCK208	79.0	889.1	67.3 ± 12.5	97.7 ± 18.8
WIRH215	56.5	1,124.4		
WIRH220	77.0	1,593.4	1,029.3 ± 633.9	357.6 ± 233.4




**FIG 4** Relationship between the most abundant water column (a) and sediment (b) taxa at the order rank level and environmental variables. Color indicates the associated Z-score. Correlations with a |Z-score| of <1.96 are not shown, and + symbols denote |Z-scores| of >3. Max, maximum.

les and Mg and TN concentrations and between *Pedosphaerales* with  $\text{NO}_2^-$  and Mg concentrations (Fig. 4b).

**Microbial communities vary according to watershed land use and stream conditions.** Stream water alpha and beta diversity differed in watersheds with high agricultural, urban, and forested land use. Sediment community structure also differed according to land use, with no change in alpha diversity. Forested streams had higher water Shannon diversity, richness, and evenness than that of agricultural (Kruskal-Wallis,  $P < 0.001$ ) and urban (Kruskal-Wallis,  $P < 0.001$ ) streams (Fig. 5). Similarly, community structure differed in both water (PERMANOVA,  $R^2 = 0.11$ ,  $P < 0.001$ ) and sediment (PERMANOVA,  $R^2 = 0.07$ ,  $P < 0.001$ ) communities. Taxa were identified that were more abundant and pervasive in streams in watersheds with high forested, agricultural, and urban land use (Tables 3 and S12). Forested streams had more taxa in the phyla *Verrucomicrobia*, *Planctomycetes*, and *Acidobacteria*. Agricul-





**FIG 5** (a to f) Alpha diversity metrics for water column (a, c, and e) and sediment (b, d, and f) samples in spring (white) and summer (gray) from streams in watersheds with highly forested (>90%), agricultural (>50%), and urban (>50%) land use. \*,  $P < 0.05$ ; \*\*,  $P < 0.01$ ; \*\*\*,  $P < 0.001$ , Dunn's test, with seasonal differences shown below and land use differences shown above.

tural streams had more taxa in the phyla *Bacteroidetes*, *Firmicutes*, and *Chloroflexi*, while urban streams had several abundant taxa in the *Proteobacteria*, *Firmicutes*, and *Chloroflexi* phyla.

Benthic macroinvertebrate Shannon diversity weakly correlated with water and sediment microbial Shannon diversity ( $\rho = 0.17$ ,  $P = 0.008$ ; and  $\rho = 0.15$ ,  $P = 0.02$ , respectively), and B-IBI scores weakly correlated with sediment microbial Shannon diversity (Spearman,  $\rho = 0.15$ ,  $P = 0.03$ ). Additionally, benthic macroinvertebrate Bray-Curtis dissimilarities correlated with sediment and water microbial Bray-Curtis dissimilarities (Mantel,  $\rho = 0.37$ ,  $P = 0.001$ ; and  $\rho = 0.35$ ,  $P = 0.001$ , respectively), and B-IBI scores weakly correlated with water (Mantel,  $\rho = 0.07$ ,  $P = 0.02$ ) and sediment microbial community structure (Mantel,  $\rho = 0.13$ ,  $P = 0.002$ ). Microbial community structure was only slightly different in streams classified as in good, fair, poor, and very poor condition using the B-IBI (PERMANOVA, water,  $R^2 = 0.02$ ,  $P = 0.006$ ; sediment,  $R^2 = 0.03$ ,  $P < 0.001$ ). *Hydrogenophaga* spp. (*Burkholderiales*), unclassified *Deltaproteobacteria* in the order *Desulfobacterales*, and heteroC45 (*Chthoniobacterales*), a *Verrucomicrobia* bacterium, were all more abundant and pervasive in streams in very poor condition than in streams in good and fair condition (Table 4).

**TABLE 3** Microbial OTUs indicative of streams in highly forested, agricultural, and urban watersheds<sup>a</sup>

Group	Substrate	Taxonomy (domain/phylum/class/order/family/genus)	Indicator <i>r</i>	<i>P</i> value	A <sup>b</sup>	B <sup>c</sup>	
Agriculture	Water	<i>Bacteria/Bacteroidetes/Flavobacteriia/Flavobacteriales/Cryomorphaeaceae/Fluviicola</i>	0.77	0.03	0.61	0.95	
	Water	<i>Bacteria/Cyanobacteria/4C0d-2/YS2/unclassified/unclassified</i>	0.70	0.006	0.54	0.90	
	Water	<i>Bacteria/Proteobacteria/Epsilonproteobacteria/Campylobacteriales/Campylobacteraceae/Sulfurospirillum</i>	0.63	0.021	0.84	0.48	
	Water	<i>Bacteria/OP3/BD4-9/unclassified/unclassified/unclassified</i>	0.60	0.028	0.58	0.62	
	Water	<i>Bacteria/Acidobacteria/BPC102/MVS-40/unclassified/unclassified</i>	0.60	0.017	0.57	0.62	
	Sediment	<i>Bacteria/Firmicutes/Bacilli/Bacillales/Bacillaceae/Bacillus</i>	0.80	0.001	0.68	0.95	
	Sediment	<i>Bacteria/Chloroflexi/Anaerolineae/GCA004/unclassified/unclassified</i>	0.78	0.002	0.62	1.00	
	Sediment	<i>Bacteria/Proteobacteria/Gammaproteobacteria/Methylococcales/Crenotrichaceae/Crenothrix</i>	0.75	0.002	0.59	0.95	
	Sediment	<i>Bacteria/Verrucomicrobia/Verrucomicrobiae/Verrucomicrobiales/Verrucomicrobiaceae/Prostheco bacter</i>	0.74	0.003	0.67	0.81	
	Sediment	<i>Bacteria/Verrucomicrobia/Verrucomicrobiae/Verrucomicrobiales/Verrucomicrobiaceae/Luteolibacter</i>	0.73	0.010	0.57	0.95	
	Forest	Water	<i>Bacteria/Verrucomicrobia/Spartobacteria/Chthoniobacteriales/Chthoniobacteraceae/DA101</i>	0.91	0.001	0.83	1.00
		Water	<i>Bacteria/Planctomycetes/Planctomycetia/Gemmatales/Gemmataceae/unclassified</i>	0.84	0.001	0.71	1.00
		Water	<i>Bacteria/Planctomycetes/Phycisphaerae/WD2101/unclassified/unclassified</i>	0.84	0.001	0.70	1.00
Water		<i>Bacteria/Elusimicrobia/Elusimicrobia/FAC88/unclassified/unclassified</i>	0.84	0.001	0.76	0.92	
Water		<i>Bacteria/Verrucomicrobia/Pedosphaerae/Pedosphaerales/auto67_4W/unclassified</i>	0.84	0.001	0.70	1.00	
Sediment		<i>Bacteria/Proteobacteria/Alphaproteobacteria/Rhizobiales/Hyphomicrobiaceae/unclassified</i>	0.80	0.001	0.67	0.96	
Sediment		<i>Bacteria/Acidobacteria/Chloracidobacteria/PK29/unclassified/unclassified</i>	0.77	0.001	0.63	0.92	
Sediment		<i>Bacteria/Verrucomicrobia/Spartobacteria/Chthoniobacteriales/Chthoniobacteraceae/DA101</i>	0.73	0.001	0.53	1.00	
Sediment		<i>Bacteria/Planctomycetes/Planctomycetia/Gemmatales/Gemmataceae/unclassified</i>	0.72	0.001	0.52	1.00	
Sediment		<i>Bacteria/Acidobacteria/Acidobacteriia/Acidobacteriales/Koribacteraceae/unclassified</i>	0.71	0.049	0.59	0.85	
Urban	Water	<i>Bacteria/Proteobacteria/Betaproteobacteria/Burkholderiales/Comamonadaceae/Hydrogenophaga</i>	0.83	0.001	0.86	0.81	
	Water	<i>Bacteria/Proteobacteria/Alphaproteobacteria/Sphingomonadales/Sphingomonadaceae/Novosphingobium</i>	0.78	0.001	0.61	1.00	
	Water	<i>Bacteria/Proteobacteria/Betaproteobacteria/Burkholderiales/Alcaligenaceae/unclassified</i>	0.78	0.001	0.81	0.75	
	Water	<i>Bacteria/Proteobacteria/Betaproteobacteria/Burkholderiales/Comamonadaceae/Rhodofera x</i>	0.74	0.021	0.55	1.00	
	Water	<i>Bacteria/Bacteroidetes/Saprosirae/Saprosirales/Chitinophagaceae/Sediminibacterium</i>	0.72	0.010	0.56	0.94	
	Sediment	<i>Bacteria/Proteobacteria/Deltaproteobacteria/Desulfuromonadales/Geobacteraceae/Geobacter</i>	0.74	0.004	0.55	1.00	
	Sediment	<i>Bacteria/Firmicutes/Clostridia/OPB54/unclassified/unclassified</i>	0.62	0.017	0.69	0.56	
	Sediment	<i>Bacteria/Chloroflexi/Anaerolineae/CFB-26/unclassified/unclassified</i>	0.61	0.037	0.53	0.69	
	Sediment	<i>Bacteria/Proteobacteria/Deltaproteobacteria/Syntrophobacteriales/Syntrophaceae/Desulfobacca</i>	0.57	0.034	0.58	0.56	
	Sediment	<i>Bacteria/Firmicutes/Clostridia/Clostridiales/unclassified/unclassified</i>	0.56	0.013	0.63	0.50	

<sup>a</sup>High defined as >90% for forest, >50% for agricultural, and >50% for urban watersheds.

<sup>b</sup>A, mean relative abundance of the OTU in each group compared to all groups.

<sup>c</sup>B, relative frequency of each OTU belonging to each group.

**DISCUSSION**

The aims of this study were to understand how stream bacteria and archaea are distributed across gradients of watershed land use and water quality, to assess how changes in microbial community composition relate to benthic macroinvertebrate diversity, and to discern how changes in stream conditions alter stream ecosystem processes, as reflected in community respiration and N<sub>2</sub>O concentrations. Bacterial and archaeal diversity significantly differed across the geographic regions of Maryland (Fig. 1 and 2), demonstrating the influence of the surrounding landscape on headwater stream microbial communities. Regional alluvium composition likely influenced stream alpha diversity, causing lower alpha diversity in Coastal Plain streams (Fig. 1). Sediments on the Coastal Plain of the eastern United States are composed of gravel, sand, silt, and clay (47), making streams more embedded (Table S2). Embeddedness was the envi-

**TABLE 4** Microbial OTUs indicative of stream conditions according to the B-IBI

Group <sup>a</sup>	Substrate	Taxonomy (domain/phylum/class/order/family/genus)	Indicator <i>r</i>	<i>P</i> value	A <sup>b</sup>	B <sup>c</sup>
Good	Sediment	<i>Archaea/unclassified/unclassified/unclassified/unclassified/unclassified</i>	0.59	0.04	0.58	0.60
Poor	Water	<i>Bacteria/Proteobacteria/Gammaproteobacteria/Methylococcales/Crenotrichaceae/Crenothrix</i>	0.71	0.03	0.58	0.86
	Water	<i>Bacteria/Proteobacteria/Betaproteobacteria/Gallionellales/Gallionellaceae/Gallionella</i>	0.53	0.04	0.53	0.54
Very poor	Water	<i>Bacteria/Proteobacteria/Betaproteobacteria/Burkholderiales/Comamonadaceae/Hydrogenophaga</i>	0.62	0.01	0.72	0.54
	Water	<i>Bacteria/Proteobacteria/Deltaproteobacteria/Desulfobacteriales/Desulfobulbaceae/unclassified</i>	0.60	0.00	0.53	0.69
	Water	<i>Bacteria/Verrucomicrobia/Spartobacteria/Chthoniobacteriales/Chthoniobacteraceae/heteroC45_4W</i>	0.59	0.00	0.56	0.62
	Water	<i>Bacteria/Proteobacteria/Alphaproteobacteria/Rhizobiales/Beijerinckiaceae/unclassified</i>	0.57	0.041	0.60	0.54
	Water	<i>Bacteria/Proteobacteria/Gammaproteobacteria/Alteromonadales/Chromatiaceae/Rheinheimera</i>	0.54	0.005	0.63	0.46
	Water	<i>Bacteria/Proteobacteria/Alphaproteobacteria/Rhodospirillales/Rhodospirillaceae/Magnetospirillum</i>	0.50	0.003	0.68	0.38

<sup>a</sup>No indicative OTUs were identified from streams in fair condition.

<sup>b</sup>A, mean relative abundance of the OTU in each group compared to all groups.

<sup>c</sup>B, relative frequency of each OTU belonging to each group.

ronmental factor that most strongly negatively correlated with Shannon diversity (Table S8), and homogeneous fine sediments have been shown to have lower diversity than that of sites with riffles, shallow turbulent sections (48). Similarly, community structure varied across the geographic regions (Fig. 2) and strongly correlated with DOC concentration, pH, and embeddedness (Table S10 and S11), all of which significantly differentiate Coastal Plain streams from the other regions (Table S2). This finding is in agreement with those from previous studies, demonstrating the strong influence of DOC concentration and pH on freshwater communities (20, 49, 50).

Despite the strong influence of stream chemistry on microbial communities (20, 49, 50), in this study, geodesic distance explained more of the variation in community composition than environmental distance. Partial Mantel test results indicated that community structure was correlated with geographic distance rather than the measured environmental variables. Geographic distance is likely a strong controlling factor in structuring headwater stream communities because there are regional differences in landscape, and stream microbes are locally seeded from the surrounding soil (20, 25, 26). Alpha diversity was greatest in spring, when water flow through the landscape is greatest and, therefore, when advection of microbes from the surrounding landscape is greatest to headwater streams. While seasonal changes in microbial diversity during fall and winter are unknown, the higher diversity in spring than in summer was likely due to higher terrestrial inputs in spring, further demonstrating the influence of landscape on stream microbial communities.

Distance-decay relationships were also observed between water column and sediment community similarity and geodesic distance (Fig. 3), further highlighting the finding that headwater stream microbes display geographic distribution patterns. Alternatively, the distance-decay relationship could be a result of spatial differences in unmeasured environmental variables. Microbial distance-decay relationships have been observed previously in streams (51, 52). Z-values represent the rate at which species similarity decreases with increasing distance; in this study, Z-values (0.12 and 0.11) are similar to microbial values from soil, salt marshes, and lakes (53–56) but lower than regional differences observed in salt marshes (57), suggesting different dispersal limitations across regional scales. In contrast to highly urban and agricultural streams, community dissimilarity in highly forested streams did not increase with distance. Neither geographic distance nor environmental distance correlated with community structure, implying that highly forested streams have a similar terrestrial microbial source.

Microbial diversity differed in streams in watersheds with high urban, agricultural, and forested land use. In contrast to previous studies, degraded streams had lower alpha diversity than that of forested streams (Fig. 5) (22–24), likely due to elevated pollution and habitat loss. Several abundant and pervasive taxa found in urban and agricultural streams (Table 3) are often associated with high-nutrient and low-oxygen environments. Members of the order *Burkholderiales* (families *Alcaligenaceae* and *Comamonadaceae*) were abundant in urban streams and correlated strongly with several anthropogenic nutrients (Fig. 4a). *Comamonadaceae* are often associated with high-nutrient conditions and are ubiquitous in many environments, including aquatic, soil, activated sludge, and wastewater (24, 58). *Comamonadaceae* have previously been associated with urban streams (24) and have been found to have the highest number of urban-tolerant taxa (23, 24). *Sulfurospirillum* spp., in the order *Campylobacteriales*, were abundant in highly agricultural streams and are often associated with microaerophilic polluted habitats, commonly growing on arsenate or selenate using  $\text{NO}_3^-$  and sulfur compounds as electron acceptors (59, 60). In contrast, an unclassified and potentially phototrophic member of *Acidobacteria* and *Hyphomicrobiaceae* (*Rhizobiales*) were more abundant in forested streams. These taxa are often associated with low-nutrient conditions and were previously identified as indicators of forested streams (21, 22) and shown to decrease in abundance with increasing watershed urbanization (23).


Only weak associations were detected between sediment and water microbial community composition and B-IBI scores. This is in contrast to findings of Simonin

et al., who found that stream microbial community structure correlates with a macroinvertebrate biotic index in North Carolina (23). Simonin et al. identified concurrent changes in microbial taxa and environmental conditions associated with the biotic index, finding a higher number of negatively responding taxa ( $n = 68$ ; taxa at higher abundance in streams in good condition) than positive responding taxa ( $n = 8$ ; taxa at higher abundance in streams in fair and poor condition) (23). Here, only one taxon (an unclassified *Archaea*) was more abundant and pervasive in streams in good condition, while several taxa were found to be abundant and pervasive in streams in very poor condition (Table 4). *Hydrogenophaga* spp. (*Burkholderiales*) and an unclassified member of *Desulfobacterales*, commonly associated with anaerobic, reducing, and contaminated environments (58, 61), were both more abundant in streams in very poor condition (Table 4). Forested water communities were more even than agricultural and urban communities, suggesting that certain taxa increase in abundance disproportionally in degraded streams, which is likely why the indicator analysis identified more taxa in streams in very poor condition. The findings here suggest that land use cover and stream chemistry are better predictors of headwater stream microbial community composition than are macroinvertebrate indices of stream conditions.

In agreement with the idea that structure determines function, in this study, water community respiration correlated with microbial community composition. Similarly, previous studies report that changes in community metabolism, specifically, the degradation of organic matter, are related to shifts in community composition and diversity (62–67). In contrast, other studies report that respiration depends on substrate availability rather than community composition due to functional redundancy (68), finding no connection between stream bacterial diversity and the activity of enzymes associated with carbon cycling (69). The weaker correlation between community respiration and community composition in sediments compared to water samples could be due to a high level of functional redundancy within sediment communities; if dominated by generalists, shifts in community composition would likely not significantly affect rates of respiration (70).

Degraded streams had lower rates of community respiration than forested streams, as evidenced by the positive correlation between water respiration and forest cover and the negative correlation between respiration and urban cover (Table S4). Rates of community respiration also negatively correlated with several physicochemical variables (Table S6 and S7), including conductivity,  $\text{Cl}^-$ , Ca, Mg, pH,  $\text{SO}_4^{2-}$ , and ANC. All signatures of anthropogenic influence, ANC,  $\text{Cl}^-$  and Zn concentrations, and pH, were found to previously correlate with benthic stream respiration across the Highlands, Piedmont, and Coastal Plain regions of the eastern United States (30), and Zn is a common urban pollutant (33, 71–73). These results suggest that environmental conditions associated with land use drive differences in community respiration, altering carbon processing in headwater streams.

In addition to altering carbon transformations, watershed modification affected stream nitrogen processing. Agricultural streams had higher  $\text{N}_2\text{O}$  concentrations than those of forested streams (Table S5), with higher  $\text{N}_2\text{O}$  concentrations being associated with elevated TN,  $\text{NO}_2^-$ ,  $\text{NO}_3^-$ , and  $\text{NH}_4^+$  concentrations (Table S5). The  $\text{N}_2\text{O}$  concentrations measured in this study, at  $0.22$  to  $4.41 \mu\text{g N}_2\text{O liter}^{-1}$  (58 to 1,217% saturated), were comparable to the values reported for agricultural streams in Illinois (74, 75) but lower than values published for agricultural drainage waters in Scotland, UK (76) and higher than values reported for other forested and agricultural streams (42, 76–78).  $\text{N}_2\text{O}$  production is known to vary by land use, with higher production from denitrification in streams in agricultural and urban basins (41, 76, 79), and changes in community composition have been shown to influence denitrification rates in agricultural (80) and urban streams (21). However, Audet et al. found no difference in  $\text{N}_2\text{O}$  concentrations measured in forested and agricultural streams in Sweden (76). Stream  $\text{N}_2\text{O}$  concentrations are often correlated with dissolved nitrogen concentrations (41, 76, 79, 81); however, variability in this relationship is often observed between sites (74, 75, 77). In this study,  $\text{N}_2\text{O}$  concentrations were not correlated with microbial community compo-



**FIG 6** Map of Maryland indicating headwater stream sampling locations. Symbol colors indicate the geographic region.

sition, but rather,  $N_2O$  production was likely elevated in streams indirectly due to high rates of denitrification in response to  $NO_3^-$  pollution.

We demonstrate that headwater stream microbial communities and ecosystem processes, such as microbial carbon and nitrogen transformations, respond to gradients in land use and stream conditions. Regional differences in stream microbial communities and the observed distance-decay relationships are further evidence that stream communities are seeded from the surrounding landscape. Across geographic regions, microbial community composition varied in streams with high urban, agricultural, and forested land use, and changes in microbial diversity and land use correlated with stream community respiration, linking changes in biodiversity to changes in ecosystem function. Our results suggest that certain microbial groups respond to land use similarly across ecosystems, making them potential candidate taxa to be used in the development of a microbial index of stream conditions.

## MATERIALS AND METHODS

**Sample collection and physicochemistry.** Stream sediment and water column samples were collected across three general geographic regions (Coastal Plain, Highlands, and Piedmont) in Maryland under baseflow conditions during spring (March and April) and summer (June to September) of 2014 and 2015 (Fig. 6). In 2014, 82 headwater streams (Table S1) were sampled for microbial diversity (Coastal Plain,  $n = 36$ ; Highlands,  $n = 29$ ; Piedmont,  $n = 17$ ), and in 2015, 23 streams across the three regions (Coastal Plain,  $n = 10$ ; Highlands,  $n = 8$ ; Piedmont,  $n = 5$ ) were resampled to assess temporal variability. Community respiration and  $N_2O$  samples were collected at a subset of sites (Tables 1 and 2) based on proximity to the laboratory to ensure proper temperature and light-controlled incubations. Sampling sites were colocated and collected in parallel with Maryland Biological Stream Survey (MBSS) sampling, a Maryland Department of Natural Resources (DNR) monitoring program that assesses the condition of Wadeable streams via physicochemical and biological variables.

One water sample for bacterial and archaeal diversity was collected from each stream in a 500-ml sterile bottle by submerging the bottle into the stream water. Water samples ( $n = 210$ ) were refrigerated until they were filtered on to 0.22- $\mu m$  pore size, 47-mm-diameter polyethersulfone filters (Mo Bio, Carlsbad, CA, USA) and stored at  $-80^\circ C$ . Three sediment samples ( $n = 630$ ) were collected within pools from each stream by inserting the open plunger end of a sterile 5-ml syringe to a depth of 1 cm, and the cores were stored in Whirl-Pak bags at  $-80^\circ C$  until extraction.

Benthic invertebrate samples were collected to calculate a Benthic Index of Biotic Integrity (B-IBI) (17), which is a legal biocriterion in the state. Covariates used as predictors of stream quality in our analyses were provided by the Maryland DNR Monitoring and Non-tidal Assessment Division (<https://dnr.maryland.gov/streams/Pages/default.aspx>), including watershed land use (urban, agricultural, and forested), substrate embeddedness, average thalweg depth, maximum depth, average stream width, average velocity, pH, specific conductance, acid-neutralizing capacity (ANC), and dissolved organic carbon (DOC), chloride ( $Cl^-$ ), sulfate ( $SO_4^{2-}$ ), total nitrogen (TN), total phosphorus (TP), orthophosphate ( $PO_4^{3-}$ ), ammonium ( $NH_4^+$ ), nitrite ( $NO_2^-$ ), nitrate ( $NO_3^-$ ), magnesium (Mg), calcium (Ca), bromide (Br), zinc (Zn), and copper (Cu) content. Land use data, benthic macroinvertebrate samples, and water chemistry samples were collected in spring of each year, while substrate embeddedness, average

thalweg depth, maximum depth, average stream width, and average velocity were measured each summer.

**Nitrous oxide concentrations.** Nitrous oxide samples were collected from a subset of Coastal Plain and Highlands streams (Table 1) in summer 2014 and summer 2015. Samples were collected in triplicate in 160-ml glass serum vials by inserting silicon tubing into the bottom of the vial and then inverting and submerging the vial into the stream water with the other end of the tube venting to the atmosphere. Samples were preserved with 100  $\mu$ l of a saturated mercuric chloride (HgCl<sub>2</sub>) solution, sealed with gray butyl septa and aluminum crimp tops, and stored at room temperature until analysis.

Nitrous oxide concentrations were measured using a headspace equilibration method, as described by Laperriere et al. (82). Each headspace was overpressurized with an addition of 2.5 or 5 ml of ultrahigh-purity (UHP) N<sub>2</sub> and equilibrated with the underlying stream water by gentle shaking at room temperature for at least 2 h. Subsamples from each headspace were analyzed using an SRI Greenhouse gas monitoring gas chromatograph (GC) equipped with an electron capture detector (ECD), dual HayeSep D packed columns, and a 1-ml sample loop (SRI Instruments, Torrance, CA, USA). The carrier gas was UHP N<sub>2</sub>, and the sample loop and column oven were heated to 60°C and 100°C, respectively. Two certified standards, 0.1 ppm and 1 ppm N<sub>2</sub>O, from Matheson Tri-Gas were used for daily calibration. The N<sub>2</sub>O concentrations ( $C_{N_2O}$ ) from the original stream sample were calculated according to Walter et al. (83), using the following equation:

$$C_{N_2O} = \frac{\left( F \times PV_w + \frac{xP}{RT} V_h \right)}{V_w}$$

where  $F$  is equal to  $\ln F = A_1 + A_2(100/T) + A_3 \ln(T/100) + A_4(T/100)^2 + S[B_1 + B_2(T/100) + B_3(T/100)^2]$ ,  $S$  is salinity,  $T$  is the equilibration temperature (K),  $A$  and  $B$  are constants from Weiss and Price (84),  $x$  is the dry gas mole fraction of N<sub>2</sub>O in the headspace,  $P$  is atmospheric pressure,  $V_w$  is the water volume,  $V_h$  is the headspace volume, and  $R$  is the gas constant (in liters atm K<sup>-1</sup> mol<sup>-1</sup>). The equilibrium N<sub>2</sub>O concentration with the atmosphere at *in situ* temperature was calculated using the Weiss and Price (84) solubility equations, using an atmospheric mole fraction of 328 ppb (<https://www.esrl.noaa.gov/gmd/hats/data.html>).

**Community respiration rates.** Sediment and water respiration rates were measured in a subset of Coastal Plain streams (Table 2) using two O<sub>2</sub> consumption methods. In 2014, O<sub>2</sub> consumption was measured using a membrane inlet mass spectrometer (MIMS), following Kana et al. (85), and in 2015 using a Fibox 3 fiber optic oxygen meter (PreSens, Regensburg, Germany). In 2014, water incubations were conducted in 12-ml Exetainer vials (Labco, Lampeter, Wales, UK). For each stream, 9 water samples were collected by inserting a piece of tubing into the bottom of the vial and inverting and submerging the vial into the stream, with the other end of the tube venting to the atmosphere. Three vials were sacrificially killed with a concentrated HgCl<sub>2</sub> solution at three time points, with the first time point immediately after collection and the remaining time points every 4 to 6 h. Vials were transported in a dark cooler back to the laboratory, where they were incubated in the dark at *in situ* temperature for the remainder of the incubation. Sediment incubations were conducted in 160-ml serum vials with butyl septa and aluminum crimp tops. Modified from the above-described collection procedure, sediment was collected by inserting the plunger end of a sterile 30-ml syringe into the sediment and collecting 5 to 10 ml of sediment. The sediment was placed into each vial and topped with stream water.

In 2015, water and sediment incubations were conducted in 60-ml glass biological demand (BOD) bottles with ground glass stoppers. Each BOD bottle contained a PSt3 oxygen sensor (PreSens, Regensburg, Germany). Five replicates were collected, with one killed control from each stream using the methods described above. Once the bottles were full, a thin layer of stream water was added to the top of each stopper to reduce gas exchange with the atmosphere. The bottles were stored in a cooler and transported back to the laboratory, where they were incubated in the dark at *in situ* temperature. O<sub>2</sub> consumption was measured using a Fibox 3 fiber optic oxygen meter (PreSens) every hour until the killed control bottle equilibrated and thereafter every 3 to 6 h for up to 24 h. All respiration rates were calculated using linear least-squares models with the function `lm` in the R package `stats v. 3.5.0` (86). Rates in 2014 were calculated by fitting a model through all 9 data points, while rates from 2015 are the mean of models from each of the replicate bottles. Water respiration rates were subtracted from sediment rates to isolate O<sub>2</sub> consumption in the sediments.

**16S rRNA gene sequencing and processing.** Water and sediment DNA were extracted using a PowerSoil-HTP 96-well soil DNA isolation kit (Mo Bio, Carlsbad, CA, USA), with modifications. For water samples, half of the filter was extracted, and filters were suspended in 925  $\mu$ l of PowerSoil-HTP bead solution and 75  $\mu$ l of solution C1 and vortexed for 10 min. Samples were digested with 20  $\mu$ l of a 20 mg ml<sup>-1</sup> proteinase K solution for 30 min at 56°C and then centrifuged for 1 min at 3,000  $\times g$ . The sediment samples were also digested with proteinase K; additionally, samples were bead beaten at 20 Hz for 20 min on a Qiagen TissueLyser. The PowerSoil-HTP 96-well soil DNA isolation kit protocol was followed for the remainder of the extractions.

16S rRNA gene amplicons were prepared using the standard Illumina protocol (San Diego, CA, USA) with primers 515F and 806R. After amplicon PCR, the three sediment core samples from each site were pooled prior to PCR cleanup. Following the second PCR cleanup, DNA was quantified using a Qubit double-stranded DNA (dsDNA) high-sensitivity kit. Illumina MiSeq 2  $\times$  150-bp (samples collected in 2014) and 2  $\times$  250-bp (samples collected in 2015) sequencing was conducted at the

University of Maryland Center for Environmental Science Institute of Marine and Environmental Technology.

Amplicon data were analyzed using the mothur software package v. 1.31.2 (87). The samples sequenced in 2014 did not have adequate read overlap to merge the reads, and for this reason, only the forward reads were used for all analyses. The mothur standard pipeline was followed, with modifications (87). Prior to quality screening, the samples contained 70,189,751 forward reads, with a median read length of 151 bp. Modified from the standard pipeline, sequences with primer mismatches were removed, and reads were trimmed using an average quality score cutoff of 35 over a 50-bp sliding window. After screening for sequencing and PCR error, 19,731,498 sequences remained, with a median read length of 132 bp. Sequences were aligned with SILVA (v. 119) and classified using Greengenes (v. 13.8.99). Sequences were binned based on taxonomy prior to clustering into operational taxonomic units (OTUs) at a 97% identity level, with a mean sample coverage of 61%. All samples were rarefied to 1,344 sequences, and after pooling, trimming, and rarefaction, 210 water and 204 sediment samples remained.

**Statistical analyses.** All statistical analyses were computed in R (v. 3.5.0) (86). OTU richness, Shannon diversity, and Pielou's evenness were used to estimate microbial alpha diversity and were calculated using the R package phyloseq (v. 1.24.2) (88). Benthic macroinvertebrate Shannon diversity was calculated using the function diversity in the vegan package. Linear least-squares models and stepwise linear regression models for Shannon diversity and the physicochemical data were fit using lm in the stats package. All colinear variables (Pearson's  $|r| \geq 0.7$ ) were removed prior to stepwise linear regression analysis.

Beta diversity was quantified using Bray-Curtis dissimilarity and Sørensen indices and was calculated, with singletons removed using vegdist in the vegan package (v. 2.5.2) (89). Community similarity was visualized using nonmetric multidimensional scaling (NMDS), which was calculated using metaMDS in the vegan package. Correlations between Bray-Curtis dissimilarity and the physicochemistry were calculated using the mantel function in vegan. Similarly, the relationships between Bray-Curtis dissimilarity, Euclidean geographic distance, and environmental variables were examined using the mantel and partial.mantel functions in vegan. The geodesic distance between two sites was calculated using gdist in the lmap package. Distance-decay was calculated using the natural-logarithm-transformed linear least-squares regressions of water and sediment community similarity (Sørensen index) and geodesic distance. The absolute values of the regression coefficients equal the exponent  $Z$  in the taxon-area relationship  $S = cA^Z$ , where  $A$  is area,  $S$  is the number of species, and  $c$  is a constant. Correlations between taxon abundance and environmental data were quantified using Spearman correlations with the function "associate" in the microbiome package (v. 1.1.10013; <http://microbiome.github.io/>). Correlations were calculated at the order rank level because a higher proportion of sequences were classified using Greengenes (v. 13.8.99) at the order rank level (73%) than the proportion classified at the family (57%) and genus (32%) rank levels. The resulting  $P$  values were converted to  $Z$ -scores, and only correlations with a  $|Z$ -score of  $\geq 1.96$  are reported. Indicator taxon analysis was used to identify microbial taxa that were both abundant and pervasive in streams of a particular land use and stream condition. Indicator taxa were identified using multipatt in indicpecies (v.1.7.6), according to Dufrène and Legendre (90) for highly urban (>50%), agricultural (>50%), and forested (>90%) streams, as well as streams classified by B-IBI scores as good (4 to 5), fair (3 to 3.9), poor (2 to 2.9), and very poor (1 to 1.9). An indicator was considered significant with an indicator statistic  $r$  of >0.5, specificity value  $A$  of >0.5, fidelity value  $B$  of >0.1, and  $P$  value of <0.05.

**Data availability.** Sequences are accessible from NCBI under BioProject accession number PRJNA545742.

## SUPPLEMENTAL MATERIAL

Supplemental material is available online only.

**SUPPLEMENTAL FILE 1**, PDF file, 1.4 MB.

## ACKNOWLEDGMENTS

We thank Jason Cessna for conducting much of the field sampling.

This work was supported by funds from a Maryland Sea Grant award (R/EH-16) to A.E.S., S.R.K., and R.H.H. S.M.L. was supported by a Maryland Sea Grant Research Fellowship and the Interdepartmental Graduate Program in Marine Science at the University of California, Santa Barbara. A.E.S. and S.M.L. were also supported by U.S. National Science Foundation award OCE-1437310.

We declare no conflicts of interest.

## REFERENCES

1. Tilman D. 1997. The influence of functional diversity and composition on ecosystem processes. *Science* 277:1300–1302. <https://doi.org/10.1126/science.277.5330.1300>.
2. Falkowski PG, Fenchel T, DeLong EF. 2008. The microbial engines that drive Earth's biogeochemical cycles. *Science* 320:1034–1039. <https://doi.org/10.1126/science.1153213>.
3. Boano F, Harvey JW, Marion A, Packman AI, Revelli R, Ridolfi L, Wörman A. 2014. Hyporheic flow and transport processes: mechanisms, models, and biogeochemical implications. *Rev Geophys* 52:603–679. <https://doi.org/10.1002/2012RG000417>.
4. Battin TJ, Besemer K, Bengtsson MM, Romani AM, Packmann AI. 2016. The ecology and biogeochemistry of stream biofilms. *Nat Rev Microbiol* 14:251–263. <https://doi.org/10.1038/nrmicro.2016.15>.
5. Schindlbacher A, Rodler A, Kuffner M, Kitzler B, Sessitsch A, Zechmeister-

- Boltenstern S. 2011. Experimental warming effects on the microbial community of a temperate mountain forest soil. *Soil Biol Biochem* 43:1417–1425. <https://doi.org/10.1016/j.soilbio.2011.03.005>.
6. Castro HF, Classen AT, Austin EE, Norby RJ, Schadt CW. 2010. Soil microbial community responses to multiple experimental climate change drivers. *Appl Environ Microbiol* 76:999–1007. <https://doi.org/10.1128/AEM.02874-09>.
  7. Sheik CS, Beasley WH, Elshahed MS, Zhou X, Luo Y, Krumholz LR. 2011. Effect of warming and drought on grassland microbial communities. *ISME J* 5:1692–1700. <https://doi.org/10.1038/ismej.2011.32>.
  8. Delong MD, Brusven MA. 1998. Macroinvertebrate community structure along the longitudinal gradient of an agriculturally impacted stream. *Environ Manage* 22:445–457. <https://doi.org/10.1007/s002679900118>.
  9. Young RG, Hury AD. 1999. Effects of land use on stream metabolism and organic matter turnover. *Ecol Appl* 9:1359–1376. [https://doi.org/10.1890/1051-0761\(1999\)009\[1359:EOLUOS\]2.0.CO;2](https://doi.org/10.1890/1051-0761(1999)009[1359:EOLUOS]2.0.CO;2).
  10. Meyer JL, Paul MJ, Taulbee WK. 2005. Stream ecosystem function in urbanizing landscapes. *J North Am Benthol Soc* 24:602–612. <https://doi.org/10.1899/04-021.1>.
  11. Moore AA, Palmer MA. 2005. Invertebrate biodiversity in agricultural and urban headwater streams: implications for conservation and management. *Ecol Appl* 15:1169–1177. <https://doi.org/10.1890/04-1484>.
  12. Whiting ER, Clifford HF. 1983. Invertebrates and urban runoff in a small northern stream, Edmonton, Alberta, Canada. *Hydrobiologia* 102:73–80. <https://doi.org/10.1007/BF00006052>.
  13. Lenat DR, Crawford JK. 1994. Effects of land use on water quality and aquatic biota of three North Carolina Piedmont streams. *Hydrobiologia* 294:185–199. <https://doi.org/10.1007/BF00021291>.
  14. Walsh CJ, Sharpe AK, Breen PF, Sonneman JA. 2001. Effects of urbanization on streams of the Melbourne region, Victoria, Australia. I. Benthic macroinvertebrate communities. *Freshw Biol* 46:535–551. <https://doi.org/10.1046/j.1365-2427.2001.00690.x>.
  15. Hall MJ, Closs GP, Riley RH. 2001. Relationships between land use and stream invertebrate community structure in a South Island, New Zealand, coastal stream catchment. *N Z J Mar Freshw Res* 35:591–603. <https://doi.org/10.1080/00288330.2001.9517025>.
  16. Bonada N, Prat N, Resh VH, Statzner B. 2006. Developments in aquatic insect biomonitoring: a comparative analysis of recent approaches. *Annu Rev Entomol* 51:495–523. <https://doi.org/10.1146/annurev.ento.51.110104.151124>.
  17. Southerland MT, Rogers GM, Kline MJ, Morgan RP, Boward DM, Kazyak PF, Klauda RJ, Stranko SA. 2007. Improving biological indicators to better assess the condition of streams. *Ecol Indic* 7:751–767. <https://doi.org/10.1016/j.ecolind.2006.08.005>.
  18. Weisberg SB, Ranasinghe JA, Dauer DM, Schaffner LC, Diaz RJ, Frithsen JB. 1997. An estuarine benthic index of biotic integrity (B-IBI) for Chesapeake Bay. *Estuaries* 20:149. <https://doi.org/10.2307/1352728>.
  19. Campbell RE, McIntosh AR. 2013. Do isolation and local habitat jointly limit the structure of stream invertebrate assemblages? *Freshw Biol* 58:128–141. <https://doi.org/10.1111/fwb.12045>.
  20. Niño-García JP, Ruiz-González C, Del Giorgio PA. 2016. Interactions between hydrology and water chemistry shape bacterioplankton biogeography across boreal freshwater networks. *ISME J* 10:1755–1766. <https://doi.org/10.1038/ismej.2015.226>.
  21. Wang S-Y, Sudduth EB, Wallenstein MD, Wright JP, Bernhardt ES. 2011. Watershed urbanization alters the composition and function of stream bacterial communities. *PLoS One* 6:e22972. <https://doi.org/10.1371/journal.pone.0022972>.
  22. Hosen JD, Febria CM, Crump BC, Palmer MA. 2017. Watershed urbanization linked to differences in stream bacterial community composition. *Front Microbiol* 8:1452. <https://doi.org/10.3389/fmicb.2017.01452>.
  23. Simonin M, Voss KA, Hassett BA, Rocca JD, Wang SY, Bier RL, Violin CR, Wright JP, Bernhardt ES. 2019. In search of microbial indicator taxa: shifts in stream bacterial communities along an urbanization gradient. *Environ Microbiol* 21:3653–3668. <https://doi.org/10.1111/1462-2920.14694>.
  24. Roberto AA, Van Gray JB, Leff LG. 2018. Sediment bacteria in an urban stream: spatiotemporal patterns in community composition. *Water Res* 134:353–369. <https://doi.org/10.1016/j.watres.2018.01.045>.
  25. Crump BC, Amaral-Zettler LA, Kling GW. 2012. Microbial diversity in arctic freshwaters is structured by inoculation of microbes from soils. *ISME J* 6:1629–1639. <https://doi.org/10.1038/ismej.2012.9>.
  26. Crump BC, Adams HE, Hobbie JE, Kling GW. 2007. Biogeography of bacterioplankton in lakes and streams of an Arctic tundra catchment. *Ecology* 88:1365–1378. <https://doi.org/10.1890/06-0387>.
  27. Claessens L, Tague CL, Groffman PM, Melack JM. 2010. Longitudinal assessment of the effect of concentration on stream N uptake rates in an urbanizing watershed. *Biogeochemistry* 98:63–74. <https://doi.org/10.1007/s10533-009-9376-y>.
  28. Groffman PM, Law NL, Belt KT, Band LE, Fisher GT. 2004. Nitrogen fluxes and retention in urban watershed ecosystems. *Ecosystems* 7:393–403. <https://doi.org/10.1007/s10021-003-0039-x>.
  29. Harbott EL, Grace MR. 2005. Extracellular enzyme response to bioavailability of dissolved organic C in streams of varying catchment urbanization. *J North Am Benthol Soc* 24:588–601. <https://doi.org/10.1899/04-023.1>.
  30. Hill BH, Herlihy AT, Kaufmann PR. 2002. Benthic microbial respiration in Appalachian Mountain, Piedmont, and Coastal Plains streams of the eastern USA. *Freshw Biol* 47:185–194. <https://doi.org/10.1046/j.1365-2427.2002.00791.x>.
  31. Battin TJ, Kaplan LA, Findlay S, Hopkinson CS, Marti E, Packman AI, Denis Newbold J, Sabater F. 2008. Biophysical controls on organic carbon fluxes in fluvial networks. *Nat Geosci* 1:95–100. <https://doi.org/10.1038/ngeo101>.
  32. Fellows CS, Clapcott JE, Udy JW, Bunn SE, Harch BD, Smith MJ, Davies PM. 2006. Benthic metabolism as an indicator of stream ecosystem health. *Hydrobiologia* 572:71–87. <https://doi.org/10.1007/s10750-005-9001-6>.
  33. Paul MJ, Meyer JL. 2001. Streams in the urban landscape. *Annu Rev Ecol Syst* 32:333–365. <https://doi.org/10.1146/annurev.ecolsys.32.081501.114040>.
  34. Wang H, Hondzo M, Xu C, Poole V, Spacie A. 2003. Dissolved oxygen dynamics of streams draining an urbanized and an agricultural catchment. *Ecol Modell* 160:145–161. [https://doi.org/10.1016/S0304-3800\(02\)00324-1](https://doi.org/10.1016/S0304-3800(02)00324-1).
  35. O'Driscoll M, Clinton S, Jefferson A, Manda A, McMillan S. 2010. Urbanization effects on watershed hydrology and in-stream processes in the southern United States. *Water* 2:605–648. <https://doi.org/10.3390/w2030605>.
  36. Hill BH, Hall RK, Husby P, Herlihy AT, Dunne M. 2000. Interregional comparisons of sediment microbial respiration in streams. *Freshw Biol* 44:213–222. <https://doi.org/10.1046/j.1365-2427.2000.00555.x>.
  37. Hill BH, Lazorchak JM, McCormick FH, Willingham WT. 1997. The effects of elevated metals on benthic community metabolism in a rocky mountain stream. *Environ Pollut* 95:183–190. [https://doi.org/10.1016/S0269-7491\(96\)00123-6](https://doi.org/10.1016/S0269-7491(96)00123-6).
  38. Newnam RM, Perry JA, Tam E, Crawford RL. 1987. Effects of chronic chlorine exposure on litter processing in outdoor experimental streams. *Freshw Biol* 18:415–428. <https://doi.org/10.1111/j.1365-2427.1987.tb01327.x>.
  39. Hosen JD, McDonough OT, Febria CM, Palmer MA. 2014. Dissolved organic matter quality and bioavailability changes across an urbanization gradient in headwater streams. *Environ Sci Technol* 48:7817–7824. <https://doi.org/10.1021/es501422z>.
  40. Seitzinger S, Harrison JA, Böhlke JK, Bouwman AF, Lowrance R, Peterson B, Tobias C, Van Drecht G. 2006. Denitrification across landscapes and waterscapes: a synthesis. *Ecol Appl* 16:2064–2090. [https://doi.org/10.1890/1051-0761\(2006\)016\[2064:DALAWA\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2006)016[2064:DALAWA]2.0.CO;2).
  41. Beaulieu JJ, Tank JL, Hamilton SK, Wollheim WM, Hall RO, Mulholland PJ, Peterson BJ, Ashkenas LR, Cooper LW, Dahm CN, Dodds WK, Grimm NB, Johnson SL, McDowell WH, Poole GC, Valett HM, Arango CP, Bernot MJ, Burgin AJ, Crenshaw CL, Helton AM, Johnson LT, O'Brien JM, Potter JD, Sheibley RW, Sobota DJ, Thomas SM. 2011. Nitrous oxide emission from denitrification in stream and river networks. *Proc Natl Acad Sci U S A* 108:214–219. <https://doi.org/10.1073/pnas.1011464108>.
  42. Beaulieu JJ, Arango CP, Hamilton SK, Tank JL. 2007. The production and emission of nitrous oxide from headwater streams in the Midwestern United States. *Glob Chang Biol* 14:878–894. <https://doi.org/10.1111/j.1365-2486.2007.01485.x>.
  43. Newcomer TA, Kaushal SS, Mayer PM, Shields AR, Canuel EA, Groffman PM, Gold AJ. 2012. Influence of natural and novel organic carbon sources on denitrification in forest, degraded urban, and restored streams. *Ecol Monogr* 82:449–466. <https://doi.org/10.1890/12-0458.1>.
  44. Mulholland PJ, Helton AM, Poole GC, Hall RO, Hamilton SK, Peterson BJ, Tank JL, Ashkenas LR, Cooper LW, Dahm CN, Dodds WK, Findlay SEG, Gregory SV, Grimm NB, Johnson SL, McDowell WH, Meyer JL, Valett HM, Webster JR, Arango CP, Beaulieu JJ, Bernot MJ, Burgin AJ, Crenshaw CL, Johnson LT, Niederlehner BR, O'Brien JM, Potter JD, Sheibley RW, Sobota DJ, Thomas SM. 2008. Stream denitrification across biomes and its response to anthropogenic nitrate loading. *Nature* 452:202–205. <https://doi.org/10.1038/nature06686>.
  45. Perryman SE, Rees GN, Walsh CJ. 2008. Analysis of denitrifying commu-



- nities in streams from an urban and non-urban catchment. *Aquat Ecol* 42:95–101. <https://doi.org/10.1007/s10452-007-9085-3>.
46. Perryman SE, Rees GN, Walsh CJ, Grace MR. 2011. Urban stormwater runoff drives denitrifying community composition through changes in sediment texture and carbon content. *Microb Ecol* 61:932–940. <https://doi.org/10.1007/s00248-011-9833-8>.
  47. Ator SW, Denver JM, Krantz DE, Newell WL, Martucci SK. 2005. A surficial hydrogeologic framework for the Mid-Atlantic Coastal Plain. U.S. Geological Survey, Reston, VA. <https://pubs.usgs.gov/pp/1680/report.pdf>.
  48. Rees GN, Watson GO, Baldwin DS, Mitchell AM. 2006. Variability in sediment microbial communities in a semipermanent stream: impact of drought. *J North Am Benthol Soc* 25:370–378. [https://doi.org/10.1899/0887-3593\(2006\)25\[370:VISMCI\]2.0.CO;2](https://doi.org/10.1899/0887-3593(2006)25[370:VISMCI]2.0.CO;2).
  49. Fierer N, Morse JL, Berthrong ST, Bernhardt ES, Jackson RB. 2007. Environmental controls on the landscape-scale biogeography of stream bacterial communities. *Ecology* 88:2162–2173. <https://doi.org/10.1890/06-1746.1>.
  50. Hullar MAJ, Kaplan LA, Stahl DA. 2006. Recurring seasonal dynamics of microbial communities in stream habitats. *Appl Environ Microbiol* 72:713–722. <https://doi.org/10.1128/AEM.72.1.713-722.2006>.
  51. Lear G, Lau K, Percec A, Buckley HL, Case BS, Neale M, Fierer N, Leff JW, Handley KM, Lewis G. 2017. Following Rapoport's Rule: the geographic range and genome size of bacterial taxa decline at warmer latitudes. *Environ Microbiol* 19:3152–3162. <https://doi.org/10.1111/1462-2920.13797>.
  52. Lear G, Washington V, Neale M, Case B, Buckley H, Lewis G. 2013. The biogeography of stream bacteria. *Glob Ecol Biogeogr* 22:544–554. <https://doi.org/10.1111/geb.12046>.
  53. Barreto DP, Conrad R, Klose M, Claus P, Enrich-Prast A. 2014. Distance-decay and taxa-area relationships for bacteria, archaea and methanogenic archaea in a tropical lake sediment. *PLoS One* 9:e110128. <https://doi.org/10.1371/journal.pone.0110128>.
  54. Horner-Devine MC, Lage M, Hughes JB, Bohannon BJM. 2004. A taxa-area relationship for bacteria. *Nature* 432:750–753. <https://doi.org/10.1038/nature03073>.
  55. Green JL, Holmes AJ, Westoby M, Oliver I, Briscoe D, Dangerfield M, Gillings M, Beattie AJ. 2004. Spatial scaling of microbial eukaryote diversity. *Nature* 432:747–750. <https://doi.org/10.1038/nature03034>.
  56. Fierer N, Jackson RB. 2006. The diversity and biogeography of soil bacterial communities. *Proc Natl Acad Sci U S A* 103:626–631. <https://doi.org/10.1073/pnas.0507535103>.
  57. Martiny JBH, Eisen JA, Penn K, Allison SD, Horner-Devine MC. 2011. Drivers of bacterial beta-diversity depend on spatial scale. *Proc Natl Acad Sci U S A* 108:7850–7854. <https://doi.org/10.1073/pnas.1016308108>.
  58. Willems A. 2014. The family Comamonadaceae, p 777–851. *In* Rosenberg E, DeLong EF, Lory S, Stackebrandt E, Thompson F (ed), *The prokaryotes: Alphaproteobacteria and Betaproteobacteria*. Springer, Berlin, Germany.
  59. Lastovica AJ, On SL, Zhang L. 2014. The family Campylobacteraceae, p 307–335. *In* Rosenberg E, DeLong EF, Lory S, Stackebrandt E, Thompson F (ed), *The prokaryotes: Deltaproteobacteria and Epsilonproteobacteria*. Springer, Berlin, Germany.
  60. Goris T, Diekert G. 2016. The genus *Sulfurospirillum*, p 209–234. *In* Löffler FE, Adrian L (ed), *Organohalide-respiring bacteria*. Springer, Berlin, Germany.
  61. Kuever J. 2014. The family Desulfobulbaceae, p 75–86. *In* Rosenberg E, DeLong EF, Lory S, Stackebrandt E, Thompson F (ed), *The prokaryotes: Deltaproteobacteria and Epsilonproteobacteria*. Springer, Berlin, Germany.
  62. Judd KE, Crump BC, Kling GW. 2006. Variation in dissolved organic matter controls bacterial production and community composition. *Ecology* 87:2068–2079. [https://doi.org/10.1890/0012-9658\(2006\)87\[2068:VIDOMC\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2006)87[2068:VIDOMC]2.0.CO;2).
  63. Ylla I, Peter H, Romani AM, Tranvik LJ. 2013. Different diversity-functioning relationship in lake and stream bacterial communities. *FEMS Microbiol Ecol* 85:95–103. <https://doi.org/10.1111/1574-6941.12101>.
  64. Peter H, Ylla I, Gudaszc C, Romani AM, Sabater S, Tranvik LJ. 2011. Multifunctionality and diversity in bacterial biofilms. *PLoS One* 6:e23225. <https://doi.org/10.1371/journal.pone.0023225>.
  65. Singer G, Besemer K, Schmitt-Kopplin P, Hödl I, Battin TJ. 2010. Physical heterogeneity increases biofilm resource use and its molecular diversity in stream mesocosms. *PLoS One* 5:e9988. <https://doi.org/10.1371/journal.pone.0009988>.
  66. Findlay SEG, Sinsabaugh RL, Sobczak WV, Hoostal M. 2003. Metabolic and structural response of hyporheic microbial communities to variations in supply of dissolved organic matter. *Limnol Oceanogr* 48:1608–1617. <https://doi.org/10.4319/lo.2003.48.4.1608>.
  67. Peter H, Beier S, Bertilsson S, Lindström ES, Langenheder S, Tranvik LJ. 2011. Function-specific response to depletion of microbial diversity. *ISME J* 5:351–361. <https://doi.org/10.1038/ismej.2010.119>.
  68. Langenheder S, Lindström ES, Tranvik LJ. 2005. Weak coupling between community composition and functioning of aquatic bacteria. *Limnol Oceanogr* 50:957–967. <https://doi.org/10.4319/lo.2005.50.3.0957>.
  69. Frossard A, Gerull L, Mutz M, Gessner MO. 2012. Disconnect of microbial structure and function: enzyme activities and bacterial communities in nascent stream corridors. *ISME J* 6:680–691. <https://doi.org/10.1038/ismej.2011.134>.
  70. Östman Ö, Drakare S, Kritzbeg ES, Langenheder S, Logue JB, Lindström ES. 2010. Regional invariance among microbial communities. *Ecol Lett* 13:118–127. <https://doi.org/10.1111/j.1461-0248.2009.01413.x>.
  71. Council TB, Duckenfield KU, Landa ER, Callender E. 2004. Tire-wear particles as a source of zinc to the environment. *Environ Sci Technol* 38:4206–4214. <https://doi.org/10.1021/es034631f>.
  72. Lettenmaier DP, Hooper ER, Wagoner C, Faris KB. 1991. Trends in stream quality in the continental United States, 1978–1987. *Water Resour Res* 27:327–339. <https://doi.org/10.1029/90WR02140>.
  73. Wilber WG, Hunter JV. 2007. The impact of urbanization on the distribution of heavy metals in bottom sediments of the Saddle River. *J Am Water Resour Assoc* 15:790–800. <https://doi.org/10.1111/j.1752-1688.1979.tb00397.x>.
  74. Davis MP, David MB. 2018. Nitrous oxide fluxes from agricultural streams in east-central Illinois. *Water Air Soil Pollut* 229:354. <https://doi.org/10.1007/s11270-018-4007-7>.
  75. Reay DS, Edwards AC, Smith KA. 2009. Importance of indirect nitrous oxide emissions at the field, farm and catchment scale. *Agric Ecosyst Environ* 133:163–169. <https://doi.org/10.1016/j.agee.2009.04.019>.
  76. Audet J, Bastviken D, Bundschuh M, Buffam I, Fleckler A, Klemmedtsson L, Laudon H, Löfgren S, Natchimuthu S, Öquist M, Peacock M, Wallin MB. 2019. Forest streams are important sources for nitrous oxide emissions. *Glob Chang Biol* 26:629–641. <https://doi.org/10.1111/gcb.14812>.
  77. Vidon P, Serchan S. 2016. Impact of stream geomorphology on greenhouse gas concentration in a New York mountain stream. *Water Air Soil Pollut* 227:428. <https://doi.org/10.1007/s11270-016-3131-5>.
  78. Höll BS, Jungkunst HF, Fiedler S, Stahr K. 2005. Indirect nitrous oxide emission from a nitrogen saturated spruce forest and general accuracy of the IPCC methodology. *Atmos Environ* 39:5959–5970. <https://doi.org/10.1016/j.atmosenv.2005.06.026>.
  79. Turner PA, Griffis TJ, Baker JM, Lee X, Crawford JT, Loken LC, Venterea RT. 2016. Regional-scale controls on dissolved nitrous oxide in the Upper Mississippi River. *Geophys Res Lett* 43:4400–4407. <https://doi.org/10.1002/2016GL068710>.
  80. Baxter AM, Johnson L, Edgerton J, Royer T, Leff LG. 2012. Structure and function of denitrifying bacterial assemblages in low-order Indiana streams. *Freshw Sci* 31:304–317. <https://doi.org/10.1899/11-066.1>.
  81. Mühlherr IH, Hiscock KM. 1997. A preliminary assessment of nitrous oxide in Chalk groundwater in Cambridgeshire, U.K. *Appl Geochem* 12:797–802. [https://doi.org/10.1016/S0883-2927\(97\)00036-X](https://doi.org/10.1016/S0883-2927(97)00036-X).
  82. Laperriere SM, Nidzicko NJ, Fox RJ, Fisher AW, Santoro AE. 2019. Observations of variable ammonia oxidation and nitrous oxide flux in a eutrophic estuary. *Estuaries Coast* 42:33–44. <https://doi.org/10.1007/s12237-018-0441-4>.
  83. Walter S, Bange HW, Breitenbach U, Wallace D. 2006. Nitrous oxide in the North Atlantic Ocean. *Biogeosci Discuss* 3:993–1022. <https://doi.org/10.5194/bgd-3-993-2006>.
  84. Weiss RF, Price BA. 1980. Nitrous oxide solubility in water and seawater. *Mar Chem* 8:347–359. [https://doi.org/10.1016/0304-4203\(80\)90024-9](https://doi.org/10.1016/0304-4203(80)90024-9).
  85. Kana TM, Darkangelo C, Duane Hunt M, Oldham JB, Bennett GE, Cornwell JC. 1994. Membrane inlet mass spectrometer for rapid high-precision determination of N<sub>2</sub>, O<sub>2</sub>, and Ar in environmental water samples. *Anal Chem* 66:4166–4170. <https://doi.org/10.1021/ac00095a009>.
  86. R Core Team. 2013. R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <https://www.r-project.org/>.
  87. Kozich JJ, Westcott SL, Baxter NT, Highlander SK, Schloss PD. 2013. Development of a dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence data on the MiSeq Illumina sequencing

- platform. *Appl Environ Microbiol* 79:5112–5120. <https://doi.org/10.1128/AEM.01043-13>.
88. McMurdie PJ, Holmes S. 2013. phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. *PLoS One* 8:e61217. <https://doi.org/10.1371/journal.pone.0061217>.
89. Oksanen JF, Blanchet G, Friendly M, Kindt R, Legendre P, McGlenn D, Minchin PR, O'Hara RB, Simpson GL, Solymos P, Stevens MHH, Szoecs E, Wagner H. 2019. vegan: community ecology package. R package version 2.5-5. <https://cran.r-project.org/package=vegan>.
90. Dufrene M, Legendre P. 1997. Species assemblages and indicator species: the need for a flexible asymmetrical approach. *Ecol Monogr* 67:345–366. <https://doi.org/10.2307/2963459>.