

# Lawrence Berkeley National Laboratory

## Recent Work

### Title

FURTHER EVIDENCE FOR STROMA LAMELLAE AS THE SOURCE OF PHOTOSYSTEM 1 FRACTIONS FROM SPINACH CHLOROPLASTS

### Permalink

<https://escholarship.org/uc/item/7162909s>

### Authors

Goodchild, David J.  
Park, Roderic B.

### Publication Date

1970-08-01

FURTHER EVIDENCE FOR STROMA  
LAMELLAE AS THE SOURCE OF  
PHOTOSYSTEM 1 FRACTIONS FROM  
SPINACH CHLOROPLASTS

David J. Goodchild and Roderic B. Park

August 1970

AEC Contract No. W-7405-eng-48

RECEIVED  
LAWRENCE  
RADIATION LABORATORY  
SEP 30 1970  
LIBRARY AND  
DOCUMENTS SECTION

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy  
which may be borrowed for two weeks.  
For a personal retention copy, call  
Tech. Info. Division, Ext. 5545*

42  
LAWRENCE RADIATION LABORATORY  
UNIVERSITY of CALIFORNIA BERKELEY

## DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

FURTHER EVIDENCE FOR STROMA LAMELLAE AS THE SOURCE OF PHOTOSYSTEM 1  
FRACTIONS FROM SPINACH CHLOROPLASTS

D. J. GOODCHILD\* AND R. B. PARK

Botany Department and Laboratory of Chemical Biodynamics, Lawrence  
Radiation Laboratory, University of California, Berkeley, California  
94720 (U.S.A.).

SUMMARY

The mechanism of digitonin action on spinach chloroplasts was investigated by thin sectioning. Evidence is presented which shows that digitonin continues to modify membranes for many minutes after the addition of the fixative glutaraldehyde. However, the action of digitonin can be stopped by simultaneous fixation and dilution of the detergent. Such experiments indicate that the initial action of digitonin is to release stroma lamellae which in turn yield the Photosystem 1 fraction. This interpretation is further supported by a significant correlation between the chlorophyll a/chlorophyll b ratio and the ratio of stroma to grana lamellae in spinach chloroplasts.

---

INTRODUCTION

Recent work from this laboratory showed that spinach chloroplasts broken in a French pressure cell can be separated into two membrane fractions consisting of stroma lamellae and grana lamellae. Assays of these fractions for the two photosystems showed that stroma lamellae

---

\*On leave from the Division of Plant Industry, CSIRO, Canberra, Australia.

contained only Photosystem 1 while the grana fraction contained both Photosystems<sup>1</sup>.

Anderson and Boardman<sup>2</sup> have used the detergent digitonin to separate spinach chloroplasts into membrane fractions which are very similar to the French pressure cell fractions in chemical composition, photochemical activity and yield. Both methods yield approximately 10% of the chlorophyll in a fraction possessing only Photosystem 1 activity and a high chlorophyll a/chlorophyll b ratio. Arntzen et al.<sup>3</sup> extended the work of Anderson and Boardman<sup>2</sup> by investigating the ultrastructure of the digitonin fractions by sectioning and freeze etching. They concluded that digitonin had split the chloroplast lamellar membranes longitudinally to yield the fraction having only Photosystem 1 activity.

The biochemical similarity between the digitonin and French pressure cell fractions has led us to re-examine the ultrastructure of the digitonin fractions to find whether our predictions or the Arntzen<sup>3</sup> interpretation is correct. We conclude from the work reported here that in the early stages of digitonin treatment, when the highest yield of Photosystem 1 fraction is obtained<sup>2</sup>, the digitonin and French pressure cell are acting in a similar manner and that in both methods the Photosystem 1 fraction is derived from stroma lamellae and not from membrane splitting.

Additional support for this distribution of photosystems between grana and stroma lamellae comes from the observation that in spinach chloroplasts there is a significant correlation between a high

chlorophyll a/chlorophyll b ratio and a high proportion of stroma lamellae.

#### MATERIALS AND METHODS

Chloroplasts were isolated from field grown spinach plants in potassium phosphate buffer (pH 7.4)-0.01 M KCl - 0.5 M sucrose and the pellet resuspended in the same buffer<sup>1</sup>. For digitonin treatment the chlorophyll concentration was adjusted to 0.3-0.4 µg/ml with buffer and to this 2% digitonin was added (2x recrystallized) to a final concentration of 0.5%<sup>3</sup>. The incubation with digitonin was carried out in an ice bath with constant stirring. Chlorophyll a and chlorophyll b were determined using the spectrophotometric method of Arnon<sup>4</sup> and a Cary Model 14R spectrophotometer with a scattered transmission accessory.

For electron microscopy of the chloroplast suspensions, samples were removed and pelleted as completely as possible in a 1 ml centrifuge tube in an adaptor in the swinging bucket rotor of a Servall centrifuge at 27,500 x g for 20 min. The resulting pellets were washed 4 times by replacing the fluid above each with phosphate buffer and then post-fixing for 1 hr in 2% osmium tetroxide in phosphate buffer. These procedures were carried out in the cold, but the samples were dehydrated in acetone and propylene oxide at room temperature. Then the end of the centrifuge tube was cut off and the pellets were extruded and embedded in epon. For sectioning, the epon embedded pellets were oriented and trimmed so that each section contained a range of material from the top to the bottom of the pellet. Sections

were stained with saturated uranyl acetate in 50% ethanol for 90 min followed by Fiske's<sup>5</sup> lead citrate for 20 min before examination in a Siemens Elmiskop 1A electron microscope.

When leaf pieces were fixed for electron microscopy a similar procedure was adopted except that pellets were replaced by mm squares of spinach leaf tissue.

## EXPERIMENTAL AND RESULTS

### 1. Digitonin treatments

Time course studies of photochemical activity of both chloroplast and algae following glutaraldehyde addition show that complete fixation may take up to 1 hr<sup>6</sup>. Therefore it is unlikely that the addition of glutaraldehyde will stop the action of digitonin immediately. Two experiments were carried out to reveal the ultrastructural state of the membranes at the time when the digitonin fractions are subjected to differential centrifugation for the separation of Photosystem 1. In the first experiment the time course of digitonin action was studied by adding fixative to the incubation mixture. In the second experiment dilution was combined with fixation in an attempt to stop the action of digitonin.

#### A. Fixative addition experiment

In this experiment 0.5 ml samples were removed from the chloroplast digitonin incubation mixture at 5, 20, and 60 min after the addition of digitonin. To each was added 0.5 ml of 4% glutaraldehyde in phosphate buffer. The sample was centrifuged and embedded as described in Materials and Methods. Figure 1 shows a section through

a pellet from 5 min digitonin treated chloroplasts. The interconnecting stroma lamellae are absent but grana stacks are still intact although most are swollen. Figure 2 shows the 60 min digitonin treatment. Grana stacks are absent and all that remains are swollen vesicles, presumably remnants of stroma lamellae and grana lamellae. The 20 min treatment was intermediate between these two and had very few grana stacks. We interpret these pictures to indicate that stroma lamellae are separated from grana during the first few minutes of digitonin action but that prolonged digitonin treatment causes grana to dissociate into vesicles.

#### B. Fixative dilution experiment

Chloroplasts were treated with digitonin as described previously, and after 5 and 30 min treatment<sup>5</sup> a 1 ml sample was removed and diluted in 100 parts of 0.05 M potassium phosphate buffer (pH 7.4) - 0.01 M KCl containing 2% glutaraldehyde. This material was sedimented at 40,000 x g in 50 ml centrifuge tubes and the pellets transferred to the 1 ml tubes for treatment as described in Materials and Methods. Figure 3 shows a section through a pellet from the 5 min digitonin treatment which shows intact, and relatively unswollen, grana stacks but no interconnecting stroma lamellae. The 30 min digitonin treatment yielded similar pictures. Both these results indicate that dilution is necessary in addition to glutaraldehyde fixation to minimize the continued action of digitonin.

A 5 min digitonin treatment followed by dilution was sufficient to separate grana and stroma lamellae. In another 5 min fixative dilution experiment the fraction that sedimented between 40,000 x g



and 160,000 x g was separated for determination of the chlorophyll a/chlorophyll b ratio. This fraction had a chlorophyll a/chlorophyll b ratio of 5.0 and represented approximately 5% of the total chlorophyll in the incubation mixture. No such fraction was obtained from a control in which distilled water replaced digitonin. This additional evidence supports the evidence from electron micrographs that chloroplast fractionation was achieved with a 5 min digitonin incubation followed by dilution. The fact that the interconnecting stroma lamellae are absent (Fig. 3) again implicates them as the source of the high chlorophyll a/chlorophyll b ratio fraction which has been shown to possess only Photosystem I activity.

## II. Chloroplast structure related to chlorophyll a/chlorophyll b ratio

In our earlier work we showed that the Photosystem I fraction obtained from spinach leaf chloroplasts by the French pressure cell technique represented the stroma lamellae of the intact chloroplasts. This fraction has a high chlorophyll a/chlorophyll b ratio, and as a consequence one would predict a simple relationship between the ratio of grana and stroma lamellae and the chlorophyll a/chlorophyll b ratio of intact spinach chloroplasts. Chloroplasts having a high chlorophyll a/chlorophyll b ratio should yield a higher percentage of the Photosystem I fraction. Young spinach obtained from the field near San Francisco in the month of April 1970 yielded a chloroplast preparation with a chlorophyll a/chlorophyll b ratio of 3.2. The yield of Photosystem I fraction from these chloroplasts following French pressure cell treatment was 26% compared to a yield of 10% from chloroplasts having a chlorophyll a/chlorophyll b ratio of 2.6.

To establish a correlation between the chlorophyll a/chlorophyll b ratio and the percentage of stroma lamellae in spinach chloroplasts, use was made of the fact that in a young spinach leaf cell division within the lamina ceases at an early stage of development in the tip region but continues for an extended period at the base of the leaf<sup>7</sup>. This finding suggested that the younger chloroplasts in the basal region may have a higher percentage of stroma lamellae and also a higher chlorophyll a/chlorophyll b ratio than the leaf as a whole.

A young leaf was selected that was obviously dark green at the tip and lighter green at the base. The terminal 0.5 cm was removed and one-half cut into \_\_\_\_\_ mm squares and fixed for electron microscopy. The other half was extracted in 80% acetone for chlorophyll determinations. The basal 0.5 cm of the leaf was treated in the same manner.

The results of two such experiments are shown in Table I. The total lengths of stroma, end grana and grana lamellae were measured on electron micrographs of single chloroplasts at a magnification of 42,000x and the results expressed as a percentage of the total lamellar length. In the first experiment, 5 chloroplasts were measured from each of four different leaf pieces and the results analyzed statistically using a t test. In the second experiment, 7 chloroplasts were measured from a leaf tip and 11 from the base. Although some variation was evident, the differences between tip and base were significant at the 0.1% level. This result indicates that there is a correlation between the chlorophyll a/chlorophyll b ratio and the percentage of stroma and end grana lamellae in spinach chloroplasts. A higher chlorophyll a/chlorophyll b ratio is significantly correlated with a greater percentage of stroma and end grana lamellae.

## DISCUSSION

The results reported here provide further support for our earlier observation that digitonin and French pressure cell treatments produce similar fractions from spinach chloroplasts<sup>1</sup>. Arntzen et al.<sup>3</sup> showed that the Photosystem 1 fraction from digitonin treated spinach chloroplasts contained only small particles on freeze fractured faces, compared with the small and large particles observed by Branton and Park<sup>8</sup> in intact spinach chloroplasts. Sane et al.<sup>1</sup> have also demonstrated small particles in a similar ultracentrifuge fraction following French pressure cell treatment. Arntzen et al.<sup>3</sup> believe that the chloroplast membranes split longitudinally to produce the small particle fraction. Their conclusion is based on fixed and sectioned digitonin treated spinach chloroplasts and their Figure 6 resembles Figure 2 of this paper.

The experiments reported here support the notion that glutaraldehyde fixation of chloroplast membranes is not rapidly achieved in the presence of 0.5% digitonin. If it were, then the images observed in the 20 and 30 min treatments from the fixative addition and fixative dilution experiments should be similar. The fact that few, if any, vestiges of grana stacks could be observed after 20 min in the fixative addition experiment, while many grana stacks were observed in the fixative dilution experiment, suggests that the mere addition of fixative is not sufficient to stop the action of digitonin. Since Arntzen et al.<sup>3</sup> added glutaraldehyde to their reaction mixture after 30 min incubation (our fixative addition procedure), their electron

micrographs probably do not reflect the true state of the membranes at that time, but an advanced stage of digitonin treatment. The differential centrifugation for the preparation of the photochemically active fractions is begun after approximately 30 min incubation in digitonin when the grana stacks are still intact, as shown by the fixative dilution experiment. The resulting rapid removal of the grana fraction leaves stroma lamellae which are subsequently precipitated as the Photosystem 1 fraction. Longer digitonin treatments (60 min, Fig. 2) show almost complete dissolution of the grana, which is consistent with the biochemical observation that prolonged incubation in digitonin followed by differential centrifugation no longer separates the two photosystems<sup>2</sup>.

Further evidence for the implication of stroma lamellae as the source of the Photosystem 1 fraction is provided by Wehrmeyer<sup>9</sup>. He used concentrations of digitonin that are lower than the 0.5% in 0.3  $\mu$ g of chlorophyll used for photosystem fractionation. However, his pictures clearly show (Figs. 3, 4 and 5) that the primary action of digitonin is on the stroma lamellae of spinach chloroplasts. These first swell, and as the digitonin concentration is increased the stroma lamellae finally form round vesicles between the grana stacks and separate from them.

Sane et al.<sup>1</sup> provided sectioning, freeze fracturing and deep etching evidence to implicate stroma lamellae and end grana membranes as the source of the Photosystem 1 fraction produced by the French pressure cell treatment of spinach chloroplasts. Further support is

provided from the measurements of chloroplasts from regions of the same leaf having different chlorophyll a/chlorophyll b ratios. In this case a higher proportion of stroma lamellae, and end grana membranes, was present in the high chlorophyll a/chlorophyll b ratio area of the leaf. Also French pressure cell fractionation of chloroplasts from high chlorophyll a/chlorophyll b ratio material produced a higher yield of Photosystem I fraction. This result may explain the findings of Ohki and Takamiya<sup>10</sup>, who obtained a Photosystem I fraction by digitonin treatment that represented 40% of the total chlorophyll in the reaction mixture. They used isolated spinach chloroplasts with a chlorophyll a/chlorophyll b ratio of 3.0 compared with the ratio of 2.6 to 2.8 of Anderson and Boardman<sup>2</sup> and Arntzen et al.<sup>3</sup> This chloroplast preparation may have had an unusually high abundance of stroma lamellae.

All these results are consistent with the interpretation suggested by Sane et al.<sup>1</sup> that both digitonin and French pressure cell treatments of spinach chloroplasts release stroma lamellae as the Photosystem I fraction.

#### ACKNOWLEDGMENTS

This work was supported by National Institute of General Medicine Grant GM-13943-14 and the U. S. Atomic Energy Commission.

REFERENCES

1. P. V. SANE, D. J. GOODCHILD AND R. B. PARK, Biochim. Biophys. Acta, in press.
2. J. M. ANDERSON AND N. K. BOARDMAN, Biochim. Biophys. Acta, 112 (1966) 403.
3. C. J. ARNTZEN, R. A. DILLEY AND F. L. CRANE, J. Cell. Biol., 43 (1969) 16.
4. D. I. ARNON, Plant Physiol., 24 (1949) 1.
5. S. FISKE, J. Microsc., 5 (1966) 355.
6. R. B. PARK, J. KELLY, S. DRURY AND K. SAUER, Proc. Nat. Acad. Sci. U.S., 55 (1966) 1056.
7. W. SAURER AND J. V. POSSINGHAM, J. Exp. Bot., 21 (1970) 151.
8. D. BRANTON AND R. B. PARK, J. Ultrastructure Res., 19 (1967) 283.
9. W. WEHRMEYER, Z. Naturforsch., 17b (1962) 54.
10. R. OHKI AND A. TAKAMIYA, Biochim. Biophys. Acta, 197 (1970) 240.

TABLE I

RELATIONSHIP BETWEEN CHLOROPHYLL a/CHLOROPHYLL b RATIO AND RELATIVE PROPORTIONS OF STROMA AND END GRANA LAMELLAE

Results from two experiments on the tip and basal regions of a young spinach leaf, showing a correlation between chlorophyll a/chlorophyll b ratio and stroma and end grana lamellae length percentages, expressed as percentages of total chloroplast lamellar length.

	Tip	Base
Ratio chlorophyll a/ chlorophyll b	2.7/2.8*	3.5/3.5
% Stroma lamellae	17.3/9.6	37.8/23.1
% Stroma and end grana lamellae	33.8/28.1	58.2/44.1

\*Values on right of / are from experiment 2.

% differences between tip and base sig. at 0.1%.

RUNNING TITLE

Origin of PS 1 Fractions





Fig. 1





Fig. 2

XBB 709-3903



XBB 709-3904

Fig. 3

LEGAL NOTICE

*This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:*

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or*
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.*

*As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.*

TECHNICAL INFORMATION DIVISION  
LAWRENCE RADIATION LABORATORY  
UNIVERSITY OF CALIFORNIA  
BERKELEY, CALIFORNIA 94720