

# UC Berkeley

## UC Berkeley Previously Published Works

### Title

Expression of a bacterial 3-dehydroshikimate dehydratase (QsuB) reduces lignin and improves biomass saccharification efficiency in switchgrass (*Panicum virgatum* L.)

### Permalink

<https://escholarship.org/uc/item/71810061>

### Journal

BMC Plant Biology, 21(1)

### ISSN

1471-2229

### Authors

Hao, Zhangying

Yogiswara, Sasha

Wei, Tong

et al.

### Publication Date

2021

### DOI

10.1186/s12870-021-02842-9

### Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>


Peer reviewed

RESEARCH ARTICLE

Open Access



# Expression of a bacterial 3-dehydroshikimate dehydratase (QsuB) reduces lignin and improves biomass saccharification efficiency in switchgrass (*Panicum virgatum* L.)

Zhangying Hao<sup>1,2</sup>, Sasha Yogiswara<sup>1,2</sup>, Tong Wei<sup>1,2,3,4</sup>, Veronica Teixeira Benites<sup>1,2</sup>, Anagh Sinha<sup>1,2</sup>, George Wang<sup>1,5</sup>, Edward E. K. Baidoo<sup>1,5</sup>, Pamela C. Ronald<sup>1,2,3</sup>, Henrik V. Scheller<sup>1,2,6</sup>, Dominique Loqué<sup>1,2</sup> and Aymerick Eudes<sup>1,2\*</sup> 

## Abstract

**Background:** Lignin deposited in plant cell walls negatively affects biomass conversion into advanced bioproducts. There is therefore a strong interest in developing bioenergy crops with reduced lignin content or altered lignin structures. Another desired trait for bioenergy crops is the ability to accumulate novel bioproducts, which would enhance the development of economically sustainable biorefineries. As previously demonstrated in the model plant *Arabidopsis*, expression of a 3-dehydroshikimate dehydratase in plants offers the potential for decreasing lignin content and overproducing a value-added metabolic coproduct (i.e., protocatechuate) suitable for biological upgrading.

**Results:** The 3-dehydroshikimate dehydratase QsuB from *Corynebacterium glutamicum* was expressed in the bioenergy crop switchgrass (*Panicum virgatum* L.) using the stem-specific promoter of an O-methyltransferase gene (*pShOMT*) from sugarcane. The activity of *pShOMT* was validated in switchgrass after observation in-situ of beta-glucuronidase (GUS) activity in stem nodes of plants carrying a *pShOMT::GUS* fusion construct. Under controlled growth conditions, engineered switchgrass lines containing a *pShOMT::QsuB* construct showed reductions of lignin content, improvements of biomass saccharification efficiency, and accumulated higher amount of protocatechuate compared to control plants. Attempts to generate transgenic switchgrass lines carrying the QsuB gene under the control of the constitutive promoter *pZmUbi-1* were unsuccessful, suggesting possible toxicity issues associated with ectopic QsuB expression during the plant regeneration process.

(Continued on next page)

\* Correspondence: [ageudes@lbl.gov](mailto:ageudes@lbl.gov)

<sup>1</sup>Joint BioEnergy Institute, Emeryville, CA 94608, USA

<sup>2</sup>Environmental Genomics and Systems Biology Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA

Full list of author information is available at the end of the article



© The Author(s). 2021 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

(Continued from previous page)

**Conclusion:** This study validates the transfer of the QsuB engineering approach from a model plant to switchgrass. We have demonstrated altered expression of two important traits: lignin content and accumulation of a co-product. We found that the choice of promoter to drive *QsuB* expression should be carefully considered when deploying this strategy to other bioenergy crops. Field-testing of engineered *QsuB* switchgrass are in progress to assess the performance of the introduced traits and agronomic performances of the transgenic plants.

**Keywords:** Switchgrass, Lignin, Shikimate, Protocatechuate, Saccharification, Bioenergy

**Background**


The development of biorefineries to reduce our dependence on nonrenewable fossil fuel resources requires production of dedicated bioenergy crops that can be grown with few inputs on marginal lands. Other desired traits for bioenergy crops include high biomass yields, stress resilience, reduced recalcitrance to conversion into biofuels and bioproducts, and the accumulation of valuable co-products [1, 2]. Switchgrass has long been recognized as an ideal crop for bioenergy purposes considering its pest and disease resistance, high biomass yields, growth performance on poor soils due to relatively low requirements for added fertilizers, carbon sequestration capacity via its extensive root system, drought tolerance, and efficient water use [3]. As a consequence, significant efforts have been implemented for the improvement of switchgrass via breeding and genetic transformation [4, 5].

Lignin is a major polymer in plant biomass that negatively impacts the conversion of cell wall polysaccharides into advanced bioproducts, and several engineering approaches have been established to modify lignin content and its monomeric composition [6, 7]. For example, the heterologous expression of a bacterial 3-dehydroshikimate dehydratase (*QsuB*) targeted to plastids resulted in strong

lignin reductions (up to 50%) in *Arabidopsis* [8]. One explanation for this observation is the possible reduction of the cytosolic shikimate pool needed for the synthesis of *p*-coumaroyl-shikimate catalyzed by hydroxycinnamoyl CoA: shikimate hydroxycinnamoyl transferase (*HCT*) during lignin biosynthesis (Fig. 1).

In switchgrass, several *HCT* gene candidates have been proposed to have a role in lignin biosynthesis based on the *HCT* activity measured with the corresponding recombinant enzymes and their expression profile in lignifying cell suspension cultures [9, 10]. In fact, more than 90% reduction in transcript levels of either *PvHCT1* or *PvHCT2* had no effect on lignin content, but simultaneous downregulation of both genes resulted in slight decreases of lignin content (5–8%) based on the yield of lignin monomers released after thioacidolysis [11]. These results not only indicate a role for *HCT* in lignin biosynthesis in switchgrass, with *PvHCT1* and *PvHCT2* being redundant, but also suggest the involvement of additional *HCTs* with similar functions.

In this work, we report on the expression of *QsuB* in switchgrass using the promoter of a sugarcane *O*-methyltransferase gene (*pShOMT*) [12]. Several switchgrass *QsuB* transformation events show reduction of



**Fig. 1** Schematic diagram of lignin biosynthesis and the conversion of 3-dehydroshikimate into protocatechuate (PCA) catalyzed by plastid-targeted *QsuB*. Grey and blue circles indicate a phenylalanine transporter and a putative shikimate transporter, respectively. Dashed arrows represent multiple enzymatic steps. E4P: Erythrose 4-phosphate; HCT: hydroxycinnamoyl CoA: shikimate hydroxycinnamoyl transferase; PEP: Phosphoenolpyruvate; PHE: Phenylalanine

lignin content and decreased cell wall recalcitrance. A significant increase in the content of protocatechuate accumulated in biomass was also observed.

## Results

### Molecular characterization of the *pShOMT::QsuB* switchgrass lines

A total of eight independent transformation events were regenerated after *Agrobacterium*-mediated transformation of switchgrass using a DNA construct that contains the plastid-targeted *QsuB* coding sequence fused downstream of the *pShOMT* promoter. The *QsuB* transgene was detected by PCR using gDNA from each transformant (Fig. 2a), and *QsuB* expression was validated by qPCR performed on cDNA synthesized from RNAs obtained from the first internode of each line at the E2 stage (Fig. 2b). A DNA construct consisting of *pShOMT* fused upstream of the *GUS* reporter gene was also transferred to switchgrass. Analysis of internodes and nodes from switchgrass plants harboring the *pShOMT::GUS* construct at the E4 stage suggested that *pShOMT* is mainly active in the nodes, whereas little activity was observed in the internodes (Figure S1). Under controlled growth conditions, all transgenic lines did not show any particular phenotype nor growth defect and were visually indistinguishable from each other or compared to non-transformed wild-type plants.

### Protocatechuate content in *pShOMT::QsuB* switchgrass


Protocatechuate (PCA), the product of *QsuB* activity, was extracted from the total aboveground biomass of


switchgrass plants at the E5 stage and quantified. Compared to control plants carrying the *pShOMT::GUS* construct, PCA was significantly increased by ~2–3-fold in four independent *pShOMT::QsuB* lines, reaching up to 380 µg/g dry weight (Fig. 3). This data shows that expression of plastid-targeted *QsuB* in transgenic switchgrass enabled the conversion of endogenous 3-dehydroshikimate into PCA.

### Lignin content and biomass saccharification efficiency in *pShOMT::QsuB* switchgrass

Total lignin content in the biomass from the *pShOMT::QsuB* switchgrass lines was measured using the Klason method. Compared to control lines containing the *pShOMT::GUS* construct, several *pShOMT::QsuB* lines showed significant reductions of lignin content ranging from 12 to 21% (Fig. 4a). Inspection of stem sections treated with phloroglucinol-HCl for the staining of lignin did not reveal any differences between the different *pShOMT::QsuB* lines and the control *pShOMT::GUS* lines (data not shown). However, on leaf blade sections, reductions in the intensity of the typical red staining were observed in the case of the *pShOMT::QsuB* lines compared to controls, especially in thick fibers located in the abaxial zone (Fig. 4b).

The recalcitrance towards enzymatic degradation of the biomass of the engineered switchgrass was evaluated by measuring the amount of sugars released from cell wall residues after pretreatment with hot water followed by a 72-h hydrolysis using a commercial cellulase







cocktail (CTec2). As shown in Fig. 5, higher amount of reducing sugars was obtained for several *pShOMT::QsuB* lines compared to the *pShOMT::GUS* control lines, with significant increases ranging between 21 and 30%.

### Discussion

Here, we describe the successful expression of the bacterial 3-dehydroshikimate dehydratase *QsuB* gene under the control of *pShOMT* in switchgrass. We show that the resulting plants display 12–21% reduction in lignin, a 2–3-fold increase in the bioaccumulation of PCA and a 5–30% increase in saccharification efficiency.

*pShOMT* was previously shown to be preferentially active in stem vascular tissues in sugarcane, rice, maize, and sorghum [12], making it a good promoter candidate to express *QsuB* specifically in lignifying tissues within vascular bundles. Similar to previous observations made in sugarcane, we were able to detect GUS activity in stem nodes from switchgrass lines carrying a *pShOMT::GUS* construct. Nevertheless, an apparent reduction of lignin content observed in some discrete regions of leaf blades (i.e., fibers on the adaxial zone) from plants carrying the *pShOMT::QsuB* construct indicate that *pShOMT* is also active in leaf cells with secondary wall accumulation (Fig. 4b). In addition to *pShOMT*, attempts to generate transgenic switchgrass lines with constructs containing *QsuB* under the control of the constitutive promoter of the maize ubiquitin1 gene (*pZmUbi-1*) was unsuccessful, whereas only a single event was obtained





with a *pZmCesa10::QsuB* construct containing the promoter of the maize cellulose synthase gene *CESA10* involved in secondary cell wall formation [13] (Figures S2, S3). This is possibly the result of toxicity occurring during the plant regeneration process when using these two *pZmUbi-1::QsuB* and *pZmCesa10::QsuB* constructs. Considering that QsuB diverts lignin biosynthesis, using the promoter of a lignin biosynthetic gene to drive *QsuB* expression may be more suited spatial-temporally during plant development. Interestingly, the single *pZmCesa10::QsuB* line showed a reduction of total lignin content as well as reduced phloroglucinol staining in leaf fibers (Figure S2E, F). Obtaining more switchgrass transgenic events with the *pZmCesa10::QsuB* construct will be essential to validate the effectiveness of *pZmCesa10* in driving QsuB expression to reduce lignin content.

The exact mechanism by which QsuB expression reduces lignin in switchgrass is still unresolved; in particular, whether the cytosolic pools of shikimate—required for HCT activity—and *p*-coumaroyl-shikimate are reduced remain to be demonstrated. Similarly, it would be interesting to determine the lignin monomeric composition in the different QsuB switchgrass lines, especially the relative amount of *p*-hydroxyphenyl (H) units, which is known to be higher in Arabidopsis QsuB plants and typically increases in HCT down-regulated dicot species [8, 14–20]. Furthermore, the recent discovery in several plant species—including switchgrass—of genes encoding putative 3-

hydroxylases (C3H) that convert *p*-coumarate to caffeate, as well as genetic evidence of their role in lignin formation in *Brachypodium distachyon*, question the exclusive role of HCT and the involvement of *p*-coumarate esters during lignin biosynthesis in monocots [21].

The overproduction of PCA in switchgrass lines expressing QsuB probably results from a partial conversion of the endogenous pool of 3-dehydroshikimate catalyzed by QsuB activity. Notably, increases in PCA titers (2–3-fold compared to control switchgrass) are smaller than those previously reported in Arabidopsis and tobacco plants containing the QsuB gene under the control of the promoter of the Arabidopsis cinnamate 4-hydroxylase gene (*pAtCAH*), which were at least two orders of magnitude higher compared to controls plants [8, 22]. In connection with these observations, it has been demonstrated in vitro that PCA acts as a competitive inhibitor of at least one HCT isoform from switchgrass (i.e., PvHCT2) [23]. Therefore, it would be informative to attempt to identify putative *p*-coumaroyl-protocatechuate conjugates in metabolite extracts from *pShOMT::QsuB* switchgrass to determine if such HCT promiscuous activity—and possibly HCT inhibition—also occurs in vivo. Finally, it is promising to observe that the QsuB engineering strategy has the potential to enhance PCA titers in switchgrass biomass because several techno-economic analyses demonstrated the benefits of producing co-products *in planta* to render bioenergy crops economically sustainable [1, 24, 25]. In fact, several studies have already reported on the use of PCA as carbon source or pathway intermediate for the biological synthesis of diverse valuable products such as beta-ketoadipic acid, muconolactone, muconic acid, 2-pyrone-4,6-dicarboxylic acid, bisabolene, and methyl ketones [22, 26–30].

## Conclusion

The QsuB engineering approach has been established in switchgrass. This work highlights the fact that selecting an adequate promoter to drive *QsuB* expression should be an important parameter for successful engineering of other crops with this gene via tissue culture-dependent transformation methods. Considering that *pShOMT* activity is induced in the leaf and root by key regulators of biotic and abiotic stress responses such as salicylic acid, jasmonic acid and methyl jasmonate [12], it will be essential to field test our engineered *pShOMT::QsuB* switchgrass to assess its agronomic performance and resilience to environmental stress.

## Methods

### Vector construction and plant transformation

The promoters *pShOMT* [12], *pZmCesa10* (2.6 kb located upstream the start codon of the maize *CESA10*

gene - GenBank: AY372244.1), and *pZmUbi-1* [31] were synthesized with the following flanking restriction sites: 5'-AscI / 3'-AvrII for *pShOMT* and 5'-HindIII / 3'-AvrII for *pZmCesa10* and *pZmUbi-1* (Genscript, Piscataway, NJ). Promoter sequences were released by enzyme digest and ligated into the binary vector pA6-GW [32] pre-digested with either AscI/AvrII or HindIII/AvrII to generate respectively the *pA6-pShOMT-GW*, *pZmCesa10-GW*, and *pA6-pZmUbi-1-GW* binary vectors. The entry vector pDONR221-schl::QsuB containing the gene encoding the 3-dehydroshikimate dehydratase QsuB from *Corynebacterium glutamicum* preceded with the nucleotide sequence of a chloroplast transit peptide [8] was LR recombined with the *pA6-pShOMT-GW*, *pA6-pZmCesa10-GW*, and *pA6-pZmUbi-1-GW* vectors using the Gateway cloning technology (Thermo Fisher Scientific, Waltham, MA) to generate the constructs *pA6-pShOMT-schl::QsuB*, *pA6-pZmCesa10-GW-schl::QsuB*, and *pA6-pZmUbi-1-GW-schl::QsuB*, respectively. A nucleotide sequence encoding the beta-glucuronidase gene (GUS) from *E. coli* was amplified from pCAM-BIA1301 using primers flanked with attB1 (5') and attB2 (3') Gateway recombination sites, and inserted into the *pA6-pShOMT-GW* and *pA6-pZmCesa10-GW* vectors by Gateway cloning to generate the constructs *pA6-pShOMT::GUS* and *pA6-pZmCesa10::GUS*, respectively. Cloning primers are listed in Table S1. The binary vectors were transformed into *Agrobacterium tumefaciens* strain AGL1 for switchgrass (*Panicum virgatum* L.) transformation which was performed at the University of Missouri's Plant Transformation Core Facility as previously described [33], where embryogenic calli used for transformation were induced from mature seeds of switchgrass cultivar Alamo-A4 (Hancock Farm & Seed Company, Dade City, FL). Hygromycin B (Life Technologies, Foster City, CA) was added to the selection medium at 50 mg/L.

### Plant growth conditions

Four transgenic switchgrass plants for each event were transferred to 2-gal pots containing Pro-Mix soil and grown in a room at 22 °C and 60% humidity using a light intensity of 250  $\mu\text{mol}/\text{m}^2/\text{s}$  and 16 h of light per day.

### PCR genotyping

Genomic DNA was extracted from leaf tissue obtained from one of the clones from each event using the Plant DNeasy plant mini kit (Qiagen, Carlsbad, CA). PCR primers specific to the *QsuB* gene were used to detect the transgene, and primers specific to the switchgrass *PvUBQ6* gene (GenBank: FE609298.1) were used to assess the quality of the gDNA. All the primers used in this study are listed in Table S1.

### RT-qPCR

Total RNAs were extracted from the first internode collected from plants at the E2 stage [34] using the TRIzol reagent (Thermo Fisher Scientific, Waltham, MA) and cDNA synthesis was conducted using the high-capacity cDNA reverse transcription kit (Applied BioSystems, Foster City, CA) as previously described [35]. RT-qPCR was performed as described previously using 40 cycles consisting of 5 s at 95 °C for denaturation and 15 s at 60 °C for annealing and amplification [35]. The relative quantification of *QsuB* transcripts was calculated using the  $2^{-\Delta\text{CT}}$  method and normalized to the reference gene *PvUBQ6* (GenBank: FE609298.1). The results are the average from two biological replicates which were each analyzed in technical replicates. RT-qPCR primers are listed in Table S1.

### Lignin assays

The Wiesner histochemical test using phloroglucinol-HCl, a reagent that reacts with coniferaldehyde groups in lignin, was performed on transverse sections of stems and leaf blades from plants at the E2 stage as previously described [36, 37]. For Klason lignin measurements, whole switchgrass plants were cut at the E5 stage (no visible flag leaf) 3 cm from the bottom, and biomass was dried in an oven at 50 °C for 7 days. Dried biomass was grinded with a Model 4 Wiley Mill equipped with a 1-mm mesh (Thomas Scientific, Swedesboro, NJ). Grinded biomass was extracted as previously described [8] and Klason lignin was measured using the standard NREL biomass protocol [38].

### Saccharification assays

Grinded and extracted biomass obtained from plants at the E5 stage was ball-milled to a fine powder using a Mixer Mill MM 400 (Retsch Inc., Newtown, PA) and stainless-steel balls. For saccharification assays, four biological replicates of 10 mg of fine biomass powder from each line was pretreated with liquid hot water followed by a 72-h enzymatic hydrolysis using 1% w/w Cellic CTec2 enzyme mixture (Novozymes, Denmark) as previously described [35]. Hydrolysates were used for measurement of reducing sugars using the 3,5-dinitrosalicylic acid (DNS) assay [39].

### Protocatechuate measurements

Whole switchgrass plants were cut 3 cm from the bottom at the E5 stage (no visible flag leaf), and biomass was dried in an oven at 50 °C for 7 days. Dried biomass was grinded with a Model 4 Wiley Mill equipped with a 1-mm mesh (Thomas Scientific, Swedesboro, NJ). An aliquot of the grinded biomass was ball-milled to a fine powder using a Mixer Mill MM 400 (Retsch Inc., Newtown, PA) and stainless-steel

balls. Metabolites were extracted from 200 mg of dried ball-milled biomass using 80% (v/v) methanol: water followed by an acid hydrolysis step as previously described [8]. Protocatechuic acid was detected in metabolite extracts using high-performance liquid chromatography (HPLC), electrospray ionization (ESI), and time-of-flight (TOF) mass spectrometry (MS) as previously described [40]. Quantification was performed using a six-point calibration curve from protocatechuic acid solutions prepared with an authentic standard (Sigma-Aldrich, St. Louis, MO).

### Histochemical GUS assays

Stem and leaf sections were obtained manually from plants at the E4 stage using a razor blade. GUS assays were conducted on plant sections using 2 mM 5-bromo-4-chloro-3-indolyl- $\beta$ -D-glucuronide (Sigma-Aldrich, St. Louis, MO) as substrate for 48 h at 37 °C as previously described [41]. After incubation, sections were dehydrated in 95% (v/v) ethanol prior to observation of the GUS staining in 70% (v/v) ethanol.

### Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12870-021-02842-9>.

**Additional file 1: Figure S1.** Representative pictures showing GUS activities in various tiller sections of switchgrass lines harboring the *pShOMT::GUS* construct. GUS expression is specifically observed in stem nodes. Scale: White bars = 2 mm, black bar = 400  $\mu$  m. N: node; IN: internode, IS: internode transverse section.

**Additional file 2: Figure S2.** Characterization of a switchgrass line harboring the *pZmCesa10::QsuB* construct. **(A)** Representative pictures showing GUS activities in various tiller sections of switchgrass lines harboring the *pZmCesa10::GUS* construct. GUS expression is mostly observed in internodes, especially in developing vascular bundles (red arrows). Scale: White bars = 2 mm, black bar = 400  $\mu$  m. N: node; IN: internode, IS: internode transverse section. **(B)** Detection of the *QsuB* gene by PCR in line *pZmCesa10::QsuB-5*. **(C)** Detection of *QsuB* transcripts by RT-qPCR. *QsuB* expression levels relative to that of *PvUBQ6* are shown. Values are means  $\pm$ SD of two biological replicates ( $n = 2$ ). **(D)** Protocatechuic acid (PCA) content measured in the biomass of the switchgrass line *pZmCesa10::QsuB-5*. A line containing the *pZmCesa10::GUS* construct was used as control. Values are means  $\pm$ SE of three biological replicates ( $n = 3$ ). Asterisks indicate a significant difference from the control using the unpaired Student's t-test ( $*P < 0.001$ ). **(E)** Klason lignin content measured in cell wall residues (CWR) obtained from the biomass of the switchgrass line *pZmCesa10::QsuB-5*. A line containing the *pZmCesa10::GUS* construct was used as control. Values are means  $\pm$ SE of four biological replicates ( $n = 4$ ). Asterisks indicate a significant difference from the control using the unpaired Student's t-test ( $*P < 0.05$ ). **(F)** Representative pictures of stem and leaf blade cross-sections stained with phloroglucinol-HCl from line *pZmCesa10::QsuB-5* and a line containing the *pZmCesa10::GUS* construct. Note in the leaves the reduction of the staining specifically in thick fibers located in both the adaxial and abaxial zones for the line *pZmCesa10::QsuB-5* (red arrows).

**Additional file 3: Table S1.** Oligonucleotides used in the study.

**Additional file 4: Figure S3.** Full length unprocessed images of PCR gels used for Figs. 2a and S2B. Note that the seven transformants obtained with the *pZmUbi-1::QsuB* construct were all false positives and did not contain the *QsuB* gene (see purple rectangle on the PCR gel).

### Abbreviations

CWR: Cell wall residue; GUS: Beta-glucuronidase; HCT: hydroxycinnamoyl-CoA: shikimate hydroxycinnamoyl transferase; HPLC-ESI-TOF-MS: High-performance liquid chromatography electrospray ionization and time-of-flight mass spectrometry; PCA: Protocatechuic acid; RT-qPCR: Real-time quantitative reverse transcription PCR

### Acknowledgements

Authors are grateful to Dr. Zhanyuan J. Zhang and the staff of the University of Missouri's Plant Transformation Core Facility for performing the switchgrass transformation work, and to Novozymes for providing Cellic CTec2.

### Authors' contributions

ZH and SY grew the plants; ZH, SY, and TW conducted the PCR genotyping and RT-qPCR; ZH and SY conducted GUS and phloroglucinol-HCl staining assays; AS and SY performed Klason lignin measurements; AS conducted metabolite extractions; GW and EEKB performed HPLC-ESI-TOF-MS analyses; VTB designed the plant binary vectors; SY performed saccharification assays; AE wrote the manuscript; ZH, PCR, HVS, and DL edited the manuscript; PCR, HVS, DL, and AE supervised the research. All authors have read and approved the manuscript.

### Funding

This work was part of the DOE Joint BioEnergy Institute (<http://www.jbeii.org>) supported by the U. S. Department of Energy, Office of Science, Office of Biological and Environmental Research, through contract DE-AC02-05CH11231 between Lawrence Berkeley National Laboratory and the U. S. Department of Energy. The United States Government retains and the publisher, by accepting the article for publication, acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this manuscript, or allow others to do so, for United States Government purposes. The funding bodies were not involved in the design of the study, data collection, interpretation of data, or in writing the manuscript.

### Availability of data and materials

The authors ensure the availability of supporting data and materials. The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

### Ethics approval and consent to participate

Not applicable.

### Consent for publication

Not applicable.

### Competing interests

AE and DL are authors of a patent related to the research (US10415052B2).

### Author details

<sup>1</sup>Joint BioEnergy Institute, Emeryville, CA 94608, USA. <sup>2</sup>Environmental Genomics and Systems Biology Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA. <sup>3</sup>Department of Plant Pathology and the Genome Center, University of California, Davis, CA 95616, USA. <sup>4</sup>Present address: State Key Laboratory of Agricultural Genomics, BGI-Shenzhen, Shenzhen 518000, China. <sup>5</sup>Biological Systems and Engineering Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA. <sup>6</sup>Department of Plant and Microbial Biology, University of California-Berkeley, Berkeley, CA 94720, USA.

Received: 31 July 2020 Accepted: 13 January 2021

Published online: 21 January 2021

### References

- Baral NR, Sundstrom ER, Das L, Gladden J, Eudes A, Mortimer JC, et al. Approaches for more efficient biological conversion of lignocellulosic feedstocks to biofuels and bioproducts. ACS Sustain Chem Eng. 2019;7:9062–79.
- Markel K, Belcher MS, Shih PM. Defining and engineering bioenergy plant feedstock ideotypes. Curr Opin Biotech. 2020;62:196–201.



3. Sanderson MA, Reed RL, McLaughlin SB, Wullschlegel SD, Conger BV, Parrish DJ, et al. Switchgrass as a sustainable bioenergy crop. *Bioresour Technol*. 1996;56:83–93.
4. Bouton J. Improvement of switchgrass as a bioenergy crop. In: Vermerris W, editor. *Genetic Improvement of Bioenergy Crops*. New York: Springer; 2008. p. 309–45.
5. Lin CY, Donohoe BS, Ahuja N, Garrity DM, Qu R, Tucker MP, et al. Evaluation of parameters affecting switchgrass tissue culture: toward a consolidated procedure for *Agrobacterium*-mediated transformation of switchgrass (*Panicum virgatum*). *Plant Methods*. 2017;13:113.
6. Eudes A, Liang Y, Mitra P, Loqué D. Lignin bioengineering. *Curr Opin Biotech*. 2014;26:189–98.
7. Halpin C. Lignin engineering to improve saccharification and digestibility in grasses. *Curr Opin Biotech*. 2019;56:223–9.
8. Eudes A, Sathitsuksanoh N, Baidoo EEK, George A, Liang Y, Yang F, et al. Expression of a bacterial 3-dehydroshikimate dehydratase reduces lignin content and improves biomass saccharification efficiency. *Plant Biotechnol J*. 2015;13:1241–50.
9. Escamilla-Treviño LL, Shen H, Hernandez T, Yin Y, Xu Y, Dixon RA. Early lignin pathway enzymes and routes to chlorogenic acid in switchgrass (*Panicum virgatum* L.). *Plant Mol Biol*. 2014;84:565–76.
10. Shen H, Mazarei M, Hisano H, Escamilla-Treviño L, Fu C, Pu Y, et al. A genomics approach to deciphering lignin biosynthesis in switchgrass. *Plant Cell*. 2013;25:4342–61.
11. Nelson RS, Stewart CN, Gou J, Holladay S, Gallego-Giraldo L, Flanagan A, et al. Development and use of a switchgrass (*Panicum virgatum* L.) transformation pipeline by the BioEnergy Science Center to evaluate plants for reduced cell wall recalcitrance. *Biotechnol Biofuels*. 2017;10:309.
12. Damaj MB, Kumpatla SP, Emani C, Beremand PD, Reddy AS, Rathore KS, et al. Sugarcane DIRIGENT and O-METHYLTRANSFERASE promoters confer stem-regulated gene expression in diverse monocots. *Planta*. 2010;231:1439–58.
13. Appenzeller L, Doblin M, Barreiro R, Wang H, Niu X, Kollipara K, et al. Cellulose synthesis in maize: isolation and expression analysis of the cellulose synthase (CesA) gene family. *Cellulose*. 2004;11:287–99.
14. Ha CM, Fine D, Bhatia A, Rao X, Martin MZ, Engle NL, et al. Ectopic defense gene expression is associated with growth defects in *Medicago truncatula* lignin pathway mutants. *Plant Physiol*. 2019;181:63–84.
15. Hoffmann L, Besseau S, Geoffroy P, Ritzenthaler C, Meyer D, Lapierre C, et al. Silencing of hydroxycinnamoyl-coenzyme a shikimate/quininate hydroxycinnamoyltransferase affects phenylpropanoid biosynthesis. *Plant Cell*. 2004;16:1446–65.
16. Liang Y, Eudes A, Yogiswara S, Jing B, Benites VT, Yamanaka R, et al. A screening method to identify efficient sgRNAs in Arabidopsis, used in conjunction with cell-specific lignin reduction. *Biotechnol Biofuels*. 2019;12:130.
17. Peng XP, Sun SL, Wen JL, Yin WL, Sun RC. Structural characterization of lignins from hydroxycinnamoyl transferase (HCT) down-regulated transgenic poplars. *Fuel*. 2014;134:485–92.
18. Shadle G, Chen F, Srinivasa Reddy MS, Jackson L, Nakashima J, Dixon RA. Down-regulation of hydroxycinnamoyl CoA: Shikimate hydroxycinnamoyl transferase in transgenic alfalfa affects lignification, development and forage quality. *Phytochemistry*. 2007;68:1521–9.
19. Vanholme B, Cesarino I, Goeminne G, Kim H, Marroni F, Van Acker R, et al. Breeding with rare defective alleles (BRDA): a natural *Populus nigra* HCT mutant with modified lignin as a case study. *New Phytol*. 2013;198:765–76.
20. Zhou X, Yang S, Lu M, Zhao S, Cai L, Zhang Y, et al. Structure and monomer ratio of lignin in C3H and HCT RNAi transgenic poplar saplings. *ChemistrySelect*. 2020;5:7164–9.
21. Barros J, Escamilla-Treviño L, Song L, Rao X, Serrani-Yarce JC, Palacios MD, et al. 4-Coumarate 3-hydroxylase in the lignin biosynthesis pathway is a cytosolic ascorbate peroxidase. *Nat Commun*. 2019;10:1994.
22. Wu W, Dutta T, Varman AM, Eudes A, Manalansan B, Loqué D, et al. Lignin valorization: two hybrid biochemical routes for the conversion of polymeric lignin into value-added chemicals. *Sci Rep*. 2017;7:8420.
23. Eudes A, Pereira JH, Yogiswara S, Wang G, Teixeira Benites V, Baidoo EEK, et al. Exploiting the substrate promiscuity of hydroxycinnamoyl-CoA: shikimate hydroxycinnamoyl transferase to reduce lignin. *Plant Cell Physiol*. 2016;57:568–79.
24. Konda NVSNM, Loqué D, Scown CD. Towards economically sustainable lignocellulosic biorefineries. In: Kumar R, Singh S, editors. *Valorization of lignocellulosic biomass in a biorefinery: from logistics to environmental and performance impact*. Balan V: Nova Publishers; 2016. p. 321–38.
25. Yang M, Baral NR, Simmons BA, Mortimer JC, Shih PM, Scown CD. Accumulation of high-value bioproducts *in planta* can improve the economics of advanced biofuels. *Proc Natl Acad Sci U S A*. 2020;117:8639–48.
26. Lin CY, Eudes A. Strategies for the production of biochemicals in bioenergy crops. *Biotechnol Biofuels*. 2020;13:71.
27. Okamura Y, Abe T, Nishimura K, Kawata Y, Sato-Izawa K, Otsuka Y, et al. Beta-ketoadipic acid and muconolactone production from a lignin-related aromatic compound through the protocatechuate 3,4-metabolic pathway. *J Biosci Bioeng*. 2016;121:652–8.
28. Otsuka Y, Nakamura M, Shigehara K, Sugimura K, Masai E, Ohara S, et al. Efficient production of 2-pyrone 4,6-dicarboxylic acid as a novel polymer-based material from protocatechuate by microbial function. *Appl Microbiol Biotechnol*. 2006;71:608–14.
29. Rodriguez A, Ersig N, Geiselman GM, Seibel K, Simmons BA, Magnuson JK, et al. Conversion of depolymerized sugars and aromatics from engineered feedstocks by two oleaginous red yeasts. *Bioresour Technol*. 2019;228:121365.
30. Dong J, Chen Y, Benites VT, Baidoo EEK, Petzold CJ, Beller HR, et al. Methyl ketone production by *Pseudomonas putida* is enhanced by plant-derived amino acids. *Biotechnol Bioeng*. 2019;116:1909–2.
31. Christensen AH, Quail PH. Ubiquitin promoter-based vectors for high-level expression of selectable and/or screenable marker genes in monocotyledonous plants. *Transgenic Res*. 1996;5:213–9.
32. Yang F, Mitra P, Zhang L, Prak L, Verherbruggen Y, Kim JS, et al. Engineering secondary cell wall deposition in plants. *Plant Biotechnol J*. 2013;11:325–35.
33. Do PT, De Tar JR, Lee H, Folta MK, Zhang ZJ. Expression of ZmGA20ox cDNA alters plant morphology and increases biomass production of switchgrass (*Panicum virgatum* L.). *Plant Biotechnol J*. 2016;14:1532–40.
34. Hardin CF, Fu C, Hisano H, Xiao X, Shen H, Stewart CN, et al. Standardization of switchgrass sample collection for cell wall and biomass trait analysis. *Bioenergy Res*. 2013;6:755–62.
35. Li G, Jones KC, Eudes A, Pidatala VR, Sun J, Xu F, et al. Overexpression of a rice BAHD acyltransferase gene in switchgrass (*Panicum virgatum* L.) enhances saccharification. *BMC Biotechnol*. 2018;18:54.
36. Hao, Monhen. A review of xylan and lignin biosynthesis: Foundation for studying Arabidopsis irregular xylem mutants with pleiotropic phenotypes. *Crit Rev Biochem Mol Biol*. 2014;49:212–41.
37. Hao Z, Avci U, Tan L, Zhu X, Glushka J, Pattathil S, et al. Loss of Arabidopsis GAUT12/IRX8 causes anther indehiscence and leads to reduced G lignin associated with altered matrix polysaccharide deposition. *Front Plant Sci*. 2014;5:357.
38. Sluiter A, Hames B, Ruiz R, Scarlata C, Sluiter J. Determination of structural carbohydrates and lignin in biomass. In: *Laboratory Analytical Procedure*. Technical Report, NREL /TP-510-42618. Golden: National Renewable Energy Laboratory; 2008.
39. Miller GL. Use of dinitrosalicylic acid reagent for determination of reducing sugar. *Anal Chem*. 1959;31:426–8.
40. Eudes A, Juminaga D, Baidoo EEK, Collins FW, Keasling JD, Loqué D. Production of hydroxycinnamoyl anthranilates from glucose in *Escherichia coli*. *Microb Cell Factories*. 2013;12:62.
41. Jefferson RA, Kavanagh TA, Bevan MW. GUS fusions: beta-glucuronidase as a sensitive and versatile gene fusion marker in higher plants. *EMBO J*. 1987;6:3901–7.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.