

UC Irvine

Clinical Practice and Cases in Emergency Medicine

Title

Steal Phenomenon with Tonsillar Arteriovenous Malformation

Permalink

<https://escholarship.org/uc/item/71q2p69q>

Journal

Clinical Practice and Cases in Emergency Medicine, 3(3)

Authors

Amin, Manish
Chaddha, Krishan
Aguñiga-Navarrete, Phillip
et al.

Publication Date

2019

DOI

10.5811/cpcem.2019.5.42882

Copyright Information

Copyright 2019 by the author(s). This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Steal Phenomenon with Tonsillar Arteriovenous Malformation

Manish Amin, DO
 Krishan Chaddha, DO
 Phillip Aguiñiga-Navarrete, RA
 Sudha Challa, MD
 Madison B. Garrett, RA

Kern Medical, Department of Emergency Medicine, Bakersfield, California

Section Editor: Christopher Sampson, MD

Submission history: Submitted February 27, 2019; Revision received May 1, 2019; Accepted May 16, 2019

Electronically published July 1, 2019

Full text available through open access at http://escholarship.org/uc/uciem_cpem

DOI: 10.5811/cpcem.2019.5.42882

Cranial vascular malformations can cause symptoms of headache, stroke, transient ischemic attack, or other cerebrovascular disorders due to steal phenomenon. Subclavian steal phenomenon is a localized change in cerebral perfusion from a cranial arteriovenous malformation (AVM). We present the only recorded case of a tonsillar AVM causing a transient ischemic attack due to steal phenomenon. [Clin Pract Cases Emerg Med. 2019;3(3):295-296.]

CASE PRESENTATION

A 35-year-old female presented to the ED with left arm and leg weakness. The patient had normal vital signs. Symptoms started 90 minutes before arrival. Past medical history included a questionable transient ischemic attack (TIA) two years prior. Physical examination noted enlargement of the right tonsillar region. The patient had four of five strength of the left upper and lower extremities with decreased light touch and pain sensation. Her National Institutes of Health Stroke Scale was one. Computed tomography (CT) of the brain was normal. CT angiogram of the brain and neck noted asymmetrical enlargement of the right pharyngeal tonsil associated with vessels and calcifications within the right tonsillar region (Image). Magnetic resonance imaging of the brain was normal. Neurology was consulted and the diagnosis of TIA from tonsillar arteriovenous malformation (AVM) was made.

DISCUSSION

This is the only reported case of tonsillar AVM with an associated TIA. Venous malformations are common types of vascular malformations that present in infancy and expand throughout a patient's lifespan. They typically present in a focal region, with 40% of them occurring in the head and neck.¹ Vascular malformations noted within the cranium can contribute to symptoms of headache, stroke, TIA, or other cerebrovascular disorders.² Intracranial AVMs are known to cause TIA symptoms due to steal phenomenon, which is a localized change




Image. Axial computed tomographic angiography of the head showing palatine tonsillar arteriovenous malformation.

in perfusion from an AVM. Use of antiplatelet agents in TIAs is evidence-based and reduces the possibility of recurrence of neurologic deficits in patients who have had TIAs. However, this poses a problem in patients known to have AVMs, as these agents can cause a higher rate of complications and rupture with AVMs.³

The steal phenomenon noted with intracranial AVMs has not been known to be associated with AVMs that are located outside the cranium. This may be an area of research for the future.

Documented patient informed consent and/or Institutional Review Board approval has been obtained and filed for publication of this case report.

Address for Correspondence: Manish Amin, DO, Kern Medical Center, Department of Emergency Medicine, 1700 Mount Vernon Ave, Bakersfield, CA 93306. Email: manish.amin@kernmedical.com.

Conflicts of Interest: By the CPC-EM article submission agreement, all authors are required to disclose all affiliations, funding sources and financial or management relationships that could be perceived as potential sources of bias. Dr. Amin serves as an Associate Editor for CPC-EM.

Copyright: © 2019 Amin et al. This is an open access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) License. See: <http://creativecommons.org/licenses/by/4.0/>

REFERENCES

1. Amato MV, Patel NA, Hu S, et al. Sporadic multifocal venous malformations of the head and neck. *Case Rep in Otolaryngol.* 2015;2015:508149.
2. Carolei A and Sacco S. Headache attributed to stroke, TIA, intracerebral haemorrhage, or vascular malformation. *Handb Clin Neurol.* 2010;97:517-28.
3. Jabbari E, Sethuraman S, Sekaran L. The clinical dilemma of treating transient ischaemic attack-like symptoms in patients with coexisting arteriovenous malformation. *BMJ Case Rep.* 2014;2014. pii: bcr2013201315.

CPC-EM Capsule

What do we already know about this clinical entity?

It is understood that vascular malformations within the cranium can result in steal phenomenon, transient ischemic attack (TIA) and stroke-like symptoms.

What is the major impact of the image(s)?

This atypical case shows arteriovenous malformation in the palatine tonsillar region, where a correlation to steal phenomenon causing TIA symptoms is not well studied.

How might this improve emergency medicine practice?

This case adds to the differential diagnosis for stroke-type presentations.