UCLA UCLA Previously Published Works

Title

The De Novo Cytosine Methyltransferase DRM2 Requires Intact UBA Domains and a Catalytically Mutated Paralog DRM3 during RNA—Directed DNA Methylation in Arabidopsis thaliana

Permalink

https://escholarship.org/uc/item/74n832mw

Journal PLOS Genetics, 6(10)

ISSN

1553-7390

Authors

Henderson, Ian R Deleris, Angelique Wong, William <u>et al.</u>

Publication Date

2010

DOI

10.1371/journal.pgen.1001182

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <u>https://creativecommons.org/licenses/by/4.0/</u>

Peer reviewed

The *De Novo* Cytosine Methyltransferase DRM2 Requires Intact UBA Domains and a Catalytically Mutated Paralog DRM3 during RNA–Directed DNA Methylation in *Arabidopsis thaliana*

Ian R. Henderson^{1,2}, Angelique Deleris¹, William Wong¹, Xuehua Zhong¹, Hang Gyeong Chin³, Gregory A. Horwitz¹, Krystyna A. Kelly², Sriharsa Pradhan³, Steven E. Jacobsen^{1,4}*

1 Department of Molecular, Cell, and Developmental Biology, University of California Los Angeles, Los Angeles, California, United States of America, 2 Department of Plant Sciences, University of Cambridge, Cambridge, United Kingdom, 3 New England Biolabs, Ipswich, Massachusetts, United States of America, 4 Howard Hughes Medical Institute, University of California Los Angeles, Los Angeles, California, United States of America

Abstract

Eukaryotic DNA cytosine methylation can be used to transcriptionally silence repetitive sequences, including transposons and retroviruses. This silencing is stable between cell generations as cytosine methylation is maintained epigenetically through DNA replication. The *Arabidopsis thaliana* Dnmt3 cytosine methyltransferase ortholog DOMAINS REARRANGED METHYLTRANSFERASE2 (DRM2) is required for establishment of small interfering RNA (siRNA) directed DNA methylation. In mammals PIWI proteins and piRNA act in a convergently evolved RNA-directed DNA methylation system that is required to repress transposon expression in the germ line. *De novo* methylation may also be independent of RNA interference and small RNAs, as in *Neurospora crassa*. Here we identify a clade of catalytically mutated DRM2 paralogs in flowering plant genomes, which in *A.thaliana* we term DOMAINS REARRANGED METHYLTRANSFERASE3 (DRM3). Despite being catalytically mutated, *DRM3* is required for normal maintenance of non-CG DNA methylation, establishment of RNA-directed DNA methylation triggered by repeat sequences and accumulation of repeat-associated small RNAs. Although the mammalian catalytically inactive Dnmt3L paralogs act in an analogous manner, phylogenetic analysis indicates that the DRM and Dnmt3 protein families diverged independently in plants and animals. We also show by site-directed mutagenesis that both the DRM2 N-terminal UBA domains and C-terminal methyltransferase domain are required for normal RNA-directed DNA methylation, supporting an essential targeting function for the UBA domains. These results suggest that plant and mammalian RNA-directed DNA methylation systems consist of a combination of ancestral and convergent features.

Citation: Henderson IR, Deleris A, Wong W, Zhong X, Chin HG, et al. (2010) The *De Novo* Cytosine Methyltransferase DRM2 Requires Intact UBA Domains and a Catalytically Mutated Paralog DRM3 during RNA–Directed DNA Methylation in *Arabidopsis thaliana*. PLoS Genet 6(10): e1001182. doi:10.1371/journal.pgen.1001182

Editor: Joseph R. Ecker, The Salk Institute for Biological Studies, United States of America

Received April 3, 2010; Accepted September 28, 2010; Published October 28, 2010

Copyright: © 2010 Henderson et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: Support was provided by a Leukemia and Lymphoma Special Fellowship (IRH), National Cancer Institute award F32CA137995 (GAH), and National Institutes of Health grant GM60398 (SEJ). SEJ is an investigator of the Howard Hughes Medical Institute. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: jacobsen@ucla.edu

Introduction

Repetitive DNA in eukaryotic genomes is often transcriptionally silent, due to genome defense mechanisms directed against transposons and other mobile DNA [1,2]. In plant genomes repetitive sequences are associated with DNA cytosine methylation [3–8], which is required for transcriptional silencing and suppression of transposon activity [9–12]. As DNA methylation can be maintained epigenetically through DNA replication, this mark serves to stably repress expression of repeated sequences following cell division [2,13]. Gene promoters in *A.thaliana* are generally not DNA methylated, though gene bodies (open reading frames) may contain DNA methylation in the CG sequence context [3–8]. Eukaryotic DNA methylation is catalyzed by cytosine methyltransferases that share ancestry with prokaryotic restriction modification enzymes and are characterized by 10 conserved catalytic motifs [13,14]. During catalysis a methyl group is transferred from the donor molecule S-adenosyl methionine to the carbon-5 position of the cytosine base, using an essential cysteine residue in catalytic motif IV [13,14].

Three functional classes of DNA methyltransferase exist in *A.thaliana*; *METHYLTRANSFERASE1* (*MET1*) (orthologous to mammalian *Dnmt1*) which maintains CG methylation, *CHROMO-METHYLASE3* (*CMT3*) (plant specific) which maintains methylation in non-CG sequence contexts and *DOMAINS REAR-RANGED METHYLTRANSFERASE2* (*DRM2*) (orthologous to *Dnmt3a/Dnmt3b*) which both *de novo* methylates DNA and maintains non-CG methylation redundantly with *CMT3* [2]. The *drm2* mutation blocks all *de novo* DNA methylation driven by repeat containing transgenes [15–17]. This phenotype is shared with mutations in a diverse set of RNA interference and chromatin proteins including *NUCLEAR RNA POLYMERASE D1* (*NRPD1*), *NUCLEAR RNA POLYMERASE (RDR2*), *DICER-LIKE3* (*DCL3*), *ARGO-*

Author Summary

Nuclear DNA quantity varies widely between species and is poorly correlated with gene number. Variation in genome size can be explained by differing amounts of repetitive DNA. Repetitive DNA may be mobile, meaning it can increase its copy number within genomes. To prevent this, plants and animals suppress expression of repeats, often by marking the repeated sequence with a methyl group on cytosine bases. DRM2 is an enzyme capable of establishing this methylation, which can be guided to target sequences by short complementary RNA guides. As repeated sequences are prone to generate short RNAs they are efficiently recognized and silenced. We show that DRM2 requires a related inactive DRM3 protein to normally silence repeated sequences. A similar situation exists in mammals, where active and inactive DNA methyltransferases act together to silence repeats. We also demonstrate that non-catalytic regions of the DRM2 enzyme are functionally important, which we speculate to be involved in targeting the enzyme to the genome. Although plant and mammal RNA-directed DNA methylation systems share key similarities, there are important mechanistic differences, meaning that they are likely to have arisen convergently.

NAUTE4 (AGO4), DEFECTIVE IN RNA DIRECTED DNA METHYLATION 1 (DRD1), DEFECTIVE IN MERISTEM SI-LENCING1 (DMS1), INVOLVED IN DE NOVO 2 (IND2), SUPPRESSOR OF VARIEGATION 3-9 HOMOLOGUE 2 (SUVH2) and SUPPRESSOR OF VARIEGATION 3-9 HOMOLOGUE 9 (SUVH9) [16-34]. These genes act in a pathway that generates short interfering RNAs (siRNAs) which can guide DRM2 DNA methyltransferase activity to homologous genomic sequences [2]. This is reflected in the strong correlation between genomic repeats, DNA methylation and siRNAs in A.thaliana [3-5,7,8]. Small RNAs are also required for DRM2 to maintain non-CG DNA methylation and silencing at a subset of endogenous loci, though this function is partially redundant with a second pathway consisting of CMT3 and the histone H3K9 methyltransferase KRYPTONITE/SUPPRESSOR OF VARIEGATION HOMO-LOGUE4 [17,35-37].

The mammalian DRM2 orthologs, Dnmt3a and Dnmt3b, are required to de novo methylate integrated retroviral sequences and imprinted genes [38,39]. Dnmt3a forms a complex with a noncatalytic paralog Dnmt3L, which is also required for normal patterns of DNA methylation in vivo [40-46]. The C-terminal domains of Dnmt3a and Dnmt3L have been co-crystallized and shown to form a hetero-tetrameric complex, which promotes Dnmt3a methyltransferase activity [42-46]. The N-terminal PHD domain of Dnmt3L interacts with unmethylated H3K4 histone tails and recruits Dnmt3a to specific loci, including imprinted genes [45,46]. Thus de novo methylation mediated by Dnmt3a requires interaction with noncatalytic Dnmt3L, which both stimulates catalysis and recruits the complex to chromatin. Animal PIWI proteins bind germ cell specific 25-30 nucleotide piRNAs, many of which are homologous to repeated sequences [47]. The mouse PIWI mutants miwi, mili and miwi2 are sterile and have defects in transposon de novo methylation in male germ cells, with a phenotype very similar to *dnmt3l* [40,48–52]. Hence, small RNAs are required to target a subset of de novo DNA methylation in mammals, though as plants lack PIWI domains and piRNA, these systems appear to have evolved independently. De novo DNA methylation can also be independent of small RNAs and RNA interference, as in Neurospora crassa [53].

Here we describe a novel protein required for RNA-directed DNA methylation in A.thaliana, DOMAINS REARRANGED METHYLTRANSFERASE3 (DRM3). Based on sequence analysis and genetic observations DRM3 appears to be catalytically mutated DNA methyltransferase paralog. A conserved clade of DRM3 genes is found throughout angiosperm genomes. Although DRM3 cannot compensate for loss of DRM2 function in vivo, it is required for normal establishment and maintenance of RNAdirected DNA methylation and accumulation of specific repeatassociated siRNA. This is analogous to the function of Dnmt3L in mammals, though phylogenetic analysis indicates that DRM3 and Dnmt3L proteins diverged independently in plant and animal lineages. We also demonstrate using site-directed mutagenesis that both the catalytic methyltransferase domain and UBA domains of DRM2 are required for RNA-directed DNA methylation. We speculate that the UBA domains are involved in targeting DRM2 activity during de novo DNA methylation, potentially by recognizing a specific chromatin state. These findings extend parallels between plant and mammalian RNA-directed DNA methylation systems, despite their independent evolution.

Results

A conserved clade of catalytically mutated DRM3 paralogs within angiosperms

The A.thaliana genome encodes a previously uncharacterized protein with high amino acid identity to DRM2, which we term DRM3 (At3g17310) (Figure 1A and 1B). Like DRM2, DRM3 contains N-terminal ubiquitin associated (UBA) domains and a Cterminal region which shares identity with the Dnmt3 cytosine methyltransferase domain, but whose catalytic motifs are rearranged such that motifs VI-X precede I-V (Figure 1A and 1B) [54]. During catalysis cytosine methyltransferases form a covalent bond between a conserved cysteine in motif IV and carbon-6 of the cytosine base [14]. The catalytic cysteine is preceded by an invariant proline, which hydrogen bonds to exo-cyclic NH2 groups of cytosine to promote specific recognition and stabilize interaction between the base and catalytic site [14]. Close inspection of the DRM3 methyltransferase domain reveals the absence of highly conserved residues and notably the invariant proline-cysteine sequence is absent from motif IV (Figure 1B). Additionally, DRM3 lacks a conserved glutamic acid within motif IX and glycine within motif X (Figure 1B). It is likely that the absence of these residues inactivate DRM3 cytosine methyltransferase activity, because a *drm2* mutation alone is sufficient to block *de novo* DNA methylation [15-17].

BLAST searches of the angiosperm Populus trichocarpa and Oryzae sativa genomes reveals a clade of DRM3 proteins shared between these species, indicating that distinct DRM2 and DRM3 proteins are conserved and were present prior to the divergence of the monocots and dicots (Figure 1A). All members of the DRM3 clade lack conserved motif IV proline-cysteine, motif IX glutamic acid and motif X glycine residues (Figure 1B). Although OsDRM3 possesses a cysteine within motif IV, it lacks a preceding proline and therefore is predicted to be catalytically inactive (Figure 1B) [14]. The mammalian Dnmt3L proteins also carry mutations in key catalytic motifs, including the catalytic proline-cysteine sequence in motif IV [36-39] (Figure 1C). However, phylogenetic analysis indicates that the Dnmt3 and DRM protein families diverged independently in plants and mammals (Figure 1A). This scenario is supported by the fact that all DRM proteins share an identical rearrangement of their catalytic motifs (Figure 1B) [54]. As Dnmt3L proteins are important for mammalian de novo DNA





(Oryzae sativa) OsDRM2 Os3g02010, OsDRM3 Os5g04330, (Homo sapiens) HsDnmt3a NP_783328, HsDnmt3b CAB53071, HsDnm3L BAA95556, (*Mus musculus*) MmDnmt3a O88508, MmDnmt3b CAM27225, MmDnmt3L AAH83147 and (Haemophilus parahaemolyticus) Hhal P05102.

doi:10.1371/journal.pgen.1001182.g001

methylation we sought to test whether *DRM3* is analogously required for RNA-directed DNA methylation in *A.thaliana*.

DRM3 is required for maintenance of non-CG DNA methylation

To test whether DRM3 is required for normal DRM2 function we obtained two independent transfer-DNA (T-DNA) insertions in this gene (drm3-1 and drm3-2) (Figure S1). We tested whether the drm3 mutations have a defect in maintenance of non-CG methylation at the MEA-ISR endogenous tandem repeat [36]. Non-CG methylation at MEA-ISR is lost in drm2 and RNAi mutants such as rdr2, which can be assayed by Southern blotting and hybridization after genomic DNA digestion with the methylsensitive restriction endonuclease MspI [36]. The higher molecular weight 4.3 kb band hybridizing to MEA-ISR represents inhibition of MspI digestion by cytosine methylation at a single CHG site [36]. The lower 2 kb band represents digested DNA and in wild type is present at levels comparable to the upper band (Figure 2A) [36]. In drm1 drm2 mutants DNA methylation at this site is eliminated and the 4.3 kb band on the Southern blot is absent (Figure 2A) [36]. Using this assay we observed that both drm3 alleles show a reduction of methylation at MEA-ISR to a level intermediate between wild type and drm1 drm2 (Figure 2A). This methylation phenotype could be complemented by transformation of drm3-1 with a genomic fragment containing the DRM3 gene (Figure 2A). As the Southern blot assay measures methylation at a single CHG site we also sequenced the MEA-ISR locus after bisulfite conversion. This independently corroborated that drm3-1 shows an intermediate level of non-CG methylation at MEA-ISR between that of wild type and drm1 drm2 (Figure 2B, Table S1 and Figure S3). Although significant differences in CG methylation are observed between genotypes, individual CG sites remain highly methylated in all samples (>70%) and reductions in the percentage of methylated sites are less than for non-CG methylation (drm3-1 shows a 16% reduction in CG versus 76% reduction in both types of non-CG methylation, while drm1 drm2 shows an 11% reduction in CG versus reduction to absence of non-CG methylation) (Figure 2B, Table S1 and Figure S3). Furthermore, drm1 drm2 and drm3 do not share developmental phenotypes characteristic of mutants that fail to maintain CG methylation such as met1, ddm1 and vim1 vim2 vim3 [55-57].

To further characterize the drm3 phenotype we performed sodium bisulfite sequencing across the promoter tandem repeats of the FWA endogene. These repeats are cytosine methylated in wild type and show an elimination of non-CG methylation in drm1 drm2 [36]. As for MEA-ISR, we observed that drm3-1 shows a strong reduction of non-CG methylation at FWA to a level intermediate between wild type and drm1 drm2 (Figure 2C, Table S2 and Figure S3). We also analyzed maintenance of DNA methylation at the SINE retroelement AtSN1 using HaeIII restriction endonuclease digestion of genomic DNA followed by quantitative PCR [18,27]. HaeIII digestion is DNA methylation sensitive, meaning that methylated sequences are protected from digestion. In wild type AtSN1 is densely methylated and resistant to HaeIII cleavage, whereas in drm1 drm2 methylation is greatly reduced and AtSN1 DNA is digested and as a consequence not amplified (Figure 2D). We repeated this assay in drm3-1 and again observed an



Figure 2. *DRM3* is required for normal maintenance of non-CG methylation. (A) Southern blots hybridized for the *MEA-ISR* repeat using DNA digested with the methyl-sensitive restriction enzyme *Mspl*. The 4.3 kb band and 2 kb bands represent digested and undigested DNA respectively. (B) Graphical representation of bisulfite sequencing at *MEA-ISR* with % cytosine methylation shown for each genotype. CG sequence contexts are shaded black, CHG are shaded grey and white represents CHH. The 95% confidence limits shown are given by the Wilson score interval. (C) Graphical representation of bisulfite sequencing at the methylated promoter repeats of the *FWA* endogene with % cytosine methylation shown for each genotype and shaded according to sequence context as in (B). (D) Quantitative PCR measurement of *AtSN1* genomic DNA following restriction endouclease digestion with DNA methylation sensitive enzyme *Hae*III. The value on the *y*-axis represents the ratio of DNA measured by quantitative PCR with and without *Hae*III digestion. The data represents three biological replicates and the error bars represent standard error. doi:10.1371/journal.pgen.1001182.g002

intermediate phenotype between wild type and *dm1 dm2*, indicating a defect in maintenance of DNA methylation at *AtSN1* (Figure 2D). We conclude from these observations that DRM3 is required for normal maintenance of non-CG DNA methylation *in vivo*. However, as non-CG methylation is not completely eliminated in *dm3*, it is not absolutely required for DRM2 activity.

DRM2 mRNA and protein expression are independent of DRM3

One explanation for dm3 phenotypes could be an influence on DRM2 expression or protein stability. To test this idea we first measured DRM2 mRNA levels using quantitative reverse-transcriptase PCR. This demonstrated that DRM2 mRNA accumulated normally in dm3 mutants relative to wild type (Figure S1). To test the accumulation of DRM2 protein we crossed a complementing DRM2-Myc transgene into dm3-1 and performed western blotting using α -Myc antibodies [58]. DRM2-Myc protein accumulated normally in dm3-1 relative to wild type (Figure S1). We conclude from these experiments that DRM2 expression is independent of DRM3.

DRM3 is required for normal establishment of RNAdirected DNA methylation

DRM2 is required for both maintenance of non-CG methylation and de novo establishment of methylation in all sequence contexts [15,35,36]. To test whether DRM3 is also required for normal establishment of RNA-directed DNA methylation we used Agrobacterium-mediated transformation with a tandem repeat containing FWA transgene [15,16,59]. In wild type endogenous FWA is transcriptionally silenced by DNA methylation at a set of promoter tandem repeats [59]. Loss of methylation at these repeats causes a dominant late-flowering phenotype, which can be quantified by counting leaf number before flowering [59]. When FWA is transformed into wild type A.thaliana the promoter repeats on the incoming transgene are efficiently methylated, causing transcriptional silencing [15,16,59]. However, FWA transformation into drm2 or RdDM pathway mutants (e.g. nrpd1, ago4, rdr2, dcl3) leads to a failure in de novo methylation, meaning incoming FWA transgenes are expressed causing a dominant late-flowering phenotype in transformed individuals [15,16]. Therefore, we transformed FWA into wild type (Col), drm3-1 and drm1 drm2



Figure 3. *DRM3* is required for normal establishment of RNA-directed DNA methylation. (A) Histograms showing flowering-time distribution of untransformed plants (black bars) or T_1 *FWA* transformant plants (grey bars). Flowering-time is measured as total leaf number and plotted along the *x*-axis, with number of plants on *y*-axis. (B) Graph showing flowering time (total leaf number) of individual T_1 plants connected to the average flowering time of their T_2 progeny. Each colour represents an individual T_1 - T_2 family. doi:10.1371/journal.pgen.1001182.g003

backgrounds and analyzed the flowering time of first generation transformed individuals (T₁) as a measure of establishment of RNA-directed DNA methylation. Untransformed plants were grown alongside +*FWA* transformants as controls and analyzed for flowering time. As previously reported *FWA* transformants in wild type flowered slightly but significantly later than untransformed controls, reflecting incomplete establishment of silencing in the T₁ generation (Figure 3A and Table 1) [15,16]. In contrast, *drm1 drm2 FWA* transformants flowered far later due to a failure to silence the

Table	1. Average	flowering-time	of FWA T ₁	transformant
popula	itions.			

	Total leaf number +/- standard error		
Genotype	Untransformed	+FWA	
Col	17.8+/-0.25	21.5+/-0.83	
drm3-1	17.0+/-0.21	30.5+/-0.82	
drm1 drm2	18.0+/-0.24	41.1+/-0.70	
DRM2-Myc drm1 drm2	15.39+/-0.18	27.8+/-0.39	
DRM2cat-Myc drm1 drm2	17.06+/-0.24	52.86+/-1.13	
DRM2uba-Myc drm1 drm2	18.09+/-0.18	51.51+/-1.06	
Col	17.1+/-0.41	20.0+/-0.23	
drm3-1	21.3+/-0.26	32.0+/-0.68	
dcl3-1	19.1+/-0.25	36.2+/-0.53	
drm3-1 dcl3-1	15.9+/-0.19	42.6+/-1.23	
drm1 drm2	16.7+/-0.23	44.7+/-0.93	

doi:10.1371/journal.pgen.1001182.t001

incoming transgenes (Figure 3A and Table 1) [15,16]. We observed that drm3-1 FWA transformants showed a flowering time distribution intermediate between the wild type and drm1 drm2 T₁ populations (Figure 3A and Table 1). This parallels our observations for maintenance of non-CG methylation with drm3-1 showing a defect in *DRM2*-mediated DNA methylation, though with a phenotype weaker than drm1 drm2.

FWA silencing is frequently incomplete in the T_1 generation and self-fertilization of transformed individuals results in an increase in FWA repeat DNA methylation and transcriptional silencing in the T_2 progeny, which as a consequence flower earlier [59]. As drm3 appears to reduce but not abolish DRM2 activity, we predicted that wild type and drm3-1 T₂ populations would become earlier flowering relative to the T1, whereas drm1 drm2 would not. Analysis of the flowering-time distribution of multiple T₂ populations relative to their T_1 parents showed this to be true (Figure 3B and Table S3). Consistent with a complete block to de novo DNA methylation in drm1 drm2 T₂ individuals did not flower significantly earlier, which they did in the case of wild type and drm3-1 (Figure 3B and Table S3). Together, we interpret this as meaning that although DRM2 is absolutely required for de novo DNA methylation in vivo, normal levels of methyltransferase activity also require non-catalytic DRM3.

Intermediate RNA-directed DNA methylation phenotypes between wild type and *drm1 drm2* have previously been observed in *dcl3* mutants [16,60,61]. DCL3 generates 24 nucleotide siRNAs with a specialized function in RNA-directed DNA methylation and transcriptional silencing, though it acts redundantly with the other DCLs in a locus-specific manner [16,60–62]. We generated *drm3-1 dcl3-1* double mutants and tested for enhancement of defects during establishment of *FWA* RNA-directed DNA methylation, relative to the single mutants. We observed that drm3-1 dcl3-1 double mutants showed a silencing defect comparable to drm1 drm2, which was greater than either drm3-1 or dcl3-1alone (Table 1). We interpret this enhancement as meaning that DRM3 and DCL3 act genetically in parallel to promote DRM2 activity *in vivo*.

Mutagenesis of DRM2 reveals a functional requirement for cytosine methyltransferase and UBA domains

DRM2 and DRM3 show a similar domain organization of Nterminal UBA domains and a C-terminal cytosine methyltransferase domain (Figure 1A). To further understand DRM2 function we performed site-directed mutagenesis of these domains and tested the consequence on RNA-directed DNA methylation. Cytosine methyltransferases require a set of 10 conserved motifs, including motif IV which contains the catalytic cysteine involved in transfer of the methyl group to cytosine (Figure 1B and 1C) [13]. We substituted the DRM2 motif IV catalytic cysteine C587 for an alanine, within an otherwise complementing DRM2 transgene containing a Myc-epitope translational fusion (hereafter termed DRM2_{cat}-Myc) (Figure 4A) [58]. DRM proteins also possess N-terminal UBA domains, which consist of three helices connected by two conserved loop regions forming an ubiquitin interaction surface [63,64]. The first loop contains a highly conserved MGF/MGY motif, which is required for correct folding and maintenance of UBA domain structure [64]. To test the

functional importance of the DRM2 UBA domains we substituting conserved phenylalanine residues (F73, F123, F206) in each loop-I region to alanines (hereafter termed $DRM2_{uba}$ -Myc) (Figure 4A). The wild type DRM2-Myc and mutant $DRM2_{cat}$ -Myc and $DRM2_{uba}$ -Myc transgenes were transformed into drm1 drm2 and tested for their ability to complement mutant RNA-directed DNA methylation phenotypes.

We first tested whether the catalytic and UBA mutations introduced into *DRM2* had any consequence on protein accumulation and localization. Intact *DRM2-Myc drm1 drm2* nuclei immunostained using α -Myc antibodies show signal throughout the nuclei with numerous bright foci and exclusion from the strongly DAPI-staining chromocentres (Figure 4B) [53]. The DRM2_{cat}-Myc and DRM2_{uba}-Myc mutant proteins showed comparable staining patterns to those observed for the unmutated DRM2-Myc protein (Figure 4B). This indicates that the UBA and methyltransferase mutations introduced do not cause a defect in DRM2 protein stability or localization to the nucleus.

We next tested the ability of the *DRM2* transgenes to complement *drm1 drm2* by assaying maintenance of non-CG methylation at *MEA-ISR* using methyl-sensitive *MspI* digestion followed by Southern blotting and hybridization as described earlier. The wild type *DRM2-Myc* transgene complemented *drm1 drm2*, whereas the *DRM2_{cat}-Myc* and *DRM2_{uba}-Myc* lines showed a complete and partial failure to remethylate *MEA-ISR* respectively



Figure 4. DRM2 catalytic and UBA domains are required for maintenance of non-CG methylation. (A) A Myc-epitope tagged *DRM2* transgene was mutated to introduce amino acid substitutions in the UBA domains or the methyltransferase domain, depicted graphically. (B) Micrographs of intact nuclei stained with DAPI and immunostained using α -Myc antibodies. (C) Southern blots hybridized for the *MEA-ISR* repeat using DNA digested with the methyl-sensitive restriction enzyme *Mspl*. (D) Graphical representation of bisulfite sequencing of *MEA-ISR* with % cytosine methylation shown for each genotype. Error bars represent 95% confidence limits (Wilson score interval). CG sequence contexts are shaded dark grey, CHG are shaded light grey and white represents CHH. doi:10.1371/journal.pgen.1001182.g004

(Figure 4C). We confirmed these observations by sequencing MEA-ISR following sodium bisulfite conversion; non-CG methylation was restored by DRM2-Myc but not by $DRM2_{cat}$ -Myc, whereas the $DRM2_{uba}$ -Myc transformants showed a low level of non-CG methylation, comparable to the level observed in drm3-I(Figure 2B, Figure 4D, Table S1 and Figure S3). Therefore, intact UBA and methyltransferase domains are required for DRM2 function during maintenance of non-CG DNA methylation.

To test whether the DRM2 mutations also blocked establishment of DNA methylation we transformed DRM2-Myc dm1 dm2, $DRM2_{cat}$ -Myc dm1 dm2 and $DRM2_{uba}$ -Myc dm1 dm2 with FWA and used the flowering-time distribution within T₁ populations as a measure of silencing. The FWA T₁ populations generated in $DRM2_{cat}$ -Myc dm1 dm2, $DRM2_{uba}$ -Myc dm1 dm2 backgrounds flowered significantly later than the wild type DRM2-Myc dm1 dm2 dm2 dm2 dm2 dm2 dm2 dm2 dm2 dm2 dm3 as meaning that intact UBA and methyltransferase domains are also required to establish RNA-directed DNA methylation and silencing at FWA.

Although *DRM2* is targeted by siRNAs, it is also required for accumulation of specific siRNA populations, for example siRNAs homologous to the 5S ribosomal DNA genes [29], which may reflect a feedback loop between transcriptional silencing and siRNA biogenesis. 5S siRNAs detected by northern blotting and hybridization are absent in *rdr2* and greatly reduced in *dm1 dm2*, relative to wild type (Figure 5B) [29]. The *DRM2-Myc* transgene is able to restore 5S siRNA accumulation in the *drm1 drm2* mutant, but this complementation is not evident with either *DRM2_{cat}-Myc* or *DRM2_{uba}-Myc* (Figure 5B). Consistent with *DRM3* functioning to promote DRM2 activity, we also observed a strong reduction in the accumulation of 5S siRNAs in *drm3-1* mutants (Figure 5B). Together these data demonstrate that both the DRM2 catalytic methyltransferase and UBA domains are required for establishment and maintenance of RNA-directed DNA methylation *in vivo*.

Discussion

A complex pathway of genes has been defined that generates siRNAs homologous to repeated sequences, which can guide DRM2 cytosine methyltransferase activity to target DNA sequences [16-34]. Here we define a new member of this pathway, the catalytically mutated DNA cytosine methyltransferase paralog DRM3, which is required for normal maintenance of non-CG DNA methylation, establishment of RNA-directed DNA methylation triggered by tandem repeats and accumulation of repeat associated siRNAs. As drm3 phenotypes are weaker than drm2 we believe that DRM3 acts to promote activity of DRM2, but is not an absolute requirement. The strength of drm3 phenotypes resembles those observed for *dcl3*, which is defective in accumulation of 24 nt siRNA known to promote RNAdirected DNA methylation. As the drm3-1 dcl3-1 double mutant shows an enhanced phenotype to the level of *drm2*, this suggests that the combined loss of 24 nt siRNA and DRM3 severely hinders DRM2 activity. The drm3 mutation does not influence DRM2 expression or protein accumulation, leading us to believe that DRM3 acts together with DRM2 at the step of catalysis. Indeed, several examples exist of catalytically inactive paralogs acting to stimulate the activity of catalytic partners [65,66]. In this respect the stimulatory interaction between Dnmt3a and Dnmt3L is particularly notable [42–46]. Despite extensive testing we have been unable to find in vitro conditions where A.thaliana DRM2 is active as a cytosine methyltransferase, which has precluded us testing a direct stimulatory effect for DRM3 on catalysis. Development of these assays will be an important next step to



Figure 5. DRM2 catalytic and UBA domains are required for *de novo* DNA methylation and siRNA accumulation. (A) Histograms showing flowering-time distribution of untransformed plants (black bars) or T_1 *FWA* transformant plants (grey bars). Flowering-time is measured as total leaf number and plotted along the *x*-axis, with number of plants on *y*-axis. (B) Northern blotting and hybridization of *5S rDNA* and *microRNA159* (*miR159*) small RNAs. doi:10.1371/journal.pgen.1001182.q005

dissect the function of DRM2 and DRM3 during RNA–directed DNA methylation.

Although an intriguing parallel exists between the functional relationships of DRM2/DRM3 and Dnmt3a/Dnmt3L, phylogenetic analysis strongly suggests that these protein families diverged independently in plants and animals (Figure 1A). There are also important differences between DRM and Dnmt3 in the mechanism of recruitment to genomic target sites. The Dnmt3L protein plays a key role in targeting Dnmt3 via an N-terminal PHD domain, which specifically recognizes histone 3 unmethylated at

lysine 4 [41-46]. By analogy the N-terminal UBA domains of DRM2 may be involved in targeting DNA methyltransferase activity. Supporting this idea we have shown that the DRM2 UBA domains are critical for RNA directed DNA methylation. We do not currently know whether DRM3 UBA domains are also functionally important. Sequence analysis shows that all DRM3 proteins possess two UBA domains, the second of which contains conserved residues known to be important for proper domain folding, including the MGF/MGY motif in loop-I (Figure S2) [64]. Interestingly, the first UBA domain of DRM3 proteins show substitution of the MGF/MGY glycine for either lysine or asparagine, which would be predicted to abolish proper folding [64]. However, as UBA domains are able to bind ubiquitin independently, DRM3 UBA domains may still play a functional role [63,64]. The precise functions of the DRM UBA domains are currently unknown, though it is interesting to speculate that they may recognize specific chromatin modifications during RdDM.

In mammals PIWI proteins together with interacting piRNA act upstream of Dnmt3a/Dnmt3L to DNA methylate homologous repeats during germ line development [40,42-48]. Plants lack orthologs of the PIWI proteins and have not so far been observed to produce piRNA [67]. Thus, small RNA-directed DNA methylation pathways appear to have evolved independently in plants and animals. Equally, the specific rearrangement of methyltransferase catalytic motifs shared between all DRM proteins indicates that DRM and Dnmt3 families have diverged independently in plants and animals (Figure 1). Here we demonstrate a further example of convergence in the parallel evolution of catalytically mutated cytosine methyltransferase paralogs (DRM3 and Dnmt3L) in both lineages. It is also important to note that information other than small RNAs must be involved in targeting DNA methylation, as this mark can be established and maintained in Neurospora crassa RNAi mutants [53], and no evidence exists of small RNAs acting upstream of Dnmt3a in the mammalian female germ line [48]. In conclusion, comparison of DRM3 and Dnmt3L function provides an example of mechanistic convergence between plant and mammalian silencing systems.

Materials and Methods

Plant materials

The drm3-1 allele is T-DNA insertion Salk_136439 and drm3-2 allele is T-DNA insertion Sail_1215_B08. The presence of the drm3-1 insertion can be genotyped by PCR amplification using JP3192 together with a primer to the T-DNA left border JP2410 (all oligonucleotide sequences are listed in Table S4). The presence of the drm3-2 insertion can be genotyped using primers JP3510 and JP2027. The drm1 drm2 and dcl3-1 T-DNA insertions have been described previously [7,16]. DRM3 complementation was performed by PCR amplification of an 8kb DRM3 genomic region from Columbia DNA using oligonucleotides JP3477 and JP3479 and cloned into pCR2.1 (Invitrogen). The DRM3 genomic region then was cloned into the pCAMBIA1300 binary vector as a Sall restriction fragment and used to transform drm3-1 plants by floral dipping using Agrobacterium strain ASE. The DRM2-Myc transgene has been described previously [58]. This construct was mutagenized using Quikchange (Stratagene) using oligonucleotides JP603/ JP604 (catalytic mutation C587A), JP2631/JP2632 (UBA domain 1 F73A), JP2633/JP2634 (UBA domain 2 F123A) and JP2635/ JP2636 (UBA domain 3 F206A).

Phylogenetic tree construction

Multiple sequence alignment was performed using the European Bioinformatics Institute CLUSTALW server (http://www.ebi. ac.uk/clustalw/). NCBI accession numbers or gene numbers for the proteins analyzed are (*Arabidopsis thaliana*) AtDRM1 At5g15380, AtDRM2 At5g14620, AtDRM3 At3g17310, (*Populus trichocarpa*) PtDRM2 DMT905, PtDRM3 DMT907, (*Oryzae sativa*) OsDRM2 Os3g02010, OsDRM3 Os5g04330, (*Homo sapiens*) HsDnmt3a NP_783328, HsDnmt3b CAB53071, HsDnm3L BAA95556, (*Mus musculus*) MmDnmt3a O88508, MmDnmt3b CAM27225, MmDnmt3L AAH83147 and (*Haemophilus parahae-molyticus*) HhaI P05102. The phylogenetic tree was constructed using the MEGA2 program using the neighbor-joining method. Bootstrap values were calculated with 1,000 replicates.

Analysis of DNA methylation

Southern blotting and hybridization were performed using 1µg of genomic DNA digested overnight using MspI and separated by gel electrophoresis using 0.8% agarose and blotted onto N+Hybond (Amersham) nitrocellulose. The MEA-ISR probe was amplified using primers JP980 and JP981. 1µg of genomic DNA was converted using a Methyleasy kit (Human Genetic Signatures) and MEA-ISR amplified using primers JP1026 and JP1027, corresponding to positions 53456-53602 of BAC clone T14P4 [15,36]. PCR products were cloned using the TOPO TA cloning kit (Invitrogen) and sequenced using the T7 primer. The FWA promoter region bisulfite sequenced corresponds to co-ordinates 46062 to 46551 of BAC clone M7J2. This region was amplified using primers JP2004 and JP2005 followed by nested amplification with primers JP2004 and JP4423. During analysis of sodium bisulfite sequencing data the percentage of methylated sites across all clones were calculated for each sequence context. The Wilson score interval was used to find the 95% confidence intervals for these percentages. Percentages were compared with the appropriate wild type percentage using the Pearson chi square test (Table S1 and Table S2). To compensate for multiple testing using the Bonferroni adjustment, a p-value of 0.002 was used as the threshold for statistical significance. To analyze cytosine methylation at AtSN1 genomic DNA was digested with HaeIII in 40 µl alongside controls without restriction endonuclease. Samples were then analyzed by quantitative real-time PCR amplifying with primers JP6699 and JP6700 combined with Biorad SYBR Green Supermix using a MX3000 Stratagene cycler. A ratio was then calculated between the amount of DNA amplified for each sample with and without HaeIII digestion.

FWA transformation and flowering-time analysis

Plants were transformed via floral dipping using Agrobacterium tumefaciens strain ASE carrying a genomic FWA clone in the binary vector pCAMBIA1300. T₁ transformant seed was selected by germination and growth on MS plates containing 50μ g/ml hygromycin followed by transfer to soil under continuous light. Vegetative and cauline leaves were counted until production of the first flowers as a measure of flowering-time. Untransformed plants were grown alongside as controls.

RNA analysis

l gram of floral buds were used to extract and purify small RNAs as described previously [60]. Small RNAs were separated by gel electrophoresis, blotted onto nitrocellulose membrane and hybridized with end-labeled oligonucleotide probes miR159 (5'-TAGAGCTCCCTTCAATCCAAA-3') and 5S siRNA (5'-AT-GCCAAGTTTGGCCTCACGGTCT-3') [61]. End-labeling was performed using T4 polynucleotide kinase (New England Biolabs) and γ -32P-ATP (Amersham). To analyze mRNA accumulation total RNA was extracted using Trizol (Invitrogen) and converted to cDNA as described previously [53]. *DRM2* expression was

analyzed using qRT-PCR with primers JP6069 and JP6070 and *DRM3* analyzed by RT-PCR using primers JP3192 and JP3193.

Immunohistochemistry

Immunostaining of fixed nuclei was performed as previously reported [68].

Supporting Information

Figure S1 DRM2 and DRM3 are expressed independently of one another. (A) Schematic diagram of At3g17310 (DRM3) with exons indicated by open boxes. The position of T-DNA insertions in the *drm3-1* and *drm3-2* alleles is indicated by open triangles. RNA was extracted from wild type (Columbia), *drm1 drm2* and *drm3-1* and used to generate cDNA. The expression of DRM3 was tested by amplifying from this cDNA using primers JP3192 and JP3193 that span the insertion site in *drm3-1*. UBIQUITIN expression was analyzed as a control using amplification with primers JP3483 and JP3484. (B) Quantitative RT-PCR was used to measure mRNA expression levels of DRM2 (dark grey) and ACTIN (light grey) in wild type (Col) and *drm3-1*. (C) Western blotting detection of DRM2-Myc accumulation probing using α -Myc antibodies. Loading was analyzed by re-probing using α -CRY1 antibodies.

Found at: doi:10.1371/journal.pgen.1001182.s001 (0.65 MB PDF)

Figure S2 Sequence analysis of DRM UBA domains. UBA domains are typically ~ 40 amino acids in length and form triple helical bundles [T. D. Mueller, J. Feigon, J Mol Biol 319, 1243 (Jun 21, 2002)]. Hydrophobic patches on the UBA surface are believed to stabilize interaction with hydrophobic surface of fivestranded ubiquitin β-sheet [T. D. Mueller, J. Feigon, J Mol Biol 319, 1243 (Jun 21, 2002)]. The most conserved residues are the MGF/MGY loop which connects helix 1 and 2 and is required for ubiquitin interaction [T. D. Mueller, J. Feigon, J Mol Biol 319, 1243 (Jun 21, 2002)]. The figure shows sequence alignments of DRM protein UBA domains performed using clustalW. Sequences are annotated with reference to solution structures of UBA domains and the presence of conserved residues [T. D. Mueller, J. Feigon, J Mol Biol 319, 1243 (Jun 21, 2002)]. The approximate helix regions are indicated above the sequences by black arrows and joining loop regions indicated by blue arrows. Conserved amino acids are highlighted in colour according to their sidechain; green indicates small (G, A), purple indicates hydrophobic (M, I, L, V), red indicates aromatic (F, Y) and blue indicates amide or acidic (K, N). AtDRM2 is exceptional in possessing three Nterminal UBA domains, whereas all other DRM proteins analyzed possess two recognizable UBA domains. In all DRM proteins the second UBA domain shows the presence of a conserved MGF motif within loop-I and the presence of other conserved hydrophobic residues. However, the first UBA domains in DRM3 proteins differ in that the conserved glycine in loop-I is replaced by either a lysine or asparagine. Mutation of this glycine in other UBA domains is sufficient to abolish interaction with ubiquitin [T. D. Mueller, J. Feigon, J Mol Biol 319, 1243 (Jun 21, 2002)]. Hence, it is possible that the first UBA domain in DRM3 proteins is inactive. However, as UBA domains have been shown to bind ubiquitin independently within a protein [T. D. Mueller, J. Feigon, J Mol Biol 319, 1243 (Jun 21, 2002)], it is possible that the second DRM3 UBA domain is functional.

Found at: doi:10.1371/journal.pgen.1001182.s002 (0.64 MB PDF)

Figure S3 Sodium bisulfite sequencing at *MEA-ISR* and *FWAM*. Graphical representation of sodium bisulfite analysis of the *MEA-ISR* and *FWA* tandem repeats. Sequences are shown with cytosines colored according to their sequence context; CG is blue, CHG is red and CHH is orange. Above the sequence are stacked blocks, each row of which represents an independent sequencing read. Methylation detected in these reads is represented by shading, again colored according to sequence context with CG blue, CHG red and CHH orange. The genotype in each case is stated at the top of each diagram.

Found at: doi:10.1371/journal.pgen.1001182.s003 (0.58 MB PDF)

 Table S1
 Sodium bisulfite sequencing analysis of MEA-ISR. The
region analyzed corresponds to positions 53456 to 53602 of BAC clone T14P4. The MEA-ISR repeat was amplified from sodium bisulfite converted DNA and the frequency of cytosine versus thymine scored. The number of independent clones analyzed is indicated, together with the number of cytosine sites scored, the number observed to be methylated and the methylation frequency expressed as a %. There are 9 CG sites, 2 CHG sites and 24 CHH sites in the amplified region. The 95% confidence limits are given by the Wilson score interval. The p-values are from Pearson chisquare tests comparing each sample with wild type (Col). In addition, drm3-1 was compared with drm1 drm2 for CG (pvalue = 0.28), CHG (p-value = 0.31) and CHH (p-value = 2.9×10^{-5}) sites. Although the percentages for Col, drm3-1 and drm1 drm2 are similar between CHG and CHH methylation, the statistical power of comparisons between the three genotypes is much lower for CHG methylation because of the smaller number of sites analyzed in the sequenced region. Hence, the confidence intervals overlap and the statistical tests fail to reach significance for the CHG context. However, the same pattern is observed for the FWA locus (Table S2), where the difference between Col and drm3-1 is statistically significant. In addition, Southern blotting and hybridization for MEA-ISR following MspI digestion (Figure 2A) measures methylation at the second CHG site and yields supporting observations of CHG methylation being intermediate in drm3-1, between wild type and drm1 drm2.

Found at: doi:10.1371/journal.pgen.1001182.s004 (0.05 MB DOC)

Table S2 Sodium bisulfite sequencing analysis of *FWA*. The region analyzed corresponds to co-ordinates 46062 to 46551 of BAC clone M7J2. The *FWA* repeats were amplified from sodium bisulfite converted DNA and the frequency of cytosine versus thymine scored. The number of independent clones analyzed is indicated, together with the number of cytosine sites scored, the number observed to be methylated and the methylation frequency expressed as a %. There are 12 CG sites, 10 CHG sites and 41 CHH sites in the amplified region. The 95% confidence limits are given by the Wilson score interval. The *p*-values are from Pearson chisquare tests comparing each sample with wild type (Col). In addition, *drm3-1* was compared with *drm1 drm2* for CG (*p*-value<0.007), CHG (*p*-value = <0.40) and CHH (*p*-value<0.002) sites.

Found at: doi:10.1371/journal.pgen.1001182.s005 (0.04 MB DOC)

Table S3Average flowering-time of T_2 FWA transformant lines.Found at:doi:10.1371/journal.pgen.1001182.s006 (0.04 MBDOC)

Table S4Oligonucleotide sequences.

Found at: doi:10.1371/journal.pgen.1001182.s007 (0.04 MB DOC)

Acknowledgments

Thank you to Max Greenberg for providing *FWA* bisulfite sequencing data in *drm1 drm2*. We thank members of the Jacobsen laboratory and anonymous reviewers for helpful comments.

References

- Aravin AA, Bourc'his D (2008) Small RNA guides for de novo DNA methylation in mammalian germ cells. Genes Dev 22: 970–975.
- Law JA, Jacobsen SE (2010) Establishing, maintaining and modifying DNA methylation patterns in plants and animals. Nat Rev Genet 11: 204–220.
- Cokus SJ, Feng S, Zhang X, Chen Z, Merriman B, et al. (2008) Shotgun bisulphite sequencing of the Arabidopsis genome reveals DNA methylation patterning. Nature 452: 215–219.
- Lippman Z, Gendrel AV, Black M, Vaughn MW, Dedhia N, et al. (2004) Role of transposable elements in heterochromatin and epigenetic control. Nature 430: 471–476.
- Lister R, O'Malley RC, Tonti-Filippini J, Gregory BD, Berry CC, et al. (2008) Highly integrated single-base resolution maps of the epigenome in Arabidopsis. Cell 133: 523–536.
- Tran RK, Henikoff JG, Zilberman D, Ditt RF, Jacobsen SE, et al. (2005) DNA methylation profiling identifies CG methylation clusters in Arabidopsis genes. Curr Biol 15: 154–159.
- Zhang X, Yazaki J, Sundaresan A, Cokus S, Chan SW, et al. (2006) Genomewide high-resolution mapping and functional analysis of DNA methylation in arabidopsis. Cell 126: 1189–1201.
- Zilberman D, Gehring M, Tran RK, Ballinger T, Henikoff S (2007) Genomewide analysis of Arabidopsis thaliana DNA methylation uncovers an interdependence between methylation and transcription. Nat Genet 39: 61–69.
- Kato M, Miura A, Bender J, Jacobsen SE, Kakutani T (2003) Role of CG and non-CG methylation in immobilization of transposons in Arabidopsis. Curr Biol 13: 421–426.
- Mirouze M, Reinders J, Bucher E, Nishimura T, Schneeberger K, et al. (2009) Selective epigenetic control of retrotransposition in Arabidopsis. Nature 461: 427–430.
- Miura A, Yonebayashi S, Watanabe K, Toyama T, Shimada H, et al. (2001) Mobilization of transposons by a mutation abolishing full DNA methylation in Arabidopsis. Nature 411: 212–214.
- Tsukahara S, Kobayashi A, Kawabe A, Mathieu O, Miura A, et al. (2009) Bursts of retrotransposition reproduced in Arabidopsis. Nature 461: 423–426.
- Goll MG, Bestor TH (2005) Eukaryotic cytosine methyltransferases. Annu Rev Biochem 74: 481–514.
- Bestor TH, Verdine GL (1994) DNA methyltransferases. Curr Opin Cell Biol 6: 380–389.
- Cao X, Jacobsen SE (2002) Role of the arabidopsis DRM methyltransferases in de novo DNA methylation and gene silencing. Curr Biol 12: 1138–1144.
- Chan SW, Zilberman D, Xie Z, Johansen LK, Carrington JC, et al. (2004) RNA silencing genes control de novo DNA methylation. Science 303: 1336.
- Henderson IR, Jacobsen SE (2008) Tandem repeats upstream of the Arabidopsis endogene SDC recruit non-CG DNA methylation and initiate siRNA spreading. Genes Dev 22: 1597–1606.
- Ausin I, Mockler TC, Chory J, Jacobsen SE (2009) IDN1 and IDN2 are required for de novo DNA methylation in Arabidopsis thaliana. Nat Struct Mol Biol 16: 1325–1327.
- El-Shami M, Pontier D, Lahmy S, Braun L, Picart C, et al. (2007) Reiterated WG/GW motifs form functionally and evolutionarily conserved ARGO-NAUTE-binding platforms in RNAi-related components. Genes Dev 21: 2539–2544.
- He XJ, Hsu YF, Pontes O, Zhu J, Lu J, et al. (2009) NRPD4, a protein related to the RPB4 subunit of RNA polymerase II, is a component of RNA polymerases IV and V and is required for RNA–directed DNA methylation. Genes Dev 23: 318–330.
- Huang L, Jones AM, Searle I, Patel K, Vogler H, et al. (2009) An atypical RNA polymerase involved in RNA silencing shares small subunits with RNA polymerase II. Nat Struct Mol Biol 16: 91–93.
- Johnson LM, Law JA, Khattar A, Henderson IR, Jacobsen SE (2008) SRAdomain proteins required for DRM2-mediated de novo DNA methylation. PLoS Genet 4: e1000280.
- Kanno T, Bucher E, Daxinger L, Huettel B, Bohmdorfer G, et al. (2008) A structural-maintenance-of-chromosomes hinge domain-containing protein is required for RNA-directed DNA methylation. Nat Genet 40: 670–675.
- Kanno T, Huettel B, Mette MF, Aufsatz W, Jaligot E, et al. (2005) Atypical RNA polymerase subunits required for RNA-directed DNA methylation. Nat Genet 37: 761–765.
- Kanno T, Mette MF, Kreil DP, Aufsatz W, Matzke M, et al. (2004) Involvement of putative SNF2 chromatin remodeling protein DRD1 in RNA–directed DNA methylation. Curr Biol 14: 801–805.
- Lahmy S, Pontier D, Cavel E, Vega D, El-Shami M, et al. (2009) PolV(PolIVb) function in RNA-directed DNA methylation requires the conserved active site and an additional plant-specific subunit. Proc Natl Acad Sci U S A 106: 941–946.

Author Contributions

Conceived and designed the experiments: IRH AD WW XZ HGC GAH SP SEJ. Performed the experiments: IRH AD WW XZ HGC GAH. Analyzed the data: IRH AD WW XZ HGC GAH KAK SP SEJ. Contributed reagents/materials/analysis tools: SP. Wrote the paper: IRH SEJ.

- Onodera Y, Haag JR, Ream T, Nunes PC, Pontes O, et al. (2005) Plant nuclear RNA polymerase IV mediates siRNA and DNA methylation-dependent heterochromatin formation. Cell 120: 613–622.
- Pontier D, Yahubyan G, Vega D, Bulski A, Sacz-Vasquez J, et al. (2005) Reinforcement of silencing at transposons and highly repeated sequences requires the concerted action of two distinct RNA polymerases IV in Arabidopsis. Genes Dev 19: 2030–2040.
- Zilberman D, Cao X, Johansen LK, Xie Z, Carrington JC, et al. (2004) Role of Arabidopsis ARGONAUTE4 in RNA-directed DNA methylation triggered by inverted repeats. Curr Biol 14: 1214–1220.
- Bies-Etheve N, Pontier D, Lahmy S, Picart C, Vega D, et al. (2009) RNA– directed DNA methylation requires an AGO4-interacting member of the SPT5 elongation factor family. EMBO Rep 10: 649–654.
- He XJ, Hsu YF, Zhu S, Liu HL, Pontes O, et al. (2009) A conserved transcriptional regulator is required for RNA-directed DNA methylation and plant development. Genes Dev 23: 2717–2722.
- He XJ, Hsu YF, Zhu S, Wierzbicki AT, Pontes O, et al. (2009) An effector of RNA-directed DNA methylation in arabidopsis is an ARGONAUTE 4- and RNA-binding protein. Cell 137: 498–508.
- Herr AJ, Jensen MB, Dalmay T, Baulcombe DC (2005) RNA polymerase IV directs silencing of endogenous DNA. Science 308: 118–120.
- Smith LM, Pontes O, Searle I, Yelina N, Yousafzai FK, et al. (2007) An SNF2 protein associated with nuclear RNA silencing and the spread of a silencing signal between cells in Arabidopsis. Plant Cell 19: 1507–1521.
- Cao X, Aufsatz W, Zilberman D, Mette MF, Huang MS, et al. (2003) Role of the DRM and CMT3 methyltransferases in RNA-directed DNA methylation. Curr Biol 13: 2212–2217.
- Cao X, Jacobsen SE (2002) Locus-specific control of asymmetric and CpNpG methylation by the DRM and CMT3 methyltransferase genes. Proc Natl Acad Sci U S A 99 Suppl 4: 16491–16498.
- Chan SW, Henderson IR, Zhang X, Shah G, Chien JS, et al. (2006) RNAi, DRD1, and histone methylation actively target developmentally important non-CG DNA methylation in arabidopsis. PLoS Genet 2: e83.
- Kaneda M, Okano M, Hata K, Sado T, Tsujimoto N, et al. (2004) Essential role for de novo DNA methyltransferase Dnmt3a in paternal and maternal imprinting. Nature 429: 900–903.
- Okano M, Bell DW, Haber DA, Li E (1999) DNA methyltransferases Dnmt3a and Dnmt3b are essential for de novo methylation and mammalian development. Cell 99: 247–257.
- Bourc'his D, Bestor TH (2004) Meiotic catastrophe and retrotransposon reactivation in male germ cells lacking Dnmt3L. Nature 431: 96–99.
- Bourc'his D, Xu GL, Lin CS, Bollman B, Bestor TH (2001) Dnmt3L and the establishment of maternal genomic imprints. Science 294: 2536–2539.
- Gowher H, Liebert K, Hermann A, Xu G, Jeltsch A (2005) Mechanism of stimulation of catalytic activity of Dnmt3A and Dnmt3B DNA-(cytosine-C5)methyltransferases by Dnmt3L J Biol Chem 280: 13341–13348.
- Suetake I, Morimoto Y, Fuchikami T, Abe K, Tajima S (2006) Stimulation effect of Dnmt3L on the DNA methylation activity of Dnmt3a2. J Biochem 140: 553–559.
- Kareta MS, Botello ZM, Ennis JJ, Chou C, Chedin F (2006) Reconstitution and mechanism of the stimulation of de novo methylation by human DNMT3L. J Biol Chem 281: 25893–25902.
- Jia D, Jurkowska RZ, Zhang X, Jeltsch A, Cheng X (2007) Structure of Dnmt3a bound to Dnmt3L suggests a model for de novo DNA methylation. Nature 449: 248–251.
- Ooi SK, Qiu C, Bernstein E, Li K, Jia D, et al. (2007) DNMT3L connects unmethylated lysine 4 of histone H3 to de novo methylation of DNA. Nature 448: 714–717.
- Aravin AA, Hannon GJ (2008) Small RNA silencing pathways in germ and stem cells. Cold Spring Harb Symp Quant Biol 73: 283–290.
- Aravin AA, Sachidanandam R, Bourc'his D, Schaefer C, Pezic D, et al. (2008) A piRNA pathway primed by individual transposons is linked to de novo DNA methylation in mice. Mol Cell 31: 785–799.
- Carmell MA, Girard A, van de Kant HJ, Bourc'his D, Bestor TH, et al. (2007) MIWI2 is essential for spermatogenesis and repression of transposons in the mouse male germline. Dev Cell 12: 503–514.
- Deng W, Lin H (2002) miwi, a murine homolog of piwi, encodes a cytoplasmic protein essential for spermatogenesis. Dev Cell 2: 819–830.
- Kuramochi-Miyagawa S, Kimura T, Ijiri TW, Isobe T, Asada N, et al. (2004) Mili, a mammalian member of piwi family gene, is essential for spermatogenesis. Development 131: 839–849.
- Kuramochi-Miyagawa S, Watanabe T, Gotoh K, Totoki Y, Toyoda A, et al. (2008) DNA methylation of retrotransposon genes is regulated by Piwi family members MILI and MIWI2 in murine fetal testes. Genes Dev 22: 908–917.

- Freitag M, Lee DW, Kothe GO, Pratt RJ, Aramayo R, et al. (2004) DNA methylation is independent of RNA interference in Neurospora. Science 304: 1939.
- Cao X, Springer NM, Muszynski MG, Phillips RL, Kaeppler S, et al. (2000) Conserved plant genes with similarity to mammalian de novo DNA methyltransferases. Proc Natl Acad Sci U S A 97: 4979–4984.
- Jeddeloh JA, Bender J, Richards EJ (1998) The DNA methylation locus DDM1 is required for maintenance of gene silencing in Arabidopsis. Genes Dev 12: 1714–1725.
- Saze H, Kakutani T (2007) Heritable epigenetic mutation of a transposonflanked Arabidopsis gene due to lack of the chromatin-remodeling factor DDM1. Embo J 26: 3641–3652.
- Woo HR, Dittmer TA, Richards EJ (2008) Three SRA-domain methylcytosinebinding proteins cooperate to maintain global CpG methylation and epigenetic silencing in Arabidopsis. PLoS Genet 4: e1000156.
- Li CF, Pontes O, El-Shami M, Henderson IR, Bernatavichute YV, et al. (2006) An ARGONAUTE4-containing nuclear processing center colocalized with Cajal bodies in Arabidopsis thaliana. Cell 126: 93–106.
- Sope WJ, Jacobsen SE, Alonso-Blanco C, Jackson JP, Kakutani T, et al. (2000) The late flowering phenotype of fwa mutants is caused by gain-of-function epigenetic alleles of a homeodomain gene. Mol Cell 6: 791–802.
- Henderson IR, Zhang X, Lu C, Johnson L, Meyers BC, et al. (2006) Dissecting Arabidopsis thaliana DICER function in small RNA processing, gene silencing and DNA methylation patterning. Nat Genet 38: 721–725.

- Xie Z, Johansen LK, Gustafson AM, Kasschau KD, Lellis AD, et al. (2004) Genetic and functional diversification of small RNA pathways in plants. PLoS Biol 2: E104.
- Gasciolli V, Mallory AC, Bartel DP, Vaucheret H (2005) Partially redundant functions of Arabidopsis DICER-like enzymes and a role for DCL4 in producing trans-acting siRNAs. Curr Biol 15: 1494–1500.
- Kozlov G, Nguyen L, Lin T, De Crescenzo G, Park M, et al. (2007) Structural basis of ubiquitin recognition by the ubiquitin-associated (UBA) domain of the ubiquitin ligase EDD. J Biol Chem 282: 35787–35795.
- Mueller TD, Feigon J (2002) Solution structures of UBA domains reveal a conserved hydrophobic surface for protein-protein interactions. J Mol Biol 319: 1243–1255.
- Kim SA, Vacratsis PO, Firestein R, Cleary ML, Dixon JE (2003) Regulation of myotubularin-related (MTMR)2 phosphatidylinositol phosphatase by MTMR5, a catalytically inactive phosphatase. Proc Natl Acad Sci U S A 100: 4492–4497.
- Willert EK, Fitzpatrick R, Phillips MA (2007) Allosteric regulation of an essential trypanosome polyamine biosynthetic enzyme by a catalytically dead homolog. Proc Natl Acad Sci U S A 104: 8275–8280.
- Shabalina SA, Koonin EV (2008) Origins and evolution of eukaryotic RNA interference. Trends Ecol Evol 23: 578–587.
- Li CF, Henderson IR, Song L, Fedoroff N, Lagrange T, et al. (2008) Dynamic regulation of ARGONAUTE4 within multiple nuclear bodies in Arabidopsis thaliana. PLoS Genet 4: e27.