

# UCLA

## UCLA Previously Published Works

### Title

Clade I-Associated Mpox Cases Associated with Sexual Contact, the Democratic Republic of the Congo.

### Permalink

<https://escholarship.org/uc/item/75x2c0mn>

### Journal

Emerging Infectious Diseases, 30(1)

### Authors

Bogoch, Isaac

Cevik, Muge

Gonsalves, Gregg

et al.

### Publication Date

2024

### DOI

10.3201/eid3001.231164

### Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

# Clade I–Associated Mpox Cases Associated with Sexual Contact, the Democratic Republic of the Congo

Emile M. Kibungu, Emmanuel H. Vakaniaki, Eddy Kinganda-Lusamaki, Thierry Kalonji-Mukendi, Elisabeth Pukuta, Nicole A. Hoff, Isaac I. Bogoch, Muge Cevik, Gregg S. Gonsalves, Lisa E. Hensley, Nicola Low, Souradet Y. Shaw, Erin Schillberg, Mikayla Hunter, Lygie Lunyanga, Sylvie Linsuke, Joule Madinga, Martine Peeters, Jean-Claude Makangara Cigolo, Steve Ahuka-Mundeke, Jean-Jacques Muyembe, Anne W. Rimoin,<sup>1</sup> Jason Kindrachuk,<sup>1</sup> Placide Mbala-Kingebeni,<sup>1</sup> Robert S. Lushima;<sup>1</sup> International Mpox Research Consortium

We report a cluster of clade I monkeypox virus infections linked to sexual contact in the Democratic Republic of the Congo. Case investigations resulted in 5 reverse transcription PCR–confirmed infections; genome sequencing suggest they belonged to the same transmission chain. This finding demonstrates that mpox transmission through sexual contact extends beyond clade IIb.

**H**uman mpox, caused by monkeypox virus (MPXV), is an emerging zoonotic viral disease first identified in the Democratic Republic of the Congo (DRC) (1). MPXV is endemic in multiple regions of Central and West Africa (2,3). The virus is subclassified into 2 clades: clade I, formerly Congo Basin (Central Africa) clade, and clade II, formerly West African clade. Clade II is further subdivided into 2 subclades, IIa and IIb; subclade IIb was responsible for the 2022 global epidemic (4; <https://www.who.int/news/item/12-08-2022-monkeypox--experts-give-virus-variants-new-names>). Clade I infections are associated with greater disease severity and more pronounced rash and had demonstrated increased

human-to-human transmission compared with clade II before the global emergence of clade IIb (5). Those difference are likely influenced by factors such as clade-specific genomic differences in host response modifier proteins, exposure type and dose, and vaccination status (<https://www.who.int/news/item/12-08-2022-monkeypox--experts-give-virus-variants-new-names>). Travel-related and animal importation–related cases have been reported in non-endemic regions (6). In 2022, rapid spread of MPXV to new geographic regions resulted in >86,000 confirmed infections in nonendemic regions and declaration of a public health emergency of international concern by the World Health Organization (7).

During the 2022 epidemic, >90% of infections were linked to secondary transmission, mainly through sexual contact among men who have sex with men (MSM) (8–11). The disease appeared to affect younger populations in endemic regions because of increased contact with zoonotic sources; the average age at infection was <25 years, and the case-fatality

Author affiliations: Ministry of Public Health, Kinshasa, Democratic Republic of the Congo (E.M. Kibungu, T. Kalonji-Mukendi, R.S. Lushima); Institut National de Recherche Biomédicale, Kinshasa (E.H. Vakaniaki, E. Kinganda-Lusamaki, E. Pukuta, L. Lunyanga, S. Linsuke, J. Madinga, J.-C. Makangara Cigolo, S. Ahuka-Mundeke, J.-J. Muyembe, P. Mbala-Kingebeni); Cliniques Universitaires de Kinshasa, Université de Kinshasa, Kinshasa (E. Kinganda-Lusamaki, J.-C. Makangara Cigolo, S. Ahuka-Mundeke, J.-J. Muyembe, P. Mbala-Kingebeni); TransVIHMI (Recherches Translationnelles sur le VIH et les Maladies Infectieuses endémiques et émergentes); University of Montpellier, French National Research Institute for Sustainable Development, INSERM, Montpellier, France (E. Kinganda-

Lusamaki, M. Peeters); University of California, Los Angeles, California, USA (N.A. Hoff, A.W. Rimoin); Toronto General Hospital, University Health Network, Toronto, Ontario, Canada (I.I. Bogoch); University of St. Andrews, St. Andrews, Scotland, UK (M. Cevik); Yale School of Public Health, New Haven, Connecticut, USA (G.S. Gonsalves); USDA Agricultural Research Service, Manhattan, Kansas, USA (L.E. Hensley); University of Bern, Bern, Switzerland (N. Low); University of Manitoba, Winnipeg, Manitoba, Canada (S.Y. Shaw, E. Schillberg, M. Hunter, J. Kindrachuk)

DOI: <https://doi.org/10.3201/eid3001.231164>

<sup>1</sup>These authors contributed equally to this article.

rate was higher among children. In nonendemic regions, however, the average age at infection was >30 years, and infection occurred predominantly in men (>95%) who self-identify as MSM (>80%) (10,11). Clinical characteristics during the 2022 epidemic included fever, physical asthenia or lethargy, and lymphadenopathy with high concentrations of papules, pustules, and vesicles on the skin of the genital and perianal organs. Rectal pain, bleeding, and purulent bloody stools were also common (12,13). A recent case series assessing data from clade IIb infections in persons living with HIV demonstrated that increased disease severity and fatal disease were also linked to CD4 counts (11).

Although clade I-related infections can occur through close contacts, including through fomites, transmission through sexual contact has not been reported previously (14,15). However, given the high sequence homology across MPXV clades and the increased disease severity associated with clade I, clarifying whether sexual contact-related infections occur across MPXV clades is critical. We report a confirmed cluster of clade I-associated mpox associated with sexual contact.

### The Study

An alert was issued by the Kwango Provincial Health Division, DRC, after reports from civil society organizations in March 2023 regarding a resident with skin rashes and pruritus. Teams from the National Programme for the Control of Monkeypox and Viral Hemorrhagic Fevers and the Institut National de


Recherche Biomédicale, accompanied by members of the senior team from the Kwango Provincial Health Division, the senior team from the Kenge Health Zone, the provincial team from the National Programme for the Control of HIV, and the provincial civil society representative, conducted an investigation. During the investigation, the national and provincial health teams instituted training for local healthcare clinics to raise awareness of the clinical signs and mode of transmission for mpox and HIV, including sexual contact in MSM.

A man from DRC in his late 20s (case-patient 1) reported having 2 sexual encounters with a man (suspected primary case-patient) in Europe 1 week before returning to DRC. The suspected primary case-patient frequently visited DRC. Case-patient 1 reported that the primary case-patient had general clinical symptoms including genital pruritus, joint pain, and physical asthenia. Nine days after contact, case-patient 1 began experiencing pruritus, vesicular skin eruptions, and genital and perianal ulcerations. He contacted a local civil organization, which initiated a preliminary investigation and alerted the National Programme for the Control of Monkeypox and Viral Hemorrhagic Fevers. He had penile lesions and penile papules (Figure 1); localized redness on the lips and in the oral buccal mucosa were also noted, and he had 1 lesion on the middle finger of the right hand. Blood samples, oropharyngeal swab samples, swab samples from rectal and genital lesions, and swab samples from vesicles on the skin of the penis and pubis were taken.



**Figure 1.** Penile lesions and papules associated with clade I monkeypox virus infection in a man in his late 20s who reported having 2 sexual encounters with a man in Europe 1 week before arriving in Democratic Republic of the Congo.

**Figure 2.** Phylogenetic analysis of MPXV sequences from a cluster of mpox cases described in Kwango Province, Democratic Republic of the Congo. A) MPXV global phylogeny showing that the Kwango Province outbreak cluster belongs to clade I MPXV. B) Phylogenetic analysis of MPXV genome sequences from the reported cases and clade I MPXV sequences from Central Africa. Posterior support values are shown at branch points. DNA was extracted at Institut National de Recherche Biomédicale using a QIAGEN DNA Mini Kit (<https://www.qiagen.com>) from blood samples and subsequently screened for MPXV with an orthopoxvirus-specific real-time PCR assay. Whole-genome sequencing was attempted on samples from the index case



by next-generation sequencing. The library preparation was performed using Illumina DNA Prep with Enrichment (<https://www.illumina.com>), and the libraries were enriched for MPXV using biotinylated custom probes synthesized by Twist Biosciences (<https://www.twistbioscience.com>). Note that 23MPX098 (or 23MPX0245V) and 23MPX099 (or 23MPX0245C) are vesicle and crust samples from case-patient 1. Scale bar indicates number of substitutions per site. DRC, Democratic Republic of the Congo; hMPXV, human MPXV; MPXV, monkeypox virus.

Case-patient 1 reported having sexual contact with 9 partners (6 men and 3 women) after arriving in DRC. A prescreening of those contacts identified suspected mpox in 3 sexual contacts (2 men and 1 woman, 30–35 years of age) who had fever and lesions in the genital and perianal region. Blood samples and oropharyngeal swab samples were taken from all 9 partners; additional rectal or vaginal swab samples were taken from the men and woman suspected to be secondary case-patients. Investigation of all sexual contacts yielded an additional 36 sexual contacts. Samples from case-patient 1 and 3 suspected secondary case-patients were confirmed to be MPXV-positive by PCR using samples from vesicles, crusts, or blood. One of the 36 sexual contacts was also PCR-positive for MPXV. We performed viral genome sequencing on PCR-positive samples; phylogenetic analysis showed tight clustering among 3 positive samples, suggesting they belong to the same chain of transmission. The closest related sequence beyond this cluster was a 2022 clade I MPXV sequence from DRC (Figure 2).

We treated the 5 PCR-positive case-patients with supportive care and pain control on an outpatient basis. We performed a follow-up investigation to further investigate transmission chains and identify additional

contacts; follow-up was conducted for each contact through individual case report forms. We monitored a total of 120 contacts over 21 days. The contacts were categorized into 3 groups: sexual contacts ( $n = 5$ ), family members ( $n = 45$ ), and persons associated with close nonsexual contact ( $n = 70$ ). None of the persons monitored during the 21 days experienced clinical signs.

## Conclusions

We describe a cluster of clade I MPXV-associated infections in DRC related to sexual contact, which has previously only been described for clade II MPXV. Of note, MPXV transmission through sexual contact is not exclusive to clade IIb and can occur during heterosexual and same-sex contact. This study demonstrates that MPXV infections can occur through additional exposure routes in MPXV-endemic regions and that the current understanding of mpox burden in clade I-endemic regions is based on classical transmission exclusively; recognizing those factors is critical. Our findings highlight additional considerations for MPXV circulation and transmission containment in endemic areas. Thus, increased MPXV surveillance, diagnostic testing access, and equitable access to both vaccines and therapeutics for persons at increased risk for infection are needed for ongoing

mitigation strategies. Given the increased disease severity associated with clade I MPXV, the potential implications of sexual transmission on broadening geographic distribution for MPXV across clades I and II must be considered. In addition, long-term immunity to mpox inferred by vaccination is unknown, including the role of mucosal immunity against clade I MPXV infections. This report highlights multiple critical global health considerations that must be addressed. Ongoing support for community engagement and educational efforts focusing on mpox recognition and reporting, including within sexual networks and for specific groups who might suffer from lack of care or experience stigma when seeking care.

Our findings highlight historically unrecognized MPXV transmission through sexual contact and indicate the need for increased routine screening in sexual health clinics in mpox-endemic and nonendemic regions. Population movement and previously unreported routes of transmission could exacerbate global distribution of MPXV, which could be compounded by the lack of routine diagnostic testing or inadequate access to rapid point-of-care testing. In view of this investigation, epidemiologic and genomic surveillance for MPXV, in both endemic and nonendemic regions, should be improved and strengthened.

### Acknowledgments

We thank the residents of Kwango Province, DRC, and the Ministry of Health, DRC, for their cooperation and assistance during this outbreak investigation. We also thank Sophie Gambia, Franc Kasongo, Jackie Muteba, and staff from the Ministry of Health, who assisted in field investigations, as well as Eric Delaporte, Ahidjo Ayoub, and all members of the AFROSCREEN Consortium (<https://www.afroscreen.org/en/network>) for their work and support on genomic surveillance in DRC.

This work was supported by the International Mpox Research Consortium (IMReC) through funding from the Canadian Institutes of Health Research and International Development Research Centre (grant no. 202209MRR-489062-MPX-CDAA-168421); Department of Defense, Defense Threat Reduction Agency, Monkeypox Threat Reduction Network grant #HDTRA1-21-1-0040; and USDA Non-Assistance Cooperative Agreement #20230048. Additional laboratory support was provided by Agence Française de Développement through the PANAFPOX and AFROSCREEN project (grant agreement CZZ3209), coordinated by ANRS Maladies infectieuses émergentes in partnership with Institut de Recherche pour le Développement (IRD).

E.L. received a PhD grant from the French Foreign Office.

### About the Author

Dr. Kibungu is a senior expert epidemiologist with a primary research interest in emerging infectious diseases, including response and containment efforts from the Ministry of Public Health, Hygiene and Prevention, Democratic Republic of the Congo. His primary research interests are emerging infectious diseases, including response and containment efforts.

### References

- Ladnyj ID, Ziegler P, Kima E. A human infection caused by monkeypox virus in Basankusu Territory, Democratic Republic of the Congo. *Bull World Health Organ*. 1972;46:593–7.
- Breman JG, Kalisa-Ruti, Steniowski MV, Zanutto E, Gromyko AI, Arita I. Human monkeypox, 1970–79. *Bull World Health Organ*. 1980;58:165–82.
- Durski KN, McCollum AM, Nakazawa Y, Petersen BW, Reynolds MG, Briand S, et al. Emergence of monkeypox – West and Central Africa, 1970–2017. *MMWR Morb Mortal Wkly Rep*. 2018;67:306–10. <https://doi.org/10.15585/mmwr.mm6710a5>
- Ulaeto D, Agafonov A, Burchfield J, Carter L, Happi C, Jakob R, et al. New nomenclature for mpox (monkeypox) and monkeypox virus clades. *Lancet Infect Dis*. 2023;23:273–5. [https://doi.org/10.1016/S1473-3099\(23\)00055-5](https://doi.org/10.1016/S1473-3099(23)00055-5)
- McCollum AM, Damon IK. Human monkeypox. *Clin Infect Dis*. 2014;58:260–7. <https://doi.org/10.1093/cid/cit703>
- Reynolds MG, Yorita KL, Kuehnert MJ, Davidson WB, Huhn GD, Holman RC, et al. Clinical manifestations of human monkeypox influenced by route of infection. *J Infect Dis*. 2006;194:773–80. <https://doi.org/10.1086/505880>
- World Health Organization. WHO Director-General declares the ongoing monkeypox outbreak a public health event of international concern [cited 2023 Nov 28]. <https://www.who.int/europe/news/item/23-07-2022-who-director-general-declares-the-ongoing-monkeypox-outbreak-a-public-health-event-of-international-concern>
- Minhaj FS, Ogale YP, Whitehill F, Schultz J, Foote M, Davidson W, et al.; Monkeypox Response Team 2022. Monkeypox outbreak – nine states, May 2022. *MMWR Morb Mortal Wkly Rep*. 2022;71:764–9. <https://doi.org/10.15585/mmwr.mm7123e1>
- Thornhill JP, Palich R, Ghosn J, Walmsley S, Moschese D, Cortes CP, et al.; Share-Net writing group. Human monkeypox virus infection in women and non-binary individuals during the 2022 outbreaks: a global case series. *Lancet*. 2022;400:1953–65. [https://doi.org/10.1016/S0140-6736\(22\)02187-0](https://doi.org/10.1016/S0140-6736(22)02187-0)
- Thornhill JP, Barkati S, Walmsley S, Rockstroh J, Antinori A, Harrison LB, et al.; SHARE-net Clinical Group. Monkeypox virus infection in humans across 16 countries – April–June 2022. *N Engl J Med*. 2022;387:679–91. <https://doi.org/10.1056/NEJMoa2207323>
- World Health Organization. Mpox (monkeypox) outbreak 2022 [cited 2023 Aug 25]. <https://www.who.int/emergencies/situations/monkeypox-outbreak-2022>
- Girometti N, Byrne R, Bracchi M, Heskin J, McOwan A, Tittle V, et al. Demographic and clinical characteristics of confirmed human monkeypox virus cases in individuals attending a sexual health centre in London, UK: an observational analysis. *Lancet Infect Dis*. 2022;22:1321–8. [https://doi.org/10.1016/S1473-3099\(22\)00411-X](https://doi.org/10.1016/S1473-3099(22)00411-X)

13. Philpott D, Hughes CM, Alroy KA, Kerins JL, Pavlick J, Asbel L, et al.; CDC Multinational Monkeypox Response Team. Epidemiologic and clinical characteristics of monkeypox cases – United States, May 17–July 22, 2022. *MMWR Morb Mortal Wkly Rep.* 2022;71:1018–22. <https://doi.org/10.15585/mmwr.mm7132e3>
14. Van Dijk C, Hoff NA, Mbala-Kingebeni P, Low N, Cevik M, Rimoin AW, et al. Emergence of mpox in the post-smallpox era – a narrative review on mpox epidemiology. *Clin Microbiol Infect.* 2023. <https://doi.org/10.1016/j.cmi.2023.08.008>
15. Titanji BK, Tegomoh B, Nematollahi S, Konomos M, Kulkarni PA. Monkeypox: a contemporary review for


healthcare professionals. *Open Forum Infect Dis.* 2022;9:ofac310.

Address for correspondence: Jason Kindrachuk, Department of Medical Microbiology & Infectious Diseases, Department of Internal Medicine, University of Manitoba, 523-745 Bannatyne Ave, Winnipeg, MB R3E 0J9, Canada; email: Jason.Kindrachuk@umanitoba.ca; Placide Mbala-Kingebeni, Département de Biologie Médicale, Université de Kinshasa, 5345, Ave. de la Démocratie, B.P. 1187 Gombe, Kinshasa, Democratic Republic of the Congo; email: mbalaplacide@gmail.com

August 2023

## Unexpected Hazards

- Clinical Characteristics of *Corynebacterium ulcerans* Infection, Japan
- Healthcare-Associated Infections Caused by *Mycobacterium neoaurum* Response to Vaccine-Derived Polioviruses Detected through Environmental Surveillance, Guatemala, 2019
- Outbreak of NDM-1– and OXA-181– Producing *Klebsiella pneumoniae* Bloodstream Infections in a Neonatal Unit, South Africa
- Spatial Epidemiologic Analysis and Risk Factors for Nontuberculous Mycobacteria Infections, Missouri, USA, 2008–2019
- Waterborne Infectious Diseases Associated with Exposure to Tropical Cyclonic Storms, United States, 1996–2018
- Elimination of *Dirofilaria immitis* Infection in Dogs, Linosa Island, Italy, 2020–2022
- Prospecting for Zoonotic Pathogens by Using Targeted DNA Enrichment
- Omicron COVID-19 Case Estimates Based on Previous SARS-CoV-2 Wastewater Load, Regional Municipality of Peel, Ontario, Canada
- Predicting COVID-19 Incidence Using Wastewater Surveillance Data, Denmark, October 2021–June 2022



- Human Fecal Carriage of *Streptococcus agalactiae* Sequence Type 283, Thailand
- Emerging *Corynebacterium diphtheriae* Species Complex Infections, Réunion Island, France, 2015–2020
- Increase of Severe Pulmonary Infections in Adults Caused by M1UK *Streptococcus pyogenes*, Central Scotland, UK
- Dengue Outbreak Response during COVID-19 Pandemic, Key Largo, Florida, USA, 2020
- SARS-CoV-2 Variants and Age-Dependent Infection Rates among Household and Nonhousehold Contacts
- Uniting for Ukraine Tuberculosis Screening Experience, San Francisco, California, USA
- *Mycobacterium abscessus* Meningitis Associated with Stem Cell Treatment During Medical Tourism
- Candidatus *Neoehrlichia mikurensis* Infection in Patient with Antecedent Hematologic Neoplasm, Spain
- Detection of Hantavirus during the COVID-19 Pandemic, Arizona, USA, 2020
- Multidrug-Resistant *Shigella sonnei* Bacteremia among Persons Experiencing Homelessness, Vancouver, British Columbia, Canada
- Multidrug-Resistant Bacterial Colonization and Infections in Large Retrospective Cohort of Mechanically Ventilated COVID-19 Patients
- Economic Evaluation of Wastewater Surveillance Combined with Clinical COVID-19 Screening Tests, Japan
- Chromosome-Borne CTX-M-65 Extended-Spectrum  $\beta$ -Lactamase–Producing *Salmonella enterica* Serovar Infantis, Taiwan
- Genome-Based Epidemiologic Analysis of VIM/IMP Carbapenemase-Producing *Enterobacter* spp., Poland
- Imported Cholera Cases, South Africa, 2023

**EMERGING  
INFECTIOUS DISEASES**

To revisit the August 2023 issue, go to:  
<https://wwwnc.cdc.gov/eid/articles/issue/29/8/table-of-contents>