

UCSF

UC San Francisco Previously Published Works

Title

What Are Patient Preferences for Integrated Behavioral Health in Primary Care?

Permalink

<https://escholarship.org/uc/item/76s783qw>

Authors

Dunn, Julia Alexandra
Garneau, H el ene Chokron
Filipowicz, Heather
et al.

Publication Date

2021

DOI

10.1177/21501327211049053

Peer reviewed

What Are Patient Preferences for Integrated Behavioral Health in Primary Care?

Julia Alexandra Dunn¹ , H el ene Chokron Garneau¹, Heather Filipowicz¹, Megan Mahoney¹, Timothy Seay-Morrison¹, Kaitlin Dent², and Mark McGovern¹

Abstract

Background: Behavioral health services, integrated into primary care practices, have become increasingly implemented. Although patient satisfaction has been studied, limited information exists about patient preferences for integrated behavioral health in primary care and how perceptions may vary. **Objective:** To determine patient preferences for integrated behavioral health within primary care and explore differences across patient groups. **Methods:** A self-report survey was distributed within a quality improvement initiative in an academic health system. A brief 8-item self-report questionnaire of perceptions and preferences for integrated behavioral health was administered to 752 primary care patients presenting before their visits at two primary care clinics. Participation was voluntary, responses were anonymous, and all patients presenting during a three-week timeframe were eligible. **Results:** In general, patients preferred to have behavioral health concerns addressed within primary care (n = 301; 41%) rather than referral to a specialist (7.5%; n = 55). There was no evidence of variation in preferences by demographic characteristics. Comfort levels to receive behavioral health services ($P < .001$) and perceived needs being met were significantly associated with preferences for receiving IBHPC ($P < .001$). **Conclusion:** This project provided valuable data to support the implementation of integrated behavioral health services in primary care clinics. In general, patients prefer to have behavioral health issues addressed within their primary care experience rather than being referred to specialty mental health care. This study adds to an expanding pool of studies exploring patient preferences for integrated behavioral health in primary care.

Keywords

behavioral health, integrated behavioral health care, patient preferences

Dates received: 19 July 2021; revised: 7 September 2021; accepted: 8 September 2021.

Introduction

Behavioral health conditions, including mental health and substance use disorders, are common and costly in primary care settings.¹⁻³ Approximately a quarter of adults in the United States meet the diagnostic criteria for a behavioral health condition annually, with a third of adults with a medical diagnosis reporting a behavioral health condition.⁴ Patients with behavioral health issues are most likely to present in primary care rather than the specialty health system, yet most providers face challenges in addressing these issues. Providers are often limited in their knowledge and training and have to identify and treat behavioral health issues with limited time and resources.¹

Integrated behavioral health in primary care (IBHPC) is designed to overcome the barriers to patients accessing and

providers offering behavioral healthcare. IBHPC combines medical and behavioral health services through integration, not simply the co-location of an embedded psychiatrist, psychologist, or social worker.⁵ Studies conducted in various settings have demonstrated that IBHPC models are associated with improved health outcomes, including depression and anxiety symptoms, patient quality of life, disability, and management of comorbid chronic diseases.⁶⁻⁸

¹Stanford University School of Medicine, Palo Alto, CA, USA

²University of Northern Colorado, Greeley, CO, USA

Corresponding Author:

Julia Alexandra Dunn, Stanford University School of Medicine, 1520 Page Mill Road, Greeley, CO 94304, USA.

Email: dunnja@stanford.edu



IBHPC is also associated with lower costs, increased treatment outcomes and adherence, enhanced patient engagement, improved patient and provider satisfaction, and reduced racial and ethnic disparities.^{5,7,9-14} IBHPC models offer an opportunity for continuity of care for patients who may not receive behavioral healthcare due to long waiting times for specialist referrals.

Although studies have documented high patient satisfaction with IBHPC post-visit, few studies have described patient preferences for IBHPC pre-visit.^{9-12,15-20} The available information on the experience with IBHPC documents a willingness to meet with an embedded behavioral health provider for a follow-up visit and that most patients prefer their mental health and medical providers co-located and in regular communication about their care.^{20,21} Patients may prefer to receive behavioral health services in primary care due to the potential for poor communication between primary and specialty care, the flexibility and comfort of care, immediate access to care, consolidated appointments, lower cost, and less perceived stigma.^{7,22} Alternatively, patients may have concerns about privacy and confidentiality and believe that primary care is for physical, not behavioral health concerns. There may also exist variation in preference by patient characteristics, such as race and ethnicity, sex, age, or prior experience with behavioral health treatment.^{23,24} If patients indicate a preference for receiving integrated behavioral healthcare in primary care settings, they may be more likely to reveal these concerns in future visits.

It remains unclear if patients prefer to have their behavioral health concerns addressed in primary care or value referrals to specialists. This question received considerable debate among primary care leaders in two clinics in California. This study addresses a major barrier around the implementation of IBHPC, whether patients prefer to receive behavioral healthcare services in primary care. A variation of primary care patients' preferences for integrated behavioral health care was explored by several patient-level factors, including demographic characteristics, attitudes, and comfort regarding behavioral healthcare. The goal of this project was to understand patients' preferences for IBHPC and help primary care providers understand how to serve patients' needs best.

Methods

Design

The question of patients preferring their behavioral health issues addressed in primary or specialty care has undergone considerable debate within a local primary care system. In this rapid quality improvement study, cross-sectional data were collected from patients at two primary care clinics in an academic healthcare system, including an internal medicine and a family medicine practice. Both of these practices are located in the same building and staffed with medical

residents, nurses, physician assistants, medical assistants, and attending faculty physicians. The clinics have a combined panel size of 21 960 patients, with approximately 1015 visits weekly. At the time of data collection, there were no integrated behavioral health services available other than referrals to a co-located social worker and external specialty care or community services. The use of standardized screening measures, such as the Patient Health Questionnaire (PHQ-9), was limited. Behavioral health screening was completed at the discretion of the individual provider. Neither patients nor providers had direct access to consulting psychiatry. Behavioral health needs were primarily addressed by primary care providers or referred to the health system's psychiatry department or mental health providers in the community. Wait times for non-acute patients for mental health specialists ranged from 3 to 9 months.

Measures

A project-specific measure, the *Integrated Behavioral Health Patient Preference survey*, was developed (Table 1). The survey is an 8-item self-report measure that focuses on patient preferences for integrated behavioral healthcare. The authors constructed the survey with input and feedback from primary care staff, leadership, and patient representatives. The instrument's psychometric properties were not investigated, and no scoring exists. The survey was designed to address the debate among primary care leaders to understand whether patients preferred IBHPC prior to implementation. The survey includes items on the patient's general medical and psychiatric treatment history, behavioral health perceptions, comfort receiving behavioral healthcare, and behavioral health services preferences. The following behavioral health conditions and comorbid chronic health outcomes were included in the survey: diabetes, hypertension, chronic obstructive pulmonary disease, asthma, depression, anxiety or stress, and alcohol or drug use. These conditions were included as they are some of the most frequent presenting conditions in primary care and have been shown to improve in settings with IBHPC models.^{8,25,26} Two items were adapted from a patient satisfaction survey and related to comfort and preference for receiving mental health services in primary care.¹⁸ Demographic information was collected and included age, sex, and race or ethnicity. The third item on the survey was the primary outcome of interest. It was operationalized as patient preferences for integrated behavioral healthcare: "If you had a behavioral, mental or emotional health concern, which of the following would you prefer?"

Data Collection

All patients presenting for their primary care visit during a 3-week data collection period in May 2019 were invited to

Table 1. Items From the Integrated Behavioral Health Patient Preference Survey Administered Among Primary Care Patients.

Items	Response options
How important is your behavioral, mental, or emotional health to your overall physical health?	Five-point Likert scale items from “not important” to “extremely important”
How comfortable would you feel being asked about or even getting help for your behavioral, mental, or emotional health issues here at the clinic?	Five-point Likert scale items from “not at all comfortable” to “extremely comfortable”
If you had a behavioral, mental, or emotional health concern, which of the following would you prefer	To get help here in the clinic from a team member; To get help here in the clinic, but only from my doctor; To get help here in the clinic from a behavioral health clinician It depends; To get a referral to a specialist outside the clinic
To what extent are your behavioral health needs met in this clinic?	Five-point Likert scale items from “not at all” to “completely”
Demographics	
Age	17 or below; 18-20; 21-29; 30-39; 40-49; 50-59; 60-69; 70 or above
Sex	Female; Male; Transgender; Other (specify)
Race/ethnicity	Asian; American Indian or Alaskan Native; Black or African American; Hispanic or Latino; Native Hawaiian or Pacific Islander; White; Other (specify)
Have you ever received treatment for any of the following problems? (Select all that apply)	Diabetes; Hypertension; COPD; Asthma; Depression; Anxiety/ stress; Alcohol or drug use
Are you currently under our care for any of the following problems? (Select all that apply)	Diabetes; Hypertension; COPD; Asthma; Depression; Anxiety/ stress; Alcohol or drug use
If you checked “yes” to diabetes or hypertension, how much of an impact do behavioral, mental, or emotional health issues have on your condition and your ability to manage it?	Five-point Likert scale items from “no impact” to “major impact”

complete a voluntary, anonymous survey when presenting at the front desk for their appointment. After providing verbal consent, participants were instructed to complete the paper survey in the waiting room. Patients were not provided any compensation for their participation. Upon completion, front desk staff placed surveys in a bin behind the front desk, and surveys were collected at the end of the week by a quality improvement team member. The survey was in English and included no identifying information. The Stanford University School of Medicine Institutional Review Board designated this project as exempt as a quality improvement project. No identifying information was collected from participants.

Statistical Analyses

The analyses were conducted using Stata Version 16. Sex was included in the survey with the options of male, female, and transgender. Very few patients identified as transgender or other; thus, these categories were set to missing for analyses. Race and ethnicity were operationalized following aggregated United States Census categories. For the analysis, American Indian/Alaska Native, Native Hawaiian, or Pacific Islander were recategorized into other. Age was collected in 10-year categories, and for analyses, the youngest group was recombined as 30 or younger, and the oldest

group as 60 or older. The importance of behavioral health to physical health was recategorized into two categories for the analysis: Not at all important to not important, and important to extremely important. Medical treatment history was categorized as: (1) chronic medical condition(s), such as diabetes, hypertension, COPD, and asthma; (2) behavioral health conditions, such as depression, anxiety/ stress, and substance use problems (alcohol, prescription medications, cannabis or illicit drugs); and (3) any patients with comorbid behavioral health and chronic medical conditions. Preference for behavioral healthcare as the primary outcome was categorized as: (1) to get help in the clinic from a team member; (2) from my doctor; (3) from a behavioral health clinician; (4) it depends; and (5) to receive a referral to a specialist outside of the clinic. The categories related to receiving any form of IBHPC were combined to address the lower frequency of patients who preferred receiving help from a non-clinical team member. Thus, this outcome was categorized into receiving help in the clinic from any staff member, it depends, and receiving a referral from a specialist.

Descriptive statistics were calculated using cross-tabulations and frequency distributions. Participants' demographic characteristics with missing data on the primary outcome were compared to those with complete data for IBHPC preferences, using Pearson chi-squared tests

and Fisher's exact test. Bivariate analyses were conducted using Pearson chi-squared tests to account for the categorical outcome with three levels of preferences for behavioral healthcare. All statistical assumptions were met for the bivariate analysis.

Results

Patient Respondent Characteristics

Of approximately 3045 patients who presented to the two clinics during the study period, a total of 752 participants completed the survey (24.8%). Table 2 presents the characteristics of the sample. A majority of participants ($n=420$; 55.9%) identified as female. Most participants were White ($n=321$; 45.0%), followed by Asian ($n=213$; 29.8%), and Hispanic or Latino ($n=66$; 9.2%). Patients' medical history of treatment included: (a) chronic diseases ($n=180$; 46.0%), including diabetes, hypertension, COPD, and asthma; (b) mental health conditions or substance use ($n=137$; 35.0%); and (c) and the comorbid conditions ($n=74$; 18.9%). The demographic characteristics that were missing included (4.3%) 32 patients on sex, 25 (3.3%) on age, and 38 (5.1%) on race and ethnicity.

Main Findings

More than half of patients reported that their behavioral health needs were mostly or completely met at the clinic ($n=322$; 53%). The majority felt comfortable receiving help for behavioral health issues ($n=673$; 90%). Patients generally preferred to have their concerns addressed within primary care ($n=301$; 41.0%) by either a behavioral health clinician (129; 17.6%), their provider (100; 13.6%), or any team member ($n=72$; 9.8%). A minority of patients desired a referral to a specialist ($n=55$; 7.5%), and half of the patients reported, "it depends" ($n=369$; 50.3%). Less than 3% of participants had missing data. Respondent missing data did not differ by patient characteristics.

Patient Factors and IBHPC Preferences

The chi-squared test results between patient-level factors and their preference for IBHPC are displayed in Table 3. There was no evidence of an association between patient preferences and patient demographic characteristics, such as age, sex, race or ethnicity, treatment history for chronic or behavioral health conditions, and behavioral health's importance to their physical health.

Patient comfort levels with receiving help for behavioral health issues were significantly associated with patient preferences for receipt of behavioral health ($P < .001$). Similarly, there was an association between patients feeling that their needs were met at the clinic and preferences for receiving behavioral healthcare ($P < .001$).

Discussion

Summary of Findings

The goal of this quality improvement project was to assess patient preferences for integrated behavioral health services and explore whether these preferences varied by patient factors. Patients preferred receiving care for their behavioral health issues within the clinics rather than receiving a referral to an outside, offsite specialist. Most patients also reported that their behavioral health needs were met at the clinic, even in the absence of onsite formal integrated behavioral health services. On average, patients within the clinic reported a higher preference for having their concerns addressed by a behavioral health clinician than their primary care doctor or other staff members. Patients reported feeling comfortable being asked about and receiving help for their behavioral health concerns. Overall, the findings from this study satisfied concerns among primary care leaders about patient perceptions for IBHPC. Some primary care leaders questioned whether patients preferred receiving behavioral healthcare at the clinic and found addressing behavioral health issues in primary care appropriate. This quality improvement project significantly impacted these primary care clinics and addressed an important question among primary care providers in two clinics in California. As a result of the findings, IBHPC models were implemented in these clinics, and the programs continue to develop.

Even before patients engage in behavioral healthcare, however, differences in individual characteristics may contribute to the prevalence of behavioral health conditions and treatment preferences.^{19,27} African Americans and older patients tend to seek treatment for behavioral health concerns in primary care preferentially but are less likely to receive evidence-based practices.²⁸ Additionally, in many older patients, behavioral health conditions largely are left undiagnosed, and there is evidence that IBHPC increases patient engagement in this population.²⁹

Despite evidence of variation in patient satisfaction with IBHPC by patient characteristics,^{19,23,27} this study showed that preferences for receiving behavioral healthcare did not vary across patient demographic characteristics. However, there was substantial evidence that patients' comfort level and the extent to which their behavioral needs were met at the clinic were associated with their preferences for receiving behavioral health services. These findings underscore that IBHPC is acceptable, if not preferred, by patients.

Limitations and Strengths

This project rapidly addressed a question in a local healthcare system by developing and deploying a brief, low-burden measure for patients to complete. In future iterations, some of the survey items should be improved. For example, 1 question included the response option "It depends" but

Table 2. Descriptive Statistics of Patients in Primary Care Settings.

Characteristics	n (%)
Sex	
Male	420 (55.9)
Female	300 (39.9)
Transgender/other	32 (4.3)
Age (years)	
20 or below	14 (1.9)
21-29	96 (13.2)
30-39	163 (22.4)
40-49	121 (16.6)
50-59	120 (16.5)
60-69	118 (16.2)
70 or above	95 (13.1)
Race or ethnicity	
Asian	213 (29.8)
American Indian/Alaska Native	5 (0.7)
Black/African American	30 (4.2)
Hispanic/Latino	66 (9.2)
Native Hawaiian or Pacific Islander	8 (1.1)
White	321 (45.0)
Other	31 (4.3)
Multiracial/multiethnic	40 (5.6)
Treatment history	
Chronic conditions	180 (46.0)
Behavioral health conditions	137 (35.0)
Comorbid conditions	74 (18.9)
Current treatment	
Chronic conditions	164 (61.2)
Behavioral health conditions	74 (27.6)
Comorbid conditions	30 (11.2)
Importance of behavioral health to physical health	
Not at all important	2 (0.3)
Somewhat important and not important	23 (3.1)
Important	113 (15.1)
Very important	237 (31.6)
Extremely important	374 (49.9)
Comfort level in the clinic	
Not at all comfortable	19 (2.5)
Somewhat comfortable and not comfortable	55 (7.4)
Comfortable	266 (35.6)
Very comfortable	214 (28.7)
Extremely comfortable	193 (25.8)
Extent of needs met	
Not at all	98 (16.1)
A little	76 (12.5)
Partially	112 (18.4)
Mostly	179 (29.4)
Completely	143 (23.5)
If you had a behavioral health concern, what would you prefer	
To get help here in the clinic from any team member	72 (9.8)
To get help here in the clinic only from my doctor	100 (13.6)
To get help here in the clinic from a behavioral health clinician	129 (17.6)
It depends	328 (44.7)
To get a referral to a specialist outside the clinic	55 (7.5)
Combination of preferences	50 (6.8)

Table 3. Chi-Squared Tests of the Relationships Between Patient-Level Factors and Preferences for Integrated Behavioral Healthcare in Primary Care Settings.

	If you had a mental health concern, where would you prefer to receive help			P-value
	In the clinic	It depends	Referral to specialty care	
	n (%)	n (%)	n (%)	
Sex				.13
Male	131 (44.3)	141 (47.6)	24 (8.1)	
Female	173 (41.9)	209 (50.6)	31 (7.5)	
Age (years)				.06
30 or below	41 (37.6)	58 (53.2)	10 (9.2)	
30-39	68 (42.5)	85 (53.1)	7 (4.4)	
40-49	50 (41.7)	60 (50.0)	10 (8.3)	
50-59	43 (36.1)	69 (58.0)	7 (5.9)	
60 or above	103 (49.5)	84 (40.4)	21 (10.1)	
Race or ethnicity				.12
White	127 (40.3)	167 (53.0)	21 (6.7)	
Asian	97 (45.8)	103 (48.6)	12 (5.7)	
Black/African American	10 (33.3)	15 (50.0)	5 (16.7)	
Hispanic/Latino	33 (50.8)	25 (38.5)	7 (10.8)	
Other	18 (42.9)	17 (40.5)	7 (16.7)	
Multiracial/multiethnic	16 (41.0)	20 (51.3)	3 (7.7)	
Treatment history				.06
Chronic conditions	80 (44.9)	82 (46.1)	16 (9.0)	
Behavioral health conditions	50 (36.8)	80 (58.8)	6 (4.4)	
Combined	29 (39.7)	34 (46.6)	10 (13.7)	
Importance of behavioral health to physical health				<.001
Not at all important to somewhat important	11 (44.0)	13 (52.0)	1 (4.0)	
Important to extremely important	298 (42.2)	355 (50.2)	54 (7.6)	
Comfort level in the clinic				
Not at all comfortable	4 (25.0)	3 (18.9)	9 (56.3)	
Somewhat comfortable and not comfortable	21 (38.2)	29 (52.7)	5 (9.1)	
Comfortable	103 (39.3)	146 (55.7)	13 (5.0)	
Very comfortable	101 (47.9)	99 (46.9)	11 (5.2)	
Extremely comfortable	81 (42.9)	91 (48.2)	17 (9.0)	
Extent of needs met				<.001
Not at all	37 (39.4)	41 (43.6)	16 (17.0)	
A little	29 (38.2)	41 (54.0)	6 (7.9)	
Partially	39 (35.5)	62 (56.4)	9 (8.2)	
Mostly	82 (46.1)	88 (49.4)	8 (4.5)	
Completely	82 (58.20)	48 (34.0)	11 (7.8)	

offered no room to elaborate. A majority of participants selected this response category. Additional qualitative data from patient interviews could have augmented these quantitative findings to explore further what types of behavioral health problems patients feel comfortable addressing in primary care and how they would like them addressed. This information could ultimately support a patient-centered implementation of IBHPC.

Additionally, many of the items focused on hypothetical perceptions of services while experiencing a behavioral health concern. More evidence is needed on how reported preferences are related to actual engagement with IBHPC.

Information on the survey's psychometric properties would also inform reliability and validity and allow future use across a range of primary care settings. To date, no validated measure exists to assess patient preferences for IBHPC.³⁰

The findings may also not generalize beyond this setting. This recruitment method did offer a naturalistic sample to answer a localized question. This study's response rate of nearly 25% was comparable to previous studies on post-visit patient satisfaction with behavioral health services.^{17,18,21} Still, no project with a survey focused on behavioral health in a primary care setting, pre- or post-visit, has featured a sample size of this magnitude. Critically, this sample may

not have been large enough to establish hypothesized differences in patient preferences for IBHPC or racially and ethnically diverse to explore these health disparities.^{27,29,31}

Future Directions

These findings suggest that relationships exist between patients' comfort level and the clinic meeting patients' behavioral health needs and preferences for IBHPC. The relationship of trust and confidence the patient has with their provider and the clinic may influence their openness to receive integrated behavioral health. Future work should investigate whether comfort level and patient needs being met may explain some of the relationships between patient demographic characteristics and preferences for IBHPC. The mechanisms that affect patients' preferences for IBHPC should be further explored in larger and more diverse samples of primary care patients. Additionally, future studies should explore the differences in patient preferences for IBHPC based on additional factors, such as the role of clinical staff, age of patients, and treatment history across more diverse primary care settings.

Conclusions

The findings from this quality improvement study in 2 primary care clinics suggest that patients prefer to have their behavioral health concerns addressed in settings familiar and comfortable to them, from individuals they know, and where they already receive care for other medical conditions. These results support the expansion of IBHPC and alleviate concerns among institutional leadership and providers that patients may not wish to have behavioral health issues raised or addressed within the context of their primary care relationship.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

ORCID iD

Julia Alexandra Dunn  <https://orcid.org/0000-0002-7892-7717>

References

- Kathol RG, Degruy F, Rollman BL. Value-based financially sustainable behavioral health components in patient-centered medical homes. *Ann Fam Med*. 2014;12(2):172-175. doi:10.1370/afm.1619
- Peek CJ, Cohen DJ, deGruy FV 3rd. Research and evaluation in the transformation of primary care. *Am Psychol*. 2014;69(4):430-442. doi:10.1037/a0036223
- Davenport S, Gray TJ, Melek S. How Do Individuals With Behavioral Health Conditions Contribute to Physical and Total Healthcare Spending? Milliman Research Report; 2020. Accessed August 25, 2021. <https://www.milliman.com/-/media/milliman/pdfs/articles/milliman-high-cost-patient-study-2020.ashx>
- Vogel ME, Kanzler KE, Aikens JE, Goodie JL. Integration of behavioral health and primary care: current knowledge and future directions. *J Behav Med*. 2017;40(1):69-84. doi:10.1007/s10865-016-9798-7
- Hunter CL, Funderburk JS, Polaha J, Bauman D, Goodie JL, Hunter CM. Primary care behavioral health (PCBH) model research: current state of the science and a call to action. *J Clin Psychol Med Settings*. 2018;25(2):127-156. doi:10.1007/s10880-017-9512-0
- Reed SJ, Shore KK, Tice JA. Effectiveness and value of integrating behavioral health into primary care. *JAMA Intern Med*. 2016;176(5):691-692. doi:10.1001/jamainternmed.2016.0804
- Balasubramanian BA, Cohen DJ, Jetelina KK, et al. Outcomes of integrated behavioral health with primary care. *J Am Board Fam Med*. 2017;30(2):130-139. doi:10.3122/jabfm.2017.02.160234
- Coventry P, Lovell K, Dickens C, et al. Integrated primary care for patients with mental and physical multimorbidity: cluster randomised controlled trial of collaborative care for patients with depression comorbid with diabetes or cardiovascular disease. *BMJ*. 2015;350:h638. doi:10.1136/bmj.h638
- O'Loughlin K, Donovan EK, Radcliff Z, Ryan M, Rybarczyk B. Using integrated behavioral healthcare to address behavioral health disparities in underserved populations. *Transl Issues Psychol Sci*. 2019;5(4):374-389. doi:10.1037/tps0000213
- Possemato K, Johnson EM, Beehler GP, et al. Patient outcomes associated with primary care behavioral health services: a systematic review. *Gen Hosp Psychiatry*. 2018;53:1-11. doi:10.1016/j.genhosppsy.2018.04.002
- Reiss-Brennan B. Mental health integration: normalizing team care. *J Prim Care Community Health*. 2014;5(1):55-60. doi:10.1177/2150131913508983
- Davis MM, Gunn R, Gowen LK, Miller BF, Green LA, Cohen DJ. A qualitative study of patient experiences of care in integrated behavioral health and primary care settings: more similar than different. *Transl Behav Med*. 2018;8(5):649-659. doi:10.1093/tbm/ibx001
- Sanchez K, Ybarra R, Chapa T, Martinez ON. Eliminating behavioral health disparities and improving outcomes for racial and ethnic minority populations. *Psychiatr Serv*. 2016;67(1):13-15. doi:10.1176/appi.ps.201400581
- Martinez LS, Lundgren L, Walter AW, et al. Behavioral health, primary care integration, and social work's role in improving health outcomes in communities of color: a systematic review. *J Soc Social Work Res*. 2019;10(3):441-457. doi:10.1086/704070
- Talen MR, & Burke Valeras A. eds *Integrated Behavioral Health in Primary Care: Evaluating the Evidence, Identifying the Essentials*. Springer; 2013.

16. Wray JM, Coulon S, Wong N, Szafranski D, Gros DF. A single integrated behavioral health appointment improves patients' perceptions of behavioral health treatment. *Prim Care Companion CNS Disord*. 2019;21(1):i: 18m02378. doi:10.4088/PCC.18m02378
17. Miller-Matero LR, Khan S, Thiem R, DeHondt T, Dubaybo H, Moore D. Integrated primary care: patient perceptions and the role of mental health stigma. *Prim Health Care Res Dev*. 2019;20:e48. doi:10.1017/S1463423618000403
18. Ede V, Okafor M, Kinuthia R, et al. An examination of perceptions in integrated care practice. *Community Ment Health J*. 2015;51(8):949-961. doi:10.1007/s10597-015-9837-9
19. Hammer JH, Perrin PB, Spiker DA. Impact of integrated care and co-location of care on mental help-seeking perceptions. *JMIR Ment Health*. 2021;30:405-410. doi:10.1080/09638237.2019.1581334
20. Mauksch LB, Tucker SM, Katon WJ, et al. Mental illness, functional impairment, and patient preferences for collaborative care in an uninsured, primary care population. *J Fam Pract*. 2001;50(1):41-47.
21. Funderburk JS, Fielder RL, DeMartini KS, Flynn CA. Integrating behavioral health services into a university health center: patient and provider satisfaction. *Fam Syst Health*. 2012;30(2):130-140.
22. Pomerantz A, Cole BH, Watts BV, Weeks WB. Improving efficiency and access to mental health care: combining integrated care and advanced access. *Gen Hosp Psychiatry*. 2008; 30(6):546-551. doi:10.1016/j.genhosppsy.2008.09.004
23. Riley AR, Walker BL, Wilson AC, Hall TA, Stormshak EA, Cohen DJ. Parents' consumer preferences for early childhood behavioral intervention in primary care. *J Dev Behav Pediatr*. 2019;40(9):669-678. doi:10.1097/DBP.0000000000000736
24. Auxier A, Farley T, Seifert K. Establishing an integrated care practice in a community health center. *Prof Psychol Res Pr*. 2011;42(5):391-397. doi:10.1037/a0024982
25. Wu LT, McNeely J, Subramaniam GA, et al. DSM-5 substance use disorders among adult primary care patients: results from a multisite study. *Drug Alcohol Depend*. 2017;179:42-46. doi:10.1016/j.drugalcdep.2017.05.048
26. Finley CR, Chan DS, Garrison S, et al. What are the most common conditions in primary care? Systematic review. *Can Fam Physician*. 2018;64(11):832-840.
27. Jimenez DE, Bartels SJ, Cardenas V, Dhaliwal SS, Alegria M. Cultural beliefs and mental health treatment preferences of ethnically diverse older adult consumers in primary care. *Am J Geriatr Psychiatry*. 2012;20(6):533-542. doi:10.1097/JGP.0b013e318227f876
28. Cook BL, Zuvekas SH, Carson N, Wayne GF, Vesper A, McGuire TG. Assessing racial/ethnic disparities in treatment across episodes of mental health care. *Health Serv Res*. 2014;49(1):206-229. doi:10.1111/1475-6773.12095
29. Bartels SJ, Coakley EH, Zubritsky C, et al. Improving access to geriatric mental health services: a randomized trial comparing treatment engagement with integrated versus enhanced referral care for depression, anxiety, and at-risk alcohol use. *Am J Psychiatry*. 2004;161(8):1455-1462. doi:10.1176/appi.ajp.161.8.1455
30. Riley AR, Walker BL, Hall TA. Development and initial validation of a measure of parents' preferences for behavioral counseling in primary care. *Fam Syst Health*. 2020;38(2):139-150. doi:10.1037/fsh0000481
31. Snowden LR, Pingitore D. Frequency and scope of mental health service delivery to African Americans in primary care. *Ment Health Serv Res*. 2002;4(3):123-130. doi:10.1023/A:1019709728333