

UC Berkeley

UC Berkeley Previously Published Works

Title

Iron addition to soil specifically stabilized lignin

Permalink

<https://escholarship.org/uc/item/7820j57w>

Authors

Hall, Steven J
Silver, Whendee L
Timokhin, Vitaliy I
[et al.](#)

Publication Date

2016-07-01

DOI

10.1016/j.soilbio.2016.04.010

Peer reviewed

Iron addition to soil specifically stabilized lignin

Author links open overlay panel [Steven J. Hall^a](#) [Whendee L. Silver^b](#) [Vitaliy I. Timokhin^c](#) [Kenneth E. Hammel^{de}](#)
Show more

<https://doi.org/10.1016/j.soilbio.2016.04.010> Get rights and content

Highlights

- Ferrous Fe addition suppressed mineralization of ¹³C-labeled lignin methoxyls.
- Ferrous Fe addition did not affect total soil respiration.
- Mineralization rates of lignin methoxyls implied decadal turnover times.

Abstract

The importance of [lignin](#) as a [recalcitrant](#) constituent of [soil organic matter](#) (SOM) remains contested. Associations with iron (Fe) [oxides](#) have been proposed to specifically protect lignin from decomposition, but impacts of Fe-lignin interactions on [mineralization](#) rates remain unclear. Oxygen (O₂) fluctuations characteristic of humid [tropical soils](#) drive reductive Fe dissolution and precipitation, facilitating multiple types of Fe-lignin interactions that could variably decompose or protect lignin. We tested impacts of Fe addition on ¹³C methoxyl-labeled lignin mineralization in soils that were exposed to static or fluctuating O₂. Iron addition suppressed lignin mineralization to 21% of controls, regardless of O₂ availability. However, Fe addition had no effect on soil CO₂ production, implying that Fe oxides specifically protected lignin methoxyls but not bulk SOM. Iron oxide-lignin interactions represent a specific mechanism for lignin stabilization, linking SOM [biochemical composition](#) to turnover via [geochemistry](#).

- [Previous article](#)
- [Next article](#)

Keywords

Carbon stabilization

Iron

Lignin

Recalcitrance

Redox

Soil organic matter

The new paradigm of [soil organic matter](#) (SOM) dynamics posits that biochemical differences among compounds that influence short-term decomposition do not impact long-term persistence in SOM ([Grandy and Neff, 2008](#), [Marschner et al., 2008](#), [Schmidt et al., 2011](#)). These findings challenged the doctrine of [humus formation](#), namely that [recalcitrant](#) biomolecules derived from [lignin](#) contribute disproportionately to SOM ([Bollag et al., 1997](#)). In support of the new paradigm, several studies reported that lignin might decompose as fast or faster than bulk SOM ([Dignac et al., 2005](#), [Gleixner et al., 1999](#), [Guggenberger et al., 1994](#), [Heim and Schmidt, 2007](#), [Kiem and Kögel-Knabner, 2003](#)). Yet, these findings are difficult to reconcile with other studies demonstrating preferential association of aromatic lignin constituents with minerals, especially Fe [oxides](#) ([Chorover and Amistadi, 2001](#), [Gu et al., 1995](#), [Hernes et al., 2013](#), [Huang et al., 1977](#), [Kaiser, 2003](#), [Riedel et al., 2013](#)). These associations would presumably decrease lignin [mineralization](#), as has been observed for coprecipitates of lignin and [ferrihydrite](#) ([Eusterhues et al., 2014](#)). Thus, a fundamental question remains: can associations with minerals, particularly Fe oxides, decrease the decomposition of lignin relative to bulk SOM?

Iron-lignin interactions in soils are also complicated by potential *stimulation* of decomposition by solid-phase and soluble Fe. Fenton reactions driven by coupled biotic/abiotic Fe redox cycling can decompose [lignin](#) ([Yelle et al., 2011](#)), and may be important in Fe-rich soils that experience fluctuating O₂ ([Hall et al., 2015a](#), [Hall and Silver, 2013](#)). Also, Fe minerals such as [goethite](#) can abiotically oxidize SOM ([Chorover and Amistadi, 2001](#)). Finally, Fe reduction dominates anaerobic metabolism in many Fe-rich soils ([Dubinsky et al., 2010](#)), solubilizing Fe and exposing adsorbed or co-precipitated organic matter to enzymatic attack.

Our previous work ([Hall et al., 2015a](#)) demonstrated that redox fluctuations increased the contribution of methoxyl groups in [lignin](#) to respiration from a [tropical forest](#) soil relative to static [aerobic conditions](#). Here, we report an additional experiment where soils were amended with aqueous Fe(II) to simulate a highly reduced microsite, and then exposed to aerobic or fluctuating aerobic/hypoxic conditions. Added Fe(II) reacted with O₂ to form Fe(III) [oxides](#), a portion of which was again reduced under subsequent [hypoxic conditions](#). We evaluated three alternative hypotheses: H1) the precipitation of Fe oxides following Fe(II) oxidation disproportionately protects lignin relative to bulk SOM; H2) precipitation of Fe oxides protects lignin and bulk SOM equally; H3) oxidation of added Fe(II) stimulates organic matter decomposition.


Soil was sampled from a tropical [montane forest](#) in the Luquillo Experimental Forest, Puerto Rico (Bisley Watershed), with 3500 mm of annual precipitation and mean annual temperature of 22 °C ([Hall and Silver, 2015](#)). This Aquic Hapludox was derived from volcanoclastic sediments and contained substantial [short-range-order](#) Fe [oxides](#) (24 mg Fe g⁻¹) and Fe(II) (0.2–0.8 mg g⁻¹). Iron(II) concentrations were much higher in reducing microsites, occasionally exceeding 5 mg g⁻¹ ([Hall and Silver, 2013](#)).

A [macromolecular](#) synthetic [guaiacyl lignin](#) labeled with ¹³C at the methoxyl position was produced ([Kirk et al., 1975](#)) to provide a sensitive index of [mineralization](#). Lignin was precipitated onto ground [leaf litter](#) and gently homogenized with soils (20 mg lignin and 1000 mg soil) following [Hall et al. \(2015a\)](#). We used a synthetic ¹³C-labeled lignin because these polymers yield similar long-term mineralization estimates as plant-derived lignins ([Haider and Martin, 1981](#)), while eliminating the labeled [polysaccharides](#) that inevitably contaminate labeled lignins produced *in planta* ([Crawford, 1981](#)). We recognize that the physical state of synthetic lignins does not completely represent that of natural lignins, which are intimately associated with other organic compounds (e.g. hemicellulose) in plant biomass. However, the fact that lignin [biodegradation](#) is a highly nonspecific oxidative process ([Ruiz-Dueñas and Martínez, 2009](#)) indicates that the precise spatial orientation of the lignin is probably not a critical factor, especially considering that extensive fragmentation of litter occurs during its breakdown in soils.

Samples were incubated under one of four headspace and Fe addition treatment combinations (n = 5 per treatment): static aerobic + aqueous Fe(II), static aerobic + water, fluctuating aerobic/hypoxic + aqueous Fe(II), fluctuating aerobic/hypoxic + water. The fluctuating aerobic/hypoxic treatment experienced three days of [aerobic conditions](#) followed by four days of [hypoxia](#)(N₂ atmosphere). Before the experiment, samples received either N₂-flushed water (controls), or 5 mg Fe(II) g soil⁻¹ as aqueous FeCl₂ (Fe addition treatment) in an anaerobic chamber. Total CO₂ and ¹³CO₂ production were measured by [gas chromatography](#) and [isotope ratio](#) mass spectrometry over 3–4 day intervals. Additional subsamples were destructively sampled for Fe(II) analysis in 0.5M HCl using a modified ferrozine method ([Hall et al., 2015a](#)).

In the Fe addition treatment, most (>70%) of added Fe(II) oxidized to Fe(III) or precipitated in non-HCl extractable phases within the first three days ([Fig. 1](#)). Iron addition increased subsequent Fe(II) production during the hypoxic phase of the fluctuating treatment 2–5 fold relative to the control ([Fig. 1](#)), likely due to the facile

reduction of newly-formed short-range-ordered Fe(III) [oxides](#) (i.e., phases with nm-scale structure). This indicated that the availability of short-range-ordered Fe phases likely limited Fe reduction even in this Fe-rich soil. Iron(II) decreased over time in all treatments, and was ultimately similar between Fe addition and control treatments.




1. [Download high-res image \(342KB\)](#)
2. [Download full-size image](#)

Fig. 1. Soil Fe(II) concentrations in 0.5 M HCl extractions by treatment and day of experiment (n = 1 per treatment/day). The grey bars represent the hypoxic phase of the fluctuating headspace treatment. In the Fe addition treatments, 5000 µg Fe(II) g⁻¹ soil were added at the beginning of the experiment.

Iron addition consistently and significantly ($p < 0.0001$) decreased the contribution of [lignin](#) methoxyl C to [soil respiration](#) (p_{lignin}) to 15–55% of the control

([Fig. 2a](#)). [Hypoxia](#) also decreased p_{lignin} in both treatments on days 7 and 14 ($p < 0.05$; [Fig. 2a](#)). However, O_2 effects were dwarfed by Fe addition, especially for cumulative [mineralization](#) ([Fig. 2b](#)). Iron-amended soils cumulatively mineralized only 21% of lignin methoxyls relative to the controls ($p < 0.0001$). Cumulative mineralization did not differ between O_2 treatments ([Fig. 2b](#)).



1. [Download high-res image \(773KB\)](#)
2. [Download full-size image](#)

Fig. 2. a) Production of CO₂ from ¹³C methoxyl [lignin](#) as a percent of total [soil respiration](#). b) Total [mineralization](#) of methoxyl lignin C. c) Soil CO₂ production by sampling date. d) Cumulative CO₂ production. Bars represent standard errors with n = 5 per treatment. See the text for a description of the treatments.

We modeled cumulative [lignin](#) methoxyl [mineralization](#) in the [aerobic treatments](#) (Fig. 2b) using first-order decay after discarding the first observation ($R^2 = 0.93$ and 0.92 , for Fe addition and controls, respectively). A two-pool model did not improve the [Akaike Information Criterion](#). Decomposition constants for Fe addition and control soils measured 0.0191 ± 0.0003 and 0.0762 ± 0.0009 y⁻¹, respectively, implying mean turnover times of 52 and 13 y. These were similar to previous estimates of total lignin turnover in other ecosystems (10–40 y; [Bahri et al., 2008](#), [Heim and Schmidt, 2007](#), [Rasse et al., 2006](#)), and of slow-pool mineral-associated C in this ecosystem (11–26 y; [Hall et al., 2015b](#)). Because we measured the most labile constituent of lignin, its methoxyl group ([Kirk et al., 1975](#)), our results suggest that the more [recalcitrant](#) aromatic C of lignin could persist in soils for decades because of interactions with Fe [oxides](#).

Trends in [soil respiration](#) differed from those of [lignin mineralization](#) (Fig. 2c,d). Total CO₂ production from Fe-amended soils was less than controls on day 3 ($p < 0.0001$), potentially due to direct toxicity of soluble Fe or [reactive oxygen species](#) produced during Fe(II) oxidation. Both mechanisms may have been important given the Fe(II) concentrations added here ([Dunning et al., 1998](#)). However, by day 10 Fe addition did not affect CO₂ production (Fig. 1c). Cumulative CO₂ production was similar between Fe addition and controls ($p = 0.15$; Fig. 1d). The similar CO₂ production from Fe addition and control treatments suggests no general suppression of organic matter mineralization upon interaction with Fe [oxides](#) as observed elsewhere ([Eusterhues et al., 2014](#)). Rather, availability of Fe(III) oxides as a terminal electron acceptor ([Sutton-Grier et al., 2011](#)), or shifts in pH, dissolved organic C, or reactive oxygen species known to be associated with Fe reduction and oxidation ([Hall and Silver, 2013](#), [Thompson et al., 2006](#)) might have sustained CO₂ production despite protective effects of Fe—C interactions.

Our data thus support H1: Fe addition disproportionately inhibited the [mineralization](#) of [lignin](#) methoxyls relative to bulk SOM, irrespective of O₂ availability. This is consistent with preferential interactions between Fe [oxide minerals](#) and aromatic lignin constituents ([Gu et al., 1995](#), [Kaiser, 2003](#), [Riedel et al., 2013](#)). Given that [short-](#)

[range-order](#) Fe [oxides](#) are present in most soils ([Reyes and Torrent, 1997](#)), Fe-lignin interactions deserve attention as contributors to “hidden lignin” pools that have potentially been obscured due to methodological artifacts associated with common lignin measurements ([Hernes et al., 2013](#)). Lignins synthesized with labeled isotopes provide one method to surmount these analytical challenges. We propose that consideration of specific interactions between [recalcitrant](#) biomolecules and soil geochemical constituents may prove useful in bridging old and new paradigms of SOM.

Acknowledgments

We thank H. Dang, C. Torrens, and the US Forest Service International Institute of Tropical Forestry. NSF grant [DEB-1457805](#) to WLS and SJH, DOE Biological and Environmental Research grant [DE-SC0006929](#) to KEH, and the NSF Luquillo Critical Zone Observatory ([EAR-0722476](#)) and LTER ([DEB-0620910](#)) supported this work.

References

[Bahri et al., 2008](#)

H. Bahri, D.P. Rasse, C. Rumpel, M.-F. Dignac, G. Bardoux, A. Mariotti **Lignin degradation during a laboratory incubation followed by ¹³C isotope analysis**

Soil Biol. Biochem., 40 (2008), pp. 1916-1922, [10.1016/j.soilbio.2008.04.002](#)

[ArticleDownload PDFView Record in Scopus](#)

[Bollag et al., 1997](#)

J.-M. Bollag, J. Dec, P.M. Huang **Formation mechanisms of complex organic structures in soil habitats**

Donald L. Sparks (Ed.), Advances in Agronomy, Academic Press (1997), pp. 237-266

[ArticleDownload PDFView Record in Scopus](#)

[Chorover and Amistadi, 2001](#)

J. Chorover, M.K. Amistadi **Reaction of forest floor organic matter at goethite, birnessite and smectite surfaces**

Geochimica et Cosmochimica Acta, 65 (2001), pp. 95-109, [10.1016/S0016-7037\(00\)00511-1](#)

[ArticleDownload PDFView Record in Scopus](#)

[Crawford, 1981](#)

R.L. Crawford **Lignin Biodegradation and Transformation**

Wiley, New York (1981)

[Dignac et al., 2005](#)

M.-F. Dignac, H. Bahri, C. Rumpel, D.P. Rasse, G. Bardoux, J. Balesdent, C. Girardin, C. Chenu, A. Mariotti **Carbon-13 natural abundance as a tool to study the dynamics of lignin monomers in soil: an appraisal at the Closeaux experimental field (France)**

Geoderma, Mech. Regul. Org. Matter Stabilisation Soils, 128 (2005), pp. 3-17, [10.1016/j.geoderma.2004.12.022](https://doi.org/10.1016/j.geoderma.2004.12.022)

[ArticleDownload](#) [PDFView](#) [Record in Scopus](#)

[Dubinsky et al., 2010](#)

E.A. Dubinsky, W.L. Silver, M.K. Firestone **Tropical forest soil microbial communities couple iron and carbon biogeochemistry**

Ecology, 91 (2010), pp. 2604-2612, [10.1890/09-1365.1](https://doi.org/10.1890/09-1365.1)

[CrossRefView](#) [Record in Scopus](#)

[Dunning et al., 1998](#)

J.C. Dunning, Y. Ma, R.E. Marquis **Anaerobic killing of oral Streptococci by reduced, transition metal cations**

Appl. Environ. Microbiol., 64 (1998), pp. 27-33

[View Record in Scopus](#)

[Eusterhues et al., 2014](#)

K. Eusterhues, J. Neidhardt, A. Hädrich, K. Küsel, K.U. Totsche **Biodegradation of ferrihydrite-associated organic matter**

Biogeochemistry, 119 (2014), pp. 45-50, [10.1007/s10533-013-9943-0](https://doi.org/10.1007/s10533-013-9943-0)

[CrossRefView](#) [Record in Scopus](#)

[Gleixner et al., 1999](#)

G. Gleixner, R. Bol, J. Balesdent **Molecular insight into soil carbon turnover**

Rapid Commun. Mass Spectrom., 13 (1999), pp. 1278-1283, [10.1002/\(SICI\)1097-0231\(19990715\)13:13<1278::AID-RCM649>3.0.CO;2-N](https://doi.org/10.1002/(SICI)1097-0231(19990715)13:13<1278::AID-RCM649>3.0.CO;2-N)

[CrossRefView](#) [Record in Scopus](#)

[Grandy and Neff, 2008](#)

A.S. Grandy, J.C. Neff **Molecular C dynamics downstream: the biochemical decomposition sequence and its impact on soil organic matter structure and function**

Sci. Total Environ., 404 (2008), pp. 297-307, [10.1016/j.scitotenv.2007.11.013](https://doi.org/10.1016/j.scitotenv.2007.11.013)

[ArticleDownload](#) [PDFView](#) [Record in Scopus](#)

[Gu et al., 1995](#)

B. Gu, J. Schmitt, Z. Chen, L. Liang, J.F. McCarthy **Adsorption and desorption of different organic matter fractions on iron oxide**

Geochim. et Cosmochim. Acta, 59 (1995), pp. 219-229, [10.1016/0016-7037\(94\)00282-Q](https://doi.org/10.1016/0016-7037(94)00282-Q)

[ArticleDownload](#) [PDFView](#) [Record in Scopus](#)

[Guggenberger et al., 1994](#)

G. Guggenberger, B.T. Christensen, W. Zech **Land-use effects on the composition of organic matter in particle-size separates of soil: I. Lignin and carbohydrate signature**

Eur. J. Soil Sci., 45 (1994), pp. 449-458, [10.1111/j.1365-2389.1994.tb00530.x](https://doi.org/10.1111/j.1365-2389.1994.tb00530.x)

[CrossRefView](#) [Record in Scopus](#)

[Haider](#)
[and](#)
[Martin,](#)
[1981](#)

K. Haider, J.P. Martin **Decomposition in soil of specifically ¹⁴C-labeled model and cornstalk lignins and coniferyl alcohol over two years as influenced by drying, rewetting, and additions of an available C substrate**

Soil Biol. Biochem., 13 (1981), pp. 447-450, [10.1016/0038-0717\(81\)90032-8](https://doi.org/10.1016/0038-0717(81)90032-8)

[ArticleDownload PDFView Record in Scopus](#)

[H](#)
[a](#)
[i](#)
[l](#)
[a](#)
[n](#)
[d](#)
[-](#)
[S](#)
[i](#)
[l](#)
[v](#)
[e](#)
[r](#)
[-](#)
[2](#)
[0](#)
[1](#)
[5](#)

S.J. Hall, W.L. Silver **Reducing conditions, reactive metals, and their interactions can explain spatial patterns of surface soil carbon in a humid tropical forest**

Biogeochemistry, 125 (2015), pp. 149-165, [10.1007/s10533-015-0120-5](https://doi.org/10.1007/s10533-015-0120-5)

[CrossRefView Record in Scopus](#)

[Hall and](#)
[Silver,](#)
[2013](#)

S.J. Hall, W.L. Silver **Iron oxidation stimulates organic matter decomposition in humid tropical forest soils**

Glob. Change Biol., 19 (2013), pp. 2804-2813, [10.1111/gcb.12229](https://doi.org/10.1111/gcb.12229)

[CrossRefView Record in Scopus](#)

[Hall et al.,
2015a](#)

S.J. Hall, W.L. Silver, V.I. Timokhin, K.E. Hammel **Lignin decomposition is sustained under fluctuating redox conditions in humid tropical forest soils**

Glob. Change Biol., 21 (2015), pp. 2818-2828, [10.1111/gcb.12908](#)

[CrossRefView Record in Scopus](#)

[Hall, 2015b](#)

S.J. Hall, G. McNicol, T. Nataka, W.L. Silver **Large fluxes and rapid turnover of mineral-associated carbon across topographic gradients in a humid tropical forest: insights from paired ¹⁴C analysis**

Biogeosciences, 12 (2015), pp. 2471-2487, [10.5194/bg-12-2471-2015](#)

[CrossRefView Record in Scopus](#)

[Heim and Schmidt](#)

A. Heim, M.W.I. Schmidt **Lignin turnover in arable soil and grassland analysed with two different labelling approaches**

Eur. J. Soil Sci., 58 (2007), pp. 599-608, [10.1111/j.1365-2389.2006.00848.x](#)

[CrossRefView Record in Scopus](#)

[Hernes et al., 20](#)

P.J. Hernes, K. Kaiser, R.Y. Dyda, C. Cerli **Molecular trickery in soil organic matter: hidden lignin**

Environ. Sci. Technol., 47 (2013), pp. 9077-9085, [10.1021/es401019n](#)

[CrossRefView Record in Scopus](#)

[Huang et al., 197](#)

P.M. Huang, T.S.C. Wang, M.K. Wang, M.H. Wu, N.W. Hsu **Retention of phenolic acids by noncrystalline hydroxy-aluminum and -iron compounds and clay minerals of soils**

Soil Sci., 123 (1977), pp. 213-219

[CrossRefView Record in Scopus](#)

[Kaiser, 2003](#)

K. Kaiser **Sorption of natural organic matter fractions to goethite (α -FeOOH): effect of chemical composition as revealed by liquid-state ¹³C NMR and wet-chemical analysis**

Org. Geochem., 34 (2003), pp. 1569-1579, [10.1016/S0146-6380\(03\)00120-7](#)

[ArticleDownload PDFView Record in Scopus](#)

[Kiem and Kögel-](#)

R. Kiem, I. Kögel-Knabner **Contribution of lignin and polysaccharides to the refractory carbon pool in C-depleted arable soils**

Soil Biol. Biochem., 35 (2003), pp. 101-118, [10.1016/S0038-0717\(02\)00242-0](#)

[ArticleDownload PDFView Record in Scopus](#)

[Kirk et al., 1975](#)

T.K. Kirk, W.J. Connors, R.D. Bleam, W.F. Hackett, J.G. Zeikus **Preparation and microbial decomposition of synthetic [¹⁴C]lignins**

Proceedings of the National Academy of Sciences (USA), 72 (1975), pp. 2515-2519

[CrossRefView Record in Scopus](#)

[Marschner et al.](#)

B. Marschner, S. Brodowski, A. Dreves, G. Gleixner, A. Gude, P.M. Grootes, U. Hamer, A. Heim, G. Jandl, R. Ji, K. Kaiser, K. Kalbitz, C. Kramer, P. Leinweber, J. Rethemeyer, A. Schäffer, M.W.I. Schmidt, L. Schwark, G.L.B. Wiesenberger **How relevant is recalcitrance for the stabilization of organic matter in soils?**

J. Plant Nutr. Soil Sci., 171 (2008), pp. 91-110, [10.1002/jpln.200700049](#)

[CrossRefView Record in Scopus](#)

[Rasse et al., 200](#)

D.P. Rasse, M.-F. Dignac, H. Bahri, C. Rumpel, A. Mariotti, C. Chenu **Lignin turnover in an agricultural field: from plant residues to soil-protected fractions**

Eur. J. Soil Sci., 57 (2006), pp. 530-538, [10.1111/j.1365-2389.2006.00806.x](#)

[CrossRefView Record in Scopus](#)

[Reyes and Torre](#)

I. Reyes, J. Torrent **Citrate-ascorbate as a highly selective extractant for poorly crystalline iron oxides**

Soil Sci. Soc. Am. J., 61 (1997), pp. 1647-1654

[CrossRefView Record in Scopus](#)

[Riedel et al., 201](#)

T. Riedel, D. Zak, H. Biester, T. Dittmar **Iron traps terrestrially derived dissolved organic matter at redox interfaces**

Proceedings of the National Academy of Sciences (2013), pp. 10101-

10105, [10.1073/pnas.1221487110](#)

[CrossRefView Record in Scopus](#)

[Ruiz-Dueñas an](#)

F.J. Ruiz-Dueñas, Á.T. Martínez **Microbial degradation of lignin: how a bulky recalcitrant polymer is efficiently recycled in nature and how we can take advantage of this**

Microb. Biotechnol., 2 (2009), pp. 164-177, [10.1111/j.1751-7915.2008.00078.x](#)

[CrossRefView Record in Scopus](#)

[Schmidt et al., 20](#)

M.W.I. Schmidt, M.S. Torn, S. Abiven, T. Dittmar, G. Guggenberger, I.A. Janssens, M. Kleber, I. Kögel-Knabner, J. Lehmann, D.A.C. Manning, P. Nannipieri, D.P. Rasse, S. Weiner, S.E. Trumbore **Persistence of soil organic matter as an ecosystem property**

Nature, 478 (2011), pp. 49-56, [10.1038/nature10386](#)

[CrossRefView Record in Scopus](#)

[Sutton-Grier et a](#)

A.E. Sutton-Grier, J.K. Keller, R. Koch, C. Gilmour, J.P. Megonigal **Electron donors and acceptors influence anaerobic soil organic matter mineralization in tidal marshes**
Soil Biol. Biochem., 43 (2011), pp. 1576-1583, [10.1016/j.soilbio.2011.04.008](https://doi.org/10.1016/j.soilbio.2011.04.008)
[ArticleDownload PDFView Record in Scopus](#)

[Thompson et al.](#)

A. Thompson, O.A. Chadwick, S. Boman, J. Chorover **Colloid mobilization during soil iron redox oscillations**
Environ. Sci. Technol., 40 (2006), pp. 5743-5749, [10.1021/es061203b](https://doi.org/10.1021/es061203b)
[CrossRefView Record in Scopus](#)

[Yelle et al., 2011](#)

D.J. Yelle, D. Wei, J. Ralph, K.E. Hammel **Multidimensional NMR analysis reveals truncated lignin structures in wood decayed by the brown rot basidiomycete Postia placenta**
Environ. Microbiol., 13 (2011), pp. 1091-1100, [10.1111/j.1462-2920.2010.02417.x](https://doi.org/10.1111/j.1462-2920.2010.02417.x)
[CrossRefView Record in Scopus](#)