

UCSF

UC San Francisco Previously Published Works

Title

Genome-wide association analysis of anti-TNF drug response in patients with rheumatoid arthritis

Permalink

<https://escholarship.org/uc/item/7895m4vm>

Journal

Annals of the Rheumatic Diseases, 72(8)

ISSN

0003-4967

Authors

Mirkov, Maša Umičević
Cui, Jing
Vermeulen, Sita H
[et al.](#)

Publication Date

2013-08-01

DOI

10.1136/annrheumdis-2012-202405

Peer reviewed

Published in final edited form as:

Ann Rheum Dis. 2013 August ; 72(8): 1375–1381. doi:10.1136/annrheumdis-2012-202405.

Genome-wide association analysis of anti-TNF drug response in rheumatoid arthritis patients

Maša Umičević Mirkov¹, Jing Cui², Sita H Vermeulen^{1,3}, Eli A. Stahl², Erik JM Toonen⁴, Remco R Makinje¹, Annette T Lee⁵, Tom WJ Huizinga⁶, Renee Allaart⁶, Anne Barton⁷, Xavier Mariette⁸, Corinne Miceli-Richard⁸, Lindsey A Criswell⁹, Paul P Tak^{10,#}, Niek de Vries¹⁰, Saedis Saevarsdottir¹¹, Leonid Padyukov¹¹, S. Louis Bridges¹², Dirk-Jan van Schaardenburg^{13,14}, Tim Jansen^{15,‡}, Ellen AJ Dutmer^{16,‡}, Mart van de Laar^{17,‡}, Pilar Barrera¹⁵, Timothy RDJ Radstake¹⁸, Piet LCM van Riel^{15,‡}, Hans Scheffer¹, Barbara Franke^{1,19}, Han G Brunner¹, Robert M Plenge^{2,*}, Peter K Gregersen^{5,*}, Henk-Jan Guchelaar^{20,*}, and Marieke JH Coenen¹

¹Department of Human Genetics, Radboud University Nijmegen Medical Centre, PO box 9101, 6500 HB Nijmegen, The Netherlands ²Brigham and Women's Hospital, Boston, Massachusetts and Broad Institute, Cambridge, Massachusetts, United States of America ³Department of Epidemiology, Biostatistics and HTA, Radboud University Nijmegen Medical Centre, PO box 3101, 6500 HB Nijmegen, The Netherlands ⁴Department of Medicine, Radboud University Nijmegen Medical Centre, PO box 9101, 6500 HB Nijmegen, The Netherlands ⁵The Feinstein Institute for Medical Research, North Shore-Long Island Jewish Health System, Manhasset, New York, United States of America ⁶Department of Rheumatology, Leiden University Medical Center, PO box 9600, 2300 RC Leiden, The Netherlands ⁷Arthritis Research UK Epidemiology Unit, Manchester Academy of Health Sciences, The University of Manchester, Manchester, United Kingdom; NIHR Manchester Musculoskeletal Biomedical Research Unit, Central Manchester University Hospitals NHS Foundation Trust, Manchester Academic Health Science Centre ⁸Université Paris-Sud, AP-HP, Hôpitaux Universitaires Paris-Sud, INSERM U1012, Le Kremlin Bicêtre, Paris, France ⁹Rosalind Russell Medical Research Center for Arthritis, Department of Medicine, University of California, San Francisco, CA, USA ¹⁰Department of Clinical Immunology and Rheumatology, AMC/University of Amsterdam, Amsterdam, The Netherlands ¹¹Rheumatology Unit, Department of Medicine, Karolinska Institutet/Karolinska University Hospital, Stockholm, Sweden ¹²Division of Clinical Immunology and Rheumatology, University of Alabama at Birmingham, Birmingham, United States of America ¹³Reade, Centre for Rehabilitation and Rheumatology (formerly Jan van Breemen Institute), Dept. of Rheumatology,

Corresponding author, Marieke Coenen, PhD, Department of Human Genetics (855), Nijmegen Centre for Evidence Based Practice and Institute for Genetic and Metabolic Disease, Radboud University Nijmegen Medical Centre, PO Box 9101, 6500 HB Nijmegen, The Netherlands, Tel: +31 (0)24 3617752, Fax: +31 (0)24 3668752, m.coenen@gen.umcn.nl.

#Currently: GlaxoSmithKline, Stevenage, U.K.

*authors contributed equally

‡on behalf of the Dutch Rheumatoid Arthritis Monitoring Registry (DREAM)

Competing Interest

The authors declare no competing interest.

Contributorship

All authors were involved in the design, analysis and interpretation of data. All authors have revised the manuscript and gave final approval for submission of the manuscript.

Amsterdam, The Netherlands ¹⁴VU University Medical Centre, Dept. of Rheumatology, Amsterdam, The Netherlands ¹⁵Department of Rheumatology, Radboud University Nijmegen Medical Centre, PO box 9101, 6500 HB Nijmegen, The Netherlands ¹⁶Department of Rheumatology, Gelderse Vallei Hospital, Ede, The Netherlands ¹⁷Arthritis Center Twente, University Twente & Medisch Spectrum Twente, Enschede, The Netherlands ¹⁸Rheumatology & Clinical Immunology, University Medical Center Utrecht & Wilhemina Children's Hospital, Utrecht, The Netherlands ¹⁹Department of Psychiatry, Radboud University Nijmegen Medical Centre, PO box 9101, 6500 HB Nijmegen, The Netherlands ²⁰Department of Clinical Pharmacy and Toxicology, Leiden University Medical Center, PO box 9600, 2300 RC Leiden, The Netherlands

Abstract

Background—Treatment strategies blocking tumor necrosis factor (anti-TNF) have proven very successful in patients with rheumatoid arthritis (RA). However, a significant subset of patients does not respond for unknown reasons. Currently there are no means of identifying these patients prior to treatment. This study was aimed at identifying genetic factors predicting anti-TNF treatment outcome in patient with RA using a genome-wide association approach.

Methods—We conducted a multi-stage, genome-wide association study with a primary analysis of 2,557,253 single nucleotide polymorphisms (SNPs) in 882 RA patients receiving anti-TNF therapy included through the Dutch Rheumatoid Arthritis Monitoring (DREAM) registry and the database of Apotheekzorg. Linear regression analysis of changes in the Disease Activity Score in 28 joints after 14 weeks of treatment was performed using an additive model. Markers with a $p < 10^{-3}$ were selected for replication in 1,821 RA patients from three independent cohorts. Pathway analysis including all SNPs with a p -value $< 10^{-3}$ was performed using Ingenuity.

Results—Seven hundred seventy two markers demonstrated evidence of association with treatment outcome in the initial stage. Eight genetic loci showed improved p -value in the overall meta-analysis compared to the first stage, three of which (rs1568885, rs1813443 and rs4411591) showed directional consistency over all four studied cohorts. We were unable to replicate markers previously reported to be associated with anti-TNF outcome. Network analysis indicated strong involvement of biological processes underlying inflammatory response and cell morphology.

Conclusion—Using a multi-stage strategy, we have identified 8 genetic loci associated with response to anti-TNF treatment. Further studies are required to validate these findings in additional patient collections.

Keywords

anti-TNF; gene polymorphism; pharmacogenetics; rheumatoid arthritis; genome-wide association study

Introduction

Rheumatoid arthritis (RA) is a chronic, systemic inflammatory disease characterized by polyarthritis, joint damage and functional disability[1]. RA cannot be cured and treatment is directed towards reducing the symptoms associated with the disease.

Tumor necrosis factor α (TNF α) is a pleiotropic, pro-inflammatory and immunoregulatory cytokine that plays a crucial role in RA [2, 3]. The introduction of TNF blocking agents, such as infliximab, etanercept and adalimumab, revolutionized therapy of RA, most notably because of the excellent clinical efficacy and ability of these agents to prevent further structural damage in patients who failed to respond to treatment with conventional disease-modifying anti rheumatic drugs (DMARDs) [4, 5]. Despite this success, a substantial proportion of RA patients (about 30%) treated with TNF inhibitors does not display any significant clinical improvement [6, 7]. Given the expensive treatment regime and the potential side effects associated with the treatment, the idea of a priori prediction of response to anti-TNF agents in RA patients is a highly relevant topic [8, 9].

Studies of clinical parameters and biomarkers have identified several factors that influence anti-TNF treatment outcome, including concurrent use of DMARDs, lower baseline health assessment questionnaire (HAQ) score, gender, smoking, serological status, TNF levels at the site of inflammation, and the synovial microarchitecture. However, these factors explain only a relatively small proportion of the observed variance in response ($R^2 = 17\text{--}29\%$) and are, therefore, not suitable to be used as predictors in clinical setting [9–12]. In addition, effort has been put into the identification of genetic markers predicting anti-TNF treatment outcome. Most of these studies are candidate gene-based studies focusing on polymorphisms in genes known to be involved in RA pathogenesis and genes implicated in TNF- α signaling pathways [13]. The most thoroughly investigated gene is *TNFA* encoding TNF α , the target of anti-TNF treatment. Initial studies suggested a role of a variant in the promoter of the gene ($-308\text{G}>\text{A}$) in anti-TNF response, although recent meta-analyses do not support this association [14, 15]. So far, using the candidate gene approaches, the most convincing evidence of association with response to anti-TNF therapy in RA patients is found for an RA risk allele at the *PTPRC* gene locus [16, 17].

A number of additional potential candidate loci have been suggested based on the results of three genome-wide association studies (GWAS) [18–20]. In a GWAS of 566 RA patients, Plant *et al.* demonstrated evidence of association at 7 genetic loci with response to TNF blockade, two of which mapped within genes: *PDZ domain-containing protein 2 (PDZD2)* and *eyes absent homolog 4 (EYA4)* [19]. In a small study ($n=89$) by Liu *et al.* association was reported for markers in the *MAFB* and *PON1* gene regions as well as in a region of chromosome 9 that contains the interferon kappa (*IFN- κ*), *MOBK2B* and *C9orf72* loci. The most compelling candidate for involvement in anti-TNF treatment response in this study is *IFN- κ* since type I IFNs play a definite role in inflammatory disease and autoimmunity [21]. However these results could not be replicated by others [20, 22]. Krintel *et al.* reported associations of SNPs within a non-coding region surrounded by the *TLR4* gene and the *DBC1* gene and a marker within the *FOXP1* gene with treatment outcome in a cohort of 196 Danish patients [20].

To determine whether the reported loci reflect true associations, and to search for novel loci that influence differential response to anti-TNF therapy, we performed a genome-wide association study in a cohort of 882 Dutch RA patients treated with anti-TNF therapy.

Materials and methods

Patients and study design

A multistage GWAS was performed including 984 RA patients treated with anti-TNF medication (stage 1) with subsequent follow up of the most significant signals in two replication cohorts (stage 2 (n=954) and 3 (n=867)).

For the initial GWAS analysis, patients were recruited through a collaborative effort in which 669 patients were included as part of the Dutch Rheumatoid Arthritis Monitoring (DREAM) registry (www.dreamregistry.nl) and 315 patients were enrolled through the database of ApotheekZorg which facilitates the Dutch distribution of adalimumab. All patients were diagnosed with RA according to the 1987 revised American College of Rheumatology (ACR) criteria and were treated with anti-TNF according to the indications in the Netherlands; Disease Activity Score 28 (DAS28) > 3.2 and previous failure on at least two disease-modifying antirheumatic drugs (DMARDs), one of which has to be methotrexate (MTX), all patients were biological naïve [23]. We used the DAS28 change at three months as outcome for our analysis. Patients that stopped treatment within the first three months, were not included in the study. All patients gave written informed consent and the study was approved by the ethical committees of the participating hospitals.

For stage 2, data from 954 RA cases treated with anti-TNF were selected from 9 different cohorts as part of the American College of Rheumatology Research and Education Foundation (REF) “Within Our Reach” project – *Autoimmune Biomarkers Collaborative Network (ABCoN)*, *Academic Medical Center Amsterdam (AMC)*, *Behandelstrategieën voor Rheumatoïde Arthritis (BeSt)*, *Biological in Rheumatoid arthritis Genetics and Genomics Study Syndicate (BRAGGSS)*, *Brigham Rheumatoid Arthritis Sequential Study (BRASS)*, *Epidemiological Investigation of Rheumatoid Arthritis (EIRA)*, *Immunex Early Rheumatoid Arthritis (ERA) study*, *Karolinska Institutet (KI) study*, *READE*, formerly *Jan van Breemen Institute (READE) study*, *Treatment of Early Aggressive RA (TEAR)* – this collection has been reported previously in [16, 24].

Finally, stage 3 included two previously described cohorts; (1) Wellcome Trust Case Control Consortium (WTCCC) comprising 595 RA patients from the UK [19] and (2) 272 RA patients from France ascertained through ReAct [25].

Genotyping and pre-imputation quality control

For stage 1, genotyping was performed using the Illumina HumanHap550-Duo Bead Chip or the Human660W-Quad BeadChips, according to the instructions of the manufacturer (Illumina, Inc, San Diego, USA).

Pre-imputation quality control procedures were applied using PLINK software [26]. Single nucleotide polymorphisms (SNPs) that had minor allele frequency (MAF) <0.05 and call rates <95% were excluded as well as SNPs with extreme departures from Hardy-Weinberg equilibrium ($p < 1 \times 10^{-5}$). Subsequently, quality control filtering was performed at the sample level. Four samples were excluded due to gender mismatch with phenotypic data and 21 samples due to a genotyping rate <95%. Cryptic relatedness between study participants was

examined by estimating IBD. Seven DNA samples were excluded based on a PI-HAT>0.125. Lastly, principal components were computed to adjust for population stratification using the EIGENSTRAT package [27]; 59 individuals were removed as outliers, based on the EIGENSTRAT default filter. After quality control (QC) 882 individuals were left for analysis. For the replication cohorts the same QC criteria were used.

Imputation

To obtain a marker set common to all studies and to increase overall coverage of the genome, imputation was performed using HapMap2 release 21 (downloaded from <http://www.sph.umich.edu/csg/abecasis/MACH/download/HapMap-r21.html>). Haplotype phasing using MaCH software (<http://www.sph.umich.edu/csg/abecasis/MACH/index.html>) [28] was followed by genotype imputation by Minimac (<http://genome.sph.umich.edu/wiki/Minimac>).

Post-imputation QC criteria were MAF ≥ 1% and good imputation quality, which was defined as RSQR ≥ 0.3. In total, 2,557,253 SNPs were included in the analysis.

Stage 1 GWAS

The additive genetic effect of each SNP allele on change in DAS28 at 3 months of treatment was estimated using linear regression analysis with adjustment for baseline DAS28 and the first three principal components (PC) derived using EIGENSTRAT. These analyses were performed using the Mach2qtl software package [29] (downloaded from <http://www.sph.umich.edu/csg/abecasis/MACH/download/HapMap-r21.html>).

The Dutch samples were not genotyped in one run therefore the results were analyzed using a meta-analysis approach that combines study-specific- β -estimates based on the fixed effect model and using the inverse of the variance of the study-specific- β -estimates to weigh the contribution of each study. Calculations were performed in the METAL package (www.sph.umich.edu/csg/abecasis/metal). Within-study genomic control (GC) correction was applied to the variance of β -estimates using lambda factors specific to each study ($\lambda_1 = 1.013$, $\lambda_2 = 1.016$, $\lambda_3 = 0.996$).

SNP selection for replication in stage 2 and 3

Markers demonstrating association with DAS28 change ($p < 10^{-3}$) in stage 1 were selected for replication. Pruning of hits based on LD was performed prior to replication: all SNPs with a HapMap CEU pair-wise correlation coefficient (r^2) > 0.8 with the most strongly associated SNPs in a locus were eliminated. 772 independent loci were left for replication. Replication analysis in stage 2 was carried out using existing GWA scan data from the REF collection. Those SNPs that passed the $p < 0.05$ threshold in stage 2 were further evaluated using GWA data from two collections (WTCCC and ReAct) in stage 3. A meta-analysis using the METAL package (www.sph.umich.edu/csg/abecasis/metal) was performed.

Explorative analysis for functional relation between genes identified in stage 1

All markers showing association with DAS change ($p < 10^{-3}$) in stage 1 were investigated for functional interactions by Ingenuity Pathway Analysis (IPA) software (Ingenuity Systems, www.ingenuity.com) using an unsupervised analysis. IPA computes a score for each network accordingly to the fit of the user's set of input genes. The score, representing the $-\log$ (p-value), indicates the likelihood of Focus Genes (genes harboring associated SNPs) in a network being found together due to a random chance.

Results

Figure 1 presents an overview of our study approach. The baseline characteristics of the patients included in the study are summarized in Table 1.

Genome-wide association analysis

In *stage 1* 2,448,996 SNPs were tested for association with anti-TNF outcome in the Dutch population. Of these SNPs 2,359 showed evidence of association with treatment response (p -value $< 10^{-3}$, Supplementary Table 1). LD pruning reduced the number of SNPs prioritized for replication analysis to 772.

We aimed to replicate the findings in 954 patients from the REF collection for *stage 2* of our study. 768 SNPs passed the QC in the second stage replication cohort. 39 markers showed nominally significant ($p < 0.05$) association with treatment outcome under an additive model, 20 of which demonstrated directionally consistent association and a resulting improvement of the association signal in a stage 1 and 2 combined meta-analysis (Supplementary Table 2).

In *stage 3* the 39 SNP stage 2 markers were further inspected for replication in two independent GWA studies, comprised of 595 patients from the UK and 272 patients from France (ReAct), separately. One SNP (rs11642036) failed QC criteria in both replication cohorts, leaving 38 SNPs for analysis. None of the tested SNPs showed nominal association with treatment outcome in these cohorts (Table 2). However, the meta-analysis including all cohorts showed improved association signals for eight SNPs compared to our stage 1 results, three of which, rs1568885, rs1813443 and rs4411591, demonstrate directional consistency over all four studied cohorts (Table 2).

None of the SNPs previously reported to be associated with treatment outcome showed evidence for association in the Dutch stage 1 cohort (Table 3) [16–20].

Explorative pathway analysis of stage 1

We explored the stage 1 dataset for potential functional relationship between genes that showed evidence of association at $p < 10^{-3}$ with treatment outcome using the Ingenuity Pathway Analysis (IPA). This resulted in the identification of 8 networks. The highest scoring network ($p = 10^{-41}$) included 26 genes identified in the GWAS analysis stage 1 and 9 additional interacting genes (Figure 2). Importantly, this network is predicted to be involved in metabolic disease and biological processes underlying inflammatory response and cell

morphology and contains genes implicated in TNF signaling (*NFκB*) and antibody formation (*IgG*).

Discussion

In this report, we described the results of the largest GWA study of response to anti-TNF treatment in RA patients conducted to date.

Using a multistage study design, we identified eight genetic loci demonstrating suggestive evidence of association (improved p-values) with treatment outcome in our overall meta-analysis, with three markers (rs4411591, rs1813443 and rs1568885) showing directional consistency over all four cohorts studied. In the combined cohort, eight identified loci together explain 3.8% of the variance in the treatment response. While no single SNP reached a genome-wide level of significance ($p < 5 \times 10^{-8}$), these variants represent excellent candidates for further investigation.

Of the eight markers with suggestive evidence of association, two map to an intergenic region in which the nearest gene is interesting in terms of its biological function. The SNPs, rs12142623 and rs4651370, are located ~400 kb downstream from the *phospholipase A2, group IVA (PLA2G4A)* gene. The protein encoded by *PLA2G4A* is a phospholipase enzyme involved in generation of eicosanoids, molecules with regulative function in inflammatory responses. TNF α is one of the first known stimuli for *PLA2G4A* activation, through action of both TNF receptor subtypes [30]. It might be possible that the identified SNPs influence long range regulatory elements.

In addition, three of the eight identified SNPs map within genes. The marker rs4411591 maps to the *Loc100130480* encoding an hypothetical protein, while rs2378945 is located in the *nucleotide binding protein-like (NUBPL)* gene. *NUBPL* encodes a protein required for the assembly of the respiratory chain NADH dehydrogenase (complex I). Finally, rs1813443 is located in the intronic region of *contactin 5 (CNTN5)* a member of the immunoglobulin superfamily which is thought to have a role in the formation of axon connections in the developing nervous system [31]. Little is known about the possible involvement of these genes in inflammatory disease and there are no apparent functional links to anti-TNF treatment outcome, yet. If association with anti-TNF response can be confirmed in additional replication studies, future functional studies are needed to prove the biological link with anti-TNF response.

We did not find evidence for association for the loci identified from previously published GWA studies on anti-TNF response [18–20] or for the *PTPRC* gene (Table 3) [16, 17] in the Dutch patients included in our stage 1 GWAS, suggesting that these genes do not play a major role in anti-TNF treatment outcome in our population. However, the power of our study to detect an association with *PTPRC* loci at $p < 0.0005$ in a set size of 882 patients was 26%.

However, the top network constructed with Ingenuity Pathway analysis indicated involvement of the genes showing suggestive association in stage 1 GWAS in following processes: cell morphology, metabolic disease and inflammatory response. This is in line

with the expected biological function of anti-TNF; TNF α is an important mediator of insulin resistance and it also impedes insulin-glucose-mediated uptake [32]. More importantly, studies showed a positive long-term effect of TNF α antagonists on insulin resistance, that was correlated with improvement in disease activity [33, 34]. The identified network also harbours two interesting interacting molecules: NF κ B and IgG. NF κ B is a transcription regulator that is preferentially activated by TNF and main downstream target of the TNF signaling pathway [35]. The finding that the genes identified through the GWAS are interacting with IgG is particularly interesting since response failure and side-effects of anti-TNF due to immunogenicity are not rare and it was found that anti-infliximab antibodies are exclusively of IgG isotype [36]. Both molecules might have a central role in determining the outcome of treatment with anti-TNF agents. Besides this network we could map the most prominent associations from stage 1 to the *VAV1* and *SPRED2* genes. *VAV1* has been found to protect T cells from Fas-mediated apoptosis in Jurkat leukemia T cells [37] and it has been confirmed that RA patients are showing differential sensitivity to apoptosis of peripheral blood lymphocytes induced by anti-TNF therapy [38]. *SPRED2* is a known RA risk loci [39]. This network might represent new leads and new additional candidates for future research.

Our study, which included 2703 RA patients treated with anti-TNF agents is the largest GWAS of treatment outcome to date. However, our sample size still remains modest compared to genetic studies of risk of RA [40, 41] and other complex traits [42, 43]. Also, there are important aspects that may affect our results. There is considerable disease heterogeneity in RA. In our dataset there is a difference in e.g. the number of women included in the studies but also type of anti-TNF treatment and co-medication use (Table 1). Furthermore, the REF collection used for replication in stage 2 consists of nine smaller cohorts from several populations. Hence, combining results across four different cohorts of RA patients that are rather diverse in subject ascertainment and assessment and previous treatment can lead to different effect estimates amongst studies and false negative results. In addition, the DAS28 score used as measure for treatment outcome in our study, is a composite score including four measures: swollen and tender joint counts, erythrocyte sedimentation rate and self-reported general health. This score is a powerful tool for measuring treatment response in a clinical setting. However, it is likely that this complex measure is influenced by genetic effects that are individually modest and would require large sample sizes to be detected.

In the present study, we have identified eight genetic loci that show suggestive evidence of influencing anti-TNF treatment response based on a multi-stage approach in a population of 2703 Caucasian RA patients. Our findings require further validation in independent cohorts and/or at a functional level.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

MJHC is supported by a grant from the Netherlands Genomics Initiative (93511014). RMP is supported by grants from the NIH (R01-AR057108, R01-AR056768, U01-GM092691, R01-AR059648), and holds a Career Award for Medical Scientists from the Burroughs Wellcome Fund. Funding for this project was provided by the American College of Rheumatology Research and Education Foundation. NdV was sponsored by CTMM, the Center for Translational Molecular Medicine (www.ctmm.nl), and the Dutch Arthritis Foundation, project TRACER (grant 04I-202). We are in debt to Prof Lars Klareskog and Prof Lars Alfredsson for providing samples from the EIRA study, as well as Doc Johan Askling, Sara Wedrén and the Swedish Rheumatology Register, for their help with the follow-up data. We thank Alejandro Arias-Vasquez for help in the design of the analysis.

References

1. McInnes IB, Schett G. Cytokines in the pathogenesis of rheumatoid arthritis. *Nat Rev Immunol.* 2007; 7:429–442. [PubMed: 17525752]
2. Tracey D, Klareskog L, Sasso EH, et al. Tumor necrosis factor antagonist mechanisms of action: a comprehensive review. *Pharmacol Ther.* 2008; 117:244–279. [PubMed: 18155297]
3. Feldmann M, Maini SR. Role of cytokines in rheumatoid arthritis: an education in pathophysiology and therapeutics. *Immunol Rev.* 2008; 223:7–19. [PubMed: 18613827]
4. Maini R, St Clair EW, Breedveld F, et al. Infliximab (chimeric anti-tumour necrosis factor alpha monoclonal antibody) versus placebo in rheumatoid arthritis patients receiving concomitant methotrexate: a randomised phase III trial. ATTRACT Study Group. *Lancet.* 1999; 354:1932–1939. [PubMed: 10622295]
5. Keystone EC, Kavanaugh AF, Sharp JT, et al. Radiographic, clinical, and functional outcomes of treatment with adalimumab (a human anti-tumor necrosis factor monoclonal antibody) in patients with active rheumatoid arthritis receiving concomitant methotrexate therapy: a randomized, placebo-controlled, 52-week trial. *Arthritis Rheum.* 2004; 50:1400–1411. [PubMed: 15146409]
6. Klareskog L, van der Heijde D, de Jager JP, et al. Therapeutic effect of the combination of etanercept and methotrexate compared with each treatment alone in patients with rheumatoid arthritis: double-blind randomised controlled trial. *Lancet.* 2004; 363:675–681. [PubMed: 15001324]
7. Maini RN, Breedveld FC, Kalden JR, et al. Therapeutic efficacy of multiple intravenous infusions of anti-tumor necrosis factor alpha monoclonal antibody combined with low-dose weekly methotrexate in rheumatoid arthritis. *Arthritis Rheum.* 1998; 41:1552–1563. [PubMed: 9751087]
8. Tak PP. A personalized medicine approach to biologic treatment of rheumatoid arthritis: a preliminary treatment algorithm. *Rheumatology (Oxford).* 2012; 51:600–609. [PubMed: 21890615]
9. Hyrich KL, Watson KD, Silman AJ, et al. Predictors of response to anti-TNF-alpha therapy among patients with rheumatoid arthritis: results from the British Society for Rheumatology Biologics Register. *Rheumatology (Oxford).* 2006; 45:1558–1565. [PubMed: 16705046]
10. Atzeni F, Antivalle M, Pallavicini FB, et al. Predicting response to anti-TNF treatment in rheumatoid arthritis patients. *Autoimmun Rev.* 2009; 8:431–437. [PubMed: 19211043]
11. Potter C, Hyrich KL, Tracey A, et al. Association of rheumatoid factor and anti-cyclic citrullinated peptide positivity, but not carriage of shared epitope or PTPN22 susceptibility variants, with anti-tumour necrosis factor response in rheumatoid arthritis. *Ann Rheum Dis.* 2009; 68:69–74. [PubMed: 18375541]
12. Klaasen R, Thurlings RM, Wijbrandts CA, et al. The relationship between synovial lymphocyte aggregates and the clinical response to infliximab in rheumatoid arthritis: a prospective study. *Arthritis Rheum.* 2009; 60:3217–3224. [PubMed: 19877042]
13. Coenen MJ, Toonen EJ, Scheffer H, et al. Pharmacogenetics of anti-TNF treatment in patients with rheumatoid arthritis. *Pharmacogenomics.* 2007; 8:761–773. [PubMed: 17638513]
14. Pavy S, Toonen EJ, Miceli-Richard C, et al. Tumour necrosis factor alpha-308G->A polymorphism is not associated with response to TNFalpha blockers in Caucasian patients with rheumatoid arthritis: systematic review and meta-analysis. *Ann Rheum Dis.* 2010; 69:1022–1028. [PubMed: 19966089]

15. Lee YH, Ji JD, Bae SC, et al. Associations between tumor necrosis factor- α (TNF- α)-308 and-238 G/A polymorphisms and shared epitope status and responsiveness to TNF- α blockers in rheumatoid arthritis: a metaanalysis update. *J Rheumatol.* 2010; 37:740–746. [PubMed: 20194454]
16. Cui J, Saevarsdottir S, Thomson B, et al. Rheumatoid arthritis risk allele PTPRC is also associated with response to anti-tumor necrosis factor alpha therapy. *Arthritis Rheum.* 2010; 62:1849–1861. [PubMed: 20309874]
17. Plant D, Prajapati R, Hyrich KL, et al. Replication of association of the PTPRC gene with response to anti-tumor necrosis factor therapy in a large UK cohort. *Arthritis Rheum.* 2012; 64:665–670. [PubMed: 21952740]
18. Liu C, Batliwalla F, Li W, et al. Genome-wide association scan identifies candidate polymorphisms associated with differential response to anti-TNF treatment in rheumatoid arthritis. *Mol Med.* 2008; 14:575–581. [PubMed: 18615156]
19. Plant D, Bowes J, Potter C, et al. Genome-wide association study of genetic predictors of anti-tumor necrosis factor treatment efficacy in rheumatoid arthritis identifies associations with polymorphisms at seven loci. *Arthritis Rheum.* 2011; 63:645–653. [PubMed: 21061259]
20. Krintel SB, Palermo G, Johansen JS, et al. Investigation of single nucleotide polymorphisms and biological pathways associated with response to TNF α inhibitors in patients with rheumatoid arthritis. *Pharmacogenetics and genomics.* 2012; 22:577–589. [PubMed: 22569225]
21. Selmi C, Lleo A, Zuin M, et al. Interferon alpha and its contribution to autoimmunity. *Curr Opin Investig Drugs.* 2006; 7:451–456.
22. Suarez-Gestal M, Perez-Pampin E, Calaza M, et al. Lack of replication of genetic predictors for the rheumatoid arthritis response to anti-TNF treatments: a prospective case-only study. *Arthritis Res Ther.* 2010; 12:R72. [PubMed: 20423481]
23. Kievit W, Fransen J, Adang EM, et al. Long-term effectiveness and safety of TNF-blocking agents in daily clinical practice: results from the Dutch Rheumatoid Arthritis Monitoring register. *Rheumatology (Oxford).* 2011; 50:196–203. [PubMed: 21078625]
24. Moreland LW, O'Dell JR, Paulus HE, et al. A randomized comparative effectiveness study of oral triple therapy versus etanercept plus methotrexate in early, aggressive rheumatoid arthritis. *Arthritis Rheum.* 2012
25. Bombardieri S, Ruiz AA, Fardellone P, et al. Effectiveness of adalimumab for rheumatoid arthritis in patients with a history of TNF-antagonist therapy in clinical practice. *Rheumatology (Oxford).* 2007; 46:1191–1199. [PubMed: 17504821]
26. Purcell S, Neale B, Todd-Brown K, et al. PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet.* 2007; 81:559–575. [PubMed: 17701901]
27. Price AL, Patterson NJ, Plenge RM, et al. Principal components analysis corrects for stratification in genome-wide association studies. *Nat Genet.* 2006; 38:904–909. [PubMed: 16862161]
28. Li Y, Willer C, Sanna S, et al. Genotype imputation. *Annu Rev Genomics Hum Genet.* 2009; 10:387–406. [PubMed: 19715440]
29. Chen WM, Abecasis GR. Family-based association tests for genomewide association scans. *Am J Hum Genet.* 2007; 81:913–926. [PubMed: 17924335]
30. Jupp OJ, Vandenabeele P, MacEwan DJ. Distinct regulation of cytosolic phospholipase A2 phosphorylation, translocation, proteolysis and activation by tumour necrosis factor-receptor subtypes. *Biochem J.* 2003; 374:453–461. [PubMed: 12786601]
31. Ogawa J, Kaneko H, Masuda T, et al. Novel neural adhesion molecules in the Contactin/F3 subgroup of the immunoglobulin superfamily: isolation and characterization of cDNAs from rat brain. *Neurosci Lett.* 1996; 218:173–176. [PubMed: 8945756]
32. Peraldi P, Hotamisligil GS, Buurman WA, et al. Tumor necrosis factor (TNF)- α inhibits insulin signaling through stimulation of the p55 TNF receptor and activation of sphingomyelinase. *J Biol Chem.* 1996; 271:13018–13022. [PubMed: 8662983]
33. Huvers FC, Popa C, Netea MG, et al. Improved insulin sensitivity by anti-TNF α antibody treatment in patients with rheumatic diseases. *Ann Rheum Dis.* 2007; 66:558–559. [PubMed: 17360784]

34. Tam LS, Tomlinson B, Chu TT, et al. Impact of TNF inhibition on insulin resistance and lipids levels in patients with rheumatoid arthritis. *Clin Rheumatol.* 2007; 26:1495–1498. [PubMed: 17237906]
35. Nagar M, Jacob-Hirsch J, Vernitsky H, et al. TNF activates a NF-kappaB-regulated cellular program in human CD45RA- regulatory T cells that modulates their suppressive function. *J Immunol.* 2010; 184:3570–3581. [PubMed: 20181891]
36. Svenson M, Geborek P, Saxne T, et al. Monitoring patients treated with anti-TNF-alpha biopharmaceuticals: assessing serum infliximab and anti-infliximab antibodies. *Rheumatology (Oxford).* 2007; 46:1828–1834. [PubMed: 18032541]
37. Yin J, Wan YJ, Li SY, et al. The distinct role of guanine nucleotide exchange factor Vav1 in Bcl-2 transcription and apoptosis inhibition in Jurkat leukemia T cells. *Acta Pharmacol Sin.* 2011; 32:99–107. [PubMed: 21151158]
38. Coury F, Ferraro-Peyret C, Le Cam S, et al. Peripheral blood lymphocytes from patients with rheumatoid arthritis are differentially sensitive to apoptosis induced by anti-tumour necrosis factor-alpha therapy. *Clin Exp Rheumatol.* 2008; 26:234–239. [PubMed: 18565243]
39. Baum L, Ng HK, Wong KS, et al. Associations of apolipoprotein E exon 4 and lipoprotein lipase S447X polymorphisms with acute ischemic stroke and myocardial infarction. *Clin Chem Lab Med.* 2006; 44:274–281. [PubMed: 16519597]
40. Okada Y, Terao C, Ikari K, et al. Meta-analysis identifies nine new loci associated with rheumatoid arthritis in the Japanese population. *Nat Genet.* 2012; 44:511–516. [PubMed: 22446963]
41. Stahl EA, Raychaudhuri S, Remmers EF, et al. Genome-wide association study meta-analysis identifies seven new rheumatoid arthritis risk loci. *Nat Genet.* 2010; 42:508–514. [PubMed: 20453842]
42. Lango Allen H, Estrada K, Lettre G, et al. Hundreds of variants clustered in genomic loci and biological pathways affect human height. *Nature.* 2010; 467:832–838. [PubMed: 20881960]
43. Chasman DI, Pare G, Zee RY, et al. Genetic loci associated with plasma concentration of low-density lipoprotein cholesterol, high-density lipoprotein cholesterol, triglycerides, apolipoprotein A1, and Apolipoprotein B among 6382 white women in genome-wide analysis with replication. *Circ Cardiovasc Genet.* 2008; 1:21–30. [PubMed: 19802338]




Figure 1. Study design of a multi-stage GWAS of response to anti-TNF medication in RA patients

We started out with meta-analysis of GWAS data from Dutch cohort comprised of 882 RA patients treated with anti-TNF medication. We selected 772 SNPs that reached $p < 0.001$ that were further followed up in stage 2 samples ($N=954$ individuals). 38 SNPs out of 768 investigated in stage 2 passed $p < 0.05$ and were further investigated in two separate cohorts in stage 3 ($N=595$ and $N=272$ individuals).

canonical information stored in the Ingenuity Knowledge Base. Nodes are displayed using various shapes that represent the functional classes of the gene product (□ cytokines, ◇ enzyme, ◎ complex/group, △ transporter, ○ transcription regulator, ○ transmembrane receptor, □ ion channel, □ ligand-dependent nuclear receptor, ▽ kinase, □ growth factor, ○ other).

Table 1

Study population characteristics

	Stage 1			Stage 2		Stage 3	
	Combined	DREAM	ApotheekZorg	REF Col*	WTCCC	ReAct	
number	984	669(69.7)	315(30.3)	954	595	272	
Gender female, %	68.6	67.8	70.4	75.6	77.3	77.9	
AntiTNF agent							
Infliximab	225(22.9)	225(33.7)		415(43.5)	268(45.0)		
Adalimumab	638(64.8)	323(48.2)	315(100)	174(18.2)	68(11.4)	272(100)	
Etanercept	121(12.3)	121(18.1)		365(38.3)	259(43.6)		
MTX co-medication, %	73.4	69.2	82.2	65.6	85.6	50	
DAS28							
baseline	5.5 ± 1.2	5.3 ± 1.3	5.8 ± 1.0	5.5 ± 1.2	6.7 ± 0.9	5.9 ± 1.0	
DeltaDAS, 14 weeks	3.6 ± 1.3	3.9 ± 1.3	3.1 ± 1.1				

Numbers are depicted as n(%) or mean ± standard deviation.

* The American College of Rheumatology Research and Education Foundation(REF) collection used for Stage 2 is composed of 9 different cohorts – Autoimmune Biomarkers Collaborative Network(ABCN), Academic Medical Center Amsterdam(AMC), Behandelstrategieën voor Rheumatoïde Arthritis(BeSt), Biological in Rheumatoid arthritis Genetics and Genomics Study Syndicate(BRAGGS), Birgham Rheumatoid Arthritis Sequential Study(BRASS),Epidemiological Investigation of Rheumatoid Arthritis(EIRA), Immunex Early Rheumatoid Arthritis(ERA) study, Karolinska Institutet(KI) study, Jan van Breemen Institute(READE) study, Treatment of Early Aggressive RA(TEAR)

Table 2

Association results for replicated SNPs in all cohorts and final meta-analysis

SNP	Chr	Position	Allele	Stage1(n=882)			Stage2(n=954)			Stage3(n=867)			Combined metaanalysis		
				PGC	β	P	PGC	β	P	WTCCC	PGC	β	P	ReAct	PGC
rs4411591	18	6540117	C	7.33×10^{-4}	0.248	0.02206	0.208	0.483	0.097	0.461	0.114	5.14×10^{-5}	0.202		
rs7767069	6	68827284	A	3.41×10^{-4}	-0.188	0.02839	-0.161	0.532	0.057	0.086	-0.203	8.34×10^{-5}	-0.144		
rs4651370	1	185505715	A	2.54×10^{-4}	0.256	5.28×10^{-3}	0.271	0.867	-0.021	0.921	0.015	1.09×10^{-4}	0.190		
rs1813443	11	99516221	C	3.50×10^{-4}	-0.195	0.04021	-0.155	0.605	-0.048	0.403	-0.097	1.37×10^{-4}	-0.148		
rs1447722	3	141037143	C	5.46×10^{-4}	0.193	7.63×10^{-4}	0.185	0.881	-0.014	0.573	-0.064	1.62×10^{-4}	0.134		
rs1568885	7	13604056	A	5.93×10^{-4}	0.255	0.0306	0.201	0.594	0.061	0.5	0.095	1.69×10^{-4}	0.185		
rs12142623	1	185557029	A	4.19×10^{-4}	0.248	7.68×10^{-3}	0.267	0.484	-0.087	0.398	0.126	2.04×10^{-4}	0.185		
rs2378945	14	31370541	A	8.61×10^{-4}	-0.166	0.01313	-0.171	0.489	0.061	0.905	0.013	6.88×10^{-4}	-0.115		

Chr, chromosome; MAF, minor allele frequency; PGC, p values with genomic control correction applied. Table lists all SNPs that have improved their p-value in meta-analysis compared to initial GWAS. rs4411591, rs1813443 and rs1568885 demonstrated directional consistency over all four cohorts studied.

Table 3

Initial GWAS association results for loci previously reported to be associated with anti-TNF treatment response

SNP	Chr	MAF	Gene	P _{GC}	β
SNPs from Liu et al.					
rs983332	1	0.18	-	0.603	0.031
rs928655	1	0.23	GBP6	0.051	0.109
rs13393173	2	0.19	LASS6	0.71	-0.006
rs437943	4	0.34	-	0.93	0.0006
rs10945919	6	0.32	AKO93144	0.44	0.03
rs854555	7	0.33	PONI	0.61	0.018
rs854548	7	0.25	PONI	0.61	0.0102
rs854547	7	0.35	PONI	0.52	0.0225
rs868856	9	0.31	MOBK2B	0.65	-0.0218
rs2814707	9	0.26	MOBK2B	0.95	-0.002
rs3849942	9	0.26	C9orf72	0.98	-0.0069
rs774359	9	0.27	C9orf72	0.89	-0.0159
rs6138150	20	0.18	-	0.66	0.0329
rs6028945	20	0.14	-	0.78	-0.0635
rs6071980	20	0.12	-	0.74	-0.0026
SNPs from Plant et al.					
rs12081765	1	0.39	-	0.73	-0.024
rs4694890	4	0.49	TEC	0.4	0.042
rs1532269	5	0.43	PDZD2	0.92	0.051
rs17301249	6	0.18	EYA4	0.36	-0.05
rs1350948	11	0.14	-	0.38	0.0642
rs7305646	12	0.50	-	0.39	0.05
rs7962316	12	0.40	BC118985	0.92	-0.026
SNPs from Krinetel et al.					
rs10520789	15	0.13	NR2F2	0.47	0.071
rs11870477	17	0.13	MAP2K6	0.67	0.073
rs16973982	15	0.14	NR2F2	0.60	0.072

SNP	Chr	MAF	Gene	P _{GC}	β
rs8046065	16	0.10	CREBBP	0.12	0.146
rs885814	1	0.32	ALPL	0.49	0.053
rs869179	1	0.34	ALPL	0.75	0.016
rs2722824	9	0.30	TLR4	0.33	-0.053
rs885813	1	0.43	ALPL	0.30	0.049
rs1875620	9	0.46	C9orf47	0.26	0.051
rs11525966	9	0.45	C9orf47	0.19	0.051
PTPRC					
rs10919563	1	0.13	PTPRC	0.384	-0.079

Chr, chromosome; MAF, minor allele frequency; PGC, p values with genomic control correction applied.