

# UC Irvine

## UC Irvine Previously Published Works

### Title

Fossil methane source dominates Cariaco Basin water column methane geochemistry

### Permalink

<https://escholarship.org/uc/item/7cw0c9mh>

### Journal

Geophysical Research Letters, 32(12)

### ISSN

0094-8276

### Authors

Kessler, JD  
Reeburgh, WS  
Southon, J  
[et al.](#)

### Publication Date

2005-06-01

### DOI

10.1029/2005gl022984

### Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

## Fossil methane source dominates Cariaco Basin water column methane geochemistry

J. D. Kessler,<sup>1</sup> W. S. Reeburgh,<sup>1</sup> J. Southon,<sup>1</sup> and R. Varela<sup>2</sup>

Received 14 March 2005; revised 16 May 2005; accepted 25 May 2005; published 25 June 2005.

[1] Natural radiocarbon measurements on methane ( $^{14}\text{C-CH}_4$ ) extracted from the Cariaco Basin water column show that 98% of the methane in Cariaco Basin waters is derived from fossil (radiocarbon-free) sources. Previous work on Cariaco Basin methane ( $\text{CH}_4$ ) considered only a diagenetic sediment source. Similar measurements of sediment  $^{14}\text{C-CH}_4$  indicate that sediment  $\text{CH}_4$  is produced from modern particulate material; thus the sediment and water column  $\text{CH}_4$  have distinct sources. Using time-dependent  $\text{CH}_4$  geochemical box models which include a fossil seep  $\text{CH}_4$  source term, we estimate 1)  $0.024\text{--}0.028 \text{ Tg y}^{-1}$  of seep  $\text{CH}_4$  are added to the Cariaco Basin water column, 2) the water column  $\text{CH}_4$  will reach a steady-state concentration by the year 2065, and 3) the seep  $\text{CH}_4$  inputs possibly began in 1967, following the July 30 Caracas earthquake. Oxidizing this  $\text{CH}_4$  to dissolved inorganic carbon does not appear to affect Cariaco Basin  $^{14}\text{C}$  chronologies. **Citation:** Kessler, J. D., W. S. Reeburgh, J. Southon, and R. Varela (2005), Fossil methane source dominates Cariaco Basin water column methane geochemistry, *Geophys. Res. Lett.*, 32, L12609, doi:10.1029/2005GL022984.

### 1. Introduction

[2] The Cariaco Basin has been an ideal site for studies of anoxic geochemistry and, because the sediments are unbioturbated, for paleoclimatic studies. The Cariaco Basin is located on the continental margin of Venezuela ( $10^\circ 30' \text{N}$ ,  $65^\circ \text{W}$ ), is almost 1400 m deep, and is separated from the tropical Atlantic by 150 m sills. Anoxic conditions are maintained below a depth between 250–300 m by thermal stratification and high flux of organic matter to the deep basin. High sedimentation rates, plus lack of bioturbation due to anoxia, have led to the extensive use of Cariaco Basin sediment cores for developing high-resolution paleoclimate records [Hughen *et al.*, 1996b; Lin *et al.*, 1997; Hughen *et al.*, 2000; Peterson *et al.*, 2000; Haug *et al.*, 2001] and  $^{14}\text{C}$  calibration chronologies [Hughen *et al.*, 2000, 2004].

[3] Anoxic conditions in the Cariaco Basin were discovered by L. V. Worthington in 1954 and were followed by studies on the stoichiometry of anaerobic degradation of organic matter [Richards and Vaccaro, 1956]. Steady-state

vertical advection-diffusion models were used to explain silica, sulfide, phosphate [Fanning and Pilson, 1972], and methane ( $\text{CH}_4$ ) [Reeburgh, 1976] distributions. Temporal changes in the hydrography, depth of the anoxic/oxic interface, and concentration of several constituents (including  $\text{CH}_4$ ) in the Cariaco Basin water column led to the development of a time-dependent box model [Scranton *et al.*, 1987; Scranton, 1988] of the anoxic region of the Cariaco Basin. Methane modeling studies [Reeburgh, 1976; Scranton, 1988] showed that the  $\text{CH}_4$  geochemistry could be explained with a diagenetic sediment  $\text{CH}_4$  source, no water column  $\text{CH}_4$  production, and water column anaerobic  $\text{CH}_4$  oxidation. Ward *et al.* [1987] measured Cariaco Basin water column aerobic and anaerobic  $\text{CH}_4$  oxidation rates using  $^{14}\text{C}$ -labelled  $\text{CH}_4$ . Although there are written and anecdotal reports of natural gas releases in this region [von Humboldt, 1900], this previous work gave no indication that seeps were an important Cariaco Basin  $\text{CH}_4$  source.

### 2. Experimental


[4] Here we present the first natural radiocarbon measurements on Cariaco Basin water column and sediment methane ( $^{14}\text{C-CH}_4$ ). Water and sediment samples were collected from January 21–24, 2004 on board the B/O *Hermano Gines* in the deepest portion of the Eastern basin ( $10.5^\circ \text{N}$ ,  $64.66^\circ \text{W}$ , 1370 m) at the time-series station used by the CARbon Retention In A Colored Ocean (CARIACO) program [Scranton *et al.*, 2001; Astor *et al.*, 2003].

[5] Seawater  $\text{CH}_4$  concentration analyses were conducted by filling 160 mL serum vials directly from Niskin bottles and inserting a 13 mL helium (He) headspace. Sediment  $\text{CH}_4$  concentration analyses were conducted by making a slurry of 3 mL of sediment and 6 mL of He purged water in 37.5 mL serum vials. All vials were poisoned with a saturated mercuric chloride solution and sealed with blue butyl rubber stoppers and crimp caps. After the samples were allowed to equilibrate, three aliquots of the headspace were removed for concentration analysis with gas chromatography (GC) and flame ionization detection. The results have been corrected for the amount of  $\text{CH}_4$  still dissolved in solution [Yamamoto *et al.*, 1976].

[6] Shipboard extraction of  $\text{CH}_4$  for radiocarbon analysis was conducted using Lamont radon stripping boards (J. D. Kessler and W. S. Reeburgh, Preparation of natural methane samples for stable isotope and radiocarbon analysis, submitted to *Limnology and Oceanography: Methods*, 2005, hereinafter referred to as Kessler and Reeburgh, submitted manuscript, 2005). Evacuated 20 L glass carboys are filled with 19 L of seawater directly from Niskin bottles, connected to the stripping

<sup>1</sup>Department of Earth System Science, University of California, Irvine, California, USA.

<sup>2</sup>Fundación La Salle de Ciencias Naturales, Estación de Investigaciones Marinas de Margarita, Isla de Margarita, Venezuela.



**Figure 1.** (a) Natural radiocarbon results for Cariaco Basin water column and (b) sediment methane ( $^{14}\text{C-CH}_4$ ) expressed as percent Modern Carbon (pMC). Note the differences in horizontal and vertical scales between the water column (Figure 1a) and sediment (Figure 1b). Error bars are  $\pm 1 \sigma$ . The dashed and solid lines in Figure 1a are predicted  $^{14}\text{C-CH}_4$  profiles based on models with and without mid-depth intrusions, respectively. The main difference between the model results is attributed to different eddy diffusion coefficients.

boards [Mathieu *et al.*, 1988], and purged with He to extract the dissolved  $\text{CH}_4$ . The extracted  $\text{CH}_4$  is trapped in a U-trap at liquid nitrogen temperature. The U-traps were fabricated with electropolished stainless steel tubing (3/8" OD, 2 feet long) bent in a "U" shape, filled with a molecular sieve (HiSiv 3000 in the 1/16" pellet form), and equipped with nonrotating-stem needle valves with PEEK stem tips (Swagelok D-Series). HiSiv 3000 (UOP Molsiv Adsorbents) was chosen as the trapping medium due to its trapping efficiency, lack of isotopic fractionation, and ability to quantitatively trap  $\text{CH}_4$  at  $-172^\circ\text{C}$ . The traps are reusable and are reactivated between each use (He flow at  $0.5 \text{ L min}^{-1}$ ,  $275^\circ\text{C}$ ) with an oven designed to heat the traps without damaging the valves.

[7] In order to extract, purify, and oxidize the  $\text{CH}_4$  collected in the U-traps, a two-stage purification-oxidation vacuum line has been developed at UC Irvine (Kessler and Reeburgh, submitted manuscript, 2005). The first stage of this vacuum line is a continuous loop designed to extract the  $\text{CH}_4$  from the U-traps and pass it through traps designed to remove  $\text{H}_2\text{O}$ ,  $\text{CO}_2$ ,  $\text{CO}$ , and non- $\text{CH}_4$  hydrocarbons. Analysis of our purified sample gas with Quadrupole Mass Spectroscopy at the National Institute of Standards and Technology [Currie *et al.*, 2000] showed no detectable traces of carbonaceous impurities. Next, the purified  $\text{CH}_4$  is injected into an adjoining oxidation loop on the same vacuum line which continuously circulates the purified  $\text{CH}_4$  through a  $975^\circ\text{C}$   $\text{CuO}$  furnace converting it to  $\text{CO}_2$  and  $\text{H}_2\text{O}$ .

[8] An aliquot of the  $\text{CO}_2$  produced from  $\text{CH}_4$  oxidation was converted to graphite with hydrogen reduction catalyzed by iron [Vogel *et al.*, 1984] and analyzed with  $^{14}\text{C}$  Accelerator Mass Spectrometry (AMS) at the Keck Carbon Cycle AMS facility at UC Irvine. We have shown that these procedures are quantitative, do not cause isotope fractionation, and that the blanks are small ( $0.477 \pm 0.24 \mu\text{moles of CH}_4$ ;  $^{14}\text{C-CH}_4 = 96.1 \pm 0.3 \text{ pMC}$  (percent

Modern Carbon)) [Stuiver and Polach, 1977; Kessler and Reeburgh, submitted manuscript, 2005].

[9] At water depths  $\leq 300 \text{ m}$ , the  $\text{CH}_4$  concentration was too low to collect enough  $\text{CH}_4$  carbon for a conventional AMS measurement. Small sample AMS measurements were conducted at these depths ( $6 \mu\text{moles of carbon}$  instead of the conventional  $83 \mu\text{moles of carbon}$ ). At depths  $\leq 250 \text{ m}$ , the amount of  $\text{CH}_4$  carbon collected required dilution with  $^{14}\text{C}$ -devoid  $\text{CO}_2$  to conduct a successful AMS measurement [Currie *et al.*, 2000]. Diluted samples have been back corrected.


### 3. Results

[10] Since previous work suggested a diagenetic sediment source of  $\text{CH}_4$  to the water column, the  $^{14}\text{C-CH}_4$  results (Figure 1a) were unexpected; most of the Cariaco Basin water column  $\text{CH}_4$  is almost completely devoid of radiocarbon (2.5 pMC), though the  $^{14}\text{C}$  content increases above 400 m. In contrast, the sediment  $\text{CH}_4$  contains significant amounts of  $^{14}\text{C}$  (86.4 pMC at 45 cm depth) indicating the  $\text{CH}_4$  is derived from relatively modern particulate carbon (Figure 1b). Since the turnover time of  $\text{CH}_4$  in the Cariaco Basin water column is short (50–60 years), as calculated from our  $\text{CH}_4$  concentration data and specific  $\text{CH}_4$  oxidation rates [Ward *et al.*, 1987], these measurements clearly indicate the presence of a large fossil source of  $\text{CH}_4$ . Since the temperature of the Cariaco Basin is too high ( $16.9^\circ\text{C}$ ) for hydrate formation, this fossil  $\text{CH}_4$  is most likely from previously unknown seeps.

[11] Primarily to quantify the input of seep  $\text{CH}_4$ , we modified Scranton's time-dependent Cariaco box model [Scranton, 1988] to include seeps. We adhered to the measured specific anaerobic  $\text{CH}_4$  oxidation rates [Ward *et al.*, 1987]. Using a procedure similar to Scranton [1988], the diffusive diagenetic sedimentary flux was calculated as  $0.20 \mu\text{moles cm}^{-2} \text{ y}^{-1}$  from our measured sediment  $\text{CH}_4$  concentration data ( $17.97 \mu\text{M}$  at 2 cm depth) and was held constant over the entire basin.

[12] The model was initiated with a water column  $\text{CH}_4$  concentration profile measured in February 1974 [Reeburgh, 1976; Wiesenburg, 1975] that was intercalibrated with other data sets by Scranton [1988]. The time-dependent model then predicted a seep-source profile, which shows large seep inputs at 870 m and at 1370 m (Figure 2). With this seep profile, the model predicts a 2004  $\text{CH}_4$  concentration profile that agrees with the measured 2004 profile to better than 1% below 650 m depth. (The GC  $\text{CH}_4$  concentrations agree with those calculated from the stripped and trapped  $\text{CH}_4$  within 3% on average below 300 m depth. If a homogeneous distribution of  $\text{CH}_4$  inputs is modeled similar to Scranton [1988], the agreement between the measured and modeled 2004  $\text{CH}_4$  concentration profiles decreases to between 30.5–42.3% on average below 650 m depth.) Above 650 m, the model over-predicts the measured concentration profile (Figure 2). Higher specific  $\text{CH}_4$  oxidation rates than those reported by Ward *et al.* [1987], as well as increased vertical resolution, are needed to achieve better agreement; this was also observed by Scranton [1988].

[13] Scranton's vertical time-dependent box model [Scranton *et al.*, 1987; Scranton, 1988] has been superseded in a strict sense by subsequent research documenting



**Figure 2.** Modeled water column methane concentration distributions excluding mid-depth intrusions unless otherwise noted: (triangles) 1974, (diamonds) 1986, (open squares) 2004, (shaded squares) 2004 (including mid-depth intrusions), (circles) 2050, (pluses) 2065. Measured water column methane concentration distributions: (dotted line) February 1974 [Reeburgh, 1976; Wiesenburg, 1975], (dashed line) February to March 1986 [Ward *et al.*, 1987], (dash-dotted line) January 2004. The solid gray and black lines represent the model-predicted seep methane input profiles with and without mid-depth intrusions, respectively. The main difference between the seep inputs is attributed to different eddy diffusion coefficients. Each box is 92 m deep.

intrusions. However the seep source is so dominant that the effects of episodic intrusions are overwhelmed and the original model can be modified to quantify the seep input. Holmén and Rooth [1990] found that in order to jointly explain temperature, salinity, and tritium distributions in the Cariaco Basin, the model presented by Scranton *et al.* [1987] must be modified to include injection of warm hypersaline shelf waters which reach the bottom of the basin and input of Caribbean thermocline waters at the sill. Scranton *et al.* [2001] observed water with sufficient density to reach the basin bottom north of the Eastern sill, however they did not observe hypersaline shelf waters reaching the bottom of the basin. Instead they proposed that a turbidity current associated with a July 1997 earthquake transported oxidized iron into the deep basin decreasing  $\text{H}_2\text{S}$  concentrations. No  $\text{CH}_4$  decrease was observed with this event and the  $\text{CH}_4$  concentration followed approximately a linear increase over a 30 year period [Scranton *et al.*, 2001] as predicted by the original time-dependent model [Scranton, 1988]. Scranton *et al.* [2001] and Astor *et al.* [2003] documented episodic mid-depth (250–350 m) intrusions of oxygenated water ( $\leq 30 \mu\text{M}$ ). These oxygen intrusions may increase the  $\text{CH}_4$  oxidation rate in the upper box (242–366 m) of the time-dependent model [Scranton, 1988] beyond what was measured by Ward *et al.* [1987]. The highest specific  $\text{CH}_4$  oxidation rate measured by Ward *et al.* [1987] was  $2.5 \times 10^{-3} \text{ day}^{-1}$  at 240 m depth, possibly corresponding to an oxygen intrusion, but decreased rapidly to  $1.5 \times$

$10^{-4} \text{ day}^{-1}$  averaged over the upper box. As an upper estimate on the effects of episodic mid-depth oxygen intrusions, we doubled the highest specific oxidation rate measured in the entire water column and applied it continuously to the upper box; the agreement between the measured and modeled 2004 water column  $\text{CH}_4$  concentration profiles decreased to only 3.6 % on average below 650 m depth. We did not model intrusions to the deep basin, because Scranton *et al.* [2001] showed that  $\text{CH}_4$  concentration has not responded to deep basin intrusions.

[14] The model is more sensitive to the values of the eddy diffusion coefficients. If the eddy diffusivities are changed from  $0.49\text{--}4.08 \text{ cm}^2 \text{ s}^{-1}$  [Scranton, 1988] to  $0.6\text{--}2.0 \text{ cm}^2 \text{ s}^{-1}$  [Holmén and Rooth, 1990], the agreement between the measured and modeled 2004 water column  $\text{CH}_4$  concentration profiles decreases to 23.7 % on average below 650 m depth. Even increasing the oxidation rate in the upper box to model oxygen intrusions, the seep  $\text{CH}_4$  fluxes must decrease by 15 % in order to cause 1 % agreement (Figure 2). We can use the sediment and seep  $\text{CH}_4$  inputs to model  $^{14}\text{C}\text{-CH}_4$  profiles (Figure 1a). Both modeled profiles agree well with the measured  $^{14}\text{C}\text{-CH}_4$  profile.

#### 4. Discussion

[15] Averaged over the anoxic region of the basin, our results indicate that  $0.024\text{--}0.028 \text{ Tg y}^{-1}$  ( $0.14\text{--}0.17 \text{ mole m}^{-2} \text{ y}^{-1}$ ) of seep  $\text{CH}_4$  and  $3 \times 10^{-4} \text{ Tg y}^{-1}$  of sediment  $\text{CH}_4$  are being added to the water column, while  $0.01 \text{ Tg y}^{-1}$  of  $\text{CH}_4$  are removed by anaerobic oxidization. Since  $\text{CH}_4$  oxidation in year 2004 only consumes one-third of the source, the bottom water  $\text{CH}_4$  concentration has more than doubled since 1974 (Figure 2). Since  $\text{CH}_4$  oxidation follows pseudo-first order kinetics [Ward *et al.*, 1987], oxidation rates will increase; both models (with and without intrusions) forecast that the Cariaco Basin will reach a steady-state ( $<0.01\%$  increase  $\text{y}^{-1}$ )  $\text{CH}_4$  concentration by year 2065 (Figure 2). Hindcasting with the models indicates the seep inputs began sometime between 1958 and 1967. If the seep inputs are initiated in 1967, then the model predicts water column  $\text{CH}_4$  concentrations that agree well with the measured 1974, 1986, and 2004  $\text{CH}_4$  concentration profiles (Figure 2). Coincidentally, a moderate-sized earthquake ( $M_w = 6.6$ ), whose epicenter was in the Caribbean Sea, 70 km NNW of Caracas (the Caracas earthquake of 1967), occurred on July 30, 1967 [Suárez and Nábělek, 1990]. This event could have initiated the release of fossil  $\text{CH}_4$  into the Cariaco Basin. The projected paths of the El Pilar and San Sebastián faults are parallel to the major axis of the Cariaco Basin [Audemard *et al.*, 2005; Suárez and Nábělek, 1990], however, their exact locations within the basin are unknown. The hindcasting results could also be interpreted as the result of a complete basin flushing event in the mid-1900s.

[16] Could oxidation of this fossil source to dissolved inorganic carbon (DIC) influence Cariaco Basin DIC  $^{14}\text{C}$  distributions and bias the marine reservoir correction used in Cariaco Basin radiocarbon chronologies [Hughen *et al.*, 2004, 2000]? Present evidence indicates that any bias is negligible. Measured modern (prebomb) Cariaco Basin reservoir corrections [Guilderson *et al.*, 2005; Hughen *et*

al., 1996a] are similar to open ocean values [Druffel et al., 1989], as are surface DIC concentrations and  $\delta^{13}\text{C}$  values [Deuser, 1973; Druffel et al., 1989]. Reservoir-corrected Cariaco Basin sediment foraminifera  $^{14}\text{C}$  data show excellent agreement with the INTCAL98 tree ring  $^{14}\text{C}$  calibration [Hughen et al., 2000; Stuiver et al., 1998] over a 2000 year overlap from 10.5 to 12.5 ka  $^{14}\text{C}$  BP (thousand radiocarbon years Before Present).

[17] Methane fluxes could have had a greater influence in the Pleistocene, when lowered sea levels sharply reduced water exchange between the basin and the Caribbean. The  $\delta^{13}\text{C}$  of the  $\text{CH}_4$  DIC source is about 60‰ lighter than atmospheric  $\text{CO}_2$ , so these episodes would leave a distinctive stable isotope signature. The planktonic foraminifera species *G. bulloides* and *G. ruber* have  $\delta^{13}\text{C}$  values that are 0.2–0.5‰ lighter than mean Holocene values for the period 14–20 ka  $^{14}\text{C}$  BP [Lin, 1992]. Ignoring any contribution to these offsets from whole-ocean glacial-interglacial  $\delta^{13}\text{C}$  shifts, this corresponds to a fossil  $\text{CH}_4$  contribution to the surface Cariaco Basin DIC pool of just 0.3–0.8%, and would bias  $^{14}\text{C}$  ages by less than 65 years.

[18] In conclusion, our radiocarbon results indicate that a large (0.024–0.028 Tg  $\text{CH}_4 \text{ y}^{-1}$ ) and previously unknown seep source of  $\text{CH}_4$  dominates Cariaco Basin water column  $\text{CH}_4$  distributions. If this seep source is localized, the possibility exists that carbonate structures, similar to those found in the Black Sea [Michaelis et al., 2002], could also occur in the Cariaco Basin.

[19] **Acknowledgments.** We acknowledge the crew of the B/O *Hermano Gines* for their enthusiasm and support at sea, Yrene Astor for help with cruise and equipment coordination, and Guaciara dos Santos for laboratory support. EDIMAR - Fundacion La Salle receives funding for the maintenance of the hydrographic time series CARIACO from the FONACIT (Fondo Nacional de Ciencia, Tecnologia e Innovacion, Venezuela), Project 2000001702. This work was supported by the National Science Foundation (NSF Grant OCE-0326928).

## References

- Astor, Y., F. Muller-Karger, and M. I. Scranton (2003), Seasonal and interannual variation in the hydrography of the Cariaco Basin: Implications for basin ventilation, *Cont. Shelf Res.*, **23**, 125–144.
- Audemard, F. A., G. Romero, H. Rendon, and V. Cano (2005), Quaternary fault kinematics and stress tensors along the southern Caribbean from fault-slip data and focal mechanism solutions, *Earth Sci. Rev.*, **69**, 181–233.
- Currie, L. A., J. D. Kessler, J. V. Marolf, A. P. McNichol, D. R. Stuart, J. C. Donoghue, D. J. Donahue, G. S. Burr, and D. Biddulph (2000), Low-level (submicromole) environmental  $^{14}\text{C}$  metrology, *Nucl. Instrum. Methods Phys. Res. B*, **172**, 440–448.
- Deuser, W. G. (1973), Cariaco Trench: Oxidation of organic matter and residence time of anoxic water, *Nature*, **242**, 601–603.
- Druffel, E. R. M., P. M. Williams, K. Robertson, S. Griffin, A. J. T. Jull, D. Donahue, L. Toolin, and T. W. Linick (1989), Radiocarbon in dissolved organic and inorganic carbon from the central North Pacific, *Radiocarbon*, **31**, 523–532.
- Fanning, K. A., and M. E. Q. Pilson (1972), A model for the anoxic zone of the Cariaco Trench, *Deep Sea Res.*, **19**, 847–863.
- Guilderson, T. P., J. E. Cole, and J. R. Southon (2005), Pre-bomb  $\Delta^{14}\text{C}$  variability and the Suess effect in Cariaco Basin surface waters as recorded in hermatypic corals, *Radiocarbon*, **47**, 57–65.
- Haug, G. H., K. A. Hughen, D. M. Sigman, L. C. Peterson, and U. Röhl (2001), Southward migration of the Intertropical Convergence Zone through the Holocene, *Science*, **293**, 1304–1308.

- Holmén, K. J., and C. G. H. Rooth (1990), Ventilation of the Cariaco Trench, a case of multiple source competition?, *Deep Sea Res., Part A*, **37**, 203–225.
- Hughen, K. A., J. T. Overpeck, L. C. Peterson, and R. F. Anderson (1996a), The nature of varved sedimentation in the Cariaco Basin, Venezuela, and its palaeoclimatic significance, in *Palaeoclimatology and Palaeoceanography From Laminated Sediments*, edited by A. E. S. Kemp, *Geol. Soc. Spec. Publ.*, **116**, 171–183.
- Hughen, K. A., J. T. Overpeck, L. C. Peterson, and S. Trumbore (1996b), Rapid climate changes in the tropical Atlantic region during the last deglaciation, *Nature*, **380**, 51–54.
- Hughen, K. A., J. R. Southon, S. J. Lehman, and J. T. Overpeck (2000), Synchronous radiocarbon and climate shifts during the last deglaciation, *Science*, **290**, 1951–1954.
- Hughen, K., S. Lehman, J. Southon, J. Overpeck, O. Marchal, C. Herring, and J. Turnbull (2004),  $^{14}\text{C}$  activity and global carbon cycle changes over the past 50,000 years, *Science*, **303**, 202–207.
- Lin, H.-L. (1992), Late Quaternary faunal and isotopic records from the anoxic Cariaco Basin, Venezuela, Ph.D. thesis, Univ. of Miami, Miami, Fla.
- Lin, H.-L., L. C. Peterson, J. T. Overpeck, S. E. Trumbore, and D. W. Murray (1997), Late Quaternary climate change from  $\delta^{18}\text{O}$  records of multiple species of planktonic foraminifera: High-resolution records from the anoxic Cariaco Basin, Venezuela, *Paleoceanography*, **12**, 415–427.
- Mathieu, G. G., P. E. Biscaye, R. A. Lupton, and D. E. Hammond (1988), System for measurement of  $^{222}\text{Rn}$  at low levels in natural waters, *Health Phys.*, **55**, 989–992.
- Michaelis, W., et al. (2002), Microbial reefs in the Black Sea fueled by anaerobic oxidation of methane, *Science*, **297**, 1013–1015.
- Peterson, L. C., G. H. Haug, K. A. Hughen, and U. Röhl (2000), Rapid changes in the hydrologic cycle of the tropical Atlantic during the last glacial, *Science*, **290**, 1947–1951.
- Reeburgh, W. S. (1976), Methane consumption in Cariaco Trench waters and sediments, *Earth Planet. Sci. Lett.*, **28**, 337–344.
- Richards, F. A., and R. F. Vaccaro (1956), The Cariaco Trench, an anaerobic basin in the Caribbean Sea, *Deep Sea Res.*, **3**, 214–228.
- Scranton, M. I. (1988), Temporal variations in the methane content of the Cariaco Trench, *Deep Sea Res., Part A*, **35**, 1511–1523.
- Scranton, M. I., F. L. Sayles, M. P. Bacon, and P. G. Brewer (1987), Temporal changes in the hydrography and chemistry of the Cariaco Trench, *Deep Sea Res., Part A*, **34**, 945–963.
- Scranton, M. I., Y. Astor, R. Bohrer, T.-Y. Ho, and F. Muller-Karger (2001), Controls on temporal variability of the geochemistry of the deep Cariaco Basin, *Deep Sea Res., Part I*, **48**, 1605–1625.
- Stuiver, M., and H. A. Polach (1977), Discussion: Reporting  $^{14}\text{C}$  data, *Radiocarbon*, **19**, 355–363.
- Stuiver, M., P. J. Reimer, E. Bard, J. W. Beck, G. S. Burr, K. A. Hughen, B. Kromer, G. McCormac, J. van der Plicht, and M. Spurk (1998), INTCAL98 radiocarbon age calibration, 24,000–0 cal BP, *Radiocarbon*, **40**, 1041–1083.
- Suárez, G., and J. Nábělek (1990), The 1967 Caracas earthquake: Fault geometry, direction of rupture propagation and seismotectonic implications, *J. Geophys. Res.*, **95**, 17,459–17,474.
- Vogel, J. S., J. R. Southon, D. E. Nelson, and T. A. Brown (1984), Performance of catalytically condensed carbon for use in accelerator mass-spectrometry, *Nucl. Instrum. Methods Phys. Res. B*, **233**, 289–293.
- von Humboldt, A. (1900), *Personal Narrative of Travels to the Equinoctial Regions of America During the Years 1799–1804*, George Bell, London.
- Ward, B. B., K. A. Kilpatrick, P. C. Novelli, and M. I. Scranton (1987), Methane oxidation and methane fluxes in the ocean surface layer and deep anoxic waters, *Nature*, **327**, 226–229.
- Wiesenburg, D. A. (1975), Processes controlling the distribution of methane in the Cariaco Trench, Venezuela, M.S. thesis, Old Dominion Univ., Norfolk, Va.
- Yamamoto, S., J. B. Alcauskas, and T. E. Crozier (1976), Solubility of methane in distilled water and seawater, *J. Chem. Eng. Data*, **21**, 78–80.

J. D. Kessler, W. S. Reeburgh, and J. Southon, Department of Earth System Science, University of California, Irvine, Irvine, CA 92697–3100, USA. (jkessler@uci.edu)

R. Varela, Fundación La Salle de Ciencias Naturales, Estación de Investigaciones Marinas de Margarita, Apartado 144 Porlamar, Isla de Margarita, Venezuela.