

Lawrence Berkeley National Laboratory

Recent Work

Title

CAPILLARY STABILITY IN AN INVERTED TECTANGULAR CHANNEL FOR FREE SURFACES WITH CURVATURE OF CHANGING SIGN

Permalink

<https://escholarship.org/uc/item/7d11d39k>

Author

Concus, Paul.

Publication Date

1964-06-23

University of California
Ernest O. Lawrence
Radiation Laboratory

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

**CAPILLARY STABILITY IN AN INVERTED RECTANGULAR CHANNEL
FOR FREE SURFACES WITH CURVATURE OF CHANGING SIGN**

Berkeley, California

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

For Am. Inst. of Aeronautic and Astronautics

UCRL-11515

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory
Berkeley, California

AEC Contract No. W-7405-eng-48

CAPILLARY STABILITY IN AN INVERTED RECTANGULAR
CHANNEL FOR FREE SURFACES WITH CURVATURE
OF CHANGING SIGN

Paul Concus

June 23, 1964

CAPILLARY STABILITY IN AN INVERTED RECTANGULAR
CHANNEL FOR FREE SURFACES WITH CURVATURE
OF CHANGING SIGN

Paul Concus

Lawrence Radiation Laboratory
University of California
Berkeley, California

June 23, 1964

SUMMARY

The stability of an incompressible inviscid fluid contained in an inverted rectangular channel is investigated, with the effects of surface tension included. The investigation covers the stability of those equilibrium free surfaces for which the curvature changes sign; the conditions for stability of free surfaces with curvature of constant sign were given previously by the author. It is found that all free surfaces with curvatures that change sign are unstable; thus surfaces with curvature of constant sign are the only possible stable ones. A graph giving the critical condition for stability as a function of contact angle is reprinted, along with the details of its derivation that were not given in the preceding report.

SYMBOLS

B	Bond number
F	Equilibrium interface
x, y	Nondimensional Cartesian coordinates
Δ	\pm Curvature
ϵ	Variational parameter
η	Perturbation to F
θ	Contact angle
λ	Undetermined parameter
ψ	Angle between tangent to F and horizontal
'	(Superscript) Denotes differentiation with respect to x

INTRODUCTION

In a previous paper,¹ an investigation of the stability of an incompressible inviscid fluid contained in an inverted rectangular channel was performed mathematically, with the effects of surface tension taken into account. The purpose was to determine the effect of contact angle and Bond number on the stability of such a fluid configuration in a low-gravity environment. The investigation was restricted to those fluid-gas equilibrium interfaces for which the curvature does not change sign. The analysis is extended in this report to include equilibrium interfaces for which the curvature does change sign. The omission of such interfaces in the original analysis is shown to be justified because they are all dynamically unstable. In addition, the details omitted in Ref. 1 of the calculation of the critical Bond number for equilibrium interfaces for which the curvature does not change sign, are also presented.

FORMULATION

The differential equation for an equilibrium free surface $y = F(x)$ (see Fig. 1) is given by Eq. (3) of Ref. 1 as

$$\left\{ F' / \left[(1 + F'^2)^{1/2} \right] \right\}' + BF - \lambda = 0 \quad , \quad -1 < x < 1 \quad (1)$$

where B is the Bond number, λ is a parameter to be determined, and the prime denotes differentiation with respect to x . The boundary conditions are given by Eq. (4) of Ref. 1, and are

$$F'(1) = -F'(-1) = \cot \theta \quad , \quad (2)$$

where θ is the contact angle between the fluid and the wall. Only values of θ for which $0 < \theta < \pi/2$ (wetting fluid) will be considered, since, as was shown in Ref. 1, the problem for $\pi/2 < \theta < \pi$ (nonwetting fluid) can be formulated in terms of an equivalent problem for $0 < \theta < \pi/2$.

Equation (1) can be solved in terms of the parameter

$$\psi = \tan^{-1} F' \quad , \quad (2a)$$

the slope of the surface, to yield,

$$B \frac{F^2}{2} - \lambda F = \cos \psi - C \quad ,$$

where F is now a function of ψ . The boundary conditions, Eq. (2), become $\psi = \pm (\pi/2 - \theta)$ at $x = \pm 1$, respectively. The constant of integration C may be conveniently evaluated by choosing the origin of coordinates properly. The coordinates shown in Fig. 1 of Ref. 1 were chosen so that the origin would lie on the fluid-vapor interface, halfway between the walls. It is more convenient here to choose the x axis to intersect the interface when $\psi = 0$ rather than when $x = 0$. That is, $F = 0$ when $\psi = 0$ and $x = x_0$ (x_0 is not necessarily 0,

see Fig. 1). The boundary conditions require that $\psi = 0$ at least once by continuity, but ψ may equal zero more than once, in which case any particular point where $\psi = 0$ and the curvature is positive (concave downward) may be singled out to define where $F = 0$. With this choice of coordinates, the constant of integration C is one, so that

$$\frac{BF^2}{2} - \lambda F = \cos \psi - 1,$$

or

$$F = \frac{\lambda \pm \Delta}{B}, \quad (3)$$

where

$$\Delta = \left[\lambda^2 + 2B(\cos \psi - 1) \right]^{1/2}. \quad (4)$$

Substitution into Eq. (1) shows that $\mp \Delta$ is the curvature. Notice that the value of $C = 1$ is not only compatible with $F = 0$ at $\psi = 0$, but also with $F = 2\lambda/B$ at $\psi = 0$, depending on whether one takes the lower or upper sign for the square root.

The corresponding solution for x passing through $x = x_0$ when $\psi = 0$ is obtained from Eqs. (2a) and (3). It is

$$x = x_0 \mp \int_0^\psi \frac{\cos \psi \, d\psi}{\Delta}. \quad (5)$$

The value of x_0 can be determined from the boundary conditions, once a choice is made concerning the sign.

In Ref. 1 (as in the first reference of Ref. 1), only the solution taking the lower sign is considered, since it is the one passing through the required value $F = \psi = 0$. The solution taking the upper sign is not considered, on the grounds that it does not pass through $F = \psi = 0$. The ensuing analysis in Ref. 1 then determines that $x_0 = 0$, and, on this basis, the critical Bond number is found.

Notice, however, that for values of ψ equal to $\pm\psi_0$ where the curvature vanishes,

$$\psi_0 = \cos^{-1} \left(1 - \frac{\lambda^2}{2E} \right), \quad 0 \leq \psi_0 < \pi, \quad (6)$$

it is possible to continuously join the two branches of the solution, one having the upper sign and the other the lower sign in Eq. (3). Since the curvature vanishes at these points, the derivatives up to the second would also join continuously. Equation (6) shows that joining can take place (for real values of ψ) only if $B \geq \lambda^2/4$. In this paper, values of $B \geq \lambda^2/4$ are considered, and equilibrium solutions made up of joined branches are allowed. The purpose is to examine the stability of such joined solutions.

STABILITY ANALYSIS FOR JOINED INTERFACES

The stability of the equilibrium interfaces consisting of joined solutions can be investigated by considering the second variation in the same manner described in Ref. 1 for unjoined solutions. The second variation for trial functions of the form

$$y = F(x) + \epsilon \eta(x),$$

where $F(x)$ is the equilibrium solution, ϵ is a parameter, and $\eta(x)$ is the perturbing function, is given by Eqs. (5) and (6) of Ref. 1. Equation (7) of Ref. 1 is the Jacobi equation, which must be satisfied by $\eta(x)$ to make the second variation vanish. It can be written in the form

$$\left[\eta' / (1 + F'^2)^{3/2} \right]' + B\eta = 0. \quad (7)$$

In the present case, care must be taken to use the appropriate signs in each of the branches of F .

No matter which sign is chosen for F and x , however, the relationship

$$\frac{dF}{dx} = \tan \psi \quad (8)$$

still holds, and Eq. (7) becomes

$$\frac{d}{dx} \left(\cos^3 \psi \frac{d\eta}{dx} \right) + B\eta = 0,$$

upon substitution of Eq. (8).

This equation, when rewritten in terms of derivatives with respect to ψ , yields

$$\frac{\frac{d}{d\psi} \left(\cos^3 \psi \frac{d\eta}{d\psi} \right)}{\left(\frac{dx}{d\psi} \right)^2} - \frac{\frac{d^2 x}{d\psi^2} \cdot \left(\cos^3 \psi \frac{d\eta}{d\psi} \right)}{\left(\frac{dx}{d\psi} \right)^3} + B\eta = 0. \quad (9)$$

Substitution of Eq. (5) into Eq. (9) shows that Eq. (9) is independent of whether the upper or lower sign is used in the solution for x . Thus, Jacobi's equation is unchanged when joined solutions, as well as nonjoined solutions, are considered. Hence, the least eigenvalue, which is the critical Bond number, is also unchanged. Therefore, the stability criterion for the critical Bond number is unaltered from that given in Ref. 1.

It is now shown that for Bond numbers less than the critical one, only the simple unjoined interfaces discussed in Ref. 1 can be stable; the joined interfaces are always unstable. To do this, it is sufficient to show that for joined interfaces a conjugate point to one of the end points $x = \pm 1$ lies within the interval $-1 < x < 1$; or, in other words, if the general solution of Eq. (7) is made to satisfy nontrivially one of the boundary conditions, Eq. (2) in Ref. 1-- that $d\eta/dx = 0$ at $x = -1$ or $x = +1$ --then $d\eta/dx$ will also vanish for some

x in this interval. Such a property makes it possible to choose a function that makes the second variation negative.

For Δ to be real, ψ must lie within the limits $-\psi_0 \leq \psi \leq \psi_0$. Thus, if F is to be an equilibrium interface satisfying the boundary conditions, Eq. (2), then $\pi/2 - \theta \leq \psi_0$. If, in addition, there is an interior joining point, then $\psi = \pm \psi_0$ for some point in $-1 < x < 1$. Then continuity in ψ requires that ψ must take on at least one of the values $-(\pi/2 - \theta)$ or $(\pi/2 - \theta)$ in the interior. That is, in order for there to be a joining point, ψ must assume one of its boundary values at least once in the interior. However, this means that a solution of Eq. (7), for which $d\eta/dx$ vanishes at a boundary point, also has $d\eta/dx$ vanish at an interior point, because, by use of $d\psi/dx$ as calculated from Eq. (5), it follows that

$$\frac{d\eta}{dx} = \mp \frac{\Delta}{\cos\psi} \frac{d\psi}{dx}.$$

a function of ψ alone. Thus, joined interfaces cannot provide an extremal solution, and hence they cannot be stable.

CONCLUSION

The above analysis shows that equilibrium interfaces possessing a curvature that changes sign are always unstable, and that the only possible stable interfaces are those with a curvature that does not change sign. Thus, the stability criteria for the critical Bond number derived in Ref. 1 by consideration of only those interfaces possessing curvature of constant sign are valid in general.

APPENDIX

Calculation of the Critical Bond Number
for the Stability of Equilibrium Interfaces

The critical Bond number can be calculated by first finding the general solution of Eq. (9) for η as a function of ψ . Substitution of $d\psi/dx$, as calculated from Eq. (5), into Eq. (9) yields

$$\frac{d}{d\psi} \left(\cos^3 \psi \frac{d\eta}{d\psi} \right) + \sin \psi \cos^2 \psi \left(1 - B \frac{\cos \psi}{\Delta^2} \right) \frac{d\eta}{d\psi} + B \eta \frac{\cos^2 \psi}{\Delta^2} = 0.$$

The general solution is

$$\eta = C \tan \psi + D \left[\frac{\lambda}{B\Delta} + \tan \psi \cdot I(\psi) \right],$$

where

$$I(\psi) = - \int_0^\psi \frac{\lambda^2 \cos \psi \, d\psi}{\Delta^3}.$$

It is always required that $-\psi_0 \leq \psi \leq \psi_0$ in order that Δ be real. The derivative of η with respect to ψ is given by $d\eta/d\psi = \sec^2 \psi [C + DI(\psi)]$, so that the derivative with respect to x is

$$\frac{d\eta}{dx} = \frac{d\eta}{d\psi} \frac{d\psi}{dx} = \frac{1}{\Delta} \sec^3 \psi \cdot [C + DI(\psi)].$$

Application of the boundary conditions at $x = \pm 1$ yields

$$\Delta_1 \sec^3 \psi_1 [C - DI(\psi_1)] = 0,$$

and

$$\Delta_1 \sec^3 \psi_1 [C + DI(\psi_1)] = 0,$$

where

where

$$\psi_1 = \pi/2 - \theta,$$

$$\psi_2 = \pi/2 - \theta.$$

$$\Delta_1 \sec^3 \psi_1 [C + DI(\psi_1)] = 0,$$

and Δ_1 is the value of Δ when ψ equals ψ_1 . Since for $0 < \psi_1 < \pi/2$ neither $\sec^3 \psi_1$ nor $I(\psi_1)$ can vanish, the equations can be satisfied only if $\Delta_1 = 0$, and the product $\Delta_1 I(\psi_1)$ may, in general, be nonzero, so that D must also be chosen zero. The requirement that $\Delta_1 = 0$ implies that

$$\lambda^2 = 2B (1 - \cos \psi_1) = 2B (1 - \sin \theta).$$

When this relationship is used in Eq. (5) and the boundary conditions that $\psi = \pm (\pi/2 - \theta)$ at $x = \pm 1$, respectively, are applied, it follows that $x_0 = 0$ and the critical Bond number relationship is

$$\frac{1}{2} \left\{ \int_0^{\pi/2-\theta} \frac{\cos\psi \, d\psi}{(\cos\psi - \sin\theta)^{1/2}} \right\}^2 = B,$$

which is Eq. (19) of Ref. 1. Figure 2 of Ref. 1 is reproduced here, which is a graph of the above relationship giving the critical Bond number as a function of contact angle.

FOOTNOTE AND REFERENCE


* This work was carried out mainly under the Independent Research Program of the Lockheed Missiles and Space Company and secondarily under the auspices of the U. S. Atomic Energy Commission.

1. Concus, P., Capillary Stability in an Inverted Rectangular Tank, in Advances in the Astronautical Sciences, (Western Periodicals Company, North Hollywood, Calif.) Vol. 14, pp. 21-37, 1963.

FIGURE LEGENDS

Fig. 1. Geometric configuration.

Fig. 2. Critical Bond number as a function of contact angle.



MUB - 3342

Fig. 1




Fig. 2.

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

