

# UC Berkeley

## UC Berkeley Previously Published Works

### Title

ClusterCAD: a computational platform for type I modular polyketide synthase design

### Permalink

<https://escholarship.org/uc/item/7h84g10b>

### Journal

Nucleic Acids Research, 46(Database issue)

### ISSN

0305-1048

### Authors

Eng, Clara H  
Backman, Tyler WH  
Bailey, Constance B  
et al.

### Publication Date

2018-01-04

### DOI

10.1093/nar/gkx893

Peer reviewed

# ClusterCAD: a computational platform for type I modular polyketide synthase design

Clara H. Eng<sup>1,†</sup>, Tyler W.H. Backman<sup>2,3,4,†</sup>, Constance B. Bailey<sup>2,3</sup>, Christophe Magnan<sup>5,6</sup>, Héctor García Martín<sup>2,3,4</sup>, Leonard Katz<sup>7</sup>, Pierre Baldi<sup>5,6</sup> and Jay D. Keasling<sup>1,2,3,4,7,8,9,\*</sup>

<sup>1</sup>Department of Chemical and Biomolecular Engineering, University of California, Berkeley, CA 94720, USA, <sup>2</sup>Joint BioEnergy Institute, 5885 Hollis Street, Emeryville, CA 94608, USA, <sup>3</sup>Biological Systems and Engineering Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA, <sup>4</sup>Department of Energy Agile BioFoundry, Emeryville, CA 94608, USA, <sup>5</sup>Department of Computer Science, University of California, Irvine, CA 92697, USA, <sup>6</sup>Institute for Genomics and Bioinformatics, University of California, Irvine, CA 92697, USA, <sup>7</sup>QB3 Institute, University of California, Berkeley, CA 94720, USA, <sup>8</sup>Department of Bioengineering, University of California, Berkeley, CA 94720, USA and <sup>9</sup>Novo Nordisk Foundation Center for Biosustainability, Technical University of Denmark, DK2970 Horsholm, Denmark

Received August 15, 2017; Revised September 17, 2017; Editorial Decision September 20, 2017; Accepted September 24, 2017

## ABSTRACT

ClusterCAD is a web-based toolkit designed to leverage the collinear structure and deterministic logic of type I modular polyketide synthases (PKSs) for synthetic biology applications. The unique organization of these megasynthases, combined with the diversity of their catalytic domain building blocks, has fueled an interest in harnessing the biosynthetic potential of PKSs for the microbial production of both novel natural product analogs and industrially relevant small molecules. However, a limited theoretical understanding of the determinants of PKS fold and function poses a substantial barrier to the design of active variants, and identifying strategies to reliably construct functional PKS chimeras remains an active area of research. In this work, we formalize a paradigm for the design of PKS chimeras and introduce ClusterCAD as a computational platform to streamline and simplify the process of designing experiments to test strategies for engineering PKS variants. ClusterCAD provides chemical structures with stereochemistry for the intermediates generated by each PKS module, as well as sequence- and structure-based search tools that allow users to identify modules based either on amino acid sequence or on the chemical structure of the cognate polyketide intermediate. ClusterCAD can be accessed at <https://clustercad.jbei.org> and at <http://clustercad.igb.uci.edu>.

## INTRODUCTION

Polyketides and their derivatives are used broadly in medicine as antibiotics, antifungals and immunosuppressants, among other applications (1), motivating research to engineer polyketide synthases (PKSs) capable of producing novel drug analogs. More recently, researchers have also harnessed the biosynthetic potential of PKSs to produce commodity chemicals in an environmentally friendly manner (2–4).

Type I modular PKSs are composed of a series of catalytic domains whose identity and order determine the structure of the final polyketide product in a predictable manner, as shown in Figure 1. A single PKS can include multiple polypeptide subunits, each comprising one or more modules. Each module, which is minimally made up of a set of ketosynthase (KS), acyltransferase (AT) and acyl carrier protein (ACP) domains, catalyzes the extension of the nascent acyl chain by two carbon units. This is accomplished via a KS-catalyzed decarboxylative condensation between an acyl starter or intermediate, and a dicarboxylic acid extender unit, which is selected by the AT (5,6). The resulting  $\beta$ -ketone may be reduced by a ketoreductase (KR) domain, which determines the stereochemistry of the resulting  $\beta$ -hydroxyl group, and that of the  $\alpha$ -substituent, if present (7–11). Additional dehydratase (DH) and enoyl reductase (ER) domains may further reduce the hydroxyl group to yield a methenyl or methylene group, respectively. The KR, DH and ER make up a set of optional reducing domains that together, are termed the reductive cassette. In addition to these reductive domains, C- or O- methyltransferases may also be present. Release of a polyketide from a PKS is carried out by a thioesterase (TE) domain that cat-

\*To whom correspondence should be addressed. Tel: +1 510 495 2620; Fax: +1 510 495 2630; Email: keasling@berkeley.edu

†These authors contributed equally to the paper as first authors.



patibility between PKS polypeptide sequences. It is thus also important to consider additional factors when designing a chimeric PKS. For example, modules that originate from well-characterized clusters, or that have been previously determined to be well-expressed in the host organism of choice, are particularly attractive choices for engineering.

## MATERIALS AND METHODS

### Data curation

The gene clusters in ClusterCAD are sourced from the Minimum Information about a Biosynthetic Gene cluster (MIBiG) database (29). To construct each ClusterCAD entry, the gene clusters in version 1.3 (3 September 2016) of the MIBiG database labeled as type I modular PKSs were first identified. These gene clusters were annotated using the antibiotics and Secondary Metabolite Analysis SHell (antiSMASH) (30), and the resulting output parsed using a list of explicitly recognized catalytic domains. Cluster analysis was truncated if a subunit containing an unrecognized catalytic domain was encountered. Domain annotations, which include predictions for AT substrate specificity and KR stereochemical outcome, informed *in silico* simulation of biocatalysis using reaction SMARTS operators (<http://www.daylight.com/dayhtml/doc/theory/>), which were implemented using the RDKit cheminformatics software library (<http://www.rdkit.org>), yielding a predicted intermediate chemical structure for each PKS module. Known final chemical structures were taken from the MIBiG database if available, or were otherwise generated from the text description of the final structure in the MIBiG database using the ChemAxon JChem Base naming tool (version 17.2.27.0, 2017) (31). Chemical structures for well-characterized polyketide products that could not be obtained by either of these methods were manually incorporated. Comparisons of the predicted and known final product chemical structures, close inspection of established amino acid sequence motifs (20,32) and an extensive literature search were used to manually curate each ClusterCAD entry by correcting subunit order, removing invalid subunits, accounting for iterating modules and correcting misannotated catalytic domain activity. The SCRATCH software suite (33,34) was used to generate secondary structure and relative solvent accessibility predictions for the PKS subunits in ClusterCAD, which were also incorporated into the database.

### Organization

*User interface.* The primary objective of ClusterCAD is to provide computational assistance to the synthetic biologist seeking to engineer PKSs. ClusterCAD provides an intuitive interface for users to browse through the clusters in the database, as well as search tools developed to assist in the design of chimeric PKSs. An example entry of a PKS cluster as displayed in the ClusterCAD web interface, is shown in Figure 2.

i) *Structure search tool:* the structure search tool was designed to enable the identification of a truncated PKS

parent to use as a starting point for PKS engineering, and takes as input either a small molecule chemical structure represented as a SMILES string, or a structure that is drawn in an interactive graphical user interface. This tool searches a database of the predicted polyketide intermediates produced by each of the PKS modules in ClusterCAD. Matches to the query structure are ranked using atom pair (AP) Tanimoto similarity scores (35), and returned matches are displayed with the maximum common substructure (36) between the query and target structures highlighted.

ii) *Sequence search tool:* the sequence search tool was designed to enable researchers to select candidate catalytic domains for domain exchange experiments, and supports a variety of queries that allow researchers to test hypotheses regarding which domain–domain interactions may be important in facilitating successful domain exchanges. The sequence search tool takes as input an amino acid sequence, and performs a protein–protein BLAST+ search against a database containing all of the subunits in ClusterCAD (37). The results page enables deeper interrogation of the returned results by highlighting catalytic domains on the basis of annotations such as AT substrate specificity, KR stereochemical outcome, and reductive domain activity.

## DISCUSSION

### Motivation behind ClusterCAD

ClusterCAD provides the first PKS database that specifically targets engineering applications. Existing PKS databases focus on identifying new metabolites through genome mining or on predicting metabolite structure from nucleotide or amino acid sequence (38). However, many of the most promising applications of PKSs are premised on harnessing their ability to generate customized molecular architectures.

From an engineering perspective, ClusterCAD offers substantial improvements over existing databases. By taking advantage of resources such as the MIBiG database and the antiSMASH command-line tool for cluster identification and annotation respectively, we were able to focus on designing tools tailored to the needs of the synthetic biologist without attempting to optimize our software for genome mining or metabolite structure prediction. Furthermore, given the strong commitment from the synthetic biology and natural products communities to update MIBiG and antiSMASH with continuous improvements, populating ClusterCAD using an end-to-end pipeline that utilizes these tools means that ClusterCAD can easily be kept up-to-date as novel biosynthetic clusters are discovered and as our understanding of PKSs develops.

While building on MIBiG and antiSMASH aids in automating the maintenance of ClusterCAD, it also means that inaccuracies in MIBiG or in the antiSMASH software may be propagated into ClusterCAD. In order to address this issue, we also employ extensive manual curation of each ClusterCAD entry in order to provide predicted polyketide intermediates that are as accurate as possible. ClusterCAD has a persistent corrections record that overwrites the output generated by antiSMASH. This enables ClusterCAD


ClusterCAD [Browse clusters](#) [Structure search](#) [Sequence search](#) [About](#)

## Borreledin

MiBiG accession  
BGC000031.1  
GenBank accession  
AJ580915.1


Show AT annotations  
 Show KR annotations

Known final product



MCS: 97.1% of atoms

Predicted pks subset




MCS: 100% of atoms

borA1

module 0

AT  ACP



**Figure 2.** ClusterCAD entry for the borreledin PKS. The ClusterCAD page for each cluster contains links to the corresponding entries in the MiBiG and NCBI Nucleotide databases. Catalytic domain annotations can be optionally displayed based on domain type. Interactive web page elements provide access to more detailed information, such as the nucleotide and amino acid sequences of each catalytic domain and subunit.

to incorporate new MiBiG entries and systematic improvements to antiSMASH, while also guaranteeing that clusters that have previously been manually corrected will remain correct. Because these records are persistent and override the automatically generated output, the manually curated clusters will only need to be further corrected if new empirical findings are generated. ClusterCAD thus combines computational annotations of AT substrate specificity and KR stereospecificity with domain expert application of heuristics obtained from the literature (20,32,39,40) to provide well-curated predictions of the polyketide intermediates naturally provided by each of the PKS modules in ClusterCAD. However, the reliance on manual curation means that some errors may be missed. We include links on each cluster page that can be used to submit corrections to the ClusterCAD maintainers for review.

ClusterCAD is further limited in that it does not support inclusion of chemistry effected by ‘on-line’ tailoring enzymes that act on ACP-tethered polyketide intermediates, but that are independent of the PKS (41). However, we believe that showing the direct product of the PKS, without taking into account ‘on-line’ tailoring enzymes provides the information that is most relevant for engineering, since ‘on-line’ tailoring enzymes are not expected to be naturally present in a heterologous host.

A substantial benefit of ClusterCAD is that it provides engineering tools that allow researchers to compare PKS modules based on amino acid sequence similarity and on cognate polyketide intermediate similarity. The substructure search in ClusterCAD is of particular utility when se-

lecting candidate modules to serve as a starting point in a PKS engineering project. Previous experimental evidence supports the assertion that the structure of the intermediate can have a significant impact on the function of individual catalytic domains. KS domains, for example, have been reported to have ‘gatekeeping’ functionality, in which they are unable to condense intermediates that are chemically dissimilar for reasons ranging from overall steric bulkiness (42) to having an opposite stereochemical configuration (43,44). Other domains, such as the KR domain (25) and the DH domain (45) have also been observed to have poor catalytic activity on substrates that are much smaller or bulkier than ones they encounter in nature.

The substructure search in ClusterCAD is one of the first that matches against polyketide intermediates associated with non-terminal PKS modules, making it more robust to false positives than similar search tools which only support substructure queries in the context of PKS final products (46). Matching directly against the polyketide intermediates produced by each module also avoids undesirable matches to structural motifs produced by tailoring enzymes, which are of limited utility for PKS engineering. To the best of our knowledge, SBSPKSv2 (47) is the only other PKS database that supports structure-based queries that match against polyketide intermediates. However, ClusterCAD boasts a larger number of type I modular PKS intermediates, and utilizes AP descriptors, which exhibit greater sensitivity to uncommon chemical motifs than the substructure fingerprint approach used by SBSPKSv2.

Further, anecdotal evidence supports the assertion that seeking catalytic domains with greater sequence similarity improves domain exchange success rates (25). While this heuristic is challenging to verify experimentally given the low throughput of domain exchange experiments, it is our hope that ClusterCAD will assist PKS researchers in elucidating the importance of sequence similarity in domain exchange experiments by streamlining the design of experiments to test their hypotheses. For instance, by facilitating chimeric PKS design, ClusterCAD will enable researchers to probe potential domain–domain interactions which may be important for engineering.

The sequence search tool in ClusterCAD highlights the domains in each returned module that overlap with the query, and also allows the user to highlight results based on catalytic domain activity. Identifying junctions in PKS chimeras that reliably leave functionally important structural elements intact (24,25) remains an open area of research. ClusterCAD provides secondary structure and relative solvent accessibility predictions that will aid in the identification and comparison of homologous structural elements across PKSs, assisting researchers in junction design. As far as we are aware, ClusterCAD is the only PKS database that provides precomputed values for the predicted secondary structure and relative solvent accessibility of each subunit. By providing support to compare PKS modules on the basis of metrics that may contribute to compatibility between PKS polypeptide sequences, ClusterCAD aims to facilitate experimental research to identify robust strategies for PKS engineering.

### Using ClusterCAD to design a chimeric PKS that produces adipic acid

While engineered PKSs have long been an attractive strategy to produce pharmaceutically relevant natural products and their analogs, more recent work has also sought to harness the chemical diversity accessible by PKS to provide an environmentally friendly alternative to chemical synthesis for commodity chemicals. Previous work by Hagen *et al.* has demonstrated that it is possible to engineer a novel PKS capable of producing adipic acid (2), a widely used commodity chemical. As a proof-of-concept, we describe how ClusterCAD could be applied to engineer this novel PKS.

- i) *Identify a truncated PKS starting point that naturally produces a polyketide intermediate that is structurally similar to adipic acid.* Hagen *et al.* selected the first module from the borrelidin PKS as a starting point based on previous biochemical studies establishing that BorMod1 tolerates succinyl-CoA as a starter, possibly because of structural similarity to the native priming unit, trans-1,2-cyclopentanedicarboxylic acid (48). Adipic acid was used to query the ClusterCAD chemical structure search tool, which returned BorMod1 as the highest-scoring result, with a naturally produced intermediate that had an AP Tanimoto similarity score of 0.32 when compared against adipic acid.
- ii) *Identify a donor reductive loop with active ER, DH and KR from a module which naturally incorporates a malonyl-CoA extender unit.* In order to produce adipic

acid using BorMod1, incorporation of an extender unit lacking a  $\alpha$ -substituent and reduction by a full complement of active reducing domains was required. Hagen *et al.* thus chose four reductive loop donors that naturally accepted malonyl-CoA, harbored a DH, ER and KR in the reductive cassette, and that came from a ‘standalone’ module, in which the corresponding open reading frame or subunit contained only a single module. Donor loops from ‘standalone’ modules were favored based on the intuition that domain exchanges using reductive cassettes derived from a ‘standalone’ module would retain more architectural similarity to BorMod1, itself a ‘standalone’ module, and thereby increase the likelihood of a functional chimera.

Using the ClusterCAD sequence search tool, we sought to identify donor modules that not only met these criteria, but also exhibited maximal sequence similarity to BorMod1. We queried the ClusterCAD sequence search tool with the amino acid sequence of BorA2, the subunit which contains BorMod1, ranking the potential donor modules by bit score and filtering for ‘standalone’ modules containing a malonyl-CoA specific AT and a complete reductive cassette. IdmO, from the indanomycin PKS, followed by SpnB from the spinosyn PKS were the only two returned results that matched all three criteria. Chimeras utilizing these two donor modules were experimentally tested by Hagen *et al.*, along with chimeras harboring reductive cassettes from the donor modules AurB, from the aureothin PKS and NanA2, from the nanchangmycin PKS. AurB, which is not contained in ClusterCAD, and SpnB were empirically determined to yield the most active chimeras. NanA2, which was not returned by ClusterCAD, but is contained in the database, yielded the least active chimera.

- iii) *Identify a DH domain that exhibits higher levels of activity on 3-hydroxy adipic acid.* The chimeras constructed by Hagen *et al.* yielded a relatively large proportion of the incompletely reduced product 3-hydroxyadipic acid, potentially due to the DH from the donor reductive cassettes having a poor substrate compatibility with the terminal carboxyl moiety, an unusual feature in PKS biosynthetic intermediates. To overcome this substrate incompatibility, Hagen *et al.* further substituted a heterologous DH into the engineered BorMod1 variant. The DH from BorMod2 was selected based on the observation that it is known to demonstrate catalytic activity on a substrate containing a terminal carboxy moiety. The intuition employed by Hagen *et al.* was consistent with the sequence-similarity based output of the ClusterCAD search tool, which returned BorMod2 as the module containing an active DH that had the highest bit score with BorMod1.

### Implications of ClusterCAD for PKS engineering

As the first PKS database that specifically targets synthetic biologists interested in PKS engineering, we also sought more generally to provide tools for computer-aided PKS design, and have released ClusterCAD under a BSD style license at <https://github.com/JBEI/clusterCAD>. This

```

{
  "description": "Borrelidin",
  "genbankAccession": "AJ580915.1",
  "mibigAccession": "BGC000031.1",
  "architecture": {
    "borA1": {
      "0": {
        "iterations": 1,
        "domains": {
          "AT": {
            "substrate": "trans-1,2-CPDA"
          }
        }
      }
    },
    .
    .
    .
    "borA6": {
      "0": {
        "iterations": 1,
        "domains": {
          "AT": {
            "substrate": "mal"
          },
          "KR": {
            "type": "B1",
            "active": true
          },
          "TE": {
            "ring": 1,
            "cyclic": true
          }
        }
      }
    }
  }
}

```

**Figure 3.** Partial JSON representation of the borrelidin PKS. This representation captures the hierarchical organization of PKSs, as well as the important catalytic properties of their components.

open source release provides Python object representations of catalytic domains, modules and clusters, that enable researchers to construct and simulate the activity of chimeric PKSs *in silico*. Furthermore, we sought to establish a method to describe PKSs in a human and computer readable manner using the JSON data-interchange format, as shown in Figure 3. The availability of such a PKS representation will aid in the development of additional software for computational PKS design by allowing researchers to easily describe novel PKSs in a computer-readable fashion.

## CONCLUSION

To the best of our knowledge, ClusterCAD is the first web-based toolkit developed specifically to aid in designing chimeric PKSs. By providing chemical structures with stereochemistry for the polyketide intermediate produced by each module, as well as relative solvent accessibility and

secondary structure predictions for each subunit, we provide rich information about each cluster to inform chimeric PKS design. Furthermore, our intuitive user interface, combined with the ability to query for modules based on either sequence- or structure-similarity provide powerful tools for PKS engineering that bring us closer to the promise of harnessing PKSs for combinatorial biosynthesis.

## ACKNOWLEDGEMENTS

We wish to acknowledge OpenEye Scientific Software and ChemAxon for academic software licenses, as well as Mark Kulawik and Yuzo Kanomata for computing support.

## FUNDING

U.S. Department of Energy Joint BioEnergy Institute and Agile BioFoundry [Contract DE-AC02-05CH11231]; DOE EERE Annual Operating Plan [BM0101020-05450-1004171, Agreement 28712, DE-AC02-05C11231]; Defense Advanced Research Projects Agency [HR0011-15-2-0045 to P.B.]; National Science Foundation [CBET-1437775 to J.D.K., EEC-0540879 to J.D.K., MCB-1341894 to J.D.K., IIS-1321053 to P.B.]; National Science Foundation Graduate Research Fellowship Program [DGE 1106400 to C.H.E.]. Funding for open access charge: Joint BioEnergy Institute.

*Conflict of interest statement.* J.D.K. has a financial interest in Amyris, Lygos, Constructive Biology and Demetrix. L.K. has a financial interest in Lygos. P.B. has a financial interest in Anivive.

## REFERENCES

- Weissman, K.J. and Leadlay, P. (2005) Combinatorial biosynthesis of reduced polyketides. *Nat. Rev. Microbiol.*, **3**, 925–936.
- Hagen, A., Poust, S., de Rond, T., Fortman, J.L., Katz, L., Petzold, C.J. and Keasling, J.D. (2015) Engineering a polyketide synthase for *in vitro* production of adipic acid. *ACS Synth. Biol.*, **5**, 21–27.
- Yuzawa, S., Deng, K., Wang, G., Baidoo, E.E., Northen, T.R., Adams, P.D., Katz, L. and Keasling, J.D. (2016) Comprehensive *in vitro* analysis of acyltransferase domain exchanges in modular polyketide synthases and its application for short-chain ketone production. *ACS Synth. Biol.*, **6**, 139–147.
- Yuzawa, S., Keasling, J.D. and Katz, L. (2017) Bio-based production of fuels and industrial chemicals by repurposing antibiotic-producing type I modular polyketide synthases: opportunities and challenges. *J. Antibiot.*, **70**, 378–385.
- Cane, D.E., Liang, T.C., Taylor, P.B., Chang, C. and Yang, C.C. (1986) Macrolide biosynthesis. 3. Stereochemistry of the chain-elongation steps of erythromycin biosynthesis. *J. Am. Chem. Soc.*, **108**, 4957–4964.
- Marsden, A.F., Caffrey, P., Aparicio, J.F., Loughran, M.S., Staunton, J. and Leadlay, P.F. (1994) Stereospecific acyl transfers on the erythromycin-producing polyketide synthase. *Science*, **263**, 378–380.
- Reid, R., Piagentini, M., Rodriguez, E., Ashley, G., Viswanathan, N., Carney, J., Santi, D.V., Hutchinson, C.R. and McDaniel, R. (2003) A model of structure and catalysis for ketoreductase domains in modular polyketide synthases. *Biochemistry*, **42**, 72–79.
- Caffrey, P. (2003) Conserved amino acid residues correlating with ketoreductase stereospecificity in modular polyketide synthases. *ChemBioChem*, **4**, 654–657.
- Garg, A., Khosla, C. and Cane, D.E. (2013) Coupled methyl group epimerization and reduction by polyketide synthase ketoreductase domains. Ketoreductase-catalyzed equilibrium isotope exchange. *J. Am. Chem. Soc.*, **135**, 16324–16327.

10. Garg, A., Xie, X., Keatinge-Clay, A., Khosla, C. and Cane, D.E. (2014) Elucidation of the cryptic epimerase activity of redox-inactive ketoreductase domains from modular polyketide synthases by tandem equilibrium isotope exchange. *J. Am. Chem. Soc.*, **136**, 10190–10193.
11. Annal, T., Paris, C., Leadlay, P.F., Jacob, C. and Weissman, K.J. (2015) Evaluating ketoreductase exchanges as a means of rationally altering polyketide stereochemistry. *ChemBioChem*, **16**, 1357–1364.
12. Khosla, C., Tang, Y., Chen, A.Y., Schnarr, N.A. and Cane, D.E. (2007) Structure and mechanism of the 6-deoxyerythronolide B synthase. *Annu. Rev. Biochem.*, **76**, 195–221.
13. Wilson, M.C. and Moore, B.S. (2012) Beyond ethylmalonyl-CoA: the functional role of crotonyl-CoA carboxylase/reductase homologs in expanding polyketide diversity. *Nat. Prod. Rep.*, **29**, 72–86.
14. Chang, C., Huang, R., Yan, Y., Ma, H., Dai, Z., Zhang, B., Deng, Z., Liu, W. and Qu, X. (2015) Uncovering the formation and selection of benzylmalonyl-CoA from the biosynthesis of splenocin and enterocin reveals a versatile way to introduce amino acids into polyketide carbon scaffolds. *J. Am. Chem. Soc.*, **137**, 4183–4190.
15. Keatinge-Clay, A.T. (2012) The structures of type I polyketide synthases. *Nat. Prod. Rep.*, **29**, 1050–1073.
16. Zheng, J., Gay, D.C., Demeler, B., White, M.A. and Keatinge-Clay, A.T. (2012) Divergence of multimodular polyketide synthases revealed by a didomain structure. *Nat. Chem. Biol.*, **8**, 615–621.
17. Weissman, K.J. (2014) The structural biology of biosynthetic megaenzymes. *Nat. Chem. Biol.*, **11**, 660–670.
18. Dutta, S., Whicher, J.R., Hansen, D.A., Hale, W.A., Chemler, J.A., Congdon, G.R., Narayan, A.R., Hakansson, K., Sherman, D.H., Smith, J.L. *et al.* (2014) Structure of a modular polyketide synthase. *Nature*, **510**, 512–517.
19. Whicher, J.R., Dutta, S., Hansen, D.A., Hale, W.A., Chemler, J.A., Dosey, A.M., Narayan, A.R., Hakansson, K., Sherman, D.H., Smith, J.L. *et al.* (2014) Structural rearrangements of a polyketide synthase module during its catalytic cycle. *Nature*, **510**, 560–564.
20. Dunn, B.J. and Khosla, C. (2013) Engineering the acyltransferase substrate specificity of assembly line polyketide synthases. *J. R. Soc. Interface*, **10**, 20130297.
21. McDaniel, R., Thamchaipenet, A., Gustafsson, C., Fu, H., Betlach, M. and Ashley, G. (1999) Multiple genetic modifications of the erythromycin polyketide synthase to produce a library of novel “unnatural” natural products. *Proc. Natl. Acad. Sci. U.S.A.*, **96**, 1846–1851.
22. Hans, M., Hornung, A., Dziarnowski, A., Cane, D.E. and Khosla, C. (2003) Mechanistic analysis of acyl transferase domain exchange in polyketide synthase modules. *J. Am. Chem. Soc.*, **125**, 5366–5374.
23. Kao, C.M., McPherson, M., McDaniel, R.N., Fu, H., Cane, D.E. and Khosla, C. (1998) Alcohol stereochemistry in polyketide backbones is controlled by the  $\beta$ -ketoreductase domains of modular polyketide synthases. *J. Am. Chem. Soc.*, **120**, 2478–2479.
24. Kellenberger, L., Galloway, I.S., Sauter, G., Bohm, G., Hanefeld, U., Cortes, J., Staunton, J. and Leadlay, P.F. (2008) A polylinker approach to reductive loop swaps in modular polyketide synthases. *ChemBioChem*, **9**, 2740–2749.
25. Eng, C.H., Yuzawa, S., Wang, G., Baidoo, E.E., Katz, L. and Keasling, J.D. (2016) Alteration of polyketide stereochemistry from *anti* to *syn* by a ketoreductase domain exchange in a type I modular polyketide synthase subunit. *Biochemistry*, **55**, 1677–1680.
26. Kao, C.M., McPherson, M., McDaniel, R.N., Fu, H., Cane, D.E. and Khosla, C. (1997) Gain of function mutagenesis of the erythromycin polyketide synthase. 2. Engineered biosynthesis of eight-membered ring tetraketide lactone. *J. Am. Chem. Soc.*, **119**, 11339–11340.
27. McDaniel, R., Kao, C.M., Hwang, S.J. and Khosla, C. (1997) Engineered intermodular and intramodular polyketide synthase fusions. *Chem. Biol.*, **4**, 667–674.
28. Gaisser, S., Kellenberger, L., Kaja, A.L., Weston, A.J., Lill, R.E., Wirtz, G., Kendrew, S.G., Low, L., Sheridan, R.M., Wilkinson, B. *et al.* (2003) Direct production of ivermectin-like drugs after domain exchange in the avermectin polyketide synthase of *Streptomyces avermitilis* ATCC31272. *Org. Biomol. Chem.*, **1**, 2840–2847.
29. Medema, M.H., Kottmann, R., Yilmaz, P., Cummings, M., Biggins, J.B., Blin, K., De Bruijn, I., Chooi, Y.H., Claesen, J., Coates, R.C. *et al.* (2015) Minimum Information about a Biosynthetic Gene cluster. *Nat. Chem. Biol.*, **11**, 625–631.
30. Weber, T., Blin, K., Duddela, S., Krug, D., Kim, H.U., Bruccoleri, R., Lee, S.Y., Fischbach, M.A., Müller, R., Wohlleben, W. *et al.* (2015) antiSMASH 3.0 – a comprehensive resource for the genome mining of biosynthetic gene clusters. *Nucleic Acids Res.*, **43**(W1), W237–W243.
31. Weber, L. (2008) JChem Base—ChemAxon. *Chem. World*, **5**, 65–66.
32. Keatinge-Clay, A.T. (2007) A tylosin ketoreductase reveals how chirality is determined in polyketides. *Chem. Biol.*, **14**, 898–908.
33. Cheng, J., Randall, A.Z., Sweredoski, M.J. and Baldi, P. (2005) SCRATCH: a protein structure and structural feature prediction server. *Nucleic Acids Res.*, **33**, W72–W76.
34. Magnan, C.N. and Baldi, P. (2014) SSpro/ACCpro 5: almost perfect prediction of protein secondary structure and relative solvent accessibility using profiles, machine learning and structural similarity. *Bioinformatics*, **30**, 2592–2597.
35. Carhart, R.E., Smith, D.H. and Venkataraghavan, R. (1985) Atom pairs as molecular features in structure-activity studies: definition and applications. *J. Chem. Inf. Model.*, **25**, 64–73.
36. Raymond, J.W., Gardiner, E.J. and Willett, P. (2002) Heuristics for similarity searching of chemical graphs using a maximum common edge subgraph algorithm. *J. Chem. Inf. Comput. Sci.*, **42**, 305–316.
37. Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K. and Madden, T.L. (2009) BLAST+: architecture and applications. *BMC Bioinformatics*, **10**, 421.
38. Zierep, P.F., Padilla, N., Yonchev, D.G., Telukunta, K.K., Klementz, D. and Günther, S. (2017) SeMPI: a genome-based secondary metabolite prediction and identification web server. *Nucleic Acids Res.*, **45**, W64–W71.
39. Zheng, J. and Keatinge-Clay, A.T. (2013) The status of type I polyketide synthase ketoreductases. *MedChemComm*, **4**, 34–40.
40. Keatinge-Clay, A.T. (2016) Stereocontrol within polyketide assembly lines. *Nat. Prod. Rep.*, **33**, 141–149.
41. Sundaram, S. and Hertweck, C. (2016) On-line enzymatic tailoring of polyketides and peptides in thio-template systems. *Curr. Opin. Chem. Biol.*, **31**, 82–94.
42. Nguyen, T., Ishida, K., Jenke-Kodama, H., Dittmann, E., Gurgui, C., Hochmuth, T., Taudien, S., Platzer, M., Hertweck, C. and Piel, J. (2008) Exploiting the mosaic structure of trans-acyltransferase polyketide synthases for natural product discovery and pathway dissection. *Nat. Biotechnol.*, **26**, 225–233.
43. Wu, J., Kinoshita, K., Khosla, C. and Cane, D.E. (2004) Biochemical analysis of the substrate specificity of the  $\beta$ -ketoacyl-acyl carrier protein synthase domain of module 2 of the erythromycin polyketide synthase. *Biochemistry*, **43**, 16301–16310.
44. Murphy, A.C., Hong, H., Vance, S., Broadhurst, R.W. and Leadlay, P.F. (2016) Broadening substrate specificity of a chain-extending ketosynthase through a single active-site mutation. *Chem. Commun.*, **52**, 8373–8376.
45. Faille, A., Gavalda, S., Slama, N., Lherbet, C., Maveyraud, L., Guillet, V., Laval, F., Quémar, A., Mourey, L. and Pedelacq, J.-D. (2017) Insights into substrate modification by dehydratases from Type I Polyketide Synthases. *J. Mol. Biol.*, **429**, 1554–1569.
46. Conway, K.R. and Boddy, C.N. (2013) ClusterMine360: a database of microbial PKS/NRPS biosynthesis. *Nucleic Acids Res.*, **41**, D402–D407.
47. Khater, S., Gupta, M., Agrawal, P., Sain, N., Prava, J., Gupta, P., Grover, M., Kumar, N. and Mohanty, D. (2017) SBSPKsv2: structure-based sequence analysis of polyketide synthases and non-ribosomal peptide synthetases. *Nucleic Acids Res.*, **45**, W72–W79.
48. Hagen, A., Poust, S., de Rond, T., Yuzawa, S., Katz, L., Adams, P.D., Petzold, C.J. and Keasling, J.D. (2014) *In vitro* analysis of carboxyacyl substrate tolerance in the loading and first extension modules of borrelidin polyketide synthase. *Biochemistry*, **53**, 5975–5977.