

UCLA

UCLA Previously Published Works

Title

Correction: Hamilton et al. Receptors for Insulin-Like Growth Factor-2 and Androgens as Therapeutic Targets in Triple-Negative Breast Cancer. *Int. J. Mol. Sci.* 2017, 18, 2305

Permalink

<https://escholarship.org/uc/item/7kt8d8bh>

Journal

International Journal of Molecular Sciences, 25(5)

ISSN

1661-6596

Authors

Hamilton, Nalo

Austin, David

Márquez-Garbán, Diana

et al.

Publication Date

2024

DOI

10.3390/ijms25052579

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed



Correction

Correction: Hamilton et al. Receptors for Insulin-Like Growth Factor-2 and Androgens as Therapeutic Targets in Triple-Negative Breast Cancer. *Int. J. Mol. Sci.* 2017, 18, 2305

Nalo Hamilton ^{1,2,*} , David Austin ³, Diana Márquez-Garbán ^{2,4}, Rudy Sanchez ⁵, Brittney Chau ⁶, Kay Foos ⁷, Yanyuan Wu ³ , Jaydutt Vadgama ^{2,3,4} and Richard Pietras ^{3,4}

¹ UCLA School of Nursing, University of California at Los Angeles, Los Angeles, CA 90095, USA

² UCLA Jonsson Comprehensive Cancer Center, University of California at Los Angeles, Los Angeles, CA 90095, USA; dmarquez@mednet.ucla.edu (D.M.-G.); jayvadgama@cdrewu.edu (J.V.)

³ Department of Medicine, Division of Cancer Research and Training, Charles Drew University School of Medicine and Science, Los Angeles, CA 90059, USA; davidAustin@cdrewu.edu (D.A.); yanyuanwu@cdrewu.edu (Y.W.); rpietras@mednet.ucla.edu (R.P.)

⁴ UCLA David Geffen School of Medicine, Department of Medicine, Division of Hematology-Oncology, University of California at Los Angeles, Los Angeles, CA 90095, USA

⁵ Department of Biology, California State University Channel Islands, Camarillo, CA 93012, USA; rudy.sanchez917@myci.csuci.edu

⁶ Department of Integrative Ecology and Evolutionary Biology and Physiology, UCLA College of Life Sciences, University of California at Los Angeles, Los Angeles, CA 90095, USA; chaubrittney@gmail.com

⁷ Department Physiological, UCLA College of Life Sciences, University of California at Los Angeles, Los Angeles, CA 90095, USA; kaymaliafoos@yahoo.com

* Correspondence: nhamilton@sonnet.ucla.edu; Tel.: +1-310-206-0589



Citation: Hamilton, N.; Austin, D.; Márquez-Garbán, D.; Sanchez, R.; Chau, B.; Foos, K.; Wu, Y.; Vadgama, J.; Pietras, R. Correction: Hamilton et al. Receptors for Insulin-Like Growth Factor-2 and Androgens as Therapeutic Targets in Triple-Negative Breast Cancer. *Int. J. Mol. Sci.* 2017, 18, 2305. *Int. J. Mol. Sci.* 2024, 25, 2579. <https://doi.org/10.3390/ijms25052579>

Received: 13 October 2023

Accepted: 8 November 2023

Published: 23 February 2024



Copyright: © 2024 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

In the original publication [1], there were mistakes in Figures 4A and 7 as published. In Figure 4A, there is an apparent duplication of flow cytometry data for the 24 h timepoint. The image for the MDA-MB-231 vehicle is duplicated for BT549 enzalutamide. In the original manuscript, Figure 7 was the result of parallel blots developed on film. As such, concerns were raised regarding the image quality and splicing of loading control.

In light of these concerns, the experiment was repeated, with additional controls, and all samples were run together. Due to the inclusion of new controls, additional information is needed for the legend associated with Figure 7. The corrected images for Figures 4A and 7 along with the updated legend for Figure 7 are presented below.

The authors state that the scientific conclusions are unaffected. This correction was approved by the Academic Editor. The original publication has also been updated.





Figure 4. IGF1R/IR antagonists induce TNBC cell death. TNBC cells were grown to 75–80% confluence in complete media, then transferred to the indicated inhibitor-conditioned media for 24 h (A)

or 48 h (B). Cells were harvested and prepared as per the manufacturer's recommended protocol for flow cytometry using 7-AAD and Annexin-V. Analyses were performed using LSRII and FloJo Software (BD FACSDiva Software v8.0.3).




Figure 7. Effect of IGF1R/IR antagonists on AR expression in mesenchymal-subtype TNBC cell line BT549. BT549 (mesenchymal-like) cultures were exposed to control media (CON), BMS-754807 (BMS; 20 μ M) or NVP-AEW807 (NVP; 8 μ M) containing media for 12 or 24 h. Total protein was isolated, processed and transferred to PVDF membranes which were probed for AR expression (1:500, Cell Signaling #5153, Cell Signaling Technology, Danvers, MA, USA). GAPDH (1:1000; h-FAB Rhodamine, BioRad, Hercules, CA, USA) and β -actin (1:500, Santa Cruz Biotechnology, Inc., Dallas, TX, USA) were used as loading controls. C2C12, a subclone of the murine myoblast cell line, was used as a negative control for AR specificity. Western immunoblot was representative of four independent experiments.

Reference

1. Hamilton, N.; Austin, D.; Márquez-Garbán, D.; Sanchez, R.; Chau, B.; Foos, K.; Wu, Y.; Vadgama, J.; Pietras, R. Receptors for Insulin-Like Growth Factor-2 and Androgens as Therapeutic Targets in Triple-Negative Breast Cancer. *Int. J. Mol. Sci.* **2017**, *18*, 2305. [[CrossRef](#)] [[PubMed](#)]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.