

UC Irvine

UC Irvine Previously Published Works

Title

Evolution of negative immune regulators

Permalink

<https://escholarship.org/uc/item/7n26h5cj>

Journal

PLOS Pathogens, 15(8)

ISSN

1553-7366

Authors

Frank, Steven A
Schmid-Hempel, Paul

Publication Date

2019

DOI

10.1371/journal.ppat.1007913

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

OPINION

Evolution of negative immune regulators

Steven A. Frank^{1*}, Paul Schmid-Hempel²

1 Department of Ecology and Evolutionary Biology, University of California, Irvine, California, United States of America, **2** Institute of Integrative Biology, ETH Zürich, Zürich, Switzerland

* safrank@uci.edu

Hosts must recognize pathogen invasion and respond rapidly. The need for speed requires a low threshold for triggering a response, causing occasional false alarms. Host immune systems therefore require strong negative regulators to shut down unnecessary responses. The evolutionary consequences of rapid trigger dynamics balanced by negative regulators have received little attention. Here, we emphasize four aspects that influence the evolutionary genetics of immunity: diverse gene families of rapidly acting triggers opposed by slower-acting negative regulators, pathogen attack against negative regulators, diversifying selection of negative regulators by pathogen pressure, and heritability of immune-related disease from imbalance between triggers and negative regulators.

Background on triggers of immunity


A broad sensor array of receptors, located on the cell surface or in the cytosol, recognize pathogen-associated molecular patterns and trigger innate immunity. Common triggers include the peptidoglycan recognition proteins, gram-negative binding proteins, Toll-like receptors (TLRs), nucleotide-binding domain leucine-rich repeats (NLRs), and the cyclic GMP-AMP synthase (cGAS) system that detects double-stranded DNA (dsDNA) [1, 2]. In the NLR family of immune triggers, some plant species have more than 1,000 genes [3], the coral *Acropora digitifera* has around 500 genes, sea urchins have approximately 200, and zebrafish have about 385 [4]. Sponges, perhaps the most ancient animals, have a diverse family of NLR receptors that can recognize a wide variety of microbes [5].

These receptors trigger immune cascades that are often evolutionarily ancient within groups or across broad domains of life. Key components of such cascades include nuclear factor- κ B (NF- κ B) transcription factors, interferons (IFNs), interleukins, and Toll and immune deficiency (IMD) pathways [6–9]. The primary dsDNA sensor cGAS triggers stimulator of IFN genes (STING), which activates a variety of fundamental immune cascades [10].

Prediction 1: Fast, error-prone triggers of innate immunity are balanced by slower-acting negative regulators that provide a more accurate assessment of invasion

The need for speed in immune response [11] demands a low threshold for triggering the immune cascade, which inevitably leads to many errors. The evolution of the trigger threshold will be influenced by the risk of attack and the costs and benefits defense [12]. The faster and more error-prone the initial triggers, the greater the need for negative regulators that can correct errors.

All organisms have an impressive array of negative regulators. Post-transcriptional modification of innate sensors and downstream molecules suppress the effect of the pattern



OPEN ACCESS

Citation: Frank SA, Schmid-Hempel P (2019) Evolution of negative immune regulators. *PLoS Pathog* 15(8): e1007913. <https://doi.org/10.1371/journal.ppat.1007913>

Editor: Carolyn B Coyne, University of Pittsburgh, UNITED STATES

Published: August 1, 2019

Copyright: © 2019 Frank, Schmid-Hempel. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: The Donald Bren Foundation supported this research. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

recognition receptor triggers [13, 14]. Tripartite motif containing 25 (TRIM25) negatively regulates the innate antiviral responses [15]. Various negative regulators suppress the Toll cascade and cytokine-signaling in mammals [16], including type I IFN [17]. The cascade triggered by the dsDNA sensor cGAS has several negative regulators, perhaps because dsDNA is nonspecific and prone to false signaling [10]. Numerous noncoding micro-RNAs (miRNAs) regulate innate immunity, including the Toll cascade in defense against gram-negative bacteria [18], modification of type I IFN regulation [17, 19], immune cell differentiation [20], and repression of the NLR triggers and their immune cascade [21, 22].

Different organisms will tune the tradeoff between faster response and lower error rates in different ways. A focus on the tuning of that tradeoff may lead to interesting comparative insights about the design of immune regulation.

Prediction 2: Negative regulators of immunity provide a point of attack for invaders

Attackers use diverse mechanisms to block the expression of host innate immunity [23, 24, 25–27]. We predict that attackers may often target stimulation of the host’s natural negative immune regulators. In fact, *Leishmania* exploits a negative TLR regulator to suppress production of key proinflammatory cytokines [28], viruses manipulate post-translational modifications that normally prevent aberrant responses [29], and many microbes interfere with (negative) receptor crosstalk that normally adjusts immune responses [30].

Invaders of insects alter the expression of an impressive variety of miRNAs, including negative regulators. Some pathogens express their own miRNAs to alter host immunity. The function of most miRNAs remains unknown, providing a rich topic for further study [31].

Tumors also evolve to down-regulate immunity against cancer cells and often manipulate the normal repressors of immunity. For example, Vinay and colleagues [32] note that “tumors can evade immune surveillance by . . . production of several immune suppressive cytokines . . . In addition to immune suppressive cytokines, other factors . . . inhibit the differentiation of progenitors . . . affecting efficient uptake and antigen presentation.” The multiple ways in which tumors manipulate negative regulators of immunity suggest that such mechanisms may be common among pathogens and parasites.

Prediction 3: Attack against negative regulators imposes diversifying selection, favoring diverse and often rapidly evolving gene families of negative regulators

Hosts gain by altering the specific sequences of their negative regulators to avoid recognition or mimicking by pathogen effectors. Negative immune regulators that are under attack may form diverse gene families and evolve relatively rapidly. Few prior studies have focused explicitly on the diversity and evolutionary rate of negative regulators.

In *Drosophila*, Lee and Ferrandon [33] note that “A striking finding of the cell line screens was the large number of IMD pathway negative regulators that were discovered, as compared with positive regulators.” Across 12 species of *Drosophila*, genes involved in the response signaling cascade tended to be more conserved than recognition proteins, but those that functioned primarily in modulating the immune response showed evidence for positive selection far above the genomic average [34].

The nature of diversifying selection depends on the specificity of mechanisms by which attackers manipulate host immunity. In human immunity, attack against negative immune regulators may be relatively nonspecific, in the sense that the attackers may not be differentially targeting molecules based on sequence variability. By contrast, for the small RNAs that

regulate immunity in plants, insects, and other organisms, the RNA sequences and the mechanisms of manipulation are likely to be highly specific. The more sequence-specific the mechanisms that attackers use to manipulate negative immune regulators, the stronger the diversifying selection will be.

Prediction 4: Strongly opposed positive and negative regulators increase the probability of misregulation and the heritability of immune-related disease

Regulation by strongly opposed triggers and repressors is prone to failure ([35] discusses opposing regulation and disease). The evolutionary tuning of immune control by opposed triggers and repressors is particularly challenging. The intensity of attack and the specific kinds of attack vary widely over time and space. Strong, fluctuating selection likely maintains genetic variability within populations for triggers and repressors.

Plants may use miRNAs to tune the threshold for the triggering of immunity by the NLR pattern recognition receptors. Canto-Pastor and colleagues [36] found that varying miRNA expression quantitatively altered the susceptibility of plants to infection. This potential for quantitative variation in the sensitivity of the triggers likely leads to heritable genetic variation in the speed and intensity of the initial immune response. It would be interesting to learn more about the later-acting negative regulators of immunity that control false alarms set off by the NLR triggers and the potential for quantitative variability in those negative regulators.

Similarly, constitutively expressed inhibitors of the type I IFN response in human immunity may “determine a threshold of activation of an antiviral response” [37]. A quantitatively regulated threshold of sensitivity is likely to be associated with heritable quantitative variability in the triggers of immunity.

Genetic variability in triggers and negative regulators can lead to mismatches in the opposing regulatory components of innate immunity and various types of immune-related disease. One possibility concerns the various immune-related disorders associated with misexpression of type I IFN [37]. Another possible example concerns the quick trigger against potentially foreign DNA by cGAS, which is susceptible to autoimmunity [10]. In the cGAS response, misregulation and immune-related disease arise in spite of the many negative regulators of the cGAS cascade.

Our hypothesis and these preliminary examples suggest that further study of genetic variability in the positive and negative regulators of immune cascades may lead to interesting insights about the heritability of immune misregulation.

Acknowledgments

SAF thanks the Theoretical Biology group of the Institute for Integrative Biology at ETH Zürich for hosting his sabbatical visit during the period of this work.

References

1. Yin Q, Fu TM, Li JX, Wu H. Structural biology of innate immunity. In: Littman DR, Yokoyama WM, editors. *Annual Review of Immunology*. Vol 33. Annual Review of Immunology. 33 2015. p. 393–416.
2. Chow KT, Gale M, Loo YM. RIG-I and Other RNA Sensors in Antiviral Immunity. In: Littman DR, Yokoyama WM, editors. *Annual Review of Immunology*, Vol 36. Annual Review of Immunology. 36 2018. p. 667–94.
3. Karasov T, Chae E, Herman J, Bergelson J. Mechanisms to mitigate the trade-off between growth and defense. *Plant Cell*. 2017; 29(4):666–80. <https://doi.org/10.1105/tpc.16.00931> PMID: 28320784
4. Jones JDG, Vance RE, Dangl JL. Intracellular innate immune surveillance devices in plants and animals. *Science*. 2016; 354(6316):aaf6395.

5. Degnan SM. The surprisingly complex immune gene repertoire of a simple sponge, exemplified by the NLR genes: A capacity for specificity? *Developmental and Comparative Immunology*. 2015; 48(2):269–74. <https://doi.org/10.1016/j.dci.2014.07.012> PMID: 25058852
6. Aggarwal K, Silverman N. Positive and negative regulation of the *Drosophila* immune response. *BMB Rep*. 2008; 41(4):267–77. Epub 2008/05/03. <https://doi.org/10.5483/bmbrep.2008.41.4.267> PMID: 18452646. *BMB Reports*. 2008;41(4):267–77.
7. Werner T, Liu G, Kang D, Ekengren S, Steiner H, Hultmark D. A family of peptidoglycan recognition proteins in the fruit fly *Drosophila melanogaster*. *Proceedings of the National Academy of Sciences USA*. 2000; 97(5):13772–7.
8. Kurata S. Peptidoglycan recognition proteins in *Drosophila* immunity. *Developmental and Comparative Immunology*. 2014; 42:36–41. <https://doi.org/10.1016/j.dci.2013.06.006> PMID: 23796791
9. Imler J-L. Overview of *Drosophila* immunity: A historical perspective. *Developmental and Comparative Immunology*. 2014; 42:3–15. <https://doi.org/10.1016/j.dci.2013.08.018> PMID: 24012863
10. Ablasser A, Chen ZJ. cGAS in action: Expanding roles in immunity and inflammation. *Science*. 2019; 363(6431):aat8657.
11. Duneau D, Ferdy JB, Revah J, Kondolf H, Ortiz GA, Lazzaro BP, et al. Stochastic variation in the initial phase of bacterial infection predicts the probability of survival in *D. melanogaster*. *Elife*. 2017; 6. Epub 2017/10/13. <https://doi.org/10.7554/eLife.28298> PMID: 29022878.
12. Metcalf CJE, Tate AT, Graham AL. Demographically framing trade-offs between sensitivity and specificity illuminates selection on immunity. *Nature Ecology & Evolution*. 2017; 1(11):1766–72. <https://doi.org/10.1038/s41559-017-0315-3> PMID: 28963489
13. Cao XT. Self-regulation and cross-regulation of pattern-recognition receptor signalling in health and disease. *Nature Reviews Immunology*. 2016; 16(1):35–50. <https://doi.org/10.1038/nri.2015.8> PMID: 26711677
14. Liu J, Qian C, Cao XT. Post-translational modification control of innate immunity. *Immunity*. 2016; 45(1):15–30. <https://doi.org/10.1016/j.immuni.2016.06.020> PMID: 27438764
15. Martin-Vicente M, Medrano LM, Resino S, Garcia-Sastre A, Martinez I. TRIM25 in the regulation of the antiviral innate immunity. *Frontiers in Immunology*. 2017; 8:1187. <https://doi.org/10.3389/fimmu.2017.01187> PMID: 29018447
16. Vidya MK, Kumar VG, Sejian V, Bagath M, Krishnan G, Bhatta R. Toll-like receptors: Significance, ligands, signaling pathways, and functions in mammals. *International Reviews of Immunology*. 2018; 37(1):20–36. <https://doi.org/10.1080/08830185.2017.1380200> PMID: 29028369
17. Arimoto KI, Miyauchi S, Stoner SA, Fan JB, Zhang DE. Negative regulation of type I IFN signaling. *Journal of Leukocyte Biology*. 2018; 103(6):1099–116. <https://doi.org/10.1002/jlb.2mir0817-342r> PMID: 29357192
18. Keck J, Gupta R, Christenson LK, Arulanandam BP. MicroRNA mediated regulation of immunity against gram-negative bacteria. *International Reviews of Immunology*. 2017; 36(5):287–99. <https://doi.org/10.1080/08830185.2017.1347649> PMID: 28800263
19. Forster SC, Tate MD, Hertzog PJ. MicroRNA as type I interferon-regulated transcripts and modulators of the innate immune response. *Frontiers in Immunology*. 2015; 6:1–9.
20. Yuan XY, Berg N, Lee JW, Le TT, Neudecker V, Jing N, et al. MicroRNA miR-223 as regulator of innate immunity. *Journal of Leukocyte Biology*. 2018; 104(3):515–24. <https://doi.org/10.1002/JLB.3MR0218-079R> PMID: 29969525
21. Li Y, Zhang QQ, Zhang JG, Wu L, Qi YJ, Zhou JM. Identification of microRNAs Involved in pathogen-associated molecular pattern-triggered plant innate immunity. *Plant Physiology*. 2010; 152(4):2222–31. <https://doi.org/10.1104/pp.109.151803> PMID: 20164210
22. Zhang Y, Xia R, Kuang HH, Meyers BC. The diversification of plant NBS-LRR defense genes directs the evolution of microRNAs that target them. *Molecular Biology and Evolution*. 2016; 33(10):2692–70. <https://doi.org/10.1093/molbev/msw154> PMID: 27512116
23. Beachboard DC, Horner SM. Innate immune evasion strategies of DNA and RNA viruses. *Current Opinion in Microbiology*. 2016; 32:113–9. <https://doi.org/10.1016/j.mib.2016.05.015> PMID: 27288760
24. Johnson DI. *Bacterial pathogens and their virulence factors*. New York, NY: Springer Science+Business Media; 2017.
25. Odendall C, Kagan JC. Activation and pathogenic manipulation of the sensors of the innate immune system. *Microbes and Infection*. 2017; 19(4–5):229–37. <https://doi.org/10.1016/j.micinf.2017.01.003> PMID: 28093320
26. Agrawal P, Nawadkar R, Ojha H, Kumar J, Sahu A. Complement evasion strategies of viruses: an overview. *Frontiers in Microbiology*. 2017; 8:1117. <https://doi.org/10.3389/fmicb.2017.01117> PMID: 28670306

27. Kobayashi SD, Malachowa N, Deleo FR. Neutrophils and Bacterial Immune Evasion. *Journal of Innate Immunity*. 2018; 10(5–6):432–41. <https://doi.org/10.1159/000487756> PMID: 29642066
28. Cecílio P, Pérez-Cabezas Ba, Santarém N, Maciel J, Rodrigues V, Cordeiro da Silva A. Deception and manipulation: the arms of *Leishmania*, a successful parasite. *Frontiers in Immunology*. 2014; 5:480. <https://doi.org/10.3389/fimmu.2014.00480> PMID: 25368612
29. Gack MU. Mechanisms of RIG-I-like receptor activation and manipulation by viral pathogens. *Journal of Virology*. 2014; 88(10):5213–6. <https://doi.org/10.1128/JVI.03370-13> PMID: 24623415
30. Hajishengallis G, Lambris JD. Microbial manipulation of receptor crosstalk in innate immunity. *Nature Reviews Immunology*. 2011; 11:187–200. <https://doi.org/10.1038/nri2918> PMID: 21350579
31. Hussain M, Asgari S. MicroRNAs as mediators of insect host–pathogen interactions and immunity. *Journal of Insect Physiology*. 2014; 70:151–8. <https://doi.org/10.1016/j.jinsphys.2014.08.003> PMID: 25152509
32. Vinay DS, Ryan EP, Pawelec G, Talib WH, Stagg J, Elkord E, et al. Immune evasion in cancer: mechanistic basis and therapeutic strategies. *Seminars in Cancer Biology*. 2015; 25:S185–S98.
33. Lee KZ, Ferrandon D. Negative regulation of immune responses on the fly. *EMBO Journal*. 2011; 30(6):988–90. <https://doi.org/10.1038/emboj.2011.47> PMID: 21407251
34. Sackton TB, Lazzaro BP, Schlenke TA, Evans JD, Hultmark D, Clark AG. Dynamic evolution of the innate immune system in *Drosophila*. *Nature Genetics*. 2007; 39(12):1461–8. <https://doi.org/10.1038/ng.2007.60> PMID: 17987029
35. Frank SA, Crespi BJ. Pathology from evolutionary conflict, with a theory of X chromosome versus auto-some conflict over sexually antagonistic traits. *Proc Natl Acad Sci U S A*. 2011; 108 Suppl 2:10886–93. Epub 2011/06/22. <https://doi.org/10.1073/pnas.1100921108> PMID: 21690397.
36. Canto-Pastor A, Santos B, Valli A, Summers W, Schornack S, Baulcombe D. Enhanced resistance to bacterial and oomycete pathogens by short tandem target mimic RNAs in tomato. *Proceedings of the National Academy of Sciences USA*. 2019; 116(7):2755–60.
37. Crow YJ. Type I interferonopathies: a novel set of inborn errors of immunity. *Ann N Y Acad Sci*. 2011; 1238:91–8. Epub 2011/12/02. <https://doi.org/10.1111/j.1749-6632.2011.06220.x> PMID: 22129056.