

UC Berkeley

UC Berkeley Previously Published Works

Title

Null structures and degenerate dispersion relations in two space dimensions

Permalink

<https://escholarship.org/uc/item/7p34x1pw>

Authors

Fu, Yuqiu

Tataru, Daniel

Publication Date

2017-12-30

Peer reviewed

NULL STRUCTURES AND DEGENERATE DISPERSION RELATIONS IN TWO SPACE DIMENSIONS

YUQIU FU AND DANIEL TATARU

ABSTRACT. For a dispersive PDE, the degeneracy of its dispersion relation will deteriorate dispersion of waves, and strengthen nonlinear effects. Such negative effects can sometimes be mitigated by some null structure in the nonlinearity.

Motivated by water-wave problems, in this paper we consider a class of nonlinear dispersive PDEs in 2D with cubic nonlinearities, whose dispersion relations are radial and have vanishing Gaussian curvature on a circle. For such a model we identify certain null structures for the cubic nonlinearity, which suffice in order to guarantee global scattering solutions for the small data problem. Our null structures in the power-type nonlinearity are weak, and only eliminate the worst nonlinear interaction. Such null structures arise naturally in some water-wave problems.

1. INTRODUCTION

We consider the following model Cauchy problem

$$(1.1) \quad \begin{cases} \partial_t u - ih(D)u = A(D)(|P_{\leq M}u|^2 P_{\leq M}u) \\ u(0, x) = u_0 \in L^2(\mathbb{R}^2). \end{cases}$$

with a cubic nonlinearity.

Here $h(\xi)$ is a radial dispersion relation on \mathbb{R}^2 which is degenerate on the unit circle, i.e. its Hessian is degenerate there. $P_{\leq M}$ is a cutoff in the frequency space selecting an annulus near the unit circle; this is where the strongest nonlinear interactions are occurring.

The interesting object here is the multiplier A , whose symbol $A(\xi)$ vanishes to some order on the same circle. We observe that without $A(D)$, the worst nonlinear interaction of waves in the power-type nonlinearity occurs when waves have equal frequencies on the unit circle. Therefore intuitively the Fourier multiplier $A(D)$ provides a null structure in (1.1) by eliminating the worst nonlinear interaction.

We will show that such types of null structures are indeed effective in controlling the long time dynamics, provided that the order of vanishing of $A(\xi)$ on the unit circle is no less than half of that of the order of degeneracy there for the Hessian of H . Under these conditions, we will prove that small data leads to global scattering solutions.

The placement of the multiplier $A(D)$ in the equation is less important. The same result holds if $A(D)$ applies instead to any of the factors in the cubic nonlinearity, e.g. as in $P_{\leq M}(A(D)uP_{\leq M}\bar{u}P_{\leq M}u)$, etc.

The second author was partially supported by the NSF grant DMS-1266182 as well as by a Simons Investigator grant from the Simons Foundation.

We now describe our model in more detail. First, we suppose there exists $\delta \in (0, 1)$ and a smooth function $\gamma : (1 - \delta, 1 + \delta) \rightarrow \mathbb{R}$ such that $h(\xi) = \gamma(|\xi|)$. We make the following assumptions on γ :

- (a) Transversality: $|\gamma'(r)| \approx 1$ for every $r \in (1 - \delta, 1 + \delta)$.
- (b) Degeneracy: There exists $\beta \in \mathbb{Z}_+$ such that $|\gamma''(r)| \approx |r - 1|^\beta$ for every $r \in (1 - \delta, 1 + \delta)$.

The first condition guarantees the transversality of waves with angularly separated frequencies, and the second condition gives a finite order β of dispersion degeneracy on the unit circle. Under these conditions, the worst cubic nonlinear interactions will turn out to occur between equal frequencies on the unit circle.

The Fourier multiplier $P_{\leq M}$ is a smooth cutoff in the frequency space selecting the region $|\xi| \in (1 - 2^{M+1}, 1 + 2^{M+1})$, where the integer $M < -100$ is chosen such that $2^{M+4} < \delta$. This is introduced so that we restrict our attention to the frequency region where the most interesting part of nonlinear interactions occur.

Last but not least, the Fourier multiplier $A(D)$ is given by a symbol $A(\xi)$ with the property that

$$(1.2) \quad |A(\xi)| \lesssim ||\xi| - 1|^{\beta/2}$$

for $|\xi| \in (1 - \delta, 1 + \delta)$, and

$$(1.3) \quad \text{supp } A(\xi) \subset \{\xi \in \mathbb{R}^2 : 1 - 2^{M-1} < |\xi| < 1 + 2^{M-1}\}.$$

The motivation for considering the Cauchy problem (1.1) comes from the study of long-term dynamics for water-wave systems. For an incompressible, inviscid and irrotational fluid occupying a time-dependent domain

$$\Omega_t = \{(x, y) \in \mathbb{R}^2 \times \mathbb{R} : -H_0 \leq y \leq h(x, t)\}$$

for some function $h(x, t)$ and for t in some interval $I_t \subset \mathbb{R}$, the water-wave problem can be reduced (see for example [Lan13]) to the following formulation for unknowns $h, \phi : \mathbb{R}_x^2 \times I_t \rightarrow \mathbb{R}$:

$$(1.4) \quad \begin{cases} \partial_t h = G(h)\phi \\ \partial_t \phi = -gh + \frac{(G(h)\phi + \nabla h \cdot \nabla \phi)^2}{2(1 + |\nabla h|^2)} - \frac{1}{2}|\nabla \phi|^2 + \sigma \operatorname{div} \left(\frac{\nabla h}{(1 + |\nabla h|^2)^{1/2}} \right), \end{cases}$$

where $\sigma \geq 0$ is the surface tension coefficient, g is the gravitational constant, and

$$G(h) := \sqrt{1 + |\nabla h|^2} \mathcal{N}(h).$$

Here $\mathcal{N}(h)$ is the Dirichlet-Neumann map associated to the domain Ω_t , and $\phi(x, t)$ is the restriction of the velocity potential to the boundary surface $y = h(x, t)$.

The dispersion relation for the linearized equation of (1.4) around the zero solution is given by

$$\Lambda(\xi) = \pm \sqrt{|\xi|(g + \sigma|\xi|^2) \tanh(H_0|\xi|)}.$$

For simplicity of computation we consider the infinite depth case $H_0 \rightarrow \infty$, in which case the dispersion relation becomes $\Lambda = \pm \sqrt{g|\xi|(1 + \sigma|\xi|^2/g)}$. If we are considering gravity-capillary water waves, that is $\sigma, g > 0$, then away from the origin, $\det(\nabla^2 \Lambda)$ vanishes exactly on a circle, which implies that we will not have optimal dispersion of waves. This motivates (and is an example of) our general dispersion relations considered above. Our nonlinearity has been chosen to be of cubic type, as for this dispersion relation, nonlinear interactions in quadratic

nonlinearities are either non-resonant or transversal. Because of this, one expects that the long-time dynamics are in fact primarily governed by cubic nonlinear interactions.

Returning to our model problem, we begin by observing that, because of the frequency localization in the nonlinearity, by Hölder's inequality and Bernstein's inequality we have

$$\|A(D)(P_{\leq M}u_1 \overline{P_{\leq M}u_2} P_{\leq M}u_3)\|_{L_x^2} \lesssim \|u_1\|_{L_x^2} \|u_2\|_{L_x^2} \|u_3\|_{L_x^2}.$$

From this estimate one can easily establish local wellposedness of (1.1) in $L^2(\mathbb{R}^2)$ using a standard fixed point argument (see Section 5):

Theorem 1 (Local well-posedness). *For every $R > 0$, there exists $T = T(R) > 0$ such that for every $u_0 \in L^2(\mathbb{R}^2)$ with $\|u_0\|_{L_x^2} \leq R$, there exists a unique strong solution $u \in C^0([0, T], L_x^2)$ to the Cauchy problem (1.1), and the solution map*

$$B_{L_x^2}(0, R) \rightarrow C^0([0, T], L_x^2), u_0 \mapsto u$$

is Lipschitz continuous.

The local well-posedness result above does not use at all the dispersive properties of the equation. This, however, becomes crucial if one consider the global in time well-posedness question. This is the main goal of this paper. Our result is as follows:

Theorem 2 (Global wellposedness for small data). *There exists $\epsilon_0 > 0$ such that for every $u_0 \in L^2(\mathbb{R}^2)$ with $\|u_0\|_{L_x^2} \leq \epsilon_0$, there exists a unique strong global solution $u \in X^0([0, \infty)) \cap C^0([0, \infty); L_x^2)$ to the Cauchy problem (1.1), and the solution map*

$$B_{L_x^2}(0, \epsilon_0) \rightarrow X^0([0, \infty)) \cap C^0([0, \infty), L_x^2), u_0 \mapsto u$$

is Lipschitz continuous.

Furthermore, the solutions are scattering, i.e. for each small data $u_0 \in L^2(\mathbb{R}^2)$ there exist $u_+ \in L^2(\mathbb{R}^2)$, small, with Lipschitz dependence on u_0 , so that

$$\lim_{t \rightarrow \infty} \left(u(t) - e^{ith(D)} u_+ \right) = 0 \quad \text{in } L^2(\mathbb{R}^2).$$

Remark 1. The above wellposedness and scattering result also holds for small solutions that are backward in time. Indeed if $u(t, x)$ is a solution to (1.1), then $\overline{u(-t, x)}$ is a solution to (1.1) with $A(D)$ replaced by $A(-D)$, and initial data $\overline{u_0}$.

The function space X^0 captures the dispersive properties of the solutions, and we have the embedding

$$X^0([0, \infty)) \subset L_t^\infty L_x^2([0, \infty) \times \mathbb{R}^2).$$

This is introduced in Section 2 and is defined using U^2 type spaces associated to the corresponding linear flow.

A key role in the proof of our result is played by localized Strichartz estimates and bilinear L^2 estimates for solutions to the linear homogeneous flow. These are derived in Section 3; we hope they will also be of independent interest due to the optimal treatment of the degeneracy.

The linear and bilinear estimates are transferred to the X^0 space in Section 7. This in turn allows us to conclude the proof of our small data result in Section 5.

Notations: In what follows \widehat{f} will denote the spatial Fourier transform of the function f ,

$$\widehat{f}(t, \xi) = \frac{1}{2\pi} \int_{\mathbb{R}^2} f(t, x) e^{-ix \cdot \xi} dx,$$

and \check{g} will denote the inverse Fourier transform, such that $(\widehat{f})^\check{=} = f$. We denote the standard inner product on L^2 by $\langle \cdot, \cdot \rangle$, that is

$$\langle f, g \rangle = \int f \bar{g}.$$

If X, Y are two subsets of \mathbb{R}^n , then $X \Subset Y$ means that X is contained in a compact subset of Y , and $X + Y$ denotes the set $\{x + y : x \in X, y \in Y\}$. 1_X will denote the indicator function of the set X . The diameter of X is defined as

$$\text{diam } X = \sup_{x, x' \in X} |x - x'|.$$

We let $A \lesssim B$ denote the statement there exists a constant $c > 0$ such that $A \leq cB$, and let $A \gtrsim B$ be the statement $B \lesssim A$. We also let $A \sim B$ be the statement $A \lesssim B$ and $B \lesssim A$. We need to be careful what those implicit constants depend on. Unless otherwise specified, they depend only on γ (and related parameters appeared in this section), and χ which is chosen in Section 2. We call those constants admissible. We will also sometimes describe the dependence explicitly.

2. FUNCTION SPACES

To prove Theorem 2, we will use a perturbative argument. Our function spaces will be of U^p, V^p type, see [KT05]. Since most results in this section concerning U^p, V^p type spaces have already been well developed, we will usually only provide references but no proofs for them. A detailed exposition of those spaces can be found in [KTV14]. Unless otherwise stated, we assume $p \in (1, \infty)$.

Let \mathcal{Z} be the collection of finite partitions of the real line

$$\mathcal{Z} = \{(t_0, \dots, t_K) : -\infty < t_0 < \dots < t_K = \infty\}.$$

Definition 1. We call the function $a : \mathbb{R} \rightarrow L^2(\mathbb{R}^2)$ a U^p -atom if

$$a = \sum_{i=1}^K 1_{[t_{i-1}, t_i)} \phi_{i-1}$$

for some partition $(t_j) \in \mathcal{Z}$, and some $\phi_i \in L^2(\mathbb{R}^2)$ satisfying $\sum_{i=0}^{K-1} \|\phi_i\|_{L^2(\mathbb{R}^2)}^p = 1$. We then define $U^p(\mathbb{R}, L^2(\mathbb{R}^2))$ to be the space of functions $u : \mathbb{R} \rightarrow L^2(\mathbb{R}^2)$ such that

$$\|u\|_{U^p} := \inf \left\{ \sum_{j=1}^{\infty} |\lambda_j| : u = \sum_{j=1}^{\infty} \lambda_j a_j, \lambda_j \in \mathbb{C}, a_j \text{ are } U^p\text{-atoms} \right\} < \infty,$$

with norm $\|\cdot\|_{U^p}$. Here we adopt the convention that $\inf \emptyset = \infty$.

Definition 2. $V^p(\mathbb{R}, L^2(\mathbb{R}^2))$ is defined to be the space of functions $v : \mathbb{R} \rightarrow L^2(\mathbb{R}^2)$ for which

$$\|v\|_{V^p} := \sup_{(t_k)_k \in \mathcal{Z}} \left(\sum_{k=1}^K \|v(t_k) - v(t_{k-1})\|_{L^2(\mathbb{R}^2)}^p \right)^{1/p} < \infty,$$

where we use the convention that $v(t_K) = 0$ if $t_K = \infty$. We let $V_{rc}^p(\mathbb{R}, L^2(\mathbb{R}^2))$ be the subspace consisting of right-continuous functions $v : \mathbb{R} \rightarrow L^2(\mathbb{R}^2)$ in V^p such that $\lim_{t \rightarrow -\infty} v(t) = 0$.

Remark 2.

- $U^p(\mathbb{R}, H), V^p(\mathbb{R}, H)$ spaces can be similarly defined for functions from the real line to any complex Hilbert space H , but for our application in this paper we will only consider functions with $H = L^2(\mathbb{R}^2)$. We will usually omit the space $L^2(\mathbb{R}^2)$ and domain \mathbb{R} in the notation, writing U^p, V^p instead.
- For $1 \leq p < q < \infty$, we have the embeddings

$$(2.1) \quad U^p \hookrightarrow V_{rc}^p \hookrightarrow U^q \hookrightarrow L_t^\infty L_x^2,$$

a proof of which can be found in [KTV14].

- U^p, V^p spaces are Banach spaces. V_{rc}^p is a closed subspace of V^p , which can be seen immediately from definitions and the embedding $V_{rc}^p \hookrightarrow L_t^\infty L_x^2$.
- U^2, V^2 spaces behave well under time truncation and frequency truncation. More precisely we have

$$(2.2) \quad \begin{aligned} \|1_{[a,b]}u\|_{U^2} &\lesssim \|u\|_{U^2} \\ \|P_k u\|_{U^2} &\lesssim \|u\|_{U^2}, \end{aligned}$$

where the implicit constants are independent of $[a, b]$ or k . For the definition of P_k see Definition 4 below. The same inequalities hold if U^2 is replaced by V^2 . These estimates can be easily checked using definitions.

Definition 3. We let $U_{h(D)}^p$ be the space

$$U_{h(D)}^p := \left\{ e^{ith(D)}u : u \in U^p \right\}$$

with the norm

$$\|u\|_{U_{h(D)}^p} = \|e^{-ith(D)}u\|_{U^p}.$$

We similarly define spaces $V_{h(D)}^p, V_{h(D),rc}^p$ corresponding to V^p, V_{rc}^p respectively.

Those spaces are Banach spaces since the original spaces are, and $e^{ith(D)}$ is unitary on L^2 .

As can be seen in the next few sections, our strategy is to perform a dyadic decomposition around the singular set $|\xi| = 1$. To make this precise, we let $\chi : \mathbb{R} \rightarrow [0, 1]$ be a smooth function satisfying

$$\chi(r) = \begin{cases} 1 & \text{if } |r| \leq 1/2, \\ 0 & \text{if } |r| \geq 3/4. \end{cases}$$

Definition 4 (Dyadic decomposition around the unit circle). For $k < 0$, we let $P_{\leq k}$ be the Fourier multipliers given by symbols

$$(2.3) \quad P_{\leq k}(\xi) = \chi(2^{-k}(|\xi| - 1)).$$

We then define $P_k = P_{\leq k} - P_{\leq k-1}$. Also expressions like $P_{k_1 < \cdot \leq k_2}$ are defined in the obvious way. We let P_0 be the Fourier multiplier with symbol

$$P_0(\xi) := 1 - P_{<0}(\xi).$$

Remark 3. We observe that $P_k(\xi)$ is supported on the disjoint union of two annuli

$$\{1 - 2^k < |\xi| < 1 - 2^{k-2}\} \cup \{1 + 2^{k-2} < |\xi| < 1 + 2^k\}.$$

We write $P_k(\xi) = P_k^+(\xi) + P_k^-(\xi)$ where P_k^+ and P_k^- are supported on $1 + 2^{k-2} \leq |\xi| \leq 1 + 2^k$ and $1 - 2^k \leq |\xi| \leq 1 - 2^{k-2}$ respectively. We can of course do the same decomposition to $P_{k_1 \leq \cdot \leq k_2}$. It is sometimes convenient for us to consider the + part and the - part separately, and usually the same argument works for both.

We also observe that projections P_k are almost orthogonal, and therefore we have

$$(2.4) \quad \|u\|_{L^2(\mathbb{R}^2)} \sim \left(\sum_{k=-\infty}^0 \|P_k u\|_{L^2(\mathbb{R}^2)}^2 \right)^{1/2}.$$

Now we can introduce the function spaces we will use in the fixed point argument.

Definition 5. Let Y^0 be the space of functions $v : \mathbb{R} \rightarrow L^2(\mathbb{R}^2)$ such that $P_k v \in V_{h(D),rc}^2$ and

$$\|v\|_{Y^0} := \left(\sum_{k=-\infty}^0 \|P_k v\|_{V_{h(D)}^2}^2 \right)^{1/2} < \infty,$$

endowed with the above norm. Let X^0 be the space of functions $u : \mathbb{R} \rightarrow L^2(\mathbb{R}^2)$ for which $P_k v \in U_{h(D)}^2$ and

$$\|u\|_{X^0} := \left(\sum_{k=-\infty}^0 \|P_k v\|_{U_{h(D)}^2}^2 \right)^{1/2} < \infty,$$

endowed with the above norm.

Remark 4. We first observe that $\|\cdot\|_{L_t^\infty L_x^2} \lesssim \|\cdot\|_{Y^0}$, $\|\cdot\|_{X^0}$, which is an immediate consequence of (2.1) and (2.4), and both spaces are Banach spaces under respective norms. Moreover using (2.4) we can check the embeddings

$$(2.5) \quad U_{h(D)}^2 \hookrightarrow X^0 \hookrightarrow Y^0 \hookrightarrow V_{h(D)}^2.$$

Also, X^0, Y^0 behave well under time truncation and frequency truncation in the sense of (2.2), since U^2, V^2 spaces do.

We can also consider the time restricted spaces $X^0([a, b))$, where we allow b to be infinity.

Definition 6. We let $X^0([a, b))$ be the space of functions $u : [a, b) \rightarrow L^2(\mathbb{R}^2)$ such that the zero extension \tilde{u}

$$\tilde{u}(t) = \begin{cases} u(t) & \text{if } t \in [a, b) \\ 0 & \text{otherwise.} \end{cases}$$

belongs to X^0 . We put norm $\|u\|_{X^0([a,b))} = \|\tilde{u}\|_{X^0}$ on the space $X^0([a, b))$.

Since $\|\cdot\|_{L_t^\infty L_x^2} \lesssim \|\cdot\|_{X^0}$, we conclude that under the identification $u \mapsto \tilde{u}$, $X^0([a, b))$ is a closed subspace of X^0 and therefore $X^0([a, b))$ is a Banach space with the norm $\|\cdot\|_{X^0([a,b))}$. If we write X^0 without specifying intervals, we always mean $X^0(\mathbb{R})$.

We also need the following duality property between U^p and $V^{p'}$ spaces. This is a consequence of Theorem 2.10 and Remark 5 in [HHK09], and Lemma 4.32 in [KTV14].

Proposition 1. *Suppose $u : \mathbb{R} \rightarrow L_x^2(\mathbb{R}^2)$ is absolutely continuous on compact intervals and $u'(t) = 0$ on $(-\infty, 0)$. Then we have*

$$(2.6) \quad \|u\|_{U^2} = \sup_{v \in V_{rc}^2: \|v\|_{V^2} = 1} \left| \int_{\mathbb{R}} \int_{\mathbb{R}^2} u'(t)v(t) dx dt \right|,$$

by which we mean if the right hand side of (2.6) is finite then $u \in U^2$ and (2.6) holds.

Suppose $T \in [0, \infty]$. We let I_T be the linear operator given by

$$(2.7) \quad I_T(f)(t, x) = \int_0^t 1_{[0, T)}(s) e^{i(t-s)h(D)} f(s) ds.$$

We claim that if $f \in L_{t,loc}^1 L_x^2$, then $e^{-ith(D)} I_T(f)$ satisfies the conditions in Proposition 1. To show that $e^{-ith(D)} I_T(f)$ is absolutely continuous on compact intervals we notice that

$$\partial_t \left(e^{-ith(D)} I_T(f) \right) = 1_{[0, T)}(t) e^{-ith(D)} f(t),$$

and by the assumption on f we have $1_{[0, T)}(t) e^{-ith(D)} f(t) \in L_{t,loc}^1 L_x^2$. Therefore by the fundamental theorem of calculus for Banach space valued functions we have that $e^{-ith(D)} I_T(f)$ is absolutely continuous on compact intervals. The previous computation also shows that

$$\partial_t \left(e^{-ith(D)} I_T(f) \right) = 0 \text{ on } (-\infty, 0).$$

As an application of Proposition 1 we prove the following estimate.

Proposition 2. *If $f \in L_{t,loc}^1 L_x^2$, and $T \in [0, \infty]$, then we have*

$$\|I_T(f)(t, x)\|_{X^0} \lesssim \sup_{\|v\|_{Y^0} \leq 1} \left| \int_0^T \int_{\mathbb{R}^2} f(t, x) \overline{v(t, x)} dx dt \right|.$$

Proof. Fix $\sigma > 0$. Observe that since

$$\|I_T(f)(t, x)\|_{X^0} = \left\| \left(\|P_k I_T(f)\|_{U_{h(D)}^2} \right)_{k \leq 0} \right\|_{l^2},$$

there exists a sequence $(b_k)_{k \leq 0}$ in \mathbb{R} such that $\|(b_k)\|_{l^2} = 1$ and

$$\|I_T(f)(t, x)\|_{X^0} \leq \sum_k b_k \|P_k I_T(f)\|_{U_{h(D)}^2} + \sigma.$$

Note that $P_k I_T(f) = I_T(P_k f)$ and $P_k f \in L_{t,loc}^1 L_x^2$. By Proposition 2, there exists $v_k \in V_{rc}^2$ with $\|v_k\|_{V^2} = 1$ such that

$$\begin{aligned} \|I_T(P_k f)\|_{U_{h(D)}^2} &\leq \int_{\mathbb{R}} \int_{\mathbb{R}^2} 1_{[0, T)}(t) e^{-ith(D)} P_k f(t) \overline{v_k(t)} dx dt + 2^k \sigma \\ &= \int_{\mathbb{R}} \int_{\mathbb{R}^2} 1_{[0, T)}(t) f(t) e^{ith(D)} P_k v_k(t) dx dt + 2^k \sigma. \end{aligned}$$

where we have used Plancherel's theorem to move the Fourier multiplier $e^{-ith(D)}P_k$ to act on $\overline{v_k}$. Now we let $v = \sum_k b_k e^{ith(D)}P_k v_k$. Then we have

$$\begin{aligned} \|I_T(f)(t, x)\|_{X^0} &\leq \sum_k b_k \|I_T(P_k f)\|_{U_{h(D)}^2} + \sigma \\ &\leq \sum_k \int_{\mathbb{R}} \int_{\mathbb{R}^2} 1_{[0, T)}(t) f(t) \overline{b_k e^{ith(D)}P_k v_k(t)} dx dt + 3\sigma \\ &= \int_0^T \int_{\mathbb{R}^2} f(t) \overline{v(t)} dx dt + 3\sigma. \end{aligned}$$

Also note that $P_N v = P_N \left(\sum_{k=N-2}^{N+2} P_k v_k \right)$ is right continuous since by assumption v_k are, and by the triangle inequality and (2.2) we have

$$\begin{aligned} \|v\|_{Y^0}^2 &= \sum_N \left\| P_N \left(\sum_{k=N-2}^{N+2} b_k e^{ith(D)} P_k v_k \right) \right\|_{V_{h(D)}^2}^2 \\ &\lesssim \sum_N \left(\sum_{k=N-2}^{N+2} b_k^2 \|e^{ith(D)} v_k\|_{V_{h(D)}^2}^2 \right) \\ &\lesssim \sum_N \left(\sum_{k=N-2}^{N+2} b_k^2 \|v_k\|_{V^2}^2 \right) \\ &\lesssim 1. \end{aligned}$$

Since we can always normalize v (in a way that is independent of σ) such that $\|v\|_{Y^0} \leq 1$ and σ is arbitrary, we obtain the desired result. \square

Remark 5. The proof also shows that if $f \in L_{t,loc}^1 L_x^2$, then for every $T \in [0, \infty]$,

$$\|I_T(f)\|_{X^0} \lesssim \|f\|_{L_t^1([0, T], L_x^2)},$$

and the implicit constant is independent of T . This is because of Hölder's inequality and the embedding $Y^0 \subset L_t^\infty L_x^2$.

3. LOCALIZED LINEAR AND BILINEAR ESTIMATES

In this section we consider linear and bilinear Strichartz estimates for our problem, localized to dyadic annuli $\{||\xi| - 1| \approx 2^k\}$. We begin with the following localized Strichartz estimate.

Proposition 3. *Suppose $k \leq M + 1$. Then we have*

$$(3.1) \quad \left\| e^{ith(D)} P_k u_0 \right\|_{L_{t,x}^4} \lesssim 2^{-\beta k/8} \|P_k u_0\|_{L_x^2}.$$

As a consequence, for $k \leq M + 1$ we have

$$(3.2) \quad \left\| e^{ith(D)} P_{k \leq \cdot \leq M+1} u_0 \right\|_{L_{t,x}^4} \lesssim 2^{-\beta k/8} \|P_{k \leq \cdot \leq M+1} u_0\|_{L_x^2}.$$

Before proving this proposition, we first establish the following localized dispersive estimate.

Lemma 1. *Let $\chi_k(D)$ be the Fourier multiplier $P_{k-2 \leq \cdot \leq k+2}$. Then we have*

$$(3.3) \quad \left\| e^{ith(D)} \chi_k^2(D) u_0 \right\|_{L_x^\infty} \lesssim \frac{2^{-\beta k/2}}{|t|} \|u_0\|_{L_x^1}.$$

Proof. We have $\chi_k(D) = \chi_k^+(D) + \chi_k^-(D)$ where $\chi_k^\pm(D)$ are Fourier multipliers with symbols $P_{k-2 \leq \cdot \leq k+2}^\pm(\xi)$. We will only establish (3.3) for $\|e^{ith(D)}(\chi_k^+(D))^2 u_0\|_{L_x^\infty}$. The other part $\|e^{ith(D)}(\chi_k^-(D))^2 u_0\|_{L_x^\infty}$ can be estimated in the same way and the triangle inequality will give the estimate (3.3), as $\chi_k^+(D)\chi_k^-(D) = 0$. For simplicity of notation we will still write $\chi_k(D)$, omitting the + sign. We first observe that

$$e^{ith(D)}\chi_k^2(D)u_0 = K * u_0$$

where

$$K(t, x) = e^{ith(\xi)}\overline{\chi_k^2(\xi)} = \int_{\mathbb{R}^2} e^{ix \cdot \xi} e^{ith(\xi)} \chi_k^2(\xi) \frac{d\xi}{(2\pi)^2}.$$

Therefore by Young's inequality it suffices to show that

$$\|K(t, x)\|_{L_x^\infty} \lesssim \frac{2^{-\beta k/2}}{|t|}.$$

Since

$$\nabla_\xi(th(\xi) + x \cdot \xi) = t\gamma'(|\xi|)\frac{\xi}{|\xi|} + x,$$

by our assumption (b) on γ in Section 1, there exists a small admissible constant $c > 0$ such that

$$(3.4) \quad |\nabla_\xi(th(\xi) + x \cdot \xi)| \geq c|x|$$

for every $|t| \leq c|x|$ and $|\xi| \in [1 - 2^{M+3}, 1 + 2^{M+3}]$, and

$$(3.5) \quad |\nabla_\xi(th(\xi) + x \cdot \xi)| \geq c|t|$$

for every $|t| > c^{-1}|x|$ and $|\xi| \in [1 - 2^{M+3}, 1 + 2^{M+3}]$.

Write $x = (x_1, x_2)$, and $\xi = (\xi_1, \xi_2)$. We consider three cases.

Case 1. $0 < |t| \leq c|x|$. We integrate by parts and obtain

$$\begin{aligned} K(t, x) &= \int_{\mathbb{R}^2} e^{ix \cdot \xi} e^{ith(\xi)} \chi_k^2(\xi) \frac{d\xi}{(2\pi)^2} \\ &= -i \int_{\mathbb{R}^2} \frac{\left(t\gamma'(|\xi|)\frac{\xi}{|\xi|} + x\right) \cdot \nabla_\xi e^{i(x \cdot \xi + th(\xi))}}{\left|t\gamma'(|\xi|)\frac{\xi}{|\xi|} + x\right|^2} \chi_k^2(\xi) \frac{d\xi}{(2\pi)^2} \\ &= i \int_{\mathbb{R}^2} e^{i(x \cdot \xi + th(\xi))} \nabla_\xi \cdot \left(\frac{\left(t\gamma'(|\xi|)\frac{\xi}{|\xi|} + x\right)}{\left|t\gamma'(|\xi|)\frac{\xi}{|\xi|} + x\right|^2} \chi_k^2(\xi) \right) \frac{d\xi}{(2\pi)^2}. \end{aligned}$$

For every $0 < |t| \leq c|x|$ and $\xi \in \text{supp } \chi_k$, by (3.4) we have for $j = 1, 2$

$$\left| \partial_{\xi_j} \frac{\left(t\gamma'(|\xi|)\frac{\xi_j}{|\xi|} + x_j\right)}{\left|t\gamma'(|\xi|)\frac{\xi}{|\xi|} + x\right|^2} \right| \lesssim \frac{|t| + |x|}{|x|^2} + \frac{|t|^2 + |x|^2}{|x|^3} \lesssim \frac{1}{|t|}.$$

Therefore for $0 < |t| \leq c|x|$,

$$(3.6) \quad \left| \int_{\mathbb{R}^2} e^{i(x \cdot \xi + th(\xi))} \left(\nabla_\xi \cdot \frac{\left(t\gamma'(|\xi|)\frac{\xi}{|\xi|} + x\right)}{\left|t\gamma'(|\xi|)\frac{\xi}{|\xi|} + x\right|^2} \right) \chi_k^2(\xi) \frac{d\xi}{(2\pi)^2} \right| \lesssim \frac{1}{|t|}.$$

By definition

$$\chi_k(\xi) = \chi(2^{-k-2}(|\xi| - 1)) - \chi(2^{-k+3}(|\xi| - 1)).$$

If we abuse our notation by writing $\chi_k(r) = \chi_k(|\xi|)$ for $|\xi| = r$ (note that χ_k is radial), then we have

$$(3.7) \quad \int |\chi'_k(r)| dr \lesssim 1,$$

which implies that

$$\int |\nabla_\xi \chi_k^2(\xi)| d\xi \lesssim 1.$$

Hence again by (3.4) we conclude that for $0 < |t| \leq c|x|$,

$$(3.8) \quad \left| \int_{\mathbb{R}^2} e^{i(x \cdot \xi + t h(\xi))} \frac{(t \gamma'(|\xi|) \frac{\xi}{|\xi|} + x)}{|t \gamma'(|\xi|) \frac{\xi}{|\xi|} + x|^2} \cdot (\nabla_\xi \chi_k^2(\xi)) \frac{d\xi}{(2\pi)^2} \right| \lesssim \frac{1}{|t|}.$$

Combining (3.6) and (3.8) we obtain

$$|K(t, x)| \lesssim \frac{1}{|t|}$$

for every $0 < |t| \leq c|x|$.

Case 2. $|t| > c^{-1}|x|$. We repeat the argument in Case 1 with slight modifications and use (3.5) in the place of (3.4) to obtain

$$|K(t, x)| \lesssim \frac{1}{|t|}$$

for every $|t| > c^{-1}|x|$.

Case 3. $c|x| < |t| \leq c^{-1}|x|$. We write the integral $K(t, x)$ in polar coordinate $\xi = r e^{i\theta}$ as

$$K(t, x) = \frac{1}{(2\pi)^2} \int_0^{2\pi} \int_0^\infty e^{i(r(x_1 \cos \theta + x_2 \sin \theta) + t \gamma(r))} \chi_k^2(r) r dr d\theta,$$

where as before we have abused our notation by writing $\chi_k(r)$ for $\chi_k(r e^{i\theta})$. Making a change of variable in θ , we have

$$K(t, x) = \frac{1}{(2\pi)^2} \int_0^{2\pi} \int_0^\infty e^{i(r|x| \cos \theta + t \gamma(r))} \chi_k^2(r) r dr d\theta.$$

Let $\psi_1(\theta) : \mathbb{S}^1 = \mathbb{R}/2\pi\mathbb{Z} \rightarrow [0, 1]$ be a smooth 2π -periodic function on \mathbb{R} such that $\psi_1|_{[-\pi, \pi]} = 1$ on $[-\pi/4, \pi/4]$ and $\text{supp } \psi_1|_{[-\pi, \pi]} \subset [-\pi/2, \pi/2]$, and let $\psi_2(\theta)$ be the function $\psi_1(\theta + \pi)$. Let $\psi_3 = 1 - \psi_1 - \psi_2$.

We now write $K(t, r)$ as the sum $(I_1 + I_2 + I_3)/(2\pi)^2$, where

$$I_i = \int_0^{2\pi} \int_{\mathbb{R}} e^{i(r|x| \cos \theta + t \gamma(r))} \chi_k^2(r) \psi_i(\theta) r dr d\theta.$$

We estimate the three integrals separately.

Estimate of $|I_1|$ and $|I_2|$. We will only estimate $|I_1|$ since the same argument applies to $|I_2|$. Suppose $(t, x) = \lambda(t_0, x_0)$ with $\lambda = |(t, x)|$. Note that $|t_0| \sim |x_0| \sim 1$ and $\lambda \sim |t|$ by our assumption $c|x| < |t| \leq c^{-1}|x|$. We write I_1 as

$$\int_{-\pi}^{\pi} \int_{\mathbb{R}} e^{i\lambda(\tilde{r}|x_0| \cos \theta + t_0 \gamma(\tilde{r}))} \chi_k^2(\tilde{r}) \psi_1(\theta) \tilde{r} d\tilde{r} d\theta.$$

Due to the support properties of χ_k and ψ_1 , we can make the change of variable

$$\begin{cases} r = \tilde{r} \\ \theta = 2 \arcsin \left(\alpha \sqrt{1/(2r)} \right) \quad (\text{or } \alpha^2 = r(1 - \cos \theta)) \end{cases}$$

in the integral and obtain

$$|I_1| = \int_{-\sqrt{1+2^{M+3}}}^{\sqrt{1+2^{M+3}}} \int_{1-2^{M+3}}^{1+2^{M+3}} e^{-i\lambda\alpha^2} e^{i\lambda(r|x_0|+t_0\gamma(r))} \chi_k^2(r) r \psi_1 \left(2 \arcsin \left(\alpha \sqrt{1/(2r)} \right) \right) \frac{2}{\sqrt{2r - \alpha^2}} dr d\alpha.$$

Therefore by the Van der Corput lemma (see e.g. [SM93]) applied to integration in α we have

$$|I_1| \leq \frac{C}{\lambda^{1/2}} \int_{-\sqrt{1+2^{M+3}}}^{\sqrt{1+2^{M+3}}} I_{1,\alpha} d\alpha \lesssim \frac{C}{\lambda^{1/2}} \sup_{|\alpha| \leq \sqrt{1+2^{M+3}}} I_{1,\alpha}$$

where C is a universal constant and

$$I_{1,\alpha} = \left| \int_{1-2^{M+3}}^{1+2^{M+3}} e^{i\lambda(r|x_0|+t_0\gamma(r))} \chi_k^2(r) r \partial_\alpha \left(\psi_1 \left(2 \arcsin \left(\alpha \sqrt{1/(2r)} \right) \right) \frac{1}{\sqrt{2r - \alpha^2}} \right) dr \right|.$$

To estimate $I_{1,\alpha}$, noting that $|\partial_r^2(r|x_0| + t_0\gamma(r))| \gtrsim 2^{\beta k}$ on $\text{supp } \chi_k$, we apply Van der Corput lemma again to the integration in r and obtain

$$I_{1,\alpha} \lesssim \frac{C 2^{-\beta k/2}}{\lambda^{1/2}} \int_{1-2^{M+3}}^{1+2^{M+3}} \left| \partial_r \left(\chi_k^2(r) r \partial_\alpha \left(\psi_1 \left(2 \arcsin \left(\alpha \sqrt{1/(2r)} \right) \right) \frac{1}{\sqrt{2r - \alpha^2}} \right) \right) \right| dr.$$

Recalling (3.7) and support properties of $\chi_k(r)$ and $\psi_1(\theta)$ we conclude

$$\sup_{|\alpha| \leq \sqrt{1+2^{M+3}}} I_{1,\alpha} \lesssim \frac{2^{-\beta k/2}}{\lambda^{1/2}},$$

which implies

$$|I_1| \lesssim \frac{2^{-\beta k/2}}{\lambda} \lesssim \frac{2^{-\beta k/2}}{|t|}.$$

Estimate of $|I_3|$. Since in the support of $\psi_3(\theta)$ we have $|\sin \theta| \gtrsim 1$, using integration by parts we obtain

$$\begin{aligned} |I_3| &= \left| \int_0^{2\pi} \int_{\mathbb{R}} \frac{\partial_\theta (e^{ir|x|\cos \theta})}{-ir|x|\sin \theta} e^{it\gamma(r)} \chi_k^2(r) \psi_3(\theta) r dr d\theta \right| \\ &= \left| \int_0^{2\pi} \int_{\mathbb{R}} e^{ir|x|\cos \theta} \partial_\theta \left(e^{it\gamma(r)} \frac{\chi_k^2(r) \psi_3(\theta)}{i|x|\sin \theta} \right) dr d\theta \right| \\ &\leq \int_0^{2\pi} \int_{\mathbb{R}} \left| \partial_\theta \left(\frac{\psi_3(\theta)}{\sin \theta} \right) \right| \frac{\chi_k^2(r)}{|x|} dr d\theta \\ &\lesssim \frac{1}{|x|} \end{aligned}$$

Therefore when $c|x| < t \leq c^{-1}|x|$ we conclude

$$|I_3| \lesssim \frac{1}{|t|}.$$

Hence we have

$$\|K(t, x)\|_{L_x^\infty} \leq |I_1| + |I_2| + |I_3| \lesssim \frac{2^{-\beta k/2}}{|t|}$$

and our proof is complete. \square

Proof of Proposition 3. We first observe that by construction we have

$$\chi_k(D)P_k(D) = P_k(D).$$

Therefore it suffice to prove the estimate

$$(3.9) \quad \left\| e^{ith(D)} \chi_k(D) u_0 \right\|_{L_{t,x}^4} \lesssim 2^{-\beta k/8} \|u_0\|_{L_x^2}$$

for every $u_0 \in L_x^2$. By the standard TT^* argument (see for example [SM93], [Tao06]) applied to the operator $e^{ith(D)} \chi_k(D)$ we conclude that the estimate (3.9) is equivalent to the following estimate

$$\left\| \int e^{i(t-s)h(D)} \chi_k^2(D) F(s) ds \right\|_{L_{t,x}^4} \lesssim 2^{-\beta k/4} \|F\|_{L_{t,x}^{4/3}}.$$

By duality it suffices to show

$$(3.10) \quad \left| \int \int \left(\int e^{i(t-s)h(D)} \chi_k^2(D) F(s) ds \right) \overline{G(t, x)} dt dx \right| \lesssim 2^{-\beta k/4} \|F\|_{L_{t,x}^{4/3}} \|G\|_{L_{t,x}^{4/3}}.$$

We can interpolate the trivial estimate

$$\left\| e^{ith(D)} \chi_k^2(D) u_0 \right\|_{L_x^2} \lesssim \|u_0\|_{L_x^2}$$

with the dispersive estimate (3.3) to obtain

$$(3.11) \quad \left\| e^{ith(D)} \chi_k^2(D) u_0 \right\|_{L_x^4} \lesssim \frac{2^{-\beta k/4}}{|t|^{1/2}} \|u_0\|_{L_x^{4/3}}.$$

Therefore by Hölder's inequality and Plancherel' theorem we have

$$\begin{aligned} & \left| \iint \left(\int e^{i(t-s)h(D)} \chi_k^2(D) F(s) ds \right) \overline{G(t, x)} dt dx \right| \\ &= \left| \iint \left(\int e^{i(t-s)h(D)} \chi_k^2(D) F(s) \overline{G(t)} dx \right) ds dt \right| \\ &\lesssim \iint \frac{2^{-\beta k/4}}{|t-s|^{1/2}} \|F(s)\|_{L_x^{4/3}} \|G(t)\|_{L_x^{4/3}} ds dt. \end{aligned}$$

Since $\frac{1}{2} + \frac{3}{4} + \frac{3}{4} = 1 + 1$, by Hardy-Littlewood-Sobolev inequality we obtain the desired inequality (3.10). (3.2) follows from (3.1) and the triangle inequality. \square

Remark 6. Observe that the weighted restriction estimate

$$(3.12) \quad \left\| e^{ith(D)} \kappa^{1/8} P_{\leq M+3} u_0 \right\|_{L_{t,x}^4} \lesssim \|P_{\leq M+3} u_0\|_{L_x^2},$$

where

$$\kappa(\xi) = \frac{\gamma'(|\xi|)\gamma''(|\xi|)}{|\xi|(1+(\gamma'(|\xi|))^2)^2}$$

is the Gaussian curvature of the surface $\tau = h(\xi)$, will imply the localized Strichartz estimates in Proposition 3. There are some prior results in this direction. If our curve γ is convex, then the weighted restriction theorem proved in [Sha07] immediately implies (3.1). Other such weighted restriction theorems have been considered for some radial surfaces in \mathbb{R}^3 [Obe04] [CKZ07]. See also [KPV91] for some weighted restriction estimates in \mathbb{R}^n . For more general hypersurfaces in \mathbb{R}^n , see [SS85] [CDMM90]. Conversely, for a radial surface in \mathbb{R}^3 given by a curve γ satisfying our assumptions, (3.12) is an easy consequence of (3.1) and the bilinear estimate (3.13) proved later. One may also prove (3.12) directly by modifying our proof for (3.1).

By the atomic structure of U^p spaces, and embedding (2.1), we obtain the following.

Corollary 1. *For every $k \leq M + 1$, we have*

$$\|P_{k \leq \cdot \leq M+1} u\|_{L_{t,x}^4} \lesssim 2^{-k\beta/8} \|P_{k \leq \cdot \leq M+1} u\|_{U_{h(D)}^4},$$

and

$$\|P_{k \leq \cdot \leq M+1} u\|_{L_{t,x}^4} \lesssim 2^{-k\beta/8} \|P_{k \leq \cdot \leq M+1} u\|_{V_{h(D)}^2}.$$

The second part of the section is devoted to bilinear L^2 estimates, which exploit transversality in the bilinear interaction of waves, rather than dispersion. As a starting point for our analysis, we utilize the following well-known bilinear estimate. The proof given here is adapted from [Bou98].

Proposition 4. *Suppose $h(\xi), h'(\xi)$ are smooth functions on some open subset Ω of \mathbb{R}^2 and $\Omega' \Subset \Omega$. Let Ω_1, Ω_2 be two open subset of Ω' . Suppose*

$$\theta := \inf_{\xi \in \Omega_1, \eta \in \Omega_2} |\nabla h(\xi) - \nabla h'(\eta)| > 0$$

and

$$l = \sup_{(\tau_0, \xi_0) \in \mathbb{R} \times \mathbb{R}^2} \text{meas}_1 \{ (h(\xi), \xi) : \xi \in \Omega_1, \xi_0 - \xi \in \Omega_2, \tau_0 - h(\xi) = h'(\xi_0 - \xi) \} < \infty.^1$$

Then we have

$$\| (e^{ith(D)}u)(e^{ith'(D)}v) \|_{L_{t,x}^2} \leq \theta^{-1/2} l^{1/2} \|u\|_{L_x^2} \|v\|_{L_x^2}$$

for every $u, v \in L_x^2$ with $\text{supp } \hat{u} \subset \Omega_1, \text{supp } \hat{v} \subset \Omega_2$.

Proof. By Plancherel's theorem and the Cauchy-Schwartz inequality (applied to the measure $\delta_0(\tau - h'(\xi - \xi') - h(\xi'))d\xi'$), we have

$$\begin{aligned} & \| (e^{ith(D)}u)(e^{ith'(D)}v) \|_{L_{t,x}^2}^2 = \left\| \int \delta_0(\tau - h'(\xi - \xi') - h(\xi')) \hat{u}(\xi - \xi') \hat{v}(\xi') d\xi' \right\|_{L_{\tau,\xi}^2}^2 \\ & \leq \left(\int \int \int |\hat{u}(\xi - \xi')|^2 |\hat{v}(\xi')|^2 \delta_0(\tau - h'(\xi - \xi') - h(\xi')) d\xi' d\xi d\tau \right) \\ & \quad \times \sup_{\tau, \xi} \left(\int 1_{\Omega_1}(\xi') 1_{\Omega_2}(\xi - \xi') \delta_0(\tau - h'(\xi - \xi') - h(\xi')) d\xi' \right) \\ & \leq \|u\|_{L_x^2}^2 \|v\|_{L_x^2}^2 \sup_{\tau, \xi} \left(\int_{h'(\xi - \xi') + h(\xi') = \tau} 1_{\Omega_1}(\xi') 1_{\Omega_2}(\xi - \xi') \frac{d\mu(\xi')}{|\nabla h'(\xi - \xi') - \nabla h(\xi')|} \right). \end{aligned}$$

By our assumption we have

$$\sup_{\tau, \xi} \left(\int_{h'(\xi - \xi') + h(\xi') = \tau} 1_{\Omega_1}(\xi') 1_{\Omega_2}(\xi - \xi') \frac{d\mu(\xi')}{|\nabla h'(\xi - \xi') - \nabla h(\xi')|} \right) \leq \theta^{-1} l.$$

Therefore we obtain the claimed estimate. \square

We now apply the above general bilinear estimate to our problem. For simplicity of exposition we assume without loss of generality that if $k_1 \leq k_2 - 10$, then $|\gamma''(r_1)| \leq |\gamma''(r_2)|/3$ whenever $|r_1 - 1| \leq 2^{k_1+2}$ and $2^{k_2-2} \leq |r_2 - 1| \leq 2^{M+3}$. In general we choose a sufficiently large admissible integer in the place of 10.

Proposition 5. *Suppose $k_1 \leq k_2 - 10$ and $k_2 \leq M + 1$. Then we have*

$$(3.13) \quad \left\| (e^{ith(D)}P_{\leq k_1}u)(e^{ith(D)}P_{k_2}v) \right\|_{L_t^2 L_x^2} \lesssim 2^{-\beta k_2/4} 2^{(k_1 - k_2)/4} \|P_{\leq k_1}u\|_{L_x^2} \|P_{k_2}v\|_{L_x^2}.$$

As a consequence we have

$$\left\| (e^{ith(D)}P_{\leq k_1}u)(e^{ith(D)}P_{k_2 \leq \cdot \leq M+1}v) \right\|_{L_t^2 L_x^2} \lesssim 2^{-\beta k_2/4} 2^{(k_1 - k_2)/4} \|P_{\leq k_1}u\|_{L_x^2} \|P_{k_2 \leq \cdot \leq M+1}v\|_{L_x^2}.$$

Remark 7. Since taking complex conjugation preserves L^2 norms, we immediately deduce that the proposition still holds if both $e^{ith(D)}$ are replaced by $e^{-ith(D)}$. The above proposition also holds if either $e^{ith(D)}$ on the left hand side is replaced by $e^{-ith(D)}$, in which case the following argument goes through with little modification.

¹The previous condition guarantees that the set being measured, which is the intersection of the graph of h over Ω_1 and an inverse translate of the graph of h' over Ω_2 , is a 1-dimensional submanifold in \mathbb{R}^3 . So here the 1-dimensional Hausdorff measure meas_1 just means the length of the curve.

Proof. The second estimate follows immediately from the first one by the triangle inequality. For (3.13) we will only estimate $\|(e^{ith(D)}P_{\leq k_1}u)(e^{ith(D)}P_{k_2}^+v)\|_{L_t^2L_x^2}$ since other combinations can be similarly estimated, and an application of the triangle inequality will give the desired result. As before we will omit $+$ in the notation. For simplicity of notation we will also abuse our notation by writing u in place of $P_{\leq k_1}u$ and v in place of $P_{k_2}v$. \hat{u} and \hat{v} are supported on Ω_1 and Ω_2 respectively where

$$(3.14) \quad \begin{aligned} \Omega_1 &= \{\xi \in \mathbb{R}^2 : 1 - 2^{k_1+2} < |\xi| < 1 + 2^{k_1+2}\} \\ \Omega_2 &= \{\xi \in \mathbb{R}^2 : 1 + 2^{k_2-2} < |\xi| < 1 + 2^{k_2+2}\}. \end{aligned}$$

We let R_k be the Fourier multiplier given by the symbol $1_{k2^{k_1} \leq |\xi| - 1 < (k+1)2^{k_1}}(\xi)$. Then we can write

$$v = \sum_{k=2^{k_2-k_1-3}}^{2^{k_2-k_1+3}} R_k v.$$

We let Ω^k be the annulus $(k-1/3)2^{k_1} < |\xi| - 1 < (k+4/3)2^{k_1}$. By Plancherel's theorem if for every $\xi, \xi' \in \Omega_1$ and $\eta \in \Omega^k, \eta' \in \Omega^{k'}$ we have $h(\xi) + h(\eta) \neq h(\xi') + h(\eta')$ then $\langle (e^{ith(D)}u)(e^{ith(D)}R_k v), (e^{ith(D)}u)(e^{ith(D)}R_{k'} v) \rangle = 0$. By our assumption (b) on γ we conclude the following almost orthogonality relation

$$(3.15) \quad \|(e^{ith(D)}u)(e^{ith(D)}v)\|_{L^2}^2 \lesssim \sum_k \|(e^{ith(D)}u)(e^{ith(D)}R_k v)\|_{L^2}^2.$$

Now we fix $k \in [2^{k_2-k_1-3}, 2^{k_2-k_1+3}] \cap \mathbb{Z}$. To estimate terms on the right hand side of (3.15), we consider the following two cases.

Case 1 $k_1 \geq (\beta+1)k_2$.

We let $L = (\beta+1)k_2$. For $L \leq m \leq -2$ we let $Q_{m,j}$ be the Fourier multiplier given by the symbol

$$1_{j2^m 2\pi \leq \xi / |\xi| < (j+1)2^m 2\pi}(\xi),$$

where we used the identification $\mathbb{S}^1 \cong \mathbb{R}/2\pi\mathbb{Z}$. Put another way, $Q_{m,j}(\xi) = 1$ if and only if $\xi = |\xi|e^{i\theta}$ for some $\theta \in [j2^m 2\pi, (j+1)2^m 2\pi)$. Then by the triangle inequality we have

$$(3.16) \quad \begin{aligned} & \left\| (e^{ith(D)}u)(e^{ith(D)}R_k v) \right\|_{L^2} \\ & \leq \sum_{m=L}^{-2} \sum_{j=0}^{2^{-L}-1} \sum_{j': 1 < |j'-j| < \min\{6, 2^{-m}-1\}} \left\| (e^{ith(D)}Q_{m,j}u)(e^{ith(D)}Q_{m,j'}R_k v) \right\|_{L^2} \\ & \quad + \sum_{j=0}^{2^{-L}-1} \sum_{j': |j'-j| \leq 1} \left\| (e^{ith(D)}Q_{m,j}u)(e^{ith(D)}Q_{m,j'}R_k v) \right\|_{L^2} \end{aligned}$$

We define

$$U_{m,j} := \{\xi \in \Omega_1 : (j-1/3)2^m 2\pi < \xi/|\xi| < (j+4/3)2^m 2\pi\}$$

$$V_{m,j'} := \{\xi \in \Omega^k : (j'-1/3)2^m 2\pi < \xi/|\xi| < (j'+4/3)2^m 2\pi\}.$$

Then $U_{m,j}, V_{m,j'}$ contain the support of $Q_{m,j}(\xi)\hat{u}(\xi)$ and $Q_{m,j'}(\xi)R_k(\xi)\hat{v}(\xi)$ respectively.

Note that when $\xi \in \Omega_1$ and $\xi' \in \Omega^k$ we have

$$\begin{aligned} |\nabla h(\xi) - \nabla h(\xi')| &= \left| \gamma'(|\xi|) \frac{\xi}{|\xi|} - \gamma'(|\xi'|) \frac{\xi'}{|\xi'|} \right| \\ &= \left| \gamma'(|\xi|) \left(\frac{\xi}{|\xi|} - \frac{\xi'}{|\xi'|} \right) + (\gamma'(|\xi|) - \gamma'(|\xi'|)) \frac{\xi'}{|\xi'|} \right| \\ &\gtrsim \left| \frac{\xi}{|\xi|} - \frac{\xi'}{|\xi'|} \right| + 2^{(\beta+1)k_2} \end{aligned}$$

because $\gamma'(|\xi|), \gamma'(|\xi'|) \sim 1$ and

$$|\gamma'(|\xi|) - \gamma'(|\xi'|)| = \left| \int_{|\xi'|}^{|\xi|} \gamma''(r) dr \right| \sim 2^{(\beta+1)k_2}.$$

So if we let θ be the parameter in Proposition 4 for $U_{m,j}$ and $V_{m,j'}$ then when $1 < |j - j'| < 6$ and $L \leq m \leq -4$,

$$\theta \gtrsim \left| \frac{\xi}{|\xi|} - \frac{\xi'}{|\xi'|} \right| \gtrsim 2^m$$

and when $|j' - j| \leq 1$ and $m = L$,

$$\theta \gtrsim 2^{(\beta+1)k_2} = 2^L.$$

We let l be the parameter in Proposition 4 for $U_{m,j}$ and $V_{m,j'}$. Since h has bounded C^3 -norm on the annulus $||\xi| - 1| \leq 2^{M+3}$, we have

$$l \sim \sup_{\tau, \eta} \text{meas}_1 \{ \xi \in U_{m,j} : \eta - \xi \in V_{m,j'}, h(\eta - \xi) + h(\xi) = \tau \}.$$

We let $\gamma_{\tau, \eta}$ be the curve

$$\{ \xi \in U_{m,j} : \eta - \xi \in V_{m,j'}, h(\eta - \xi) + h(\xi) = \tau \}.$$

Fix $\eta \in \mathbb{R}^2$ and $\tau \in \mathbb{R}$. To estimate the length of $\gamma_{\tau, \eta}$ we examine its defining function

$$f(\xi) := h(\eta - \xi) + h(\xi).$$

Then

$$\begin{aligned} (3.17) \quad \nabla f(\xi) &= \gamma'(|\xi|) \frac{\xi}{|\xi|} - \gamma'(|\eta - \xi|) \frac{\eta - \xi}{|\eta - \xi|} \\ &= (\gamma'(|\xi|) - \gamma'(|\eta - \xi|)) \frac{\xi}{|\xi|} + \gamma'(|\eta - \xi|) \left(\frac{\xi}{|\xi|} - \frac{\eta - \xi}{|\eta - \xi|} \right). \end{aligned}$$

We write $\nabla f = \nabla_r f + \nabla_\theta f$ where

$$\nabla_r f(\xi) = \left(\nabla f \cdot \frac{\xi}{|\xi|} \right) \xi, \quad \nabla_\theta f = \nabla f - \nabla_r f$$

are the radial and tangential parts of ∇f respectively. Note that when $\xi \in U_{m,j}$ and $\eta - \xi \in V_{m,j'}$,

$$-\gamma'(|\xi|) + \gamma'(|\eta - \xi|) \sim 2^{(\beta+1)k_2}, \quad \gamma'(|\eta - \xi|) \sim 1$$

So by (3.17) and our assumption $k_1 \geq (\beta + 1)k_2$ we know that when $1 < |j - j'| < 6$

$$\frac{|\nabla_r f(\xi)|}{|\nabla_\theta f(\xi)|} \begin{cases} \lesssim 1 & \text{if } k_1/2 + (\beta + 1)k_2/2 < m \leq -4, \\ \gtrsim \frac{2^{k_2(\beta+1)}}{2^m} & \text{if } L \leq m \leq k_1/2 + (\beta + 1)k_2/2. \end{cases}$$

for every $\xi \in U_{m,j}$ satisfying $\eta - \xi \in V_{m,j'}$. So if we let l be the parameter in Proposition 4 for $U_{m,j}$ and $V_{m,j'}$, then² when $1 < |j - j'| < 6$,

$$l \lesssim \begin{cases} 2^{k_1} & \text{if } k_1/2 + (\beta + 1)k_2/2 < m \leq -4, \\ \frac{2^m}{2^{k_2(\beta+1)}} 2^m & \text{if } L \leq m \leq k_1/2 + (\beta + 1)k_2/2 \end{cases}$$

and when $|j' - j| \leq 1$ and $m = L = (\beta + 1)k_2$,

$$l \lesssim 2^L = 2^{(\beta+1)k_2}.$$

Applying Proposition 4 we obtain when $1 < |j - j'| < 6$,

$$\begin{aligned} & \left\| (e^{ith(D)} Q_{m,j} u) (e^{ith(D)} Q_{m,j'} R_k v) \right\|_{L^2} \\ & \lesssim \begin{cases} 2^{k_1/2} 2^{-m/2} \|Q_{m,j} u\|_{L^2} \|Q_{m,j'} R_k v\|_{L^2} & \text{if } k_1/2 + (\beta + 1)k_2/2 < m \leq -4, \\ 2^{m-k_2(\beta+1)/2} 2^{-m/2} \|Q_{m,j} u\|_{L^2} \|Q_{m,j'} R_k v\|_{L^2} & \text{if } L \leq m \leq k_1/2 + (\beta + 1)k_2/2 \end{cases} \\ & = \begin{cases} 2^{k_1/2} 2^{-m/2} \|Q_{m,j} u\|_{L^2} \|Q_{m,j'} R_k v\|_{L^2} & \text{if } k_1/2 + (\beta + 1)k_2/2 < m \leq -4, \\ 2^{m/2} 2^{-k_2(\beta+1)/2} \|Q_{m,j} u\|_{L^2} \|Q_{m,j'} R_k v\|_{L^2} & \text{if } L \leq m \leq k_1/2 + (\beta + 1)k_2/2 \end{cases} \end{aligned}$$

and when $|j' - j| \leq 1$ and $m = L$,

$$\begin{aligned} \left\| (e^{ith(D)} Q_{m,j} u) (e^{ith(D)} Q_{m,j'} R_k v) \right\|_{L^2} & \lesssim 2^{L/2} 2^{-L/2} \|Q_{m,j} u\|_{L^2} \|Q_{m,j'} R_k v\|_{L^2} \\ & \lesssim \|Q_{m,j} u\|_{L^2} \|Q_{m,j'} R_k v\|_{L^2} \\ & \lesssim 2^{-\beta k_2/4} 2^{(k_1 - k_2)/4} \|Q_{m,j} u\|_{L^2} \|Q_{m,j'} R_k v\|_{L^2} \end{aligned}$$

where we have used our assumption $k_1 \geq (\beta + 1)k_2$.

When $1 < |j - j'| < \min\{6, 2^{-m} - 1\}$ and $-3 \leq m \leq -2$ we can easily see that $\theta \gtrsim 1$. By Lemma 2 below we have $l \lesssim 2^{(k_1 - k_2)/2}$. So Proposition 4 gives

$$(3.18) \quad \left\| (e^{ith(D)} Q_{m,j} u) (e^{ith(D)} Q_{m,j'} R_k v) \right\|_{L^2} \lesssim 2^{(k_1 - k_2)/4} \|Q_{m,j} u\|_{L^2} \|Q_{m,j'} R_k v\|_{L^2}.$$

Therefore combining the above estimates, (3.16), and the Cauchy-Schwartz inequality yields

$$\begin{aligned} & \left\| (e^{ith(D)} u) (e^{ith(D)} R_k v) \right\|_{L^2} \\ & \lesssim \left(\sum_{k_1/2 + (\beta+1)k_2/2 < m \leq -4} 2^{k_1/2} 2^{-m/2} + \sum_{L \leq m \leq k_1/2 + (\beta+1)k_2/2} 2^{m/2} 2^{-k_2(\beta+1)/2} \right. \\ & \quad \left. + 2^{(k_1 - k_2)/4} + 2^{-\beta k_2/4} 2^{(k_1 - k_2)/4} \right) \sup_m \left(\sum_j \|Q_{m,j} u\|_{L^2}^2 \right)^{1/2} \left(\sum_{j'} \|Q_{m,j'} R_k v\|_{L^2}^2 \right)^{1/2} \\ & \lesssim \left(2^{k_1/2} 2^{-(\beta+1)k_2/4 - k_1/4} + 2^{(\beta+1)k_2/4} 2^{k_1/4} 2^{-k_2(\beta+1)/2} + 2^{-\beta k_2/4} 2^{(k_1 - k_2)/4} \right) \|u\|_{L^2} \|R_k v\|_{L^2} \\ & \lesssim 2^{-\beta k_2/4} 2^{(k_1 - k_2)/4} \|u\|_{L^2} \|R_k v\|_{L^2}. \end{aligned}$$

²To be completely rigorous we need to show that the number of components of $\gamma_{\tau,\xi}$ is bounded by some admissible constant. See the proof of Lemma 2 below.

Case 2.2 $k_1 < (\beta + 1)k_2$. We let $L' = k_1/2 + (\beta + 1)k_2/2$ (without loss of generality we may assume that k_2 is even). By the triangle inequality we have

$$(3.19) \quad \left\| (e^{ith(D)}u)(e^{ith(D)}R_kv) \right\|_{L^2} \\ \leq \sum_{m=L'}^{-2} \sum_{j=0}^{2^{-m}-1} \sum_{j': 1 < |j' - j| < \min\{6, 2^{-m}-1\}} \left\| (e^{ith(D)}Q_{m,j}u)(e^{ith(D)}Q_{m,j'}R_kv) \right\|_{L^2} \\ + \sum_{j=0}^{2^{-L'}-1} \sum_{j': |j' - j| \leq 1} \left\| (e^{ith(D)}Q_{m,j}u)(e^{ith(D)}Q_{m,j'}R_kv) \right\|_{L^2}.$$

Arguing as before and noting the assumption $k_1 < (\beta + 1)k_2$, we have when $1 < |j - j'| < 6$ and $L' \leq m \leq -4$,

$$\left\| (e^{ith(D)}Q_{m,j}u)(e^{ith(D)}Q_{m,j'}R_kv) \right\|_{L^2} \\ \lesssim \left((2^{k_1})^{1/2} (2^m)^{-1/2} + \left(\frac{2^{(\beta+1)k_2}}{2^m} 2^{k_1} \right)^{1/2} 2^{-(\beta+1)k_2/2} \right) \|Q_{m,j}u\|_{L^2} \|Q_{m,j'}R_kv\|_{L^2} \\ \lesssim 2^{k_1/2} 2^{-m/2} \|Q_{m,j}u\|_{L^2} \|Q_{m,j'}R_kv\|_{L^2},$$

and when $|j' - j| \leq 1$ and $m = L' = k_1/2 + (\beta + 1)k_2/2$,

$$\left\| (e^{ith(D)}Q_{m,j}u)(e^{ith(D)}Q_{m,j'}R_kv) \right\|_{L^2} \\ \lesssim \left(2^{(\beta+1)k_2/2} 2^{k_1/2} \right)^{1/2} \left(2^{(\beta+1)k_2} \right)^{-1/2} \|Q_{m,j}u\|_{L^2} \|Q_{m,j'}R_kv\|_{L^2} \\ \lesssim 2^{-\beta k_2/4} 2^{(k_1 - k_2)/4} \|Q_{m,j}u\|_{L^2} \|Q_{m,j'}R_kv\|_{L^2}.$$

Therefore combining the above estimates, (3.18) and the Cauchy-Schwarz inequality, we deduce from (3.19) that

$$\left\| (e^{ith(D)}u)(e^{ith(D)}R_kv) \right\|_{L^2} \\ \lesssim \left(2^{(k_1 - k_2)/4} + 2^{-\beta k_2/4} 2^{(k_1 - k_2)/4} + \sum_{L' \leq m \leq -4} 2^{k_1/2} 2^{-m/2} \right) \|u\|_{L^2} \|R_kv\|_{L^2} \\ \lesssim \left(2^{k_2/2} 2^{-(\beta+1)k_2/4 - k_1/4} + 2^{-\beta k_2/4} 2^{(k_1 - k_2)/4} \right) \|u\|_{L^2} \|R_kv\|_{L^2} \\ \lesssim 2^{-\beta k_2/4} 2^{(k_1 - k_2)/4} \|u\|_{L^2} \|R_kv\|_{L^2}.$$

Combining Case 1 and Case 2 we therefore conclude that

$$\left\| (e^{ith(D)}u)(e^{ith(D)}R_kv) \right\|_{L^2} \lesssim 2^{-\beta k_2/4} 2^{(k_1 - k_2)/4} \|u\|_{L^2} \|R_kv\|_{L^2}$$

for every k . Now the almost orthogonality relation (3.15) implies

$$\left\| (e^{ith(D)}u)(e^{ith(D)}v) \right\|_{L^2} \lesssim 2^{-\beta k_2/4} 2^{(k_1 - k_2)/4} \|u\|_{L^2} \|v\|_{L^2},$$

as desired. \square

Lemma 2. *Suppose $k_1 \leq k_2 - 10$ and $k_2 \leq M + 1$. Let Ω_1, Ω_2 be the annuli defined by (3.14). For every $\tau_0 \in \mathbb{R}, \xi \in \mathbb{R}^2$ define*

$$\gamma_{\tau_0, \xi_0} := \{\xi : \xi \in \Omega_1, \xi_0 - \xi \in \Omega_2, \tau_0 - h(\xi) = h(\xi_0 - \xi)\}.$$

Then

$$\Theta := \{\theta \in [0, 2\pi] : \text{there exists } r \text{ such that } re^{i\theta} \in \gamma_{\tau_0, \xi_0}\}$$

is a disjoint union of at most c many intervals, each of which has size less than $C2^{(k_1 - k_2)/2}$ for some admissible constants c, C .

Proof. We fix ξ_0, τ_0 . Without loss of generality we may suppose that in polar coordinates $\xi_0 = r_0 e^{i\theta_0} = r_0$. Let Ω be the set

$$\Omega := \{\xi : \xi \in \Omega_1, \xi_0 - \xi \in \Omega_2\}.$$

and let Ω' be the set

$$\Omega' := \{re^{i\theta} \in \Omega : \theta \neq 0, \pi\}.$$

So using the above notation we have

$$\gamma_{\tau_0, \xi_0} = \{\xi \in \Omega : \tau_0 - h(\xi) = h(\xi_0 - \xi)\}.$$

Writing down the inequalities defining annuli Ω_1 and $\xi_0 - \Omega_2$ we see that, depending on r_0 , Ω' can have at most two components, which we will denote by U' and U'' . They satisfy

$$U' = \{re^{i\theta} \in \Omega : 0 < \theta < \pi\}, \quad U'' = \{re^{i\theta} \in \Omega : -\pi < \theta < 0\}.$$

In the case when there is only one component, we put the other one to be the empty set. If Ω' has no component, we put both to be the empty set.

We first observe that U' satisfies

$$(3.20) \quad U'_\theta := \{r : re^{i\theta} \in U'\}, \quad U'_r := \{\theta : re^{i\theta} \in U'\}$$

are intervals, for every θ, r . U'' has the same properties (3.20) with U' replaced by U'' . From now on we will only consider the part of γ_{τ_0, ξ_0} that lies in U' , and the same argument will apply to the part lying in U'' .

Let $f(r, \theta)$ be the defining function of γ_{τ_0, ξ_0}

$$f(r, \theta) := \gamma(|re^{i\theta} - r_0|) + \gamma(r).$$

Noting that

$$|re^{i\theta} - r_0| = \sqrt{r^2 + r_0^2 - 2rr_0 \cos \theta},$$

we differentiate f with respect to r, θ respectively and obtain

$$(3.21) \quad \partial_r f(r, \theta) = \gamma'(|re^{i\theta} - r_0|) \frac{r - r_0 \cos \theta}{|re^{i\theta} - r_0|} + \gamma'(r),$$

$$(3.22) \quad \partial_\theta f(r, \theta) = \gamma'(|re^{i\theta} - r_0|) \frac{rr_0 \sin \theta}{|re^{i\theta} - r_0|}.$$

Recall that under the assumption $\xi = re^{i\theta} \in \Omega$, we always have

$$(3.23) \quad \gamma'(r), \gamma'(|re^{i\theta} - r_0|) \sim 1.$$

Also note that in order for γ_{τ_0, ξ_0} to be nonempty, we must have

$$(3.24) \quad 2^{k_2} \lesssim r_0 \lesssim 1.$$

When $re^{i\theta} \in U'$, we have

$$\partial_\theta f(r, \theta) > 0.$$

Therefore due to property (3.20), for every $r \in [1 - 2^{k_1+2}, 1 + 2^{k_1+2}]$, there exists at most one $\theta \in (0, \pi)$ such that $re^{i\theta} \in U'$ and $f(r, \theta) = \tau_0$. So we can parametrize

$$\gamma_1 := \gamma_{\tau_0, \xi_0} \cap U'$$

by $\theta = \theta(r)$. By the implicit function theorem we know that $\theta(r)$ is a smooth function defined on a union of disjoint intervals $I := \bigcup_i I_i$ in $(1 - 2^{k_1+2}, 1 + 2^{k_1+2})$, and

$$(3.25) \quad \theta'(r) = \frac{\gamma'(|re^{i\theta} - r_0|)(r - r_0 \cos \theta) + \gamma'(r)|re^{i\theta} - r_0|}{\gamma'(|re^{i\theta} - r_0|)rr_0 \sin \theta}.$$

We write $I_i = (a_i, b_i)$ where $1 - 2^{k_1+2} \leq a_i < b_i \leq 1 + 2^{k_1+2}$.

We fix i in the index set J of I_i and $r_i \in I_i$. By (3.25) we have

$$(\cos \theta)'(r) = -\frac{\gamma'(|re^{i\theta} - r_0|)(r - r_0 \cos \theta) + \gamma'(r)|re^{i\theta} - r_0|}{rr_0 \gamma'(|re^{i\theta} - r_0|)},$$

which implies for $r \in (a_i, b_i)$

$$\cos \theta(r) = \cos \theta(r_i) - \int_{r_i}^r \frac{\gamma'(|re^{i\theta} - r_0|)(r - r_0 \cos \theta) + \gamma'(r)|re^{i\theta} - r_0|}{rr_0 \gamma'(|re^{i\theta} - r_0|)} dr.$$

Since for every $r \in (a_i, b_i)$

$$\left| \frac{\gamma'(|re^{i\theta} - r_0|)(r - r_0 \cos \theta) + \gamma'(r)|re^{i\theta} - r_0|}{rr_0 \gamma'(|re^{i\theta} - r_0|)} \right| \lesssim 2^{-k_2}$$

and $b_i - a_i \lesssim 2^{k_1}$, we therefore conclude that for every $r \in (a_i, b_i)$

$$|\theta(r) - \theta(r_i)| \lesssim 2^{(k_1 - k_2)/2}.$$

To finish the proof we are left to show that the cardinality of the index set J is bounded by some admissible constant c . Applying implicit function theorem to $f(r, \theta)$ on $\Omega' + B(0, 2^{k_1})$, the 2^{k_1} neighborhood of Ω' , we see that for every $j \in J$, $a_j e^{i\theta(a_j)}, b_j e^{i\theta(b_j)}$ are contained in $\partial\Omega'$, and for distinct $j, k \in J$, $\{a_j e^{i\theta(a_j)}, b_j e^{i\theta(b_j)}\} \neq \{a_k e^{i\theta(a_k)}, b_k e^{i\theta(b_k)}\}$. So it suffices to show that

$$K := \{re^{i\theta} \in \partial\Omega' : \tau_0 - \gamma(r) = \gamma(|r_0 - re^{i\theta}|)\}$$

has cardinality no more than c for some admissible constant c . Note that $\partial\Omega'$ is a subset of $\partial\Omega_1 \cup (r_0 + \partial\Omega_2) \cup \{re^{i\theta} \in \overline{\Omega_1} : \theta = 0, \pi\}$. Previous argument has already shown that

$$|K \cap \partial\Omega_1| \leq 2.$$

Similarly we have³

$$|K \cap (r_0 + \partial\Omega_2)| \leq 2.$$

Setting $\theta = 0$ in (3.21) yields

$$\partial_r f(r, 0) = \gamma'(|r - r_0|) \frac{r - r_0}{|r - r_0|} + \gamma'(r).$$

Therefore for $re^{i\theta} \in \partial\Omega'$

$$\partial_{rr} f(r, 0) = \gamma''(r) + \gamma''(|r_0 - r|)$$

³One way to see this is to repeat the previous argument in polar coordinates centered at ξ_0 .

is either always positive or always negative, as $|\gamma''(|r_0 - r|)| \geq |\gamma''(r)|/3$. Noting that $\{r : re^{i\theta} \in \partial\Omega'\}$ is an interval, we therefore conclude that

$$|K \cap \{re^{i\theta} \in \overline{\Omega}_1 : \theta = 0\}| \leq 2.$$

Similarly we have

$$|K \cap \{re^{i\theta} \in \overline{\Omega}_1 : \theta = \pi\}| \leq 2.$$

Therefore $|K| \leq 8$ and the proof is complete. \square

Using the extension property of U^p spaces (Proposition 2.16 in [HHK09]), we obtain

Corollary 2. *Suppose $k_1 \leq k_2 - 10$ and $k_2 \leq M + 1$. Then we have*

$$(3.26) \quad \|P_{\leq k_1} u P_{k_2 \leq \cdot \leq M+1} v\|_{L_{t,x}^2} \lesssim 2^{-\beta k_2/4} 2^{(k_1 - k_2)/4} \|P_{\leq k_1} u\|_{U_{h(D)}^2} \|P_{k_2 \leq \cdot \leq M+1} v\|_{U_{h(D)}^2}.$$

The right hand side of (3.26) is evaluated in U^2 -based spaces, which, as we shall see later, is insufficient for our fixed point argument. However, we can use the following interpolation result from [HHK09] to transfer from U^p -based estimates to V^p -based estimates, provided we have an additional U^q -based estimate for some $q > p$. Another way of transferring from U^p to V^p can be found in [CH16], which is not easy to implement here.

Proposition 6. *Suppose $1 < p < q$. Let E be a Banach space and $T : U^q \rightarrow E$ be a linear bounded operator with operator norm C_q . Suppose in addition there exists $C_p \in (0, C_q]$ such that the estimate $\|Tu\|_E \leq C_p \|u\|_{U^p}$ holds for all $u \in U^p$. Then we have for every $u \in V_{rc}^p$*

$$\|Tu\|_E \leq C_{p,q} C_p \left(\log \frac{C_q}{C_p} + 1 \right) \|u\|_{V^p},$$

where $C_{p,q}$ is a constant depending only on p, q .

In our case we already have a U^2 -based estimate in Corollary 2. We can also obtain a U^4 -based estimate from Proposition 3. Indeed, under the same condition as in Corollary 2, by Hölder's inequality we have

$$\begin{aligned} \|P_{k_1} u P_{k_2 \leq \cdot \leq M+1} v\|_{L_{t,x}^2} &\lesssim \|P_{k_1} u\|_{L_{t,x}^4} \|P_{k_2 \leq \cdot \leq M+1} v\|_{L_{t,x}^4} \\ &\lesssim 2^{-\beta(k_1+k_2)/8} \|P_{k_1} u\|_{U_{h(D)}^4} \|P_{k_2 \leq \cdot \leq M+1} v\|_{U_{h(D)}^4}. \end{aligned}$$

Therefore applying Proposition 6 twice⁴, and noticing the assumption that $k_1 \leq k_2 - 10$, we obtain

$$\|P_{k_1} u P_{k_2 \leq \cdot \leq M+1} v\|_{L_{t,x}^2} \lesssim 2^{-\beta k_2/4} 2^{(k_1 - k_2)/4} |k_2 - k_1|^2 \|P_{k_1} u\|_{V_{h(D)}^2} \|P_{k_2 \leq \cdot \leq M+1} v\|_{V_{h(D)}^2}$$

if $P_{k_1} u, P_{k_2 \leq \cdot \leq M+1} v \in V_{h(D),rc}^2$. Summing the above inequality over k_1 and using the triangle inequality we obtain the following corollary:

Corollary 3. *Suppose $k_1 \leq k_2 - 10$ and $k_2 \leq M + 1$. If $P_{\leq k_1} u, P_{k_2 \leq \cdot \leq M+1} v \in V_{rc,h(D)}^2$, then*

$$(3.27) \quad \|P_{\leq k_1} u P_{k_2 \leq \cdot \leq M+1} v\|_{L_{t,x}^2} \lesssim 2^{-\beta k_2/4} \|P_{\leq k_1} u\|_{V_{h(D)}^2} \|P_{k_2 \leq \cdot \leq M+1} v\|_{V_{h(D)}^2}$$

⁴In fact we apply Proposition 6 twice to obtain the estimate for enlarged projections:

$$\|P_{k_1-2 \leq \cdot \leq k_1+2} u P_{k_2-2 \leq \cdot \leq M+3} v\|_{L_{t,x}^2} \lesssim 2^{-\beta k_2/4} 2^{(k_1 - k_2)/4} |k_2 - k_1|^2 \|u\|_{V_{h(D)}^2} \|v\|_{V_{h(D)}^2}.$$

Then we substitute u and v by $P_{k_1} u$ and $P_{k_2 \leq \cdot \leq M+1} v$ respectively.

The localized linear and bilinear estimates discussed above allow us to obtain the following crucial estimate on the nonlinear term, whose proof will occupy the next section. We recall that I_T is the operator given by

$$I_T(f) = \int_0^t 1_{[0,T)}(s) e^{i(t-s)h(D)} f(s) ds.$$

Proposition 7. *For $T \in [0, \infty]$ we have*

$$(3.28) \quad \left\| I_T \left(A(D) \left(P_{\leq M} u_1 \overline{P_{\leq M} u_2} P_{\leq M} u_3 \right) \right) \right\|_{X^0} \lesssim \|u_1\|_{X^0} \|u_2\|_{X^0} \|u_3\|_{X^0}.$$

4. PROOF OF PROPOSITION 7

If k_1, k_2, k_3 are three nonpositive integers, then we let $\tilde{k}_1, \tilde{k}_2, \tilde{k}_3$ be the increasing rearrangement of k_1, k_2, k_3 , that is, $\tilde{k}_1 \geq \tilde{k}_2 \geq \tilde{k}_3$. Observe that when $u_1, u_2, u_3 \in X^0$, we have

$$A(D) \left(P_{\leq M} u_1(s) \overline{P_{\leq M} u_2(s)} P_{\leq M} u_3(s) \right) \in L_{t,loc}^1 L_x^2$$

because of Hölder's inequality, the embedding $X^0 \subset L_t^\infty L_x^2$, and Bernstein's inequality. Therefore by Proposition 2 we have

$$\begin{aligned} & \left\| I_T \left(A(D) \left(P_{\leq M} u_1 \overline{P_{\leq M} u_2} P_{\leq M} u_3 \right) \right) \right\|_{X^0} \\ &= \sup_{\|v\|_{Y^0} \leq 1} \int_0^T \int_{\mathbb{R}^2} \left| (A(D) \left(P_{\leq M} u_1 \overline{P_{\leq M} u_2} P_{\leq M} u_3 \right)) \bar{v} dx dt \right|. \end{aligned}$$

We fix $v \in Y^0$ with $\|v\|_{Y^0} \leq 1$. By Plancherel's theorem we can move the Fourier multiplier $A(D)$ to act on v . Therefore we have

$$(4.1) \quad \begin{aligned} & \left| \int_0^T \int_{\mathbb{R}^2} (A(D) \left(P_{\leq M} u_1 \overline{P_{\leq M} u_2} P_{\leq M} u_3 \right)) \bar{v} dx dt \right| \\ &= \left| \int_0^T \int_{\mathbb{R}^2} (P_{\leq M} u_1 \overline{P_{\leq M} u_2} P_{\leq M} u_3) \overline{A(D)v} dx dt \right| \\ &\leq \sum_{k \leq M} \left| \int_0^T \int_{\mathbb{R}^2} (P_{\leq M} u_1 \overline{P_{\leq M} u_2} P_{\leq M} u_3) \overline{A(D)P_k v} dx dt \right|. \end{aligned}$$

Above we have used the support property of $A(\xi)$ to restrict the sum over $k \in \mathbb{Z}$ to $k \leq M$. To shorten notation, we let

$$f_{k_1, k_2, k_3, k} := P_{k_1} u_1 \overline{P_{k_2} u_2} P_{k_3} u_3 \overline{A(D)P_k v}.$$

We similarly define $f_{k_1 \leq M; k_2 \leq M; k_3 \leq M; k}$ and others in the obvious way.

We can split the last sum in (4.1) into four parts I_1, I_2, I_3, I_4 , where

$$\begin{aligned} I_1 &= \sum_{k \leq M} \left| \int_0^T \int_{\mathbb{R}^2} \sum_{k-10 \leq \tilde{k}_1, \tilde{k}_2, \tilde{k}_3 \leq M} f_{k_1, k_2, k_3, k} dx dt \right| \\ I_2 &= \sum_{k \leq M} \left| \int_0^T \int_{\mathbb{R}^2} \sum_{k-10 \leq \tilde{k}_1, \tilde{k}_2 \leq M; \tilde{k}_3 < k-10} f_{k_1, k_2, k_3, k} dx dt \right| \\ I_3 &= \sum_{k \leq M} \left| \int_0^T \int_{\mathbb{R}^2} \sum_{k-10 \leq \tilde{k}_1 \leq M; \tilde{k}_2, \tilde{k}_3 < k-10} f_{k_1, k_2, k_3, k} dx dt \right| \\ I_4 &= \sum_{k \leq M} \left| \int_0^T \int_{\mathbb{R}^2} \sum_{\tilde{k}_1, \tilde{k}_2, \tilde{k}_3 < k-10} f_{k_1, k_2, k_3, k} dx dt \right|. \end{aligned}$$

We estimate the four sums separately. Our general strategy will be to use Corollary 1 and Corollary 3 to establish the estimates

$$I_i \leq \|u_1\|_{X^0} \|u_2\|_{X^0} \|u_3\|_{X^0}$$

for $i = 1, 2, 3, 4$, and therefore conclude Proposition 7.

Estimate of I_1 :

First we consider the near diagonal sum, that is, the sum over $k_1, k_2, k_3 \in [k-10, k+10]$. In this case we use Hölder's inequality and Corollary 1 four times to conclude that

$$\begin{aligned} \left| \int_0^T \int_{\mathbb{R}^2} f_{k_1, k_2, k_3, k} dx dt \right| &= \left| \int_0^T \int_{\mathbb{R}^2} P_{k_1} u_1 \overline{P_{k_2} u_2} P_{k_3} u_3 \overline{P_k A(D) v} dx dt \right| \\ &\lesssim \|P_{k_1} u_1\|_{L^4_{t,x}} \|P_{k_2} u_2\|_{L^4_{t,x}} \|P_{k_3} u_3\|_{L^4_{t,x}} \|P_k A(D) v\|_{L^4_{t,x}} \\ &\lesssim 2^{-4k\beta/8} 2^{\beta k/2} \|P_{k_1} u_1\|_{V^2_{h(D)}} \|P_{k_2} u_2\|_{V^2_{h(D)}} \|P_{k_3} u_3\|_{V^2_{h(D)}} \|P_k v\|_{V^2_{h(D)}} \\ &\lesssim \|P_{k_1} u_1\|_{V^2_{h(D)}} \|P_{k_2} u_2\|_{V^2_{h(D)}} \|P_{k_3} u_3\|_{V^2_{h(D)}} \|P_k v\|_{V^2_{h(D)}}. \end{aligned}$$

We then need to estimate the sum over the near diagonal region $k_1, k_2, k_3 \in [k-10, k+10]$. Using the Cauchy-Schwartz inequality and the embedding $U^2 \subset V^2_{rc}$, we have

$$\begin{aligned} &\sum_k \sum_{k_1, k_2, k_3 = k-10}^{k+9} \left| \int_0^T \int_{\mathbb{R}^2} f_{k_1, k_2, k_3, k} dx dt \right| \\ &\lesssim \left(\sum_k \|P_k u_1\|_{U^2_{h(D)}}^2 \right)^{1/2} \left(\sum_k \|P_k u_2\|_{U^2_{h(D)}}^2 \right)^{1/2} \left(\sum_k \|P_k u_3\|_{U^2_{h(D)}}^2 \right)^{1/2} \left(\sum_k \|P_k v\|_{V^2_{h(D)}}^2 \right)^{1/2} \\ &\lesssim \|u_1\|_{X^0} \|u_2\|_{X^0} \|u_3\|_{X^0} \|v\|_{Y^0} \\ &\lesssim \|u_1\|_{X^0} \|u_2\|_{X^0} \|u_3\|_{X^0}. \end{aligned}$$

Now we consider the off-diagonal sum, that is, the sum over $\tilde{k}_1 \geq k+10$. We will only consider the sum over $k_1 = \tilde{k}_1 \geq k+10$, since the remaining sum can be split into finitely many similar cases and be similarly treated. Applying Hölder's inequality, Corollary 3 once

and Corollary 1 twice, we obtain when $k + 10 \leq k_1 \leq M$

$$\begin{aligned}
& \left| \int_0^T \int_{\mathbb{R}^2} f_{k_1; k-10 \leq k_2, k_3 \leq M; k} dx dt \right| \\
&= \left| \int_0^T \int_{\mathbb{R}^2} P_{k_1} u_1 \overline{P_{k-10 \leq k_2 \leq M} u_2} P_{k-10 \leq k_3 \leq M} u_3 \overline{P_k A(D) v} dx dt \right| \\
&\lesssim \|P_{k_1} u_1 \overline{P_k A(D) v}\|_{L_{t,x}^2} \|P_{k-10 \leq k_2 \leq M} u_2\|_{L_{t,x}^4} \|P_{k-10 \leq k_3 \leq M} u_3\|_{L_{t,x}^4} \\
&\lesssim 2^{\beta k/2} 2^{-\beta k_1/4} 2^{-2\beta k/8} \|P_{k_1} u_1\|_{V_{h(D)}^2} \|P_k v\|_{V_{h(D)}^2} \|u_2\|_{V_{h(D)}^2} \|u_3\|_{V_{h(D)}^2} \\
&\lesssim 2^{(k-k_1)\beta/4} \|P_{k_1} u_1\|_{U_{h(D)}^2} \|P_k v\|_{V_{h(D)}^2} \|u_2\|_{X^0} \|u_3\|_{X^0}.
\end{aligned}$$

Here we have also used the embedding (2.5). Therefore by the Cauchy-Schwartz inequality we conclude that

$$\begin{aligned}
& \sum_{k \leq M} \sum_{k+10 \leq k_1 \leq M} \left| \int_0^T \int_{\mathbb{R}^2} f_{k_1; k-10 \leq k_2, k_3 \leq M; k} dx dt \right| \\
&\lesssim \|u_2\|_{X^0} \|u_3\|_{X^0} \left(\sum_{k \leq M} \sum_{k+10 \leq k_1 \leq M} 2^{\beta(k-k_1)/4} \|P_{k_1} u_1\|_{U_{h(D)}^2}^2 \right)^{1/2} \\
&\quad \times \left(\sum_{k \leq M} \sum_{k+10 \leq k_1 \leq M} 2^{\beta(k-k_1)/4} \|P_k v\|_{V_{h(D)}^2}^2 \right)^{1/2} \\
&\lesssim \|u_1\|_{X^0} \|u_2\|_{X^0} \|u_3\|_{X^0} \|v\|_{Y^0} \\
&\lesssim \|u_1\|_{X^0} \|u_2\|_{X^0} \|u_3\|_{X^0}.
\end{aligned}$$

Estimate of I_2 . We will only consider the sum over $k_3 = \tilde{k}_3 < k - 10$. Applying Hölder's inequality, Corollary 1 twice and Corollary 3 once, we obtain when $k_3 < k - 10$,

$$\begin{aligned}
& \left| \int_0^T \int_{\mathbb{R}^2} f_{k-10 \leq k_1, k_2 \leq M; k_3; k} dx dt \right| \\
&\lesssim \|P_{k_3} u_3 P_k A(D) v\|_{L_{t,x}^2} \|P_{k-10 \leq k_1 \leq M} u_1\|_{L_{t,x}^4} \|P_{k-10 \leq k_2 \leq M} u_2\|_{L_{t,x}^4} \\
&\lesssim 2^{-\beta k_3/4} 2^{-2\beta k/8} 2^{\beta k/2} \|u_1\|_{X^0} \|u_2\|_{X^0} \|P_{k_3} u_3\|_{V_{h(D)}^2} \|P_k v\|_{V_{h(D)}^2} \\
&\lesssim 2^{\beta(k-k_3)/4} \|P_{k_3} u_3\|_{U_{h(D)}^2} \|P_k v\|_{V_{h(D)}^2} \|u_1\|_{X^0} \|u_2\|_{X^0},
\end{aligned}$$

Here we have again used the embedding (2.5). Now we sum the above estimate over $k \leq M, k_3 < k - 10$ using the Cauchy-Schwartz inequality as before to conclude that

$$I_2 \lesssim \|u_1\|_{X^0} \|u_2\|_{X^0} \|u_3\|_{X^0}.$$

Estimate of I_3 . We will only consider the sum over $k_2 = \tilde{k}_2 < k - 10$, $k_3 = \tilde{k}_3 < k - 10$. Applying Hölder's inequality and Corollary 1, 3, we obtain when $k - 10 \leq k_1 \leq M$,

$$\begin{aligned} & \left| \int_0^T \int_{\mathbb{R}^2} f_{k_1; k_2 < k-10; k_3 < k-10; k} dx dt \right| \\ & \lesssim \left(\|P_{k_1} u_1 \overline{P_{<k-20} u_2}\|_{L_{t,x}^2} + \|P_{k_1} u_1 \overline{P_{k-20 \leq \cdot < k-10} u_2}\|_{L_{t,x}^2} \right) \|P_{<k-10} u_3 \overline{P_k A(D)v}\|_{L_{t,x}^2} \\ & \lesssim \left(2^{-\beta(k+k_1)/4} + 2^{-\beta(k_1+3k)/8} \right) 2^{\beta k/2} \|u_2\|_{X^0} \|u_3\|_{X^0} \|P_{k_1} u_1\|_{V_{h(D)}^2} \|P_k v\|_{V_{h(D)}^2} \\ & \lesssim 2^{\beta(k-k_1)/8} \|P_{k_1} u_1\|_{U_{h(D)}^2} \|P_k v\|_{V_{h(D)}^2} \|u_2\|_{X^0} \|u_3\|_{X^0}. \end{aligned}$$

We sum the above estimate over $k - 10 \leq k_1 \leq M$, and use the Cauchy-Schwartz inequality as before to conclude

$$\sum_{k \leq M} \sum_{k-10 \leq k_1 \leq M} \left| \int_0^T \int_{\mathbb{R}^2} f_{k_1; k_2 < k-10; k_3 < k-10; k} dx dt \right| \leq \|u_1\|_{X^0} \|u_2\|_{X^0} \|u_3\|_{X^0}.$$

Hence we obtain

$$I_3 \lesssim \|u_1\|_{X^0} \|u_2\|_{X^0} \|u_3\|_{X^0}.$$

Estimate of I_4 . Without loss of generality we only consider the sum over $k_1 \leq k_2 \leq k_3 < k - 10$. We let $N > 0$ be an admissible integer which will be chosen later. We let T_i be the Fourier multiplier with symbol $1_{2\pi i/N \leq \xi/|\xi| < 2\pi(i+1)/N}(\xi)$ where we again use the identification $\mathbb{S}^1 \cong \mathbb{R}/2\pi\mathbb{Z}$. Then

$$\int_0^T \int_{\mathbb{R}^2} f_{k_1, k_2, k_3, k} dx dt = \sum_{i_1, i_2, i_3, i=0}^{N-1} \int_0^T \int_{\mathbb{R}^2} (T_{i_1} P_{k_1} u_1) (\overline{T_{i_2} P_{k_2} u_2}) (T_{i_3} P_{k_3} u_3) (\overline{T_i P_k A(D)v}) dx dt.$$

We let $O_{i,k}$ be the open set

$$O_{i,k} = \{ \xi \in \mathbb{R}^2 : 2\pi(i-1/3)/N < \xi/|\xi| < 2\pi(i+4/3)/N, \quad 2^{k-2} < ||\xi| - 1| < 2^{k+2} \}$$

By Plancherel's theorem, in order for the integral

$$\int_0^T \int_{\mathbb{R}^2} (T_{i_1} P_{k_1} u_1) (\overline{T_{i_2} P_{k_2} u_2}) (T_{i_3} P_{k_3} u_3) (\overline{T_i P_k A(D)v}) dx dt$$

not to vanish, we must have

$$(4.2) \quad \{ \xi_1 + \xi_3 : \xi_1 \in O_{i_1, k_1}, \xi_3 \in O_{i_3, k_3} \} \cap \{ \xi_2 + \xi : \xi_2 \in O_{i_2, k_2}, \xi \in O_{i, k} \} \neq \emptyset.$$

When $k_1, k_2, k_3 < k - 10$, (4.2) implies that there exists an admissible integer $N > 0$ satisfying $N \sim 2^{k/2}$ and an admissible constant $d > 0$ such that

$$(4.3) \quad \int_0^T \int_{\mathbb{R}^2} f_{k_1, k_2, k_3, k} dx dt = \sum_{i_2, i=0}^{N-1} \left(\sum_{|i_1 - i_2| \leq d} \sum_{|i_3 - i| \leq d} + \sum_{|i_1 - i| \leq d} \sum_{|i_3 - i_2| \leq d} \right) \int_0^T \int_{\mathbb{R}^2} (T_{i_1} P_{k_1} u_1) (\overline{T_{i_2} P_{k_2} u_2}) (T_{i_3} P_{k_3} u_3) (\overline{T_i P_k A(D)v}) dx dt.$$

In particular (4.2) implies that if we let θ_1, θ_2 be the angles between ξ_1, ξ_3 and ξ_2, ξ respectively, then

$$2(\cos \theta_1 - \cos \theta_2) |\xi_1| |\xi_3| = |\xi_1| + |\xi_3| - |\xi_2| - |\xi| - 2 \cos \theta_2 (|\xi_1| |\xi_3| - |\xi_2| |\xi|).$$

The above constraint implies that if we choose N large enough (still satisfying $N \sim 2^{k/2}$) then

$$\int_0^T \int_{\mathbb{R}^2} (T_{i_1} P_{k_1} u_1) \overline{(T_{i_2} P_{k_2} u_2)} (T_{i_3} P_{k_3} u_3) \overline{(T_i P_k A(D) v)} dx dt = 0$$

if $|i_1 - i_3| \leq 2$, $N/2 - 2 \leq |i_1 - i_2| \leq N/2 + 2$, $N/2 - 2 \leq |i_3 - i_2| \leq N/2 + 2$ all hold.

We let l, θ be the parameters in Proposition 4 for $O_{i', k'}, O_{i'', k''}$ with $k' \leq k'', k' < k - 10$ and $2 < |i' - i''| \leq d$. Then the same argument in the proof of Proposition 5 combined with Lemma 2 shows that

$$l \lesssim 2^{k'} + 2^{(k' - k'')/2} \lesssim 2^{(k' - k'')/2}, \quad \theta \gtrsim 2^{k/2}.$$

So Proposition (4) implies that

$$\|(e^{ith(D)} T_{i'} P_{k'} u)(e^{ith(D)} T_{i''} P_{k''} v)\|_{L_{t,x}^2} \lesssim 2^{(k' - k'')/4} 2^{-k/4} \|T_{i'} P_{k'} u\|_{L_x^2} \|T_{i''} P_{k''} v\|_{L_x^2}$$

when $2 < |i' - i''| \leq d$. Similarly we have

$$\|(e^{ith(D)} T_{i'} P_{k'} u) \overline{(e^{ith(D)} T_{i''} P_{k''} v)}\|_{L_{t,x}^2} \lesssim 2^{(k' - k'')/4} 2^{-k/4} \|T_{i'} P_{k'} u\|_{L_x^2} \|T_{i''} P_{k''} v\|_{L_x^2}.$$

when $-N/2 + 2 < |i' - i''| < N/2 - 2$. In particular $2^{(k' - k'')/4} 2^{-k/4} < 2^{(k' - k)/4} 2^{-\beta k/4}$. Therefore arguing as in the proof of (5) and the argument deriving Corollary 3, we conclude from the above analysis that when $k_1 \leq k_2 \leq k_3 < k - 10$,

$$\begin{aligned} & \left| \int_0^T \int_{\mathbb{R}^2} (P_{k_1} u_1) \overline{(P_{k_2} u_2)} (P_{k_3} u_3) \overline{(P_k A(D) v)} dx dt \right| \\ & \lesssim \left(2^{(k_1 - k)/8} 2^{(k_2 - k_3)/8} + 2^{(k_2 - k)/8} 2^{(k_1 - k_3)/8} + 2^{(k_3 - k)/8} 2^{(k_1 - k_2)/8} \right) \\ & \quad \times 2^{-\beta k/2} 2^{\beta k/2} \|P_{k_1} u_1\|_{V_{h(D)}^2} \|P_{k_2} u_2\|_{V_{h(D)}^2} \|P_{k_3} u_3\|_{V_{h(D)}^2} \|P_k v\|_{V_{h(D)}^2}. \end{aligned}$$

Summing the above estimate over $k_1 \leq k_2 \leq k_3 < k - 10$ and using the Cauchy-Schwartz inequality yield

$$I_4 \lesssim \|u_1\|_{X^0} \|u_2\|_{X^0} \|u_3\|_{X^0}.$$

5. PROOF OF THE MAIN THEOREM

We recall that I_T is the linear operator given by

$$I_T(f)(t, x) = \int_0^t 1_{[0, T)}(s) e^{i(t-s)h(D)} f(s) ds$$

where $T \in [0, \infty]$. By Duhamel's formula we know that $u \in C^0([0, T], L_x^2)$ is a strong solution to the Cauchy problem (1.1) if and only if it satisfies the following integral equation

$$u(t) = e^{ith(D)} u_0 + i \int_0^t e^{i(t-s)h(D)} A(D) (|P_{\leq M} u(s)|^2 P_{\leq M} u(s)) ds$$

on $[0, T)$, which is implied by

$$(5.1) \quad u(t) = 1_{[0, T)}(t) e^{ith(D)} u_0 + 1_{[0, T)}(t) I_T (A(D) (|P_{\leq M} u|^2 P_{\leq M} u)).$$

The identity (5.1) also implies that $u(t) = 0$ outside $[0, T)$.

We let N_T be the (nonlinear) operator given by

$$(5.2) \quad N_T(u) = 1_{[0, T)}(t) I_T (A(D) (|P_{\leq M} u|^2 P_{\leq M} u)).$$

We will use a fixed point argument to the map⁵

$$\phi_T : u \mapsto 1_{[0,T)}(t)e^{ith(D)}u_0 + N_T(u)$$

to obtain a strong solution $u \in C_t^0([0, T), L_x^2)$. Such fixed point argument of establishing wellposedness results has been a standard method (see for example [Tsu87] [Tao06]).

Proof of Theorem 2. Let $\epsilon_0, \delta_0 > 0$ be small constants to be chosen later, and let $T = \infty$. We let $B_{L_x^2}(0, \epsilon_0)$ be the closed ball of radius ϵ_0 in the space $L_x^2(\mathbb{R}^2)$, and similarly we let $B_{X^0}(0, \delta_0)$ be the closed ball of radius δ_0 in X^0 . Because of the embedding $X^0 \hookrightarrow L_t^\infty L_x^2$, $B_{X^0}(0, \delta_0) \cap C^0([0, \infty), L_x^2)$ is a closed subspace of X^0 , and we put X^0 -norm on it which turns it into a Banach space.⁶

Since the nonlinearity is of algebraic power type, we can show using $\|1_{[0,\infty)} \cdot\|_{X^0} \lesssim \|\cdot\|_{X^0}$, Proposition 7 and the triangle inequality that

$$\|N_\infty(u) - N_\infty(v)\|_{X^0} \leq c_1(\|u\|_{X^0}^2 + \|v\|_{X^0}^2)\|u - v\|_{X^0},$$

for some admissible constant $c_1 > 0$. We also have

$$(5.3) \quad \|\phi_\infty(u)\|_{X^0} \leq c_2(\|u_0\|_{L_x^2} + \|u\|_{X^0}^3)$$

for some admissible constant $c_2 > 0$, since $\|1_{[0,\infty)}(t)e^{ith(D)}u_0\|_{X^0} \lesssim \|1_{[0,\infty)}u_0\|_{U^2} \lesssim \|u_0\|_{L_x^2}$. Therefore we can choose $\delta_0, \epsilon_0 > 0$ small enough depending only on c_1, c_2 such that when $u_0 \in B_{L_x^2}(0, \epsilon_0)$,

$$\phi_\infty : B_{X^0}(0, \delta_0) \cap C^0([0, \infty), L_x^2) \rightarrow B_{X^0}(0, \delta_0) \cap C^0([0, \infty), L_x^2)$$

and we have a strict contraction

$$\|\phi_\infty(u) - \phi_\infty(v)\|_{X^0} \leq \frac{1}{2}\|u - v\|_{X^0}.$$

Therefore for every $u_0 \in B_{L_x^2}(0, \epsilon_0)$, there exists a unique fixed point $u \in B_{X^0}(0, \delta_0) \cap C^0([0, \infty), L_x^2)$ of ϕ_∞ , which by Duhamel's formula is a strong global solution to the Cauchy problem (1.1).

For fixed points u, v of ϕ_∞ with initial data $u_0, v_0 \in B_{L_x^2}(0, \epsilon_0)$ respectively, we have

$$\|u - v\|_{X^0} \leq c_2\|u_0 - v_0\|_{L_x^2} + 2c_1\delta_0^2\|u - v\|_{X^0}.$$

If we choose δ_0, ϵ_0 sufficiently small then we can conclude the map

$$B_{L_x^2}(0, \epsilon_0) \rightarrow B_{X^0}(0, \delta_0) \cap C^0([0, \infty), L_x^2), u_0 \mapsto u$$

is Lipschitz continuous. Hence the solution map

$$B_{L_x^2}(0, \epsilon_0) \rightarrow B_{X^0([0,\infty))}(0, \delta_0) \cap C^0([0, \infty), L_x^2), u_0 \mapsto u|_{[0,\infty)}$$

is Lipschitz continuous. Scattering of solutions is an immediate consequence of the fact that the limit $\lim_{t \rightarrow \infty} g$ exists for any function $g \in U^2$.

To establish unconditional uniqueness of solutions in $X^0([0, \infty)) \cap C^0([0, \infty), L_x^2)$, we will show that if $u, v \in X^0([0, \infty)) \cap C^0([0, \infty), L_x^2)$ are two strong solutions to our Cauchy problem (1.1) with $u(0) = v(0)$, then $u = v$ on $[0, \infty)$. By the time-translation invariance of our Cauchy problem (1.1) and a continuity argument, it suffices to show that if $u(0) = v(0)$,

⁵Note that the map here depends on u_0 , although not indicated explicitly in the notation.

⁶By $B_{X^0}(0, \delta_0) \cap C^0([0, \infty), L_x^2)$ we mean the space of functions $u \in B_{X^0}(0, \delta_0)$ such that u is continuous on $[0, \infty)$ as a function into L_x^2 .

then the solutions u, v agree on a short time interval $[0, T')$ for some $T' > 0$. However this is immediate from local wellposedness of our Cauchy problem (1.1). \square

REFERENCES

- [Bou98] Jean Bourgain. Refinements of Strichartz inequality and applications to 2D-NLS with critical nonlinearity. *International Mathematics Research Notices*, 1998(5):253–283, 1998.
- [CDMM90] Michael Cowling, Shaun Disney, Giancarlo Mauzeri, and Detlef Müller. Damping oscillatory integrals. *Inventiones mathematicae*, 101(1):237–260, 1990.
- [CH16] Timothy Candy and Sebastian Herr. Transference of bilinear restriction estimates to quadratic variation norms and the Dirac-Klein-Gordon system. *arXiv preprint arXiv:1605.04882*, 2016.
- [CKZ07] Anthony Carbery, Carlos Kenig, and Sarah Ziesler. Restriction for flat surfaces of revolution in \mathbb{R}^3 . *Proceedings of the American Mathematical Society*, 135(6):1905–1914, 2007.
- [HHK09] Martin Hadac, Sebastian Herr, and Herbert Koch. Well-posedness and scattering for the KP-II equation in a critical space. *Annales de l'Institut Henri Poincaré (C) Nonlinear Analysis*, 26(3):917–941, 2009.
- [KPV91] Carlos Kenig, Gustavo Ponce, and Luis Vega. Oscillatory integrals and regularity of dispersive equations. *Indiana University Mathematics Journal*, 40(1):33–69, 1991.
- [KT05] Herbert Koch and Daniel Tataru. Dispersive estimates for principally normal pseudodifferential operators. *Communications on pure and applied mathematics*, 58(2):217–284, 2005.
- [KTV14] Herbert Koch, Daniel Tataru, and Monica Visan. *Dispersive Equations and Nonlinear Waves: Generalized Korteweg-de Vries, Nonlinear Schrödinger, Wave and Schrödinger Maps*, volume 45 of *Oberwolfach Seminars*. 2014.
- [Lan13] David Lannes. *The water waves problem*, volume 188 of *Mathematical surveys and monographs*. American Mathematical Society, 2013.
- [Obe04] Daniel Oberlin. A uniform Fourier restriction theorem for surfaces in \mathbb{R}^3 . *Proceedings of the American Mathematical Society*, pages 1195–1199, 2004.
- [Sha07] Bassam Shayya. An affine restriction estimate in \mathbb{R}^3 . *Proceedings of the American Mathematical Society*, 135(4):1107–1113, 2007.
- [SM93] Elias Stein and Timothy Murphy. *Harmonic Analysis: Real-Variable Methods, Orthogonality, and Oscillatory Integrals*, volume 43 of *PMS*. 1993.
- [SS85] Christopher Sogge and Elias Stein. Averages of functions over hypersurfaces in \mathbb{R}^n . *Inventiones mathematicae*, 82(3):543–556, 1985.
- [Tao06] Terence Tao. *Nonlinear dispersive equations: local and global analysis*, volume 106 of *CBMS*. 2006.
- [Tsu87] Y. Tsutsumi. L^2 solutions for nonlinear Schrödinger equations and nonlinear groups. *Funk. Ekv.*, 30:115–125, 1987.

DEPARTMENT OF MATHEMATICS, 970 EVANS HALL, BERKELEY, CA 94720
E-mail address: yuqiu@outlook.com, tataru@math.berkeley.edu