UC Agriculture & Natural Resources

Nutrition and Health

Title

Nutrition and Health Info Sheet: Omega-3 Fatty Acids

Permalink

https://escholarship.org/uc/item/7qs416b1

ISBN

978-1-60107-599-4

Authors

Heneman, Karrie Zidenberg-Cherr, Sheri

Publication Date

2008-10-01

DOI

10.3733/ucanr.8316

Peer reviewed



UNIVERSITY OF CALIFORNIA

Division of Agriculture and Natural Resources http://anrcatalog.ucdavis.edu

NUTRITION AND HEALTH INFO SHEET

Omega-3 Fatty Acids

KARRIE HENEMAN, Assistant Project Scientist, Department of Nutrition, University of California, Davis; **SHERI ZIDENBERG-CHERR**, UC Cooperative Extension Nutrition Science Specialist, Department of Nutrition, University of California, Davis

What are omega-3 fatty acids?

Omega-3 fatty acids are a group of fatty acids important for human health. There are three main types of omega-3 fatty acids: eicosapentaenoic acid (EPA), docosahexaenoic acid (DHA), and alpha linolenic acid (ALA), which is converted in our bodies to EPA or DHA after consumption. High concentrations of omega-3 fatty acids are found in fish and plant oils (1), in addition to human milk and eggs from chickens fed a fishmeal diet.

What is the omega-3 fatty acid content of commonly consumed fish and plant oils? (2)

Fish	Omega-3 fatty acids (mg/3oz)	Plant oils	Omega-3 fatty acids (g/tbsp)
cod	134	pumpkin seeds	0.051
catfish	151	olive oil	0.103
haddock	203	walnuts, black	0.156
clams	241	soybean oil	1.231
shrimp	267	rapeseed oil	1.302
flounder	426	walnut oil	1.414
pollock	460	flaxseeds	2.350
tuna, canned	733	walnuts, English	2.574
salmon	1825	flaxseed oil	7.249

What are the beneficial effects associated with consumption of omega-3 fatty acids found in fish?

Consumption of omega-3 fatty acids from fish has been associated with a variety of beneficial effects, including a reduction in overall mortality. This is largely due to protection against coronary heart disease, in part through lowered heart rate and blood pressure (3). Other beneficial effects associated with fish consumption include reduced risk of stroke, depression, and mental decline with aging (3). Current research also suggests that a deficit in omega-3 fatty acids could negatively impact brain development in a fetus (4).

What are the current recommendations for fish consumption?

The American Heart Association recommends eating a variety of fish (particularly fatty fish) at least two times a week for people without coronary heart disease. (A serving is 3 ounces of cooked fish. See the metric conversion table at the end of this publication.) For people with coronary heart disease, the recommendation is 1 gram of omega-3 fatty acids per day (preferably from fatty fish) (5). However, some species of fish are high in mercury, a metal which can negatively impact brain development in young children and developing fetuses (see table below).



What is the omega-3 fatty acid and mercury content of fish and shellfish commonly eaten in the United States? (6)

Fish	Omega-3 fatty acids (g/3 oz)	Mercury (ppm)
catfish*	0.15-0.20	0.05
clams*	0.24	below the level of detection
cod*	0.13-0.24	0.11
crabs*	0.34-0.40	0.06
flounder or sole*+	0.43	0.05
grouper	0.21	0.55
halibut [†]	0.40-1.00	0.26
herring*†	1.71–1.81	0.04
lobster	0.07-0.41	0.31
mahi mahi	0.12	0.19
orange roughy	0.002	0.54
oysters*†	0.37–1.17	below the level of detection
pollock*†	0.46	0.06
red snapper	0.27	0.60
salmon (fresh, frozen)*+	0.68–1.83	0.01
scallops*	0.17	0.05
shrimp*	0.27	below the level of detection
tuna (canned, light)*+	0.26-0.73	0.12
tuna (fresh, frozen)†	0.24–1.28	0.38

Note: Be sure to check local advisories regarding the safety of the fish you catch (7). Advisories for California water bodies can be found at http://oehha.ca.gov/fish/so_cal/index.html.

Because of the mercury danger, the United States Food and Drug Administration and the United States Environmental Protection Agency recommend not eating swordfish, tilefish, king mackerel, or shark for women who are pregnant, breastfeeding, or of childbearing age and for children and teens less than 17 years old (see table below). These groups may eat up to 12 ounces per week of a variety of fish low in mercury, but they should limit the amount of albacore tuna and tuna steaks or fillets to no more than 6 ounces per week (7). Overall, the health benefits of eating fish outweigh the potential risks, especially when the above safety recommendations are followed (9).

What fish have high levels of mercury? (6)

Fish	Mercury (ppm)	Omega-3 fatty acids (g/3 oz)
king mackerel	0.73	0.34
swordfish	0.97	0.70
shark	0.99	0.90
tilefish	1.45	0.80

^{*} Indicates fish and shellfish with lower levels of mercury (8).

[†] Indicates fish and shellfish highest in omega-3 fatty acids.

METRIC CONVERSIONS

English	Conversion factor for English to metric	Conversion factor for metric to English	Metric
grain	64.80	0.015	milligram (mg)
fluid ounce (fl oz)	29.57	0.034	milliliter (ml)
fluid ounce (fl oz)	2.96	0.0034	deciliter (dl)
ounce (oz)	28.35	0.035	gram (g)
cup	236.6	0.004	milliliter (ml)

REFERENCES

- 1. Jenkins, D. J, and A. R. Josse. 2008. Fish oil and omega-3 fatty acids. Can Med Assoc J 178:150.
- 2. Gebauer, S. K., T. L. Psota, W. S. Harris, and P. M. Kris-Etherton. 2006. n-3 fatty acid dietary recommendations and food sources to achieve essentiality and cardiovascular benefits. Am J Clin Nutr 83:1526S-1535S.
- 3. Torpy, J. M., C. Lynm, and R. M. Glass. 2006. Eating fish: Health benefits and risks. JAMA 296:1926.
- 4. McNamara, R. K., and S. E. Carlson. 2006. Role of omega-3 fatty acids in brain development and function: Potential implications for the pathogenesis and prevention of psychopathology. Prostaglandins Leukot Essent Fatty Acids 75:329-49.
- 5. Kris-Etherton, P. M., W. S. Harris, and L. J. Appel. 2002. Fish consumption, fish oil, omega-3 fatty acids, and cardiovascular disease. Circulation 106:2747-57.
- 6. American Heart Association. 2008. Fish, levels of mercury and omega-3 fatty acids. American Heart Association Web site, http://www.americanheart.org/presenter.jhtml?identifier=3013797.
- 7. The United States Food and Drug Administration and The United States Environmental Protection Agency. 2004. What you need to know about mercury in fish and shellfish. Environmental Protection Agency Web site, http://www.epa.gov/waterscience/fish/files/MethylmercuryBrochure.pdf.
- 8. U.S. Department of Health and Human Services and U.S. Environmental Protection Agency. 2006. Mercury levels in commercial fish and shellfish. Center for Food Safety and Applied Nutrition Web site, http://www.cfsan.fda.gov/~frf/sea-mehg.html.
- 9. Mozaffarian, D., and E. B. Rimm. 2006. Fish intake, contaminants, and human health: Evaluating the risks and the benefits. JAMA 296:1885-99.

FOR FURTHER INFORMATION

To order or obtain printed ANR publications and other products, visit the ANR Communication Services online catalog at http://anrcatalog.ucdavis.edu. You can also place orders by mail, phone, or FAX, or request a printed catalog of our products from:

University of California Agriculture and Natural Resources Communication Services 6701 San Pablo Avenue, 2nd Floor Oakland, California 94608-1239 Telephone: (800) 994-8849 or (510) 642-2431

FAX: (510) 643-5470

E-mail inquiries: danrcs@ucdavis.edu

© 2008 The Regents of the University of California Division of Agriculture and Natural Resources

All rights reserved.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise, without the written permission of the publisher and the authors.

Publication 8316

ISBN-13: 978-1-60107-599-4

Production of this material was supported by a grant from the Vitamin Cases Consumer Settlement Fund, created as a result of an antitrust class action. One of the purposes of the fund is to improve the health and nutrition of California consumers.

The University of California prohibits discrimination or harassment of any person on the basis of race, color, national origin, religion, sex, gender identity, pregnancy (including childbirth, and medical conditions related to pregnancy or childbirth), physical or mental disability, medical condition (cancer-related or genetic characteristics), ancestry, marital status, age, sexual orientation, citizenship, or service in the uniformed services (as defined by the Uniformed Services Employment and Reemployment Rights Act of 1994: service in the uniformed services includes membership, application for membership, performance of service, application for service, or obligation for service in the uniformed services) in any of its programs or activities.

University policy also prohibits reprisal or retaliation against any person in any of its programs or activities for making a complaint of discrimination or sexual harassment or for using or participating in the investigation or resolution process of any such complaint.

University policy is intended to be consistent with the provisions of applicable State and Federal laws.

Inquiries regarding the University's nondiscrimination policies may be directed to the Affirmative Action/Equal Opportunity Director, University of California, Agriculture and Natural Resources, 1111 Franklin Street, 6th Floor, Oakland, CA 94607, (510) 987-0096. For information about ordering this publication, telephone 1-800-994-8849. For assistance in downloading this publication, telephone 530-754-3927.

An electronic copy of this publication can be found at the ANR Communication Services catalog Web site, http://anrcatalog.ucdavis.edu.

This publication has been anonymously peer reviewed for technical accuracy by University of California scientists and other qualified professionals. This review process was managed by the ANR Associate Editor for Food and Nutrition.

pr-10/08-LR/CR

