

UCLA

Proceedings of the UCLA Department of Medicine

Title

Oral Allergy Syndrome

Permalink

<https://escholarship.org/uc/item/7r06682n>

Journal

Proceedings of the UCLA Department of Medicine, 17(1)

Author

Morris, Brian S

Publication Date

2012-12-03

CLINICAL VIGNETTE

Oral Allergy Syndrome

By Brian S. Morris, MD

Case Report

The patient is a 35-year-old male with a history of childhood asthma and allergic rhinitis who began to experience episodes of oral angioedema over the past few ten years. Starting in his mid-20's, symptoms began to occur once or twice per year. Over the course of the past five years, symptoms have occurred more frequently and now occur weekly. Episodes can affect various parts of the face and can vary in intensity and duration. Some involve the upper right lip while other episodes involve the lower left lip. Some involve the cheek while others involve the tissues surrounding one of the eyes. The patient has been to the emergency room several times and received courses of steroids and anti-histamines. Testing revealed a CBC with a slight eosinophilia of 12% but other labs including chemistries and thyroid function were within normal limits. Complement levels and the C1-esterase inhibitor level was normal. Total IgE was normal.

His past medical history is remarkable for GERD and multiple episodes of otitis media as a child. He takes no medications and has no drug allergies. He does not smoke or drink alcohol. His family history is unremarkable.

Vital signs included blood pressure of 120/78 mm hg., pulse of 64 beats/minute and temperature of 36.7 C. Physical examination was unremarkable except for nasal and sinus congestion consistent with allergic rhinitis.

General Discussion

Oral allergy syndrome is a common IgE-mediated syndrome typically found in pollen-sensitized adults. Episodes (usually contact urticaria) usually occur in or near the mouth from cross-reactivity to protein founds in certain fruits, vegetables, nuts, or other foods¹. It is one of the most common allergy-mediated processes in adults and has been described under various names including "pollen-food allergy syndrome", "pollen-food syndrome," and "pollen-associated food allergy syndrome"². These reactions are believed to be due to cross-reactions between allergens in pollen and homologous allergens in plant foods³. Non-plant foods such as eggs, seafood, and cow's milk tend not to be

associated with this syndrome. Patients usually report edema and itching of the mouth and surrounding areas shortly after ingesting uncooked fruits, vegetables, or nuts⁴. While cooking often eliminates the possibility of cross-reactivity, freezing foods does not appear to affect the risk⁵. Unlike other food allergy syndromes, most reactions in oral allergy syndrome are limited to the mouth, lips, tongue and throat⁶. Anaphylaxis, intestinal symptoms, and hemodynamic changes are rarely seen in these patients⁷. Clinical manifestations usually include itching, tingling, and swelling of the lips, throat, palate, nasal passages or oral mucosa shortly after the ingestion of the offending food. Most of these patients have seasonal pollen allergies, but some may not appear to react to pollen⁶. Birch pollen-allergic patients are more likely to develop oral allergy syndrome than those who are sensitized to other types of pollen⁸.

Epidemiology

The prevalence of this syndrome is not clearly defined but it is felt to be fairly common occurring in at least 5% of the population⁴. Risk factors include sensitization to tree pollen or other pollens, coexistent allergy-related problems, and living where environmental exposure to pollen is high⁹. Symptoms related to oral allergy syndrome can occur any time of year but are more common during the pollen season(s)¹⁰. Most patients with food allergy syndrome have a history of atopic disorders including asthma, eczema, and allergic rhinitis¹¹.

Etiology and Pathogenesis

Fruit and vegetable proteins have a high degree of structural homology to pollen allergens¹². The symptoms associated with this disorder are believed to result from contact-related urticaria caused by IgE-mediated reactions to pollen-related proteins found in certain foods¹³. Patients with this disorder have produced IgE antibodies against pollen and these antibodies cross-react with other structurally similar proteins found in ingested foods¹⁴. A complex cascade involving mast cells, T-cells, and inflammatory mediators result in the signs and symptoms of this syndrome¹⁵. Some individuals react to

only one food type while others react to many foods¹⁶. The allergenic proteins felt to trigger these reactions are usually destroyed during the cooking process. Thus, cooked foods (with a few exceptions such as nuts and celery) are much less likely than raw foods to be problematic⁵. Antibodies that trigger oral allergy syndrome react to the linear amino acid sequence of the protein or to a conformational epitope of the protein. Cooking food tends to alter the conformational epitope but not the amino acid sequence¹⁷. If the immune response is to the conformational epitope, then the person with oral allergy syndrome may be able to eat the food if it is cooked, but not when it is raw. If the response is to the linear sequence (common in tree pollen/nut allergies), then cooking usually has no effect on the allergic process¹⁸. In most cases patients can tolerate the culprit food in any cooked form, with tree nuts and peanuts being two exceptions¹⁹.

Food-allergy patients who are allergic to certain types of pollens (tree, weed, and grass) tend to react to certain types of foods. For example, patients who are allergic to grass pollen are more likely to react to melon, tomatoes, and oranges²⁰. Patients who are allergic to alder pollen are more likely to react to almonds, apples, peaches, and strawberries¹⁴. Patients with ragweed pollen allergies are more likely to react to bananas, green peppers, and honey¹⁴. Patients with birch pollen allergies are more likely to react to celery, cherries, plums, soy, and wheat¹⁴. Although there are statistical correlations between various types of pollen and types of foods, some people react to only one food while others react to many foods.

Interestingly, latex allergens (like pollen) can also sensitize patient to cross-react to protein found in some foods including banana, kiwi, tomato, and avocado²¹. Many of these reactions can be systemic and severe in nature²². In addition, some patients developed oral allergy syndrome through cross reactions to nonspecific lipid transfer proteins which belong to the prolamin superfamily of plant proteins²³.

Diagnosis and Testing

The evaluation of these patients centers on a detailed, accurate historical record of the circumstances surrounding the patient's symptoms, physical examination, objective allergy testing to confirm pollen sensitization (usually with IgE RAST testing, skin testing, and patch testing), and an elimination diet leading to a reduced incidence of symptoms²⁴. A food challenge is often advocated by specialists to more accurately determine whether a certain food predictably elicits a reaction. Prick testing, patch testing, and RAST IgE testing are used as adjuncts to history taking,

elimination protocols, and food challenges²⁵. In addition, prick testing with samples of fresh foods may be more reliable and less likely to result in false negative testing than more unstable commercially available extracts²⁶. A diagnostic questionnaire for oral allergy syndrome patients has been developed and validated²⁷.

Treatment and Prognosis

Once proper diagnostic testing has been performed, treatment of oral allergy syndrome involves avoiding the offending foods and treating the pollen-induced allergic process. Avoidance of the specific fruits and vegetables that have triggered reactions is the mainstay of treatment. Severe reactions on first exposure to an allergen are rare but can occur²⁸. Severe reactions to foods that were previously tolerated can also occur²⁹. The decision as to whether to simply avoid the uncooked versions of certain foods or all versions of such foods needs to be made case by case. Desensitization to the pollen with immunotherapy is recommended in some cases and can sometimes help minimize cross-reactions³⁰. Peeling or cooking foods may also help minimize or eliminate reactions to foods to which the patient is sensitive⁹. When the offending agents cannot be isolated or avoided or when symptoms persist despite the above measures, anti-histamines can help prevent and control episodes³¹. Oral steroids are also sometimes prescribed in certain clinical circumstances³². An epinephrine pen is also often prescribed to patients to be used in the rare circumstance of systemic anaphylaxis. This is particularly important for patients with a history of a reaction to cooked foods, a history of systemic or pharyngeal reactions, reactions to foods with a higher rate of severe reactions (peanuts, celery, tree nuts, peaches, and mustard), or those at higher risk for pharyngeal occlusion³³. Immunotherapy has also been used with some level of effectiveness in cases of food allergy syndrome. For example, immunotherapy with extracts of birch pollen has been shown to reduce the incidence of apple related episodes³⁴. Some allergists caution these patients to avoid medications that increase gastric pH as preliminary studies suggest that a higher gastric pH may reduce the digestion of some proteins and could increase the risk of some allergic reactions³⁵. Treatment with anti-IgE therapy is as of yet unproven but may hold some promise³⁶.

Unfortunately, there are limited data to assess the prognosis for these patients. The long term consequences of continuing to ingest foods that trigger such reactions are also unknown. Systemic reactions are uncommon but can occur even in those without a history of severe reactions³⁷.

Clinical Course and Follow-Up

The 35-year old patient was diagnosed with food allergy syndrome. After a careful review of his food inventory, it appeared that most of his episodes occurred about the ingestion of apples, plums, and cherries. On skin testing, the patient tested positive for birch pollen. The patient was instructed to avoid apples, plums, and cherries and immunotherapy for birch pollen was initiated. He was also instructed to fully cook fruits and vegetables whenever possible. After the above measures were instituted and the frequency of episodes dramatically decreased over the course of several months.

REFERENCES

1. **Ma S, Sicherer SH, Nowak-Wegrzyn A.** A survey on the management of pollen-food allergy syndrome in allergy practices. *J Allergy Clin Immunol.* 2003 Oct;112(4):784-8. PubMed PMID: 14564362.
2. **NIAID-Sponsored Expert Panel, Boyce JA, Assa'ad A, Burks AW, Jones SM, Sampson HA, Wood RA, Plaut M, Cooper SF, Fenton MJ, Arshad SH, Bahna SL, Beck LA, Byrd-Bredbenner C, Camargo CA Jr, Eichenfield L, Furuta GT, Hanifin JM, Jones C, Kraft M, Levy BD, Lieberman P, Luccioli S, McCall KM, Schneider LC, Simon RA, Simons FE, Teach SJ, Yawn BP, Schwanger JM.** Guidelines for the diagnosis and management of food allergy in the United States: report of the NIAID-sponsored expert panel. *J Allergy Clin Immunol.* 2010 Dec;126(6 Suppl):S1-58. doi: 10.1016/j.jaci.2010.10.007. PubMed PMID: 21134576.
3. **Egger M, Mutschlechner S, Wopfner N, Gadermaier G, Briza P, Ferreira F.** Pollen-food syndromes associated with weed pollinosis: an update from the molecular point of view. *Allergy.* 2006 Apr;61(4):461-76. Review. PubMed PMID: 16512809.
4. **Zuidmeer L, Goldhahn K, Rona RJ, Gislason D, Madsen C, Summers C, Sodergren E, Dahlstrom J, Lindner T, Sigurdardottir ST, McBride D, Keil T.** The prevalence of plant food allergies: a systematic review. *J Allergy Clin Immunol.* 2008 May;121(5):1210-1218.e4. doi: 10.1016/j.jaci.2008.02.019. Epub 2008 Apr 18. PubMed PMID: 18378288.
5. **Bégin P, Des Roches A, Nguyen M, Masse MS, Paradis J, Paradis L.** Freezing does not alter antigenic properties of fresh fruits for skin testing in patients with birch tree pollen-induced oral allergy syndrome. *J Allergy Clin Immunol.* 2011 Jun;127(6):1624-6.e3. doi: 10.1016/j.jaci.2011.01.028. Epub 2011 Mar 16. PubMed PMID: 21411128.
6. **Ortolani C, Ispano M, Pastorello E, Bigi A, Ansaloni R.** The oral allergy syndrome. *Ann Allergy.* 1988 Dec;61(6 Pt 2):47-52. PubMed PMID: 3264668.
7. **Pastorello EA, Ortolani C, Farioli L, Pravettoni V, Ispano M, Borga A, Bengtsson A, Incorvaia C, Berti C, Zanussi C.** Allergenic cross-reactivity among peach, apricot, plum, and cherry in patients with oral allergy syndrome: an in vivo and in vitro study. *J Allergy Clin Immunol.* 1994 Oct;94(4):699-707. PubMed PMID: 7930303.
8. **Enrique E, Cisteró-Bahíma A, Bartolomé B, Alonso R, San Miguel-Moncín MM, Bartra J, Martínez A.** *Platanus acerifolia* pollinosis and food allergy. *Allergy.* 2002 Apr;57(4):351-6. PubMed PMID: 11906368.
9. **Mari A, Wallner M, Ferreira F.** Fagales pollen sensitization in a birch-free area: a respiratory cohort survey using Fagales pollen extracts and birch recombinant allergens (rBet v 1, rBet v 2, rBet v 4). *Clin Exp Allergy.* 2003 Oct;33(10):1419-28. PubMed PMID: 14519150.
10. **Bohle B.** The impact of pollen-related food allergens on pollen allergy. *Allergy.* 2007 Jan;62(1):3-10. Review. PubMed PMID: 17156335.
11. **Ebo DG, Hagedorens MM, Bridts CH, Schuerwegh AJ, De Clerck LS, Stevens WJ.** Flow cytometric analysis of in vitro activated basophils, specific IgE and skin tests in the diagnosis of pollen-associated food allergy. *Cytometry B Clin Cytom.* 2005 Mar;64(1):28-33. PubMed PMID: 15668987.
12. **Valenta R, Kraft D.** Type 1 allergic reactions to plant-derived food: a consequence of primary sensitization to pollen allergens. *J Allergy Clin Immunol.* 1996 Apr;97(4):893-5. Review. PubMed PMID: 8655883.
13. **Sicherer SH.** Clinical implications of cross-reactive food allergens. *J Allergy Clin Immunol.* 2001 Dec;108(6):881-90. PubMed PMID: 11742262.
14. **Santos A, Van Ree R.** Profilins: mimickers of allergy or relevant allergens? *Int Arch Allergy Immunol.* 2011;155(3):191-204. doi: 10.1159/000321178. Epub 2011 Feb 2. Review. PubMed PMID: 21293140.
15. **Hirschwehr R, Valenta R, Ebner C, Ferreira F, Sperr WR, Valent P, Rohac M, Rumpold H, Scheiner O, Kraft D.** Identification of common allergenic structures in hazel pollen and hazelnuts: a possible explanation for sensitivity to hazelnuts in patients allergic to tree pollen. *J Allergy Clin Immunol.* 1992 Dec;90(6 Pt 1):927-36. PubMed PMID: 1281178.
16. **Bohle B, Zwölfer B, Heratizadeh A, Jahn-Schmid B, Antonia YD, Alter M, Keller W, Zuidmeer L, van Ree R, Werfel T, Ebner C.** Cooking birch pollen-related food: divergent consequences for IgE- and T cell-mediated reactivity in vitro and in vivo. *J Allergy Clin Immunol.* 2006 Jul;118(1):242-9. Epub 2006 May 11. PubMed PMID: 16815162.
17. **Sancho AI, Rigby NM, Zuidmeer L, Asero R, Mistrello G, Amato S, González-Mancebo E, Fernández-Rivas M, van Ree R, Mills EN.** The effect of thermal processing on the IgE reactivity of the non-specific lipid transfer protein from apple. *Mal d 3. Allergy.* 2005 Oct;60(10):1262-8. PubMed PMID: 16134992.
18. **Asero R, Mistrello G, Roncarolo D, de Vries SC, Gautier MF, Ciurana CL, Verbeek E, Mohammadi T, Knul-Brettlova V, Akkerdaas JH, Bulder I, Aalberse RC, van Ree R.** Lipid transfer protein: a pan-allergen in plant-derived foods that is highly resistant to pepsin digestion. *Int Arch Allergy Immunol.* 2001 Jan-Mar;124(1-3):67-9. PubMed PMID: 11306929.
19. **Hansen KS, Ballmer-Weber BK, Lüttkopf D, Skov PS, Wüthrich B, Bindslev-Jensen C, Vieths S, Poulsen LK.** Roasted hazelnuts--allergenic activity evaluated by double-blind, placebo-controlled food challenge. *Allergy.* 2003 Feb;58(2):132-8. PubMed PMID: 12622744.
20. **Cuesta-Herranz J, Lázaro M, Martínez A, Figueredo E, Palacios R, de-Las-Heras M, Martínez J.** Pollen allergy in peach-allergic patients: sensitization and cross-reactivity to taxonomically unrelated pollens. *J Allergy Clin Immunol.* 1999 Sep;104(3 Pt 1):688-94. PubMed PMID: 10482847.
21. **García Ortiz JC, Moyano JC, Alvarez M, Bellido J.** Latex allergy in fruit-allergic patients. *Allergy.* 1998 May;53(5):532-6. PubMed PMID: 9636814.
22. **Blanco C, Carrillo T, Castillo R, Quiralte J, Cuevas M.** Latex allergy: clinical features and cross-reactivity with fruits. *Ann Allergy.* 1994 Oct;73(4):309-14. PubMed PMID: 7943998.
23. **Krause S, Reese G, Randow S, Zennaro D, Quarantino D, Palazzo P, Ciardiello MA, Petersen A, Becker WM, Mari A.**

- Lipid transfer protein (Ara h 9) as a new peanut allergen relevant for a Mediterranean allergic population. *J Allergy Clin Immunol.* 2009 Oct;124(4):771-8.e5. doi: 10.1016/j.jaci.2009.06.008. Epub 2009 Aug 8. PubMed PMID: 19665774.
24. **Vieths S, Scheurer S, Ballmer-Weber B.** Current understanding of cross-reactivity of food allergens and pollen. *Ann N Y Acad Sci.* 2002 May;964:47-68. Review. PubMed PMID: 12023194.
25. **Kazemi-Shirazi L, Pauli G, Purohit A, Spitzauer S, Fröschl R, Hoffmann-Sommergruber K, Breiteneder H, Scheiner O, Kraft D, Valenta R.** Quantitative IgE inhibition experiments with purified recombinant allergens indicate pollen-derived allergens as the sensitizing agents responsible for many forms of plant food allergy. *J Allergy Clin Immunol.* 2000 Jan;105(1 Pt 1):116-25. PubMed PMID: 10629461.
26. **Ricci G, Righetti F, Menna G, Bellini F, Miniaci A, Masi M.** Relationship between Bet v 1 and Bet v 2 specific IgE and food allergy in children with grass pollen respiratory allergy. *Mol Immunol.* 2005 Jun;42(10):1251-7. Epub 2005 Jan 16. PubMed PMID: 15829314.
27. **Skypala IJ, Calderon MA, Leeds AR, Emery P, Till SJ, Durham SR.** Development and validation of a structured questionnaire for the diagnosis of oral allergy syndrome in subjects with seasonal allergic rhinitis during the UK birch pollen season. *Clin Exp Allergy.* 2011 Jul;41(7):1001-11. doi: 10.1111/j.1365-2222.2011.03759.x. Epub 2011 Apr 25. PubMed PMID: 21518043.
28. **Vila L, Sánchez G, Sanz ML, Diéguez I, Martínez A, Palacios R, Martínez J.** Study of a case of hypersensitivity to lettuce (*Lactuca sativa*). *Clin Exp Allergy.* 1998 Aug;28(8):1031-5. PubMed PMID: 9756210.
29. **Kosma P, Sjölander S, Landgren E, Borres MP, Hedlin G.** Severe reactions after the intake of soy drink in birch pollen-allergic children sensitized to Gly m 4. *Acta Paediatr.* 2011 Feb;100(2):305-6. doi: 10.1111/j.1651-2227.2010.02049.x. Epub 2010 Dec 1. PubMed PMID: 20942860.
30. **Niederberger V, Pauli G, Grönlund H, Fröschl R, Rumpold H, Kraft D, Valenta R, Spitzauer S.** Recombinant birch pollen allergens (rBet v 1 and rBet v 2) contain most of the IgE epitopes present in birch, alder, hornbeam, hazel, and oak pollen: a quantitative IgE inhibition study with sera from different populations. *J Allergy Clin Immunol.* 1998 Oct;102(4 Pt 1):579-91. PubMed PMID: 9802365.
31. **Gotoda H, Maguchi S, Kawahara H, Terayama Y, Fukuda S.** Springtime pollinosis and oral allergy syndrome in Sapporo. *Auris Nasus Larynx.* 2001 May;28 Suppl:S49-52. PubMed PMID: 11683343.
32. **Maeda N, Inomata N, Morita A, Kirino M, Ikezawa Z.** Correlation of oral allergy syndrome due to plant-derived foods with pollen sensitization in Japan. *Ann Allergy Asthma Immunol.* 2010 Mar;104(3):205-10. doi: 10.1016/j.ana.2009.11.049. PubMed PMID: 20377109.
33. **Cuesta-Herranz J, Lázaro M, Figueredo E, Igea JM, Umpiérrez A, De-Las-Heras M.** Allergy to plant-derived fresh foods in a birch- and ragweed-free area. *Clin Exp Allergy.* 2000 Oct;30(10):1411-6. PubMed PMID: 10998017.
34. **Asero R.** Effects of birch pollen-specific immunotherapy on apple allergy in birch pollen-hypersensitive patients. *Clin Exp Allergy.* 1998 Nov;28(11):1368-73. PubMed PMID: 9824409.
35. **Schöll I, Untersmayr E, Bakos N, Roth-Walter F, Gleiss A, Boltz-Nitulescu G, Scheiner O, Jensen-Jarolim E.** Antiulcer drugs promote oral sensitization and hypersensitivity to hazelnut allergens in BALB/c mice and humans. *Am J Clin Nutr.* 2005 Jan;81(1):154-60. PubMed PMID: 15640475.
36. **McWilliams L, Mousallem T, Burks W.** Future therapies for food allergy. *Hum Vaccin Immunother.* 2012 Aug 16;8(10). [Epub ahead of print] PubMed PMID: 22894951.
37. **Codreanu F, Moneret-Vautrin DA, Morisset M, Guénard L, Rancé F, Kanny G, Lemerdy P.** The risk of systemic reactions to skin prick-tests using food allergens: CICBAA data and literature review. *Eur Ann Allergy Clin Immunol.* 2006 Feb;38(2):52-4. Review. PubMed PMID: 16711536.

Submitted on December 3, 2012