

UC Irvine

UC Irvine Previously Published Works

Title

Superlinear problems

Permalink

<https://escholarship.org/uc/item/7rq240bd>

Journal

Pacific Journal of Mathematics, 214(1)

ISSN

0030-8730

Authors

Schechter, M

Zou, W M

Publication Date

2004-03-01

Peer reviewed

Superlinear Problems

Martin Schechter *

Department of Mathematics, University of California

Irvine, CA 92697-3875, U. S. A

E-mail: mschecht@math.uci.edu

Wenming Zou[†]

Department of Mathematical Sciences, Tsinghua University,

Beijing 100084, P. R. China

E-mail: wzou@math.tsinghua.edu.cn

Abstract

We solve elliptic semilinear boundary value problems in which the nonlinear term is superlinear. By weakening the hypotheses, we are able to include more equations than hitherto permitted. In particular, we do not require the superquadracity condition imposed by most authors, and it is not assumed that the region is bounded.

1 Introduction

Consider the problem

$$(1) \quad -\Delta u = f(x, u), \quad x \in \Omega; \quad u = 0 \quad \text{on} \quad \partial\Omega,$$

where $\Omega \subset \mathbb{R}^n$ is a bounded domain whose boundary is a smooth manifold, and $f(x, t)$ is a continuous function on $\bar{\Omega} \times \mathbb{R}$. This semilinear Dirichlet problem has been studied by many authors. It is called sublinear if there is a constant C such that

$$|f(x, t)| \leq C(|t| + 1), \quad x \in \Omega, \quad t \in \mathbb{R}.$$

Otherwise, it is called superlinear. Beginning in [1], almost all researchers studying the superlinear problem assumed

(a₁) There are constants $c_1, c_2 \geq 0$ such that

$$|f(x, t)| \leq c_1 + c_2|t|^s,$$

*Supported in part by an NSF grant.

[†]Supported by NSFC grant (GP10001019).

where $0 \leq s < (n+2)/(n-2)$ if $n > 2$.

(a₂) $f(x, t) = o(|t|)$ as $t \rightarrow 0$.

(a₃) there are constants $\mu > 2$, $r \geq 0$ such that

$$(2) \quad 0 < \mu F(x, t) \leq t f(x, t), \quad |t| \geq r,$$

where

$$F(x, t) = \int_0^t f(x, s) ds.$$

They proved

Theorem 1.1. *Under hypotheses (a₁)–(a₃), problem (1) has a nontrivial weak solution.*

The condition (a₃) is convenient, but it is very restrictive. In particular, it implies that there exist positive constants c_3, c_4 such that

$$(3) \quad F(x, t) \geq c_3 |t|^\mu - c_4, \quad x \in \Omega, t \in \mathbb{R}.$$

This eliminates many superlinear problems.

A much weaker condition that implies superlinearity is

(a'₃) Either

$$F(x, t)/t^2 \rightarrow \infty \text{ as } t \rightarrow \infty$$

or

$$F(x, t)/t^2 \rightarrow \infty \text{ as } t \rightarrow -\infty$$

The purpose of the present paper is to explore what happens when (a₃) is replaced with (a'₃). Surprisingly, we find the following to be true.

Theorem 1.2. *Under hypotheses (a₁), (a₂), (a'₃) the boundary value problem*

$$(4) \quad -\Delta u = \beta f(x, u), \quad x \in \Omega; \quad u = 0 \text{ on } \partial\Omega,$$

has a nontrivial solution for almost every positive β .

Unfortunately, this theorem does not give any information for any specific β . We then turned our attention to solving (1) under a weaker assumption than (a₃). For this purpose we introduced

(a''₃) There are constants $\mu > 2$, $r \geq 0$ such that

$$(5) \quad \mu F(x, t) - t f(x, t) \leq C(t^2 + 1), \quad |t| \geq r.$$

Note that (a₃) implies both (a'₃) and (a''₃), but they are much weaker. We prove

Theorem 1.3. *Under hypotheses (a₁), (a₂), (a'₃), (a''₃) problem (1) has a nontrivial solution.*

We also have

Theorem 1.4. *If we replace hypothesis (a_3'') with*

(a_3''') *The function*

$$(6) \quad H(x, t) := tf(x, t) - 2F(x, t)$$

is convex in t .

then the problem (1) has at least one nontrivial solution.

Costa-Magalhães [2] solved (1) under the following assumptions:

$$|f(x, t)| \leq a_0|t|^{p-1} + b_0, \quad x \in \Omega, \quad t \in \mathbb{R},$$

$$\limsup_{|t| \rightarrow \infty} \frac{F(x, t)}{|t|^q} \leq b < \infty \quad \text{uniformly for a.e. } x \in \Omega,$$

$$\liminf_{|t| \rightarrow \infty} \frac{H(x, t)}{|t|^\mu} \geq a > 0 \quad \text{uniformly for a.e. } x \in \Omega,$$

$$\limsup_{t \rightarrow 0} 2F(x, t)/t^2 \leq \alpha < \lambda_0 \quad \text{uniformly for a.e. } x \in \Omega,$$

$$F(x, t)/t^2 \rightarrow \infty \quad \text{as } |t| \rightarrow \infty,$$

where $1 \leq p < 2n/(n-2)$, $\mu > n(q-2)/2$.

Willem-Zou [11] proved a weaker form of Theorem 1.2 for a special case. They do not require hypothesis (a_3) , but they do assume

$$tf(x, t) \geq 0, \quad t \in \mathbb{R}$$

and

$$tf(x, t) \geq c_0|t|^\mu, \quad |t| \geq r$$

for some constants $c_0 > 0$, $r \geq 0$, $\mu > 2$. Some authors have replaced hypothesis (a_3) with (3). Although (3) is a more natural assumption, it is still too restrictive to be desirable. It is for this reason that we introduced assumptions (a_3') and (a_3'') .

Stronger versions of Theorems 1.2 – 1.4 will be given in the next section. In them we are not restricted to any particular boundary value problem, and it is not assumed that the region Ω is bounded.

2 The main theorems

Many elliptic semilinear problems can be described in the following way. Let Ω be a domain in \mathbf{R}^n , and let A be a selfadjoint operator on $L^2(\Omega)$. We assume that $A \geq \lambda_0 > 0$ and that

$$(7) \quad C_0^\infty(\Omega) \subset D := D(A^{1/2}) \subset H^{m,2}(\Omega)$$

for some $m > 0$, where $C_0^\infty(\Omega)$ denotes the set of test functions in Ω (i.e., infinitely differentiable functions with compact supports in Ω) and $H^{m,2}(\Omega)$ denotes the Sobolev space. If m is an integer, the norm in $H^{m,2}(\Omega)$ is given by

$$(8) \quad \|u\|_{m,2} := \left(\sum_{|\mu| \leq m} \|D^\mu u\|^2 \right)^{1/2}.$$

Here D^μ represents the generic derivative of order $|\mu|$ and the norm on the right hand side of (8) is that of $L^2(\Omega)$. If m is not an integer, there are several ways of defining the space $H^{m,2}(\Omega)$, all of which are equivalent. We shall not assume that m is an integer.

Let q be a number satisfying

$$(9) \quad \begin{aligned} 2 &< q \leq 2n/(n-2m), & 2m < n, \\ 2 &< q < \infty, & n \leq 2m, \end{aligned}$$

and let $f(x, t)$ be a Carathéodory function on $\Omega \times \mathbf{R}$. This means that $f(x, t)$ is continuous in t for a.e. $x \in \Omega$ and measurable in x for every $t \in \mathbf{R}$.

We consider the problem

$$(10) \quad Au = f(x, u), \quad u \in D.$$

By a solution of (10) we shall mean a function $u \in D$ such that

$$(11) \quad (u, v)_D = (f(\cdot, u), v), \quad v \in D.$$

If u is a solution of (11) and $f(x, u)$ is in $L^2(\Omega)$, then u is in $D(A)$ and solves (10) in the classical sense. Otherwise we call it a weak (or semistrong) solution.

We make the following assumptions

(A) The function $f(x, t)$ satisfies

$$(12) \quad |f(x, t)| \leq V(x)^q (|t|^{q-1} + 1)$$

and

$$(13) \quad f(x, t)/V(x)^q = o(|t|^{q-1}) \text{ as } |t| \rightarrow \infty,$$

where $V(x) > 0$ is a function in $L^q(\Omega)$ such that

$$(14) \quad \|Vu\|_q \leq C\|u\|_D, \quad u \in D.$$

Here

$$(15) \quad \|u\|_q := \left(\int_{\Omega} |u(x)|^q dx \right)^{1/q},$$

$$(16) \quad \|u\|_D := \|A^{1/2}u\|,$$

and $q' = q/(q-1)$. If Ω and $V(x)$ are bounded, then (14) will hold automatically by the Sobolev inequality. However, there are functions $V(x)$ which are unbounded and such that (14) holds even on unbounded regions Ω (cf., e.g., [4]). With the norm (16), D becomes a Hilbert space.

(B) The point λ_0 is an isolated simple eigenvalue with a bounded eigenfunction $\varphi_0(x) \neq 0$ a.e. in Ω .

(C) There is a $\delta > 0$ such that

$$2F(x, t) \leq \lambda_0 t^2, \quad |t| \leq \delta, \quad x \in \Omega,$$

where

$$(17) \quad F(x, t) := \int_0^t f(x, s) ds.$$

(D) There is a function $W(x) \in L^1(\Omega)$ such that either

$$-W(x) \leq F(x, t)/t^2 \rightarrow \infty \text{ as } t \rightarrow \infty, \quad x \in \Omega.$$

or

$$-W(x) \leq F(x, t)/t^2 \rightarrow \infty \text{ as } t \rightarrow -\infty, \quad x \in \Omega.$$

(E) There are constants $\mu < 1, C \geq 0$ such that

$$[2F(x, t) - \mu t f(x, t)]/(t^2 + 1) \leq C, \quad t \in \mathbb{R}, \quad x \in \Omega.$$

We shall prove

Theorem 2.1. *Under the above hypotheses, the problem*

$$(18) \quad Au = f(x, u), \quad u \in D$$

has at least one nontrivial solution.

We also have

Theorem 2.2. *If we replace hypothesis (E) with*

(E') *The function*

$$(19) \quad H(x, t) := tf(x, t) - 2F(x, t)$$

is convex in t .

then the problem (18) has at least one nontrivial solution.

Problem (18) is called sublinear if $f(x, t)$ satisfies

$$(20) \quad |f(x, t)| \leq C(|t| + 1), \quad x \in \Omega, t \in R.$$

Otherwise it is called superlinear. Hypothesis (D) requires (18) to be superlinear.

The problem (18) has been studied by many people. The vast majority of results obtained concern sublinear problems. Much less has been proved for the superlinear case. In [1] the basic assumption was

$$(21) \quad 0 < \mu F(x, t) \leq tf(x, t), \quad |t| \geq r$$

for some $\mu > 2$ and $r \geq 0$. This is a very convenient hypothesis since it readily achieves mountain pass geometry as well as satisfaction of the Palais-Smale condition. However it is a severe restriction; it strictly controls the growth of $f(x, t)$ as $|t| \rightarrow \infty$. Almost every author discussing superlinear problems has made this assumption. We have been able to weaken this assumption considerably, but not to our complete satisfaction. We assume either that

$$\mu F(x, t) - tf(x, t) \leq C(t^2 + 1), \quad |t| \geq r$$

for some $\mu > 2$ and $r \geq 0$ or that (19) is convex in t . These allow much more freedom for the function $f(x, t)$. But they do not allow as much freedom as we would like.

If we drop hypothesis (E) completely, then we are able to prove the following theorems.

Theorem 2.3. *If we replace hypotheses (C), (D) with*

(C') *There are a $\delta > 0$ and a $\tilde{\lambda} > \lambda_0$ such that*

$$2F(x, t) \geq \tilde{\lambda}t^2, \quad |t| \leq \delta, x \in \Omega.$$

and

(D') *There is a function $W(x) \in L^1(\Omega)$ such that*

$$W(x) \geq P(x, t) \rightarrow -\infty \text{ as } |t| \rightarrow \infty, \quad x \in \Omega,$$

where

$$(22) \quad P(x, t) := F(x, t) - \frac{1}{2}\lambda_0 t^2.$$

and drop hypothesis **(E)**, then problem (18) has at least one nontrivial solution.

We also have

Theorem 2.4. *Assume that **(A)**–**(D)** hold. Then for almost every $\beta \in (0, 1)$, the equation*

$$(23) \quad Au = \beta f(x, u)$$

has a nontrivial solution. In particular, the eigenvalue problem (23) has infinitely many solutions.

Theorem 2.5. *If we replace hypothesis **(C)** in Theorem 2.4 with*

(C'') *There are a $\delta > 0$ and a $\tilde{\lambda} \leq \lambda_0$ such that*

$$2F(x, t) \leq \tilde{\lambda} t^2, \quad |t| \leq \delta, \quad x \in \Omega.$$

and **(D)** with

(D'') *Either*

$$\int_{\Omega} F(x, R\varphi_0) dx / R^2 \rightarrow \infty \text{ as } R \rightarrow \infty$$

or

$$\int_{\Omega} F(x, -R\varphi_0) dx / R^2 \rightarrow \infty \text{ as } R \rightarrow \infty.$$

then (23) has a nontrivial solution for almost every $\beta \in (0, \lambda_0/\tilde{\lambda})$.

Corollary 2.6. *If we replace hypothesis **(C'')** in Theorem 2.5 with*

(C''') *$F(x, t)/t^2 \rightarrow 0$ uniformly as $t \rightarrow 0$.*

then (23) has a nontrivial solution for almost every $\beta \in (0, \infty)$.

The method (called the monotonicity trick) which allows one to solve (23) for almost all values of β in some interval was first introduced by Struwe [8] for minimization problems. It was applied by Jeanjean [3] and others for various types of problems.

3 Preliminaries

Define

$$(24) \quad G(u) := \|u\|_D^2 - 2 \int_{\Omega} F(x, u) dx.$$

Under hypothesis **(A)**, it is known that G is a continuously differentiable functional on the whole of D . In fact, the following were proved in [5, pp. 56–58]

Proposition 3.1. Under hypothesis **(A)**, $F(x, u(x))$ and $v(x)f(x, u(x))$ are in $L^1(\Omega)$ whenever $u, v \in D$.

Proposition 3.2. $G(u)$ has a Fréchet derivative $G'(u)$ on D given by

$$(25) \quad (G'(u), v)_D = 2(u, v)_D - 2(f(\cdot, u), v).$$

Proposition 3.3. The derivative $G'(u)$ given by (25) is continuous in u .

Theorem 3.4. Under hypotheses **(A)**–**(C)**, the following alternative holds:

Either

(a) *there is an infinite number of $y(x) \in D(A) \setminus \{0\}$ such that*

$$(26) \quad Ay = f(x, y) = \lambda_0 y$$

or

(b) *for each $\rho > 0$ sufficiently small, there is an $\varepsilon > 0$ such that*

$$(27) \quad G(u) \geq \varepsilon, \quad \|u\|_D = \rho.$$

4 Proofs

We now give the proof of Theorem 2.1.

Proof. We take

$$(28) \quad G(u) = \|u\|_D^2 - 2 \int_{\Omega} F(x, u) dx.$$

Under our hypotheses, Propositions 3.1–3.3 apply, and

$$(29) \quad (G'(u), v) = 2(u, v)_D - 2(f(\cdot, u), v), \quad u, v \in D.$$

By Theorem 3.4 we see that there are positive constants ε, ρ such that

$$(30) \quad G(u) \geq \varepsilon, \quad \|u\|_D = \rho$$

unless

$$(31) \quad Au = \lambda_0 u = f(x, u), \quad u \in D \setminus \{0\}$$

has a solution. This would give a nontrivial solution of (18). We may therefore assume that (30) holds. Next we note that

$$G(\pm R\varphi_0)/R^2 = \|\varphi_0\|_D^2 - 2 \int_{\Omega} \{F(x, \pm R\varphi_0)/R^2 \varphi_0^2\} \varphi_0^2 dx \rightarrow -\infty \text{ as } R \rightarrow \infty$$

by hypothesis **(D)**, since $\varphi_0 \neq 0$ a.e. Since $G(0) = 0$ and (30) holds, we can now apply the usual mountain pass theorem (cf., e.g., [5, p. 22]) to conclude that there is a sequence $\{u_k\} \subset D$ such that

$$G(u_k) \rightarrow c \geq \varepsilon, \quad G'(u_k) \rightarrow 0.$$

Then

$$(32) \quad G(u_k) = \rho_k^2 - 2 \int_{\Omega} F(x, u_k) dx \rightarrow c$$

and

$$(33) \quad (G'(u_k), u_k) = 2\rho_k^2 - 2(f(\cdot, u_k), u_k) = o(\rho_k),$$

where $\rho_k = \|u_k\|_D$. Assume that $\rho_k \rightarrow \infty$, and let $\tilde{u}_k = u_k/\rho_k$. Since $\|\tilde{u}_k\|_D = 1$, there is a renamed subsequence such that $\tilde{u}_k \rightarrow \tilde{u}$ weakly in D , strongly in $L^2_{loc}(\Omega)$ and a.e. in Ω . By (32),

$$\int_{\Omega} \frac{2F(x, u_k)}{u_k^2} \tilde{u}_k^2 dx \rightarrow 1.$$

Let

$$\Omega_1 = \{x \in \Omega : \tilde{u}(x) \neq 0\}, \quad \Omega_2 = \Omega \setminus \Omega_1.$$

Then

$$\frac{2F(x, u_k)}{u_k^2} \tilde{u}_k^2 \rightarrow \infty, \quad x \in \Omega_1$$

by hypothesis **(D)**. If Ω_1 has positive measure, then

$$\int_{\Omega} \frac{2F(x, u_k)}{u_k^2} \tilde{u}_k^2 dx \geq \int_{\Omega_1} \frac{2F(x, u_k)}{u_k^2} \tilde{u}_k^2 dx + \int_{\Omega_2} [-W(x)] dx \rightarrow \infty.$$

Thus, the measure of Ω_1 must be 0, i.e., we must have $\tilde{u} \equiv 0$ a.e. Moreover,

$$\int_{\Omega} \frac{2F(x, u_k) - \mu u_k f(x, u_k)}{u_k^2} \tilde{u}_k^2 dx \rightarrow 1 - \mu.$$

But by hypothesis **(E)**,

$$\limsup \frac{2F(x, u_k) - \mu u_k f(x, u_k)}{u_k^2} \tilde{u}_k^2 \leq \limsup C \frac{u_k^2 + 1}{u_k^2} \tilde{u}_k^2 = 0,$$

which implies that $1 - \mu \leq 0$, contrary to assumption. Hence, the ρ_k are bounded. We can now follow the usual procedures to obtain a weak solution of (18) satisfying $G(u) = c \geq \varepsilon$ (cf., e.g., [5, p. 64]). Since $G(0) = 0$, we see that $u \neq 0$. This completes the proof. \square

We postpone the proof of Theorem 2.2 until the next section.

In proving Theorem 2.3, we shall make use of

Lemma 4.1. *Under hypothesis (C'), there is an $\alpha \neq 0$ such that $G(\alpha\varphi_0) < 0$.*

Proof. We can assume that

$$(34) \quad \|\varphi_0\|_D = 1.$$

Thus,

$$\begin{aligned} G(\alpha\varphi_0) &= \alpha^2 - 2 \int_{\Omega} F(x, \alpha\varphi_0) dx \\ &\leq \alpha^2 - \tilde{\lambda}\alpha^2 \int_{|\alpha\varphi_0(x)| < \delta} \varphi_0(x)^2 dx \\ &\quad + \int_{|\alpha\varphi_0(x)| > \delta} V^q(|\alpha\varphi_0|^q + |\alpha\varphi_0|) \\ &\leq \alpha^2 - \tilde{\lambda}\alpha^2 \|\varphi_0\|^2 + C|\alpha|^q \|V\varphi_0\|_q^q \\ &\leq \alpha^2 [1 - (\tilde{\lambda}/\lambda_0) + C'|\alpha|^{q-2}]. \end{aligned}$$

This can be made negative by taking α sufficiently small. □

Lemma 4.2. *Under hypothesis (D'),*

$$(35) \quad G(u) \rightarrow \infty \quad \text{as} \quad \|u\|_D \rightarrow \infty.$$

Proof. Suppose there is a sequence $\{u_k\} \subset D$ such that $\rho_k = \|u_k\| \rightarrow \infty$ and

$$G(u_k) \leq K.$$

Write

$$u_k = w_k + \alpha_k \varphi_0, \quad \tilde{u}_k = u_k / \rho_k, \quad \tilde{w}_k = w_k / \rho_k, \quad \tilde{\alpha}_k = \alpha_k / \rho_k,$$

where $w_k \perp \varphi_0$. If $\lambda_1 > \lambda_0$ is the next point in the spectrum of A , then

$$\lambda_1 \|w\|^2 \leq \|w\|_D^2, \quad w \perp \varphi_0.$$

Thus

$$\begin{aligned} G(u_k) &= \|u_k\|_D^2 - \lambda_0 \|u_k\|^2 - 2 \int_{\Omega} P(x, u_k) dx \\ &\geq (1 - \frac{\lambda_0}{\lambda_1}) \|w_k\|_D^2 - 2 \int_{\Omega} P(x, u_k) dx \\ &\geq (1 - \frac{\lambda_0}{\lambda_1}) \|w_k\|_D^2 - 2 \int_{\Omega} W(x) dx. \end{aligned}$$

The only way this would not converge to ∞ is if $\|w_k\|_D$ is bounded. But then $\|\tilde{w}_k\|_D \rightarrow 0$, and $|\tilde{\alpha}_k| \rightarrow 1$. Since $\|\tilde{u}_k\|_D = 1$, there is a renamed subsequence such that $\tilde{u}_k \rightarrow \tilde{u}$ weakly in D , strongly in $L^2_{loc}(\Omega)$ and a.e. in Ω . Since $\tilde{w} = 0$ and $|\tilde{\alpha}| = 1$, we have $\tilde{u}(x) = \tilde{\alpha}\varphi_0(x) \neq 0$ a.e. Hence, $|u_k(x)| = \rho_k|\tilde{u}_k(x)| \rightarrow \infty$ a.e. Consequently,

$$\int_{\Omega} P(x, u_k) dx \rightarrow -\infty,$$

showing that $G(u_k) \rightarrow \infty$. This completes the proof. \square

We can now give the proof of Theorem 2.3.

Proof. Let

$$m = \inf_D G.$$

Then there is a sequence $\{u_k\} \subset D$ such that $G(u_k) \rightarrow m$. In view of Lemma 4.2, we must have $\|u_k\|_D \leq C$. Thus, there is a renamed subsequence such that $u_k \rightarrow u$ weakly in D , strongly in $L^2_{loc}(\Omega)$ and a.e. in Ω . Now,

$$\begin{aligned} G(u) &= \|u\|_D^2 - 2 \int_{\Omega} F(x, u) dx \\ &= \|u_k\|_D^2 - 2([u_k - u], u)_D - \|u_k - u\|_D^2 \\ &\quad - 2 \int_{\Omega} F(x, u_k) dx + 2 \int_{\Omega} [F(x, u_k) - F(x, u)] dx \\ &\leq G(u_k) - 2([u_k - u], u)_D + 2 \int_{\Omega} [F(x, u_k) - F(x, u)] dx. \end{aligned}$$

From our hypotheses, it follows that

$$\int_{\Omega} F(x, u_k) dx \rightarrow \int_{\Omega} F(x, u) dx$$

(cf., e.g., [5, p. 64]). We therefore have in the limit $G(u) \leq m$, from which we conclude that $G(u) = m$ and $G'(u) = 0$. Hence, u is a weak solution of (10). We see from Lemma 4.1 that $m < 0$. Since $G(0) = 0$, we see that $u \neq 0$. This completes the proof. \square

5 The eigenvalue problem

In this section we shall give the proofs of Theorems 2.4, 2.5 and 2.2. They will be based on the following result given in [7]. Let E be a reflexive Banach space with norm $\|\cdot\|$, and let A, B be two closed subsets of E . Suppose that $G \in C^1(E, \mathbb{R})$ is of the form: $G(u) := I(u) - J(u)$, $u \in E$, where $I, J \in C^1(E, \mathbb{R})$ map bounded sets to bounded sets. Define

$$G_{\lambda}(u) = \lambda I(u) - J(u), \quad \lambda \in \Lambda,$$

where Λ is an open interval contained in $(0, +\infty)$. Assume one of the following alternatives holds.

(H₁) $I(u) \geq 0$ for all $u \in E$ and either $I(u) \rightarrow \infty$ or $|J(u)| \rightarrow \infty$ as $\|u\| \rightarrow \infty$.

(H₂) $I(u) \leq 0$ for all $u \in E$ and either $I(u) \rightarrow -\infty$ or $|J(u)| \rightarrow \infty$ as $\|u\| \rightarrow \infty$.

Furthermore, we suppose that

(H₃) $a_0(\lambda) := \sup_A G_\lambda \leq b_0(\lambda) := \inf_B G_\lambda$, for any $\lambda \in \Lambda$.

We let Φ be the set of mappings $\Gamma(t) \in C(E \times [0, 1], E)$ with the following properties:

- a) for each $t \in [0, 1]$, $\Gamma(t)$ is a homeomorphism of E onto itself and $\Gamma(t)^{-1}$ is continuous on $E \times [0, 1]$
- b) $\Gamma(0) = I$
- c) for each $\Gamma(t) \in \Phi$ there is a $u_0 \in E$ such that $\Gamma(1)u = u_0$ for all $u \in E$ and $\Gamma(t)u \rightarrow u_0$ as $t \rightarrow 1$ uniformly on bounded subsets of E .

We have

Theorem 5.1. *Assume that (H₁) (or (H₂)) and (H₃) hold.*

- (1) *If A links B and A is bounded, then for almost all $\lambda \in \Lambda$ there exists $u_k(\lambda) \in E$ such that $\sup_k \|u_k(\lambda)\| < \infty$, $G'_\lambda(u_k(\lambda)) \rightarrow 0$ and*

$$G_\lambda(u_k(\lambda)) \rightarrow a(\lambda) := \inf_{\Gamma \in \Phi} \sup_{s \in [0, 1], u \in A} G_\lambda(\Gamma(s, u)), \quad k \rightarrow \infty.$$

Furthermore, if $a(\lambda) = b_0(\lambda)$, then $\text{dist}(u_k(\lambda), B) \rightarrow 0$, $k \rightarrow \infty$.

- (2) *If B links A and B is bounded, then for almost all $\lambda \in \Lambda$ there exists $v_k(\lambda) \in E$ such that $\sup_k \|v_k(\lambda)\| < \infty$, $G'_\lambda(v_k(\lambda)) \rightarrow 0$ and*

$$G_\lambda(v_k(\lambda)) \rightarrow b(\lambda) := \sup_{\Gamma \in \Phi} \inf_{s \in [0, 1], v \in B} G_\lambda(\Gamma(s, v)), \quad k \rightarrow \infty.$$

Furthermore, if $a_0(\lambda) = b(\lambda)$, then $\text{dist}(v_k(\lambda), A) \rightarrow 0$, $k \rightarrow \infty$.

We shall also need the following extension of Theorem 3.4.

Theorem 5.2. *Let λ be a parameter satisfying $1 < \lambda \leq K < \infty$. Under hypotheses **(A)**–**(D)**, for each $\rho > 0$ sufficiently small (not depending on λ), we have*

$$(36) \quad G_\lambda(u) := \lambda \|u\|_D^2 - 2 \int_\Omega F(x, u) dx \geq (\lambda - 1)\rho^2, \quad \|u\|_D = \rho.$$

*If we replace hypothesis **(C)** with hypothesis **(C'')**, assuming $1 < \tilde{\lambda}/\lambda_0 < \lambda \leq K < \infty$, then we have*

$$(37) \quad G_\lambda(u) \geq \left(\lambda - \frac{\tilde{\lambda}}{\lambda_0} \right) \rho^2, \quad \|u\|_D = \rho.$$

Proof. Let $\lambda_1 > \lambda_0$ be the next point in the spectrum of A , and let N_0 denote the eigenspace of λ_0 . We take $M = N_0^\perp \cap D$. By hypothesis **(B)**, there is a $\rho > 0$ such that

$$\|y\|_D \leq \rho \Rightarrow |y(x)| \leq \delta/2, \quad y \in N_0.$$

Now suppose $u \in D$ satisfies

$$(38) \quad \|u\|_D \leq \rho \text{ and } |u(x)| \geq \delta$$

for some $x \in \Omega$. We write

$$(39) \quad u = w + y, \quad w \in M, \quad y \in N_0.$$

Then for those $x \in \Omega$ satisfying (38) we have

$$\delta \leq |u(x)| \leq |w(x)| + |y(x)| \leq |w(x)| + (\delta/2).$$

Hence

$$(40) \quad |y(x)| \leq \delta/2 \leq |w(x)|,$$

and consequently,

$$(41) \quad |u(x)| \leq 2|w(x)|$$

for all such x . Now we have by (12) and (14)

$$\begin{aligned} G_\lambda(u) &\geq \lambda \|u\|_D^2 - \lambda_0 \int_{|u| < \delta} u^2 dx - C \int_{|u| > \delta} (|Vu|^q + V^q|u|) dx \\ &\geq \lambda \|u\|_D^2 - \lambda_0 \|u\|^2 - C' \int_{|u| > \delta} |Vu|^q dx \\ &\geq (\lambda - 1) \|y\|_D^2 + \lambda \|w\|_D^2 - \lambda_0 \|w\|^2 - C'' \int_{2|w| > \delta} |Vw|^q dx \end{aligned}$$

in view of the fact that $\|y\|_D^2 = \lambda_0 \|y\|^2$ and (41) holds. Thus, by (14),

$$(42) \quad G_\lambda(u) \geq (\lambda - 1) \|y\|_D^2 + \left(\lambda - \frac{\lambda_0}{\lambda_1} - C''' \|w\|_D^{q-2} \right) \|w\|_D^2, \quad \|u\|_D \leq \rho.$$

We take $\rho > 0$ to satisfy

$$1 - \frac{\lambda_0}{\lambda_1} > C''' \rho^{q-2}$$

This gives

$$\begin{aligned} G_\lambda(u) &\geq (\lambda - 1) \rho^2 + \left(\lambda - \frac{\lambda_0}{\lambda_1} - C''' \rho^{q-2} - \lambda + 1 \right) \|w\|_D^2 \\ &\geq (\lambda - 1) \rho^2, \quad \|u\|_D = \rho. \end{aligned}$$

Hence, (36) holds.

To prove (37) under hypothesis (\mathbf{C}'') , let $\eta = \tilde{\lambda}/\lambda_0$ and $\Lambda = (\eta, K)$. Under hypothesis (\mathbf{C}'') we have in place of (42)

$$(43) \quad G_\lambda(u) \geq (\lambda - \eta)\|y\|_D^2 + \left(\lambda - \frac{\tilde{\lambda}}{\lambda_1} - C'''\|w\|_D^{q-2} \right) \|w\|_D^2, \quad \|u\|_D \leq \rho.$$

We take $\rho > 0$ to satisfy

$$\eta - \frac{\tilde{\lambda}}{\lambda_1} > C'''\rho^{q-2}$$

Consequently,

$$\begin{aligned} G_\lambda(u) &\geq (\lambda - \eta)\rho^2 + \left(\lambda - \frac{\tilde{\lambda}}{\lambda_1} - C'''\rho^{q-2} - \lambda + \eta \right) \|w\|_D^2 \\ &\geq (\lambda - \eta)\rho^2, \quad \|u\|_D = \rho. \end{aligned}$$

This gives (37), and the proof is complete. \square

We now turn to the proofs of Theorems 2.4 and 2.5. We prove the latter first. We shall prove Theorem 2.5 by applying Theorems 5.1 and 5.2.

Proof. We take $E = D$, $\Lambda = (\eta, K)$, where $\eta = \tilde{\lambda}/\lambda_0$, $K > 1$ is a finite number, and

$$I(u) = \|u\|_D^2, \quad J(u) = 2 \int_{\Omega} F(x, u) dx.$$

For the purpose of this application, it is sufficient to know that the sets

$$A_{\pm} = [0, \pm R\varphi_0], \quad B = \{x \in D : \|x\|_D = \rho\}$$

link each other if $R > \rho$ (cf., e.g., [5]). In our case hypothesis (\mathbf{H}_1) is satisfied. We now check that (\mathbf{H}_3) holds. We observe that

$$G'_\lambda(u) = 0$$

is equivalent to (23) with $\beta = 1/\lambda$. Now, at least one of the expressions

$$J(\pm R\varphi_0)/R^2 = 2 \int_{\Omega} F(x, \pm R\varphi_0) dx/R^2 \rightarrow \infty \text{ as } R \rightarrow \infty$$

by hypothesis (\mathbf{D}'') . Hence, for R sufficiently large, one of the inequalities

$$G_\lambda(\pm R\varphi_0)/R^2 \leq K\|\varphi_0\|_D^2 - 2 \int_{\Omega} \{F(x, \pm R\varphi_0)/R^2\} \leq 0$$

holds. Thus,

$$a_0(\lambda) \leq 0, \quad \lambda \in \Lambda.$$

Moreover, it follows from Theorem 5.2 that (37) holds. Hence,

$$b_0(\lambda) \geq (\lambda - \eta)\rho^2, \quad \lambda \in \Lambda.$$

This shows that hypothesis **(H₃)** holds. We can now apply Theorem 5.1 to conclude that for almost all $\lambda \in \Lambda$, there exists $u_k(\lambda) \in D$ such that $\sup_k \|u_k(\lambda)\| < \infty$, $G'_\lambda(u_k(\lambda)) \rightarrow 0$ and

$$G_\lambda(u_k(\lambda)) \rightarrow a(\lambda) \geq b_0(\lambda).$$

Once it is known that the sequence $\{u_k\}$ is bounded, we can apply the usual theory to conclude that there is a solution of

$$G'_\lambda(u) = 0, \quad G_\lambda(u) = a(\lambda)$$

(cf., e.g., [5, p. 64]). Moreover, from the definition, we see that $a(\lambda) \geq (\lambda - \eta)\rho^2$. Hence, the equation $G'_\lambda(u) = 0$ has a nontrivial solution for almost every $\lambda \in \Lambda$. This is equivalent to (23) having a nontrivial solution for almost every $\beta \in (K^{-1}, \eta^{-1})$. Since K was arbitrary, the result follows. \square

To prove Theorem 2.4, it suffices to take $\tilde{\lambda} = \lambda_0$ and show that hypothesis **(D)** implies hypothesis **(D'')**. To see this, we note that

$$\int_{\Omega} F(x, \pm R\varphi_0) dx / R^2 = \int_{\Omega} \frac{F(x, \pm R\varphi_0)}{R^2 \varphi_0^2} \varphi_0^2 dx \rightarrow \infty$$

by hypothesis **(D)** and the fact that $\varphi_0(x) \neq 0$ a.e.

To prove Corollary 2.6, we let ε be any positive number. By hypothesis **(C''')**, there is a $\delta > 0$ such that

$$F(x, t)/t^2 \leq \varepsilon, \quad |t| \leq \delta, \quad x \in \Omega.$$

By Theorem 2.5, equation (23) has a nontrivial solution for a.e. $\beta \in (0, \lambda_0/\varepsilon)$. Since ε was arbitrary, the result follows.

We now give the proof of Theorem 2.2.

Proof. By Theorem 2.4, for each arbitrary $K > 1$, and a.e. $\lambda \in (1, K)$, there exists u_λ such that $G'_\lambda(u_\lambda) = 0$, $G_\lambda(u_\lambda) = a(\lambda) \geq (\lambda - 1)\rho^2$. Choose $\lambda_n \rightarrow 1$, $\lambda_n > 1$. Then there exists u_n such that

$$G'_{\lambda_n}(u_n) = 0, \quad G_{\lambda_n}(u_n) = a(\lambda_n) \geq a(1) \geq b_0(1).$$

By Theorem 3.4, we may assume $b_0(1) \geq \varepsilon > 0$. Therefore,

$$\int_{\Omega} \frac{2F(x, u_n)}{\|u_n\|_D^2} dx \leq c.$$

Now we prove that $\{u_n\}$ is bounded. If $\|u_n\|_D \rightarrow \infty$, let $w_n = u_n/\|u_n\|_D$, then $w_n \rightarrow w$ weakly in D , strongly in $L^2_{loc}(\Omega)$ and a.e. in Ω .

Case 1: $w \neq 0$ in D . We get a contradiction as follows:

$$\begin{aligned} c &\geq \int_{\Omega} \frac{2F(x, u_n)}{\|u_n\|_D^2} dx = \int_{\Omega} \frac{2F(x, u_n)}{u_n^2} |w_n|^2 dx \\ &\geq \int_{w \neq 0} \frac{2F(x, u_n)}{u_n^2} |w_n|^2 dx - \int_{w=0} W_1(x) dx \rightarrow \infty. \end{aligned}$$

Case 2: $w = 0$ in D . We define $t_n \in [0, 1]$ by

$$G_{\lambda_n}(t_n u_n) = \max_{t \in [0, 1]} G_{\lambda_n}(t u_n).$$

For any $c > 0$ and $\bar{w}_n = c w_n$, we have

$$\int_{\Omega} F(x, \bar{w}_n) dx \rightarrow 0$$

(cf., e.g., [5, p. 64]). Thus,

$$G_{\lambda_n}(t_n u_n) \geq G_{\lambda_n}(c w_n) = c^2 \lambda_n - 2 \int_{\Omega} F(x, \bar{w}_n) dx \geq c^2/2$$

for n large enough. That is, $\lim_{n \rightarrow \infty} G_{\lambda_n}(t_n u_n) = \infty$ and $(G'_{\lambda_n}(t_n u_n), u_n) = 0$. Therefore,

$$\begin{aligned} G_{\lambda_n}(t_n u_n) &= \int_{\Omega} \left(f(x, t_n u_n) t_n u_n - 2F(x, t_n u_n) \right) dx \\ &= \int_{\Omega} H(x, t_n u_n) dx \rightarrow \infty. \end{aligned}$$

By hypothesis **(E')**,

$$G_{\lambda_n}(u_n) = \int_{\Omega} H(x, u_n) dx \geq \int_{\Omega} H(x, t_n u_n) dx \rightarrow \infty.$$

But

$$\begin{aligned} G_{\lambda_n}(u_n) = a(\lambda_n) &\leq \sup_{s \in [0, 1], u \in A} G_{\lambda_n}((1-s)u) \\ &\leq \sup_{s \in [0, 1], u \in A} G_K((1-s)u) \\ &< c, \end{aligned}$$

a contradiction. Thus, $\|u_n\|_D \leq C$. It now follows that

$$G'(u_n) \rightarrow 0, \quad G(u_n) \rightarrow a(1) \geq b_0(1).$$

We can now apply Theorem 3.4.1 in [5, p. 64] to obtain the desired solution. \square

References

- [1] A. Ambrosotti and P. H. Rabinowitz, Dual variational methods in critical point theory and applications, *J. Func. Anal.*, 14 (1973) 349–381.
- [2] D. G. Costa and C. A. Magalhães, Variational elliptic problems which are nonquadratic at infinity, *Nonlinear Analysis TMA*, 23(1994) 1401–1412.
- [3] L. Jeanjean, On the existence of bounded Palais-Smale sequences and applications to a Landesman-Lazer type problem set on \mathbb{R}^N , *Proc. Roy. Soc. Edinburgh A* 129 (1999) 787– 809.
- [4] M. Schechter, Superlinear elliptic boundary value problems, *Manuscripta Math.*, 86 (1995) 253–265.
- [5] M. Schechter, *Linking Methods in Critical Point Theory*, Birkhäuser Boston, 1999.
- [6] M. Schechter, *Spectra of Partial Differential Operators*, North-Holland, Amsterdam, 1986.
- [7] M. Schechter and W. Zou, Double linking theorem and multiple solutions, preprint.
- [8] M. Struwe, The existence of surfaces of constant mean curvature with free boundaries, *Acta Math.* 160 (1988) 19–64.
- [9] M. Struwe, *Variational Methods*, Springer-Verlog, 1996.
- [10] M. Willem, *Minimax Theorems*, Birkhäuser Boston, 1996.
- [11] M. Willem and W. Zou, On a semilinear Dirichlet problem and a nonlinear Schrödinger equation with periodic potential, preprint.