

Lawrence Berkeley National Laboratory

Recent Work

Title

QUANTUM CORRECTIONS (WITHIN THE CLASSICAL PATH APPROXIMATION) TO THE BOLTZMANN DENSITY MATRIX

Permalink

<https://escholarship.org/uc/item/7s43g175>

Author

Hornstein, Steven M.

Publication Date

1971-11-01

c.2

QUANTUM CORRECTIONS
(WITHIN THE CLASSICAL PATH APPROXIMATION)
TO THE BOLTZMANN DENSITY MATRIX

Steven M. Hornstein and William H. Miller

November 29, 1971

AEC Contract No. W-7405-eng-48

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*



34

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

QUANTUM CORRECTIONS (WITHIN THE CLASSICAL PATH APPROXIMATION)

TO THE BOLTZMANN DENSITY MATRIX

by

Steven M. Hornstein and William H. Miller*

Inorganic Materials Research Division, Lawrence Berkeley Laboratory
and Department of Chemistry; University of California,
Berkeley, California 94720

ABSTRACT

A previously derived classical path approximation for diagonal matrix elements of the Boltzmann density matrix (i.e., the particle density) is expanded in a power series in \hbar about the classical density. Comparing term by term with Wigner's expansion of the exact quantum mechanical density, one sees that the first quantum correction to the classical density (which involves first and second derivatives of the potential energy) is given correctly by the classical path approximation. Wigner's expression for the second correction term, however, involves the first four derivatives of the potential, and the classical path approximation gives only the part of this term that involves the first three derivatives.

I. Introduction

Recent work has demonstrated the value of the classical path approximation to quantum mechanical problems. This classical-limit approach, in which the Feynman sum over all paths is replaced by a sum over only the classical path (or paths), has yielded accurate results in scattering problems¹ and quantum statistical mechanics.²

In this Letter we investigate in more detail a recently derived classical path approximation for the Boltzmann density matrix². The particle density (i.e., the diagonal elements of the density matrix) given by this approximation was seen to be considerably more accurate than the ordinary classical density, being qualitatively correct (and reasonably quantitative) even in the low temperature limit where the classical density is completely inadequate. Furthermore, in regions where the potential energy function has only first and second non-vanishing derivatives, the classical path approximation was seen to give the exact quantum mechanical result for the density.

To gain more insight into the nature of the classical path approximation and its level of accuracy, this paper systematically expands the classical-path density in a power series in \hbar , and explicit expressions are obtained for the first two "quantum correction" terms to the ordinary classical density. Comparing with the exact quantum expressions for these correction terms obtained by Wigner³, one sees that the first quantum correction to the classical density (which involves first and second derivatives of the potential) is given correctly by the classical path approximation. The second correction term, however, involves the first four derivatives of the potential energy and the classical path

approximation gives only the part involving its first three derivatives.

II. Expansion of the Classical Path Approximation

For a single particle in a one-dimensional potential well the particle density $\rho_\beta(x) \equiv \langle x | e^{-\beta H} | x \rangle$ is given in the classical path approximation by² ($\beta \leq 1/kT$)

$$\rho_\beta(x) = D_\beta(x)^{-\frac{1}{2}} \exp[-\phi_\beta(x)] \quad (1)$$

where

$$\phi_\beta(x) = \beta V(x_0) + 2(2m/\hbar^2)^{\frac{1}{2}} \int_{x_0}^x dx' [V(x') - V(x_0)]^{\frac{1}{2}} \quad (2)$$

$$D_\beta(x) = 4\pi(\hbar^2/2m)^{\frac{1}{2}} [V(x_0) - V(x)] \int_{x_0}^x dx' [V(x') - V(x_0)]^{-\frac{3}{2}}. \quad (3)$$

The turning point x_0 is the function of x and β defined by the equation

$$\lambda^{\frac{1}{2}} = \int_{x_0}^x dx' [V(x') - V(x_0)]^{-\frac{1}{2}}, \quad (4)$$

where the "quantum parameter" (dimensions = length²/energy) is

$$\lambda = \hbar^2 \beta^2 / 2m. \quad (5)$$

It is easy to show² that in the limit $\lambda \rightarrow 0$ Equations (1)-(4) reduce to the usual classical density

$$\rho_\beta^{CL}(x) = \left(\frac{2m}{\hbar^2} / 4\pi\beta \right)^{\frac{1}{2}} \exp[-\beta V(x)]. \quad (6)$$

Here we wish to expand Equations (2)-(4) in a power series in λ to obtain

Equation (1) in the form

$$\rho_{\beta}(x) = \rho_{\beta}^{CL}(x) [1 + \lambda C_1(x, \beta) + \lambda^2 C_2(x, \beta) + \dots]; \quad (7)$$

i.e., the goal is to obtain explicit expressions for the "quantum correction" functions $C_k(x, \beta), k = 1, 2, \dots$.

The first step is to change variables of integration in Equations (2)-(4) from x' to the dimensionless variable z , $x' = x_0 + z \Delta x$, where $\Delta x \equiv x - x_0$; the limits of all the integrals then become 0 to 1. The integrands in Equations (2)-(4) are then expanded in a power series in Δx and integrated term by term; Equation (4), for example, becomes

$$\lambda^{\frac{1}{2}} = 2(\Delta x/V')^{\frac{1}{2}} \left\{ 1 + \Delta x(5/12)(V''/V') + \Delta x^2 [(43/160)(V''/V')^2 - (11/60)(V'''/V')] + o(\Delta x^3) \right\}, \quad (4')$$

where $V' = V'(x)$, $V'' = V''(x)$, etc. Equation (4'), a power series in Δx for λ , can be inverted to obtain Δx as a power series in λ :

$$\Delta x = (\lambda V'/4) \left\{ 1 - \lambda(5/24)V'' + \lambda^2 [(61/1440)(V'')^2 + (11/480)V'V'''] + o(\lambda^3) \right\} \quad (8)$$

Inserting Equation (8) into the Δx -expansions of Equations (2) and (3) then gives a λ -expansion for these quantities:

$$\phi_{\beta}(x) = \beta [V - \lambda(V')^2/12 + \lambda^2(V')^2V''/60 + o(\lambda^3)] \quad (2')$$

$$D_{\beta}(x) = 4\pi\beta(\hbar^2/2m) \left\{ 1 + \lambda V''/3 + \lambda^2 [(V'')^2/30 - V'V'''/15] + o(\lambda^3) \right\}, \quad (3')$$

or

$$D_{\beta}(x)^{-\frac{1}{2}} = \left(\frac{2m/4\pi\beta}{\hbar^2} \right)^{\frac{1}{2}} \left\{ 1 - \lambda V''/6 + \lambda^2 [(V'')^2/40 + V'V''''/30] + O(\lambda^3) \right\} \quad (3'')$$

Combining Equations (2') and (3'') in with Equation (1) gives the density in the form of Equation (7), where the first two quantum correction functions are identified as

$$C_1(x, \beta) = \beta [V'(x)]^2/12 - V''(x)/6 \quad (9)$$

$$C_2(x, \beta) = \beta^2 [V'(x)]^4/288 - (11/360)\beta [V'(x)]^2 V''(x) + [V''(x)]^2/40 + V'(x)V''''(x)/30 \quad (10)$$

Higher terms can be similarly generated.

The exact quantum expressions for these correction terms have been determined by Wigner³, and one sees that Equation (9) for the first quantum correction is the same as Wigner's result. Wigner's expression for $C_2(x, \beta)$, however, is the same as that in Equation (10) but with the term $-V''''(x)/60\beta$ added to it.

III. Discussion

Knowing from Wigner's work³ that the first quantum correction to the classical density involves only first and second derivatives of the potential, one could have predicted beforehand that the classical path approximation would give the exact quantum result for this first correction term - for it has been noted above that the classical path approximation is exact (for all values of λ) if the potential has only first and second non-zero derivatives. Thus it is not surprising, too, that the part of the second correction term which involves the first two derivatives of the potential is given correctly. The surprising feature, which could not have been anticipated, is that the part of

$C_2(x,\beta)$ which involves the third derivative is also given correctly.

The classical path approximation was seen not to be capable, however, of producing the fourth derivative term in the quantum expression for $C_2(x,\beta)$.

These observations are another example of the fact that the classical path approximation is not a high temperature (small β) approximation in the present statistical mechanical application, nor a short time approximation in scattering situations; rather it is an approximation that the potential energy is a slowly varying function. The second correction term $C_2(x,\beta)$, therefore, will be given quite accurately by Equation (10) if the fourth derivative of the potential is small compared to the three lower derivatives. This notion that the classical path approximation is basically an approximation of a slowly varying potential is also a well-known idea from ordinary WKB theory⁴.

Acknowledgment

This work was supported by the U. S. Atomic Energy Commission.

References

* Alfred P. Sloan Fellow

1. See, for example, W. H. Miller, J. Chem. Phys., 53, 1949; 3578 (1970); Chem. Phys. Letters, 7, 431 (1970); J. Chem. Phys., 54, 5386 (1971); 55, 3150 (1971).
2. W. H. Miller, J. Chem. Phys., 55, 3146 (1971).
3. E. P. Wigner, Phys. Rev., 40, 749 (1932).
4. See, for example, L. I. Schiff, Quantum Mechanics, McGraw-Hill, New York, 1968, p. 271.

LEGAL NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

TECHNICAL INFORMATION DIVISION
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720