

UCSF

UC San Francisco Previously Published Works

Title

M. tuberculosis Ser/Thr Protein Kinase D Phosphorylates an Anti-Anti-Sigma Factor Homolog

Permalink

<https://escholarship.org/uc/item/7vh3737z>

Journal

PLOS Pathogens, 3(4)

ISSN

1553-7366

Authors

Greenstein, Andrew E

MacGurn, Jason A

Baer, Christina E

et al.

Publication Date

2007-04-01

DOI

10.1371/journal.ppat.0030049

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

M. tuberculosis Ser/Thr Protein Kinase D Phosphorylates an Anti-Anti-Sigma Factor Homolog

Andrew E. Greenstein¹, Jason A. MacGurn², Christina E. Baer³, Arnold M. Falick⁴, Jeffery S. Cox², Tom Alber^{1*}

1 Department of Molecular and Cell Biology, University of California, Berkeley, California, United States of America, **2** Department of Microbiology and Immunology, University of California San Francisco, San Francisco, California, United States of America, **3** Biophysics Graduate Group, University of California, Berkeley, California, United States of America, **4** Howard Hughes Medical Institute Mass Spectrometry Laboratory, University of California, Berkeley, California, United States of America

Receptor Ser/Thr protein kinases are candidates for sensors that govern developmental changes and disease processes of *Mycobacterium tuberculosis* (*Mtb*), but the functions of these kinases are not established. Here, we show that *Mtb* protein kinase (Pkn) D overexpression alters transcription of numerous bacterial genes, including Rv0516c, a putative anti-anti-sigma factor, and genes regulated by sigma factor F. The PknD kinase domain directly phosphorylated Rv0516c, but no other sigma factor regulator, in vitro. In contrast, the purified PknB and PknE kinase domains phosphorylated distinct sigma regulators. Rather than modifying a consensus site, PknD phosphorylated Rv0516c in vitro and in vivo on Thr2 in a unique N-terminal extension. This phosphorylation inhibited Rv0516c binding in vitro to a homologous anti-anti-sigma factor, Rv2638. These results support a model in which signals transmitted through PknD alter the transcriptional program of *Mtb* by stimulating phosphorylation of a sigma factor regulator at an unprecedented control site.

Citation: Greenstein AE, MacGurn JA, Baer CE, Falick AM, Cox JS, et al. (2007) *M. tuberculosis* Ser/Thr protein kinase D phosphorylates an anti-anti-sigma factor homolog. PLoS Pathog 3(4): e49. doi:10.1371/journal.ppat.0030049

Introduction

Mycobacterium tuberculosis (*Mtb*) is among the world's most harmful pathogens, causing approximately two million deaths annually [1]. In addition to the emergence of multi-drug-resistant strains, *Mtb* evades current therapeutics by shifting from active infection to a persistent, metabolically dormant state [2]. This transition exemplifies the distinctive *Mtb* life cycle, which encompasses unique developmental adaptations to distinct environments [3]. Little is known about the signaling mechanisms that mediate the biochemical changes that initiate and maintain the stages of *Mtb* development.

Candidate regulators of *Mtb* development include receptor Ser/Thr protein kinases (STPKs) that modulate intracellular events in response to external stimuli. In eukaryotes, homologous STPKs sense environmental cues and transduce signals that regulate virtually all aspects of cell physiology. The *Mtb* genome encodes 11 such Hanks-type (also called "eukaryotic-like") STPKs, including nine putative transmembrane receptor kinases [4]. Although the activating stimuli for these kinases have not been identified, the extracellular C-terminal sensor domains include a β -propeller interaction motif, a PASTA repeat thought to bind cell wall structures, and a redox-sensitive DsbG homolog [5–8]. The intracellular, N-terminal kinase domains structurally resemble eukaryotic homologs, and similar receptor STPKs are widely distributed in bacterial genera. The first reported bacterial STPK substrates include pThr-binding forkhead-associated (FHA) domains [9], metabolic enzymes [10], and apparent regulators of cell division [11,12], but the mechanisms of signaling in vivo are not established. Genetic studies suggest that two of the 11 *Mtb* STPKs are essential for growth [13] and that the STPKs regulate characteristics such as cell shape [11], virulence [14], and nitrogen balance [15]. Identifying the

intracellular targets of *Mtb* STPKs is essential to understanding their mechanistic roles in *Mtb* biology.

A second class of bacterial Ser/Thr kinases, the anti-sigma factors, regulates gene expression by controlling alternative sigma factors [16]. Alternative sigma factors, such as sigma B (SigB) and sigma F (SigF) in *Bacillus subtilis*, mediate transcriptional responses to environmental cues by binding RNA polymerase and mediating promoter recognition. Work on *B. subtilis* has established the paradigm by which complex regulatory cascades influence alternative sigma factor activity (reviewed by Hughes and Mathee [16]). Anti-sigma factor proteins (e.g., RsbW) directly sequester cognate alternative sigma factors and prevent RNA polymerase binding. Anti-anti-sigma factors (e.g., RsbV) relieve this transcriptional repression by binding the anti-sigma factor. The anti-sigma factors phosphorylate anti-anti-sigma factors on a conserved Ser or Thr, and this modification promotes dissociation of the complex.

This basic regulatory organization is recapitulated for multiple layers in which paralogs of anti-sigma factors and anti-anti-sigma factors switch partners and ultimately determine the concentration of the active sigma factor [17]. In this

Editor: Lalita Ramakrishnan, University of Washington, United States of America

Received: August 10, 2006; **Accepted:** February 15, 2007; **Published:** April 6, 2007

Copyright: © 2007 Greenstein et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abbreviations: MBP, maltose binding protein; *Mtb*, *Mycobacterium tuberculosis*; MyBP, myelin basic protein; Pkn[letter], protein kinase [letter]; SigF, sigma factor F; STPK, Ser/Thr protein kinase; TEV, tobacco etch virus; WT, wild-type

* To whom correspondence should be addressed. E-mail: tom@ucray.berkeley.edu

Author Summary

Many bacteria, including *Mycobacterium tuberculosis* (*Mtb*), sense the environment using a family of signaling proteins called Ser/Thr protein kinases (STPKs), but the functions of these sensors are not well understood. This study shows that the *Mtb* protein kinase (Pkn) D STPK attaches a phosphate group to one and only one member of a family of regulators of “alternative” sigma factors, which activate sets of genes in numerous bacteria. Phosphorylation of the regulator at an unprecedented position abolished binding in vitro to a putative partner. Remarkably, increasing PknD activity in *Mtb* not only strongly activated the gene encoding the specific regulatory protein phosphorylated by PknD, but also altered the expression of genes controlled by an alternative sigma factor. By providing evidence for a mechanistic link between PknD and gene regulation, this work supports a new model in which STPKs in numerous microorganisms transduce environmental signals by controlling expression of specific groups of genes. Thus, certain bacterial STPKs may orchestrate aspects of the coordinate control of gene expression essential for adaptation in the environment and in host infections.

“partner switching” mechanism, anti-sigma factor paralogs play two distinct roles. Some anti-sigma factors (e.g., RsbW) antagonize transcription by directly sequestering alternative sigma factors. In contrast, other anti-sigma factors (e.g., RsbT) act upstream to stimulate transcription by binding and activating the master “environmental sensing” phosphatase (RsbU in *B. subtilis* [18]). This phosphatase reactivates anti-anti-sigma factors, which bind the cognate anti-sigma factor, thus increasing the concentration of free sigma factor. Environmental cues affect the phosphorylation state of upstream anti-anti-sigma factor paralogs (such as RsbS and the RsbRA-D proteins), and these proteins form a complex (termed the “stressosome”) that binds the positive regulator of the phosphatase [19]. The central role of Ser/Thr phosphorylation in anti-anti-sigma factor regulation and the established role of eukaryotic kinases in gene regulation led us to test the hypothesis that the eukaryotic-like STPKs may impinge on transcription regulated by alternative sigma factors.

Here, we demonstrate that increasing the activity of the PknD STPK in *Mtb* resulted in specific phosphorylation of a single anti-anti-sigma factor homolog, Rv0516c. Simultaneously, the *Rv0516c* gene was activated and transcription of genes regulated by the SigF alternative sigma factor was coordinately altered. PknD phosphorylated Rv0516c at a novel site, Thr2, distinct from conserved Ser/Thr phosphorylation sites in the anti-anti-sigma factor family. Thr2 phosphorylation abolished binding to another anti-anti-sigma factor. These results demonstrate that PknD phosphorylates a putative sigma factor regulator in *Mtb*, alters binding of a cognate regulator, and, by a mechanism that has not been determined, changes the expression of SigF-dependent genes.

Results

To investigate the pathways regulated by receptor STPK signaling, we constructed *Mtb* strains expressing either wild-type (WT) or kinase-dead (Asp138Asn) PknD under the control of an acetamide-inducible promoter [20]. The Asp138Asn mutation in the catalytic site reduced the in vitro

activity of kinase domain ~2,600-fold (Figure S1). In this approach, excess kinase substituted for an activating signal to stimulate downstream pathways. Western blotting with anti-PknD and anti-pThr antibodies showed that the WT or mutant kinases accumulated after induction and produced a concomitant increase in Thr phosphorylation (Figure 1A). Expression of the attenuated Asp138Asn mutant produced a much smaller increase in phosphorylation of cellular targets. Consistent with the idea that the expressed PknD (directly or indirectly) mediated the observed phosphorylation in vivo, cellular-protein phosphorylation was blocked when the PknD variants were induced in the presence of SP600125 (Figure S1B), a c-Jun N-terminal kinase (JNK) inhibitor that shows specificity for PknD over other *Mtb* STPKs (C. Mieczkowski and T. Alber, unpublished data).

Transcriptional profiling using microarrays confirmed the induction of *PknD* transcripts and revealed a set of genes regulated by PknD overexpression in a kinase-dependent manner (Figure 1B; Table S1). Remarkably, the transcripts most differentially expressed in the strain expressing WT PknD (Figure 1B) included the genes with the largest reductions in transcription during log phase growth of an *Mtb* mutant harboring a deletion of *sigF* [21]. Moreover, the *Rv0516c* gene, which is homologous to anti-anti-sigma factors, was dramatically induced by PknD overexpression. The established role of phosphorylation in anti-anti-sigma factor regulation supported the hypothesis that PknD specifically phosphorylates Rv0516c.

To test this idea, we measured the phosphorylation by the purified PknD kinase domain of all predicted *Mtb* homologs of the *B. subtilis* alternative sigma factor regulators. Potential regulators were identified using iterative PSI-BLAST searches for homologs of Rv0516c, SpoIIAA, and SpoIIAB, and putative homologs were confirmed using 3D-PSSM [22] to verify that the predicted fold resembled anti- or anti-anti-sigma factors (Figure 2). All of the identified sigma factor regulators were cloned, expressed in *Escherichia coli*, and purified. Using a [γ - 32 P]ATP transfer assay, we found that the PknD kinase domain (PknD₁₋₃₇₈) efficiently phosphorylated Rv0516c, but not any of the other sigma factor regulator homologs (Figure 3A). The SP600125 inhibitor blocked this Rv0516c phosphorylation (Figure 3B) in a dose-dependent manner, indicating that PknD catalyzed the observed phosphorylation. Rv0516c phosphorylation was reversed by PstP (Figure 3C), the *Mtb* protein Ser/Thr phosphatase that dephosphorylates all *Mtb* STPK substrates tested to date [23]. These results showed that PknD and PstP act on the putative anti-anti-sigma factor Rv0516c in vitro.

To determine if *Mtb* UsfX (Rv3287c), the anti-sigma factor Ser kinase that controls SigF [24], also phosphorylates Rv0516c, we incubated these proteins under conditions in which UsfX phosphorylated the model substrate, myelin basic protein (MyBP). In contrast to PknD, UsfX failed to phosphorylate Rv0516c (Figure S2). The anti-sigma factor paralogs RshA [25] and Rv0941c also failed to phosphorylate Rv0516c (unpublished data). Thus, unlike anti-anti-sigma factors that are phosphorylated by anti-sigma factors, Rv0516c is phosphorylated by a eukaryotic-like STPK, PknD.

To determine if sigma factor regulator phosphorylation is a general function of *Mtb* STPKs, we assayed the ability of four other *Mtb* kinase domains (PknA, PknB, PknE, and PknK) to phosphorylate Rv0516c and the eight other purified *Mtb*










Gene number	Domain Architecture	Function	Name
Rv0516c		unknown	
Rv0941c		unknown	
Rv1364c		unknown	
Rv1365c		Anti-anti SigF	RsfA
Rv1904		unknown	
Rv2638		unknown	
Rv3221		Anti-SigH	RshA
Rv3287c		Anti-SigF	UsfX
Rv3687c		Anti-anti SigF	RsfB

Figure 2. *Mtb* Proteins Containing Domains Homologous to the *B. subtilis* Alternative Sigma Factor Regulators SpoIIAA and SpoIIAB

Four of the *Mtb* homologs of the *B. subtilis* SpoIIAB (anti-sigma factor) or SpoIIAA (anti-anti-sigma factor) encode proteins that have been shown to regulate SigF or SigH. *Rv0941c* contains both types of domain, and *Rv1364c* contains both domains as well as an RsbU-like, Ser/Thr phosphatase domain. Three of the seven anti-anti-sigma factor domains are preceded by a Ser/Thr-rich N-terminal extension (blue). This extension in Rv0516c contains the site of PknD phosphorylation.

doi:10.1371/journal.ppat.0030049.g002

absence of a Ser or Thr at the consensus phosphorylation site and our failure to observe Rv0516c phosphorylation by any anti-sigma factor in vitro are consistent with PknD phosphorylation at the distinct site, Thr2.

A yeast two-hybrid analysis of interactions among *Mtb* sigma factor regulators has suggested that Rv0516c can bind the homologous predicted anti-anti-sigma factor, Rv2638 [18]. To test this association and investigate the role of Thr2 phosphorylation in regulating the interaction, we used affinity chromatography to compare binding of purified Rv2638 to Rv0516c before and after PknD phosphorylation. Rv2638 bound Rv0516c, and Rv0516c phosphorylation by PknD abolished this association (Figure 5B). These results showed that Thr2 phosphorylation regulates the interaction in vitro between Rv0516c and the anti-anti-sigma factor paralog, Rv2638.

Discussion

Because the identity of activating environmental signals remains unknown, we stimulated PknD receptor kinase activity in *Mtb* by overexpressing the protein (Figure 1A). Overexpression was expected to stimulate phosphorylation of PknD substrates directly by increasing the concentration of the kinase and indirectly by favoring dimerization (by mass action), which leads to allosteric activation [26]. PknD activity produced a transcriptional response that altered genes activated by SigF during log phase growth (Figure 1B), including the anti-anti-sigma factor homolog Rv0516c [21]. Strikingly, the PknD kinase domain also directly phosphorylated the Rv0516c protein (but none of the other *Mtb* sigma factor regulators) in vitro and upon overexpression in vivo. In contrast to the conserved internal phosphorylation sites found in many anti-anti-sigma factors [16], PknD phosphorylated Rv0516c on Thr2 in an N-terminal extension similar to that found in two additional mycobacterial anti-anti-sigma factors, Rv1904 and Rv2638. Phosphorylation directly blocked Rv0516c binding to Rv2638, indicating that Thr2 phosphorylation has a direct functional consequence.

Although the roles of the Rv0516c:Rv2638 complex are unknown, alternative phosphorylation sites and functional interactions between upstream anti-anti-sigma factor homologs have been demonstrated in *B. subtilis* for the RsbS and RsbRA-RsbRD regulators [17,27–29]. These *B. subtilis* regulators form a large complex that controls the environmental sensing phosphatase RsbU, which dephosphorylates anti-anti-sigma factors [17,27–29].

The correlation between SigF-responsive genes [21] and genes that are transcriptionally sensitive to PknD activity (Figure 1) is specific to PknD; overexpression of *Mtb* PknB produces a distinct transcriptional response (T. Lombana, J. MacGurn, J. Cox, and T. Alber, unpublished data). Nonetheless, these data do not demonstrate a direct mechanistic link between Rv0516c and SigF. To the contrary, the lack of Rv0516c phosphorylation by the *Mtb* anti-SigF (UsfX, Rv3287c) or any other anti-sigma factor hints that PknD indirectly influences SigF-mediated transcription by phosphorylating Rv0516c or other substrates. The present data do not distinguish models in which repression of the SigF response is caused by Rv0516c phosphorylation or by a distinct signal generated by phosphorylation of one or more other proteins in vivo.

The substrates of the *Mtb* STPKs are not restricted to transcriptional regulators. Rough estimates suggest that the number of Ser/Thr phosphoproteins in *Mtb* may exceed 100 [30], and proposed *Mtb* substrates include metabolic enzymes [31], regulatory proteins [12,32], and membrane channels [33]. Using proteomic methods to analyze lysates phosphorylated in vitro by the PknD kinase domain, Perez and coworkers recently reported that PknD phosphorylates MmpL7, the transporter for phthiocerol dimycocerosate (PDIM), a lipid essential for virulence [34]. These studies, however, did not test whether MmpL7 is phosphorylated in vivo, whether phosphorylation altered the function of MmpL7, or whether MmpL7 is a better substrate of other STPKs. Perez and coworkers did not detect PknD phosphorylation of Rv0516c, perhaps because this regulatory protein




Figure 3. PknD Specifically Phosphorylates Rv0516c In Vitro
 (A) Each anti-sigma factor or anti-anti-sigma factor (gene [Rv] number listed along the top) fused to MBP was cloned, expressed, purified, and incubated with (+) or without (-) PknD in a [γ - 32 P]ATP transfer assay. The assay was quenched with EDTA, and TEV protease was added to separate the MBP tag from the sigma factor regulator. The two dark bands in the center of each PknD lane correspond to the autophosphorylated kinase and the MBP tag. The sigma factor regulators released by TEV protease are between 12 and 20 kDa. Only Rv0516c was phosphorylated by PknD. (B) Dose-dependent attenuation by the PknD inhibitor, SP600125, shows that PknD phosphorylates Rv0516c. (C) PstP, a protein Ser/Thr phosphatase from *Mtb*, dephosphorylates Rv0516c. PknD (lane 1), but not heat-inactivated PknD (lane 2), phosphorylates Rv0516c. PstP was added to PknD-phosphorylated Rv0516c, and the phosphatase reaction was quenched immediately (lane 3) or after 30 min (lane 4). (D) In vitro kinase assays were performed to investigate the specificity of five *Mtb* kinases for the eight sigma factor regulators (Figure S3). Assays were carried out in a manner identical to the assay shown in Figure 3A. Kinase concentrations were adjusted to catalyze equal phosphorylation of myelin basic protein in vitro. PknB, PknD, and PknE phosphorylated a specific subset of substrates.
 doi:10.1371/journal.ppat.0030049.g003

may not be sufficiently abundant in the *Mtb* lysates or because proteins <20 kDa (such as Rv0516c) were run off the two-dimensional gels used to identify potential substrates [34]. Similar reasons may explain the failure of Perez and coworkers to detect phosphorylation of small proteins containing FHA domains previously found to be in vitro substrates of PknD [9].

In contrast, the complementary approach used here, based on assaying the biochemical and transcriptional effects of kinase activation in vivo, is sensitive to changes in the activity of regulatory factors, even proteins present in small amounts. By assaying in vitro PknD phosphorylation of all the *Mtb* homologs of the *B. subtilis* SpoIIAA and SpoIIAB sigma factor




Figure 4. PknD Phosphorylates Rv0516c on Thr2 In Vitro
 (A) Mass spectrometry of trypsin-digested, PknD-phosphorylated Rv0516c indicated the presence of a single phosphorylation site within the N-terminal nine residues. The figure shows the mass spectrum of an HPLC fraction of the digest. The measured mass (monoisotopic MH^+) of the N-terminal peptide (1384.60 Da [arrow]) was 79.97 Da larger than the expected mass of 1,304.63 Da, corresponding to addition of a single phosphate group. Tandem MS and N-terminal sequencing of the peptide at m/z 1,384.60 revealed Thr2 as the site of phosphorylation. (B) Rv0516c Thr2 occurs in a unique, 28-residue Ser/Thr-rich, N-terminal extension, not in the anti-anti-sigma factor domain. (C) Mutations in the N-terminus of Rv0516c confirm that PknD phosphorylates Rv0516c on Thr2. The eight indicated mutants of Rv0516c were purified, and in vitro phosphorylation by the PknD kinase domain was assayed after cleavage with the TEV protease (left). Non-native residues before the start of Rv0516c left after TEV protease cleavage are shaded in gray. Protein-stained loading controls (center) confirmed that an equal amount of each Rv0516c variant was assayed. Controls lacking kinase showed that the phosphorylation was due to PknD (far right). Thr2Ala (T2A) and deletions of the N-terminal four and eight residues caused the most dramatic reductions in phosphorylation.
 doi:10.1371/journal.ppat.0030049.g004

regulators, we found that only Rv0516c was efficiently phosphorylated (Figure 3). Nonetheless, overexpressing PknD may cause abnormal phosphorylation or physiological changes that result in indirect transcriptional changes unrelated to normal kinase functions. The striking correlation between PknD phosphorylation of Rv0516c and activation of the *Rv0516c* gene, however, suggests a potential autoregulatory loop and sets the stage to explore the biological roles of this sigma factor regulator and the consequences of Rv0516c phosphorylation in vivo.

Although bacterial STPKs phosphorylate many types of




Figure 5. PknD Phosphorylates Rv0516c on Thr2 In Vivo, and This Phosphorylation Blocks Binding to Another Sigma Factor Regulator

(A) Western blot of lysates of *Mtb* strains expressing full-length PknD under the control of an acetamide-inducible promoter and constitutively expressing Rv0516c with a C-terminal, FLAG epitope tag. As controls, the kinase was rendered inactive (D138N) or the Rv0516c phosphorylation site was mutated (Thr2Ala [T2A]). The active (but not the mutant) PknD phosphorylated the overexpressed WT Rv0516c. The levels of endogenous proteins were too low to detect phosphorylated Rv0516c in this experiment. KatG was detected with antibodies as a loading control. Rv0516c (but not the Thr2Ala mutant) was phosphorylated upon WT PknD induction.

(B) Phosphorylation on Thr2 abolished binding to Rv2638. Equal amounts of cell lysates overexpressing Rv2638 were incubated with pre-phosphorylated (lane 4) or unphosphorylated (lane 3) Rv0516c. As a control, Rv2638 binding to the purification tag alone was evaluated (lane 2). Protein staining was used to insure that equal amounts of recombinant protein were used as bait in each reaction, and bound Rv2638 was visualized by Western blotting with antibodies against the His₆ tag. Total soluble protein before incubation with the amylose-Sepharose (lane 1) indicated the presence of Rv2638. Only unphosphorylated Rv0516c bound the anti-anti-sigma factor, Rv2638. doi:10.1371/journal.ppat.0030049.g005

proteins [30], alternative sigma factor regulators may be substrates of STPKs in diverse genera. In addition to the activity of PknD, the PknB and PknE kinase domains phosphorylated sigma factor regulators in vitro (Figures 3D and S3). In contrast, some kinase domains (e.g., PknA and PknK; Figure 3D) apparently do not phosphorylate these sigma factor regulators. With up to 12 candidate alternative sigma factors in the *Mtb* genome, it is unlikely that each kinase controls a completely autonomous pathway. Instead,

our data suggest that phosphorylation pathways may converge on overlapping sets of regulators (Figure 3D). The specific phosphorylation of Rv0516c on a novel functional site by PknD suggests that STPK phosphorylation of sigma factor regulators goes beyond the paradigm established to date in *B. subtilis*.

Materials and Methods

Bacterial strains, media, and growth conditions. The strains and plasmids used in this study are listed in Table 1. *M. tuberculosis* (Erdman) cultures were grown in 7H9 medium and transformed as previously described [35]. Plasmids were maintained episomally by growth in medium containing antibiotics.

Microarrays: Data collection and analysis. Strains were grown to mid-log phase in 7H9 media before induction of PknD by addition of acetamide (0.2%). RNA was isolated from cultures at indicated time points as previously described [36] and quantified by measuring OD₂₆₀. RNA was random-primed and reverse transcribed in the presence of amino-allyl dUTP. Residual RNA was hydrolyzed by addition of 0.2 N NaOH, 0.1 M EDTA, and incubation at 65 °C for 15 min, followed by addition of 0.2 N HCl to neutralize. The cDNA was purified with Zymo binding columns (Zymo Research, <http://www.zymoresearch.com>) and conjugated to either Cy3 (individual cDNA samples) or Cy5 (common reference pool cDNA). An equal quantity of each RNA sample within an experiment (representing both *Mtb* strains at all time points) was used to make a common cDNA reference pool. Dye-conjugated cDNA from each individual sample was mixed and co-hybridized with dye-conjugated cDNA from the common reference pool on microarray slides containing oligonucleotide spots representing every gene in *M. tuberculosis* (Qiagen, <http://www.qiagen.com>). After 2 d of hybridization at 63 °C, arrays were washed and scanned using a GenePix 3000B scanner (Axon Instruments, <http://www.moleculardevices.com>). Array gridding was performed in GenePix Pro 4.1, and Nomad 2.0 was used to select high quality spots. For each spot, the ratio of medians (Rm) was averaged from repeat hybridizations and normalized to $t = 0$ (uninduced). Cluster analysis was performed using Cluster 3.0. Two biological replicates were performed, and each biological replicate was averaged over two hybridizations.

Protein expression and purification for in vitro assays. Using *Mtb* H37Rv genomic DNA as a template for PCR amplification, gene segments encoding PknB₁₋₃₀₈, PknE₁₋₂₈₆, and PknK₁₋₂₈₉ were cloned into pET-28b vectors (Novagen, <http://www.emdbiosciences.com>). PknD₁₋₃₇₈ was cloned into pET-24b (Novagen). PknA₁₋₃₃₇ and full-length clones of each anti-sigma factor or anti-anti-sigma factor were inserted into the Gateway vector pHMGWA [37], which included NH₂-terminal 6X-His and maltose binding protein (MBP) tags, followed by a tobacco etch virus (TEV) protease site. All constructs were confirmed by DNA sequencing.

Proteins were expressed in *E. coli* BL21 CodonPlus (Stratagene, <http://www.stratagene.com>) at 18 °C. The kinase-domain constructs and Rv0516c were purified to homogeneity (as assayed by SDS-PAGE) by immobilized metal affinity chromatography (IMAC) using nickel-equilibrated HiTrap chelating Sepharose (Amersham, <http://www.amershambiosciences.com>), size-exclusion chromatography using HiLoad 26/60 Superdex 75 (Amersham), and anion-exchange chromatography using HiTrap Q Sepharose (Amersham). The sigma factor regulators prepared for kinase-activity screens were purified by nickel-IMAC (Rv0516c was purified by IMAC only for these assays as well). The molecular weight of each sigma factor regulator, as assayed by SDS-PAGE, corresponded to the mass predicted by the gene sequence. Because the kinase-domain constructs autophosphorylated during expression, migration on SDS-PAGE was slightly retarded.

In vitro kinase assays. The sigma factor regulators were dialyzed into the reaction buffer (80 mM NaCl, 20 mM Tris [pH 7.5], 0.5 mM Tris(2-carboxyethyl)phosphine hydrochloride [TCEP], 250 μM MnCl₂). In a total reaction volume of 19 μl, the final concentration of each IMAC-purified 6X-His-MBP-tagged sigma factor regulator was 20 μM, and the final concentration of kinase was 1.2 μM. The reaction was initiated with the simultaneous addition of 1 μL of [γ -³²P]ATP (800 Ci/mmol and 10 mCi/ml; ICN, <http://www.mpbio.com>) and ATP (Sigma, <http://www.sigmaaldrich.com>) to final concentrations of 250 nCi/μl and 50 μM, respectively. The reaction was allowed to proceed for 2 h at room temperature and quenched by the simultaneous addition of EDTA to 20 mM and 7.2 μg of TEV protease.

Table 1. Plasmids and Strains Used in This Study

Category	Plasmid/Strain Name	Description/Genotype	Source	
Plasmids	Original Gateway vector	pDONR207, GentR, CmR	Invitrogen	
	pGWdest3.kan	oriE, oriM, attR1, CmR, ccdB, attR2, acetamidase promoter, KanR	This study	
	pAG1	pGWdest3 with PknD (WT), KanR	This study	
	pAG2	pGWdest3 with PknD (D138N), KanR	This study	
	pGWdest1.hyg	oriE, oriM, attR1, CmR, ccdB, attR2, GroEL promoter, KanR	This study	
	pAG3	pGWdest1 with Rv0516c (WT, C' 3XFLAG), HygR	This study	
	pAG4	pGWdest1 with Rv0516c (T2A, C' 3XFLAG), HygR	This study	
	pAG5	pGWdest1 with Rv0516c (T2E, C' 3XFLAG), HygR	This study	
	M. tuberculosis strains	Erdman	WT	W. R. Jacobs, Jr.
		JMM55	Erdman + pAG1, KanR	This study
JMM57		Erdman + pAG2, KanR	This study	
AGM1		JMM55 + pAG3, KanR HygR	This study	
AGM2		JMM57 + pAG3, KanR HygR	This study	
AGM31		JMM55 + pAG4, KanR HygR	This study	
AGM32		JMM55 + pAG5, KanR HygR	This study	
AGM33		JMM57 + pAG4, KanR HygR	This study	
AGM34		JMM57 + pAG5, KanR HygR	This study	

doi:10.1371/journal.ppat.0030049.t001

The TEV cleavage was allowed to proceed 2 h or overnight at room temperature, resulting in efficient separation of each sigma factor regulator from the tag. The sequence GlyHisMet was left at the NH₂-terminus after TEV cleavage of the tag. The cleavage reactions were separated by SDS-PAGE on 4%–12% NuPage Novex BisTris gels (Invitrogen, <http://www.invitrogen.com>), and the gels were dried. Radioactivity was quantified with a Molecular Dynamics Typhoon 8600 phosphoimager.

Activity of the Asp138Asn mutant. To assess the activity of the Asp138Asn mutant of PknD, we incubated 0.036 nM WT kinase or 3.6 nM Asp138Asn kinase with 0.5 mg/ml MyBP or 0.5 mg/ml Rv0516c. The reaction was carried out for 30 min under buffer, metal, and ATP concentrations similar to those described above, and then quenched with either 5X SDS-PAGE loading dye (for MyBP) or the TEV/EDTA mixture described above. Phosphorylation was quantified with ImageQuant (GE Healthcare, <http://www.gehealthcare.com>) after electrophoresis, drying, and phosphoimager data collection.

PknD inhibition by SP600125. Untagged PknD_{1–378} was purified by IMAC, cleaved with TEV, purified on HiLoad 26/60 Superdex 75 (Amersham), and concentrated from the flow-through fraction of a second IMAC column. Each reaction was set up with or without 38 nM kinase, 20 μM Rv0516c, 250 nCi/μl [γ -³²P]ATP, and 25 μM unlabeled ATP in 50 mM NaCl, 50 mM HEPES (pH 7.5), 0.5 mM TCEP, 10 mM MnCl₂, and 10 mM MgCl₂. SP600125 was diluted into water and added to a concentration of 20 nM to 20 μM. Reactions were carried out and analyzed as described above.

In vitro dephosphorylation of Rv0516c by PstP. PknD, Rv0516c, and PstP were purified to homogeneity [30]. Heat-inactivated PknD was prepared by incubation at 95 °C for 1 h. Phospho-Rv0516c prepared in a 2-h incubation with PknD and [γ -³²P]ATP was treated with 2.3 μg of PstP. The reaction was quenched with EDTA and TEV after zero or 30 min. Separation and quantification were carried out as described above.

Phosphorylation site mapping. Purified Rv0516c was phosphorylated using 6X-His-PknD_{1–378} and 2 mM ATP. The reaction proceeded overnight, and the kinase was removed by IMAC. The flow-through fraction was diluted with water and rocked at room temperature for 2 d to induce precipitation. Supernatant was removed, and the resulting pellet was dissolved in 6 M guanidinium hydrochloride.

The mass of the intact protein was determined by electrospray ionization–ion-trap mass spectrometry. Rv0516c was digested with trypsin; the resulting digest mixture was separated on a reversed-phase C-18 column (0.15 × 150 mm), and fractions were collected. The MALDI TOF spectrum of each fraction was obtained, and the phosphorylated peptide was identified using MALDI-tandem TOF (MS/MS). The MS/MS spectrum was used, along with Edman sequencing, to identify the phosphorylation site.

Mutations to the N-terminus of Rv0516c were created with QuikChange (Stratagene). Proteins were purified and phosphorylated

as described, except that the Rv0516c variants were treated with TEV protease and quenched with the protease inhibitor, aminoethylbenzene sulfonyl fluoride (AEBSF) (MP Biomedicals, <http://www.mpbio.com>), prior to phosphorylation.

In vivo phosphorylation state analysis. Full-length PknD was cloned into an acetamide-inducible *M. tuberculosis* expression vector (pGWdest3.kan) [38]. Full-length Rv0516c (WT or mutant) was cloned into a tuberculosis expression vector under the control of the constitutive GroEL promoter (pGWdest1.hyg). The vector also encoded a C-terminal antigenic FLAG (DDDDK) tag. Mutants were made using Quikchange (Stratagene) on the *Rv0516c* gene in the entry vector. *Mtb* Erdman was transformed by electroporation [35], grown for 3–6 wk on solid rich medium, and single colonies were picked and grown to mid-log phase. Large (50 mL) cultures were inoculated and adjusted to an optical density (OD) at 600 nm of 0.3 after 5 d. To induce PknD expression, acetamide was added to a final concentration of 0.2% at 24, 8, 4, or 2 h before harvesting. Induced and uninduced cultures were grown to a final OD₆₀₀ of ~0.6. Then, 10 mL of each culture were harvested by centrifugation, and resuspended in 200 μL of extraction buffer (1% SDS, 20 mM EDTA, 50 mM HEPES (pH 7.5), 1 mM AEBSF). Samples were immediately boiled for 25 min. Next, 200 μL of 100-μm glass beads were added, and samples were bead-beaten twice for 5 min. Samples were boiled a second time for 10 min and centrifuged. The soluble fraction was removed and diluted with SDS-PAGE loading dye. After electrophoresis in 10%–20% Tris-glycine gels, proteins were transferred to PVDF membrane, and detected with anti-phosphoThreonine (Invitrogen), anti-DDDDK (AbCam), anti-KatG, or anti-PknD antibodies (Pacific Immunology, <http://www.pacificimmunology.com>). HRP-conjugated secondary antibody was used with Kodak BioMax MR film to develop the Western blots. In cases of multiple antibody detection, blots were stripped in 2% SDS/100 mM DTT, 62 mM Tris (pH 7), for 30 min at 50 °C.

In vitro interaction assays. Rv2638 was expressed in *E. coli* in the pHxGWA (N-terminal His₆-thioredoxin) vector [37]. Phosphorylated Rv0516c was prepared by incubation of purified His₆-MBP-tagged protein with 5 mM MnCl₂, 2 mM ATP, and 10:1 (w:w) PknD_{1–378}. Phospho-Rv0516c was purified by IMAC, and both phosphorylated and unphosphorylated Rv0516c were dialyzed into the pull-down buffer (70 mM NaCl, 20 mM HEPES 7.5, 0.5 mM TCEP).

Pull-downs were performed by lysing (by sonication) *E. coli* that expressed Rv2638 in the presence of 200 μg MBP-tagged Rv0516c (phosphorylated or unphosphorylated) or His₆-MBP control in 70 mM NaCl, 20 mM HEPES 7.5, 0.5 mM TCEP, 1 mM AEBSF, and 10 mM MnCl₂. After rocking the lysate for 30 min at 4 °C, samples were centrifuged for 10 min at 14,000 rpm in a microcentrifuge at 4 °C. A small amount of the supernatant was retained for analysis, and the majority was applied to 50 μL of amylose-Sepharose (New England Biolabs, <http://www.neb.com>) pre-equilibrated in the buffer. After rocking for 10 min at 4 °C, resin was washed three times in buffer,

Table 2. GenBank Accession Numbers of Genes Described in This Study

Gene Rv Number	Name	Synonym	Accession	Organism
Rv0516c			NP_215030	<i>M. tuberculosis</i>
Rv0941c			NP_215456	<i>M. tuberculosis</i>
Rv1365c	RsfA		NP_215881	<i>M. tuberculosis</i>
Rv1904			NP_216420	<i>M. tuberculosis</i>
Rv2638			NP_217154	<i>M. tuberculosis</i>
Rv3221c	RshA		NP_217737	<i>M. tuberculosis</i>
Rv3287	UsfX		NP_217804	<i>M. tuberculosis</i>
Rv3687	RsfB		NP_218204	<i>M. tuberculosis</i>
Rv0015c	PknA		NP_214529	<i>M. tuberculosis</i>
Rv0014c	PknB		NP_214528	<i>M. tuberculosis</i>
Rv0931c	PknD		NP_215446	<i>M. tuberculosis</i>
Rv1743	PknE		NP_216259	<i>M. tuberculosis</i>
Rv3080c	PknK		NP_217596	<i>M. tuberculosis</i>
Rv0018c	PstP		NP_214532	<i>M. tuberculosis</i>
Rv1364c			YP_177802	<i>M. tuberculosis</i>
Rv3286c	SigF		NP_217803	<i>M. tuberculosis</i>
	SigB		NP_388354	<i>B. subtilis</i>
	SigF		NP_390226	<i>B. subtilis</i>
	SpolIAA		NP_390228	<i>B. subtilis</i>
	SpolIAB		NP_390227	<i>B. subtilis</i>
	RsbS		NP_388349	<i>B. subtilis</i>
	RsbT		NP_388350	<i>B. subtilis</i>
	RsbRA	RsbR	NP_388348	<i>B. subtilis</i>
	RsbRB	YkoB	NP_389203	<i>B. subtilis</i>
	RsbRC	YojH	NP_389827	<i>B. subtilis</i>
	RsbRD	YqhA	NP_390356	<i>B. subtilis</i>
	RsbU		NP_388351	<i>B. subtilis</i>
	RsbV		NP_388352	<i>B. subtilis</i>
	RsbW		NP_388353	<i>B. subtilis</i>

doi:10.1371/journal.ppat.0030049.t002

resuspended in 40 μ L of 2X SDS-PAGE loading dye, and boiled for 10 min. Samples were separated on 12% Tris-glycine gels (Invitrogen), transferred to PVDF, blocked in 4% non-fat dry milk, and incubated overnight at room temperature with 1:2000 monoclonal anti-His₆ clone HIS-1 (Sigma). Blots were washed and imaged as described above. For loading controls, the same reactions were separated on 12% Tris-glycine gels and stained with Coomassie blue.

Supporting Information

Figure S1. Inhibition of PknD

(A) The D138N mutant of PknD is 2,600-fold less active than WT *in vitro*. Using a 100-fold excess of PknD D138N, we measured a 27.2-fold difference in Rv0516c phosphorylation, and a 25.3-fold difference in MyBP phosphorylation compared to the WT PknD kinase domain. Values were averaged over three replicates.

(B) The PknD inhibitor SP600125 delays changes in phosphorylation upon PknD expression in *Mtb*. *Mtb* strains expressing WT (left) or D138N PknD (right) were induced with or without the simultaneous addition of 60 μ M SP600125 and grown for 8 or 24 h. All samples were lysed at the same time, clarified by centrifugation, separated by SDS-PAGE, and probed with anti-pThr antibodies. The presence of the inhibitor, which is a green compound, turned the cells green, indicating that it was likely taken up. The WT PknD, but not the D138N mutant, was present and phosphorylated in the uninduced cells (compare the 0 time points). At 8 h, SP600125 inhibited cellular protein phosphorylation by the expressed PknD. The D138N mutant PknD, however, was phosphorylated at 8 h in the presence of the inhibitor. This increased phosphorylation suggests that the overexpressed PknD is phosphorylated by other cellular kinases that are not inhibited by SP600125. By 24 h, perhaps due to overexpression of PknD or efflux of the compound, *Mtb* cellular proteins are highly phosphorylated.

Found at doi:10.1371/journal.ppat.0030049.sg001 (239 KB PDF).

Figure S2. Rv3287c (*Mtb* UsfX) Phosphorylates the Model Substrate MyBP, but Not Rv0516c

Autoradiogram showing substrate phosphorylation in the presence (+) and absence (–) of the anti-sigma factor kinase, Rv3287.

Found at doi:10.1371/journal.ppat.0030049.sg002 (44 KB PDF).

Figure S3. Multiple *Mtb* Ser/Thr Protein Kinases Phosphorylate a Unique Set of Putative Sigma-Regulators

The phosphorylation of the putative anti- and anti-anti-sigma factors by PknA, PknB, PknE, and PknK was assessed. PknD did not phosphorylate Rv1364c (unpublished data), but this regulator was omitted from the analysis due to the presence of a phosphatase domain. Each kinase domain was incubated in assay buffer with MBP-tagged substrate protein. The tag was removed with TEV protease and the phosphoproteins were visualized by autoradiography of SDS gels. The MBP tag was phosphorylated by the PknA, PknB, and PknE kinase domains. The kinase-domain concentrations were adjusted to produce similar phosphorylation of the nonspecific substrate, MyBP. Sigma factor regulators migrated in the bottom half of each gel. (A) PknA does not phosphorylate any sigma factor regulators. (B) PknB phosphorylates Rv0516c and Rv3221A (RshA). (C) PknE phosphorylates Rv0516c, Rv1365c (RsfA), Rv1904, and Rv3221A (RshA). (D) PknK does not phosphorylate any of these sigma factor regulators.

Found at doi:10.1371/journal.ppat.0030049.sg003 (624 KB PDF).

Figure S4. Multiple Sequence Alignment around the Conserved Anti-Anti-Sigma Factor Phosphorylation Site

Rv0516c and other putative *Mtb* anti-anti-sigma factors are highly homologous to known anti-anti-sigma factors. For example, Rv0516c and Rv2638 align to the well-characterized RsfA (anti-anti-sigma factor F, Rv1365c) with *e*-values of 2×10^{-5} and 1×10^{-29} , respectively.

(A) A multiple sequence alignment of anti-anti-sigma factors and Rv0516c around the phosphorylation site in *B. subtilis* SpoIIAA (shaded). Phosphorylation of all of these anti-anti-sigma factors has been observed in the respective bacteria. While Rv0516c contains a serine adjacent to the conserved phosphorylation site, the conserved Ser/Thr is replaced with a Gly in Rv0516c.

(B) Multiple sequence alignment showing that the phosphorylation site observed in SpoIIAA and RsbV from *B. subtilis* is conserved only in two of the six *Mtb* anti-anti-sigma factor domains. Oxidation of the highlighted cysteine in Rv1365c has been shown to mimic the physiological role of Ser phosphorylation.

Found at doi:10.1371/journal.ppat.0030049.sg004 (21 KB PDF).

Table S1. Genes Differentially Regulated in Response to Overexpression of WT or Kinase-Dead PknD

For each gene at each time point, the value in the table represents the (averaged) ratio of medians (Rm) which has been normalized to the Rm at $t = 0$. Rm is defined as the median pixel intensity for the reference pool divided by the median pixel intensity of the sample. Genes depicted in this table exhibit at least a 2-fold difference in Rm between WT and kinase-dead samples at any time point. The data in the table were ordered by hierarchical clustering using Gene Cluster 3.0. N.D. indicates no data available. Genes under the control of SigF during log phase growth (e.g., *acpM*, *Rv1592c*, and *fabD*) were expressed at a lower level at $t = 0$ in the strain harboring the WT PknD expression plasmid compared to the kinase-dead PknD plasmid. These genes showed little fluctuation in transcriptional levels as WT PknD was overexpressed for 2, 8, or 24 h. In contrast, as expression of the kinase-dead PknD increased over time, these genes under SigF control during log phase growth were progressively repressed to levels similar to those in the strain expressing WT PknD. This pattern coincided with the leaky expression of active phosphorylated WT PknD at $t = 0$, while kinase activity was stimulated only after 4 h of kinase-dead PknD overexpression (Figure 1A). The idea that active WT PknD may have been signaling in uninduced cells is supported by the finding that the average Rm value for five representative SigF regulated genes (*acpM*, *Rv1592c*, *fabD*, *atpE* and *sodA*) at time zero in WT versus the Asp138Asn mutant strain was 1.9. By comparison, the mean and median Rm ratios at time zero for all genes were 1.07 and 0.96, respectively. These data suggest that some effects of WT PknD overexpression may have been manifest at $t = 0$ in these microarray experiments.

Found at doi:10.1371/journal.ppat.0030049.st001 (415 KB DOC).

Accession Numbers

Please see Table 2 for a list of genes described in this study.

Acknowledgments

We thank Kristi Pullen for providing PstP, Carl Mieczkowski for sharing unpublished observations about the specificity of the SP600125 inhibitor, T. Noelle Lombana for PknB, Christoph Grundner for PknK, Laurie Gay for PknE, and Janis Allen for preliminary microarray analysis. We thank Pei-Yi Sung, Brian Abe, and Danielle Tomkiel for assistance with cloning. We are indebted to David King of the HHMI mass spectrometry laboratory. We were

aided by the use of Gateway expression vectors provided by Bénédicte Delagoutte, Didier Busso, and Dino Moras.

Author contributions. AEG, JAM, JSC, and TA conceived and designed the experiments. AEG, JAM, CEB, and AMF performed the experiments. AEG, JAM, AMF, JSC, and TA analyzed the data. AEG, JAM, and AMF contributed reagents/materials/analysis tools. AEG, JAM, CEB, AMF, JSC, and TA wrote the paper.

Funding. This work was supported by National Institutes of Health grants to JSC and TA and a National Science Foundation Graduate Research Fellowship to AEG.

Competing interests. The authors have declared that no competing interests exist.

References

- Dye C, Scheele S, Dolin P, Pathania V, Ravigione MC (1999) Consensus statement. Global burden of tuberculosis: Estimated incidence, prevalence, and mortality by country. WHO Global Surveillance and Monitoring Project. *JAMA* 282: 677–686.
- Zhang Y (2004) Persistent and dormant tubercle bacilli and latent tuberculosis. *Front Biosci* 9: 1136–1156.
- Russell DG (2001) *Mycobacterium tuberculosis*: Here today, and here tomorrow. *Nat Rev Mol Cell Biol* 2: 569–577.
- Cole ST, Brosch R, Parkhill J, Garnier T, Churcher C, et al. (1998) Deciphering the biology of *Mycobacterium tuberculosis* from the complete genome sequence. *Nature* 393: 537–544.
- Av-Gay Y, Everett M (2000) The eukaryotic-like Ser/Thr protein kinases of *Mycobacterium tuberculosis*. *Trends Microbiol* 8: 238–244.
- Gay LM, Ng HL, Alber T (2006) A conserved dimer and global conformational changes in the structure of apo-PknE Ser/Thr protein kinase from *Mycobacterium tuberculosis*. *J Mol Biol* 360: 409–420.
- Good MC, Greenstein AE, Young TA, Ng HL, Alber T (2004) Sensor domain of the *Mycobacterium tuberculosis* receptor Ser/Thr protein kinase, PknD, forms a highly symmetric beta propeller. *J Mol Biol* 339: 459–469.
- Jones G, Dyson P (2006) Evolution of transmembrane protein kinases implicated in coordinating remodeling of gram-positive peptidoglycan: inside versus outside. *J Bacteriol* 188: 7470–7476.
- Grundner C, Gay LM, Alber T (2005) *Mycobacterium tuberculosis* serine/threonine kinases PknB, PknD, PknE, and PknF phosphorylate multiple FHA domains. *Protein Sci* 14: 1918–1921.
- Rajagopal L, Vo A, Silvestroni A, Rubens CE (2005) Regulation of purine biosynthesis by a eukaryotic-type kinase in *Streptococcus agalactiae*. *Mol Microbiol* 56: 1329–1346.
- Kang CM, Abbott DW, Park ST, Dascher CC, Cantley LC, et al. (2005) The *Mycobacterium tuberculosis* serine/threonine kinases PknA and PknB: Substrate identification and regulation of cell shape. *Genes Dev* 19: 1692–1704.
- Thakur M, Chakraborti PK (2006) GTPase activity of mycobacterial FtsZ is impaired due to its transphosphorylation by the eukaryotic type Ser/Thr kinase, PknA. *J Biol Chem* 281: 40107–40113.
- Sasseti CM, Boyd DH, Rubin EJ (2003) Genes required for mycobacterial growth defined by high density mutagenesis. *Mol Microbiol* 48: 77–84.
- Papavinasundaram KG, Chan B, Chung JH, Colston MJ, Davis EO, et al. (2005) Deletion of the *Mycobacterium tuberculosis* pknH gene confers a higher bacillary load during the chronic phase of infection in BALB/c mice. *J Bacteriol* 187: 5751–5760.
- Cowley S, Ko M, Pick N, Chow R, Downing KJ, et al. (2004) The *Mycobacterium tuberculosis* protein serine/threonine kinase PknG is linked to cellular glutamate/glutamine levels and is important for growth in vivo. *Mol Microbiol* 52: 1691–1702.
- Hughes KT, Mathee K (1998) The anti-sigma factors. *Annu Rev Microbiol* 52: 231–286.
- Kim TJ, Gaidenko TA, Price CW (2004) In vivo phosphorylation of partner switching regulators correlates with stress transmission in the environmental signaling pathway of *Bacillus subtilis*. *J Bacteriol* 186: 6124–6132.
- Parida BK, Douglas T, Nino C, Dhandayuthapani S (2005) Interactions of anti-sigma factor antagonists of *Mycobacterium tuberculosis* in the yeast two-hybrid system. *Tuberculosis (Edinb)* 85: 347–355.
- Delumeau O, Chen CC, Murray JW, Yudkin MD, Lewis RJ (2006) High-molecular-weight complexes of RsbR and paralogues in the environmental signaling pathway of *Bacillus subtilis*. *J Bacteriol* 188: 7885–7892.
- Mahenthiralingam E, Draper P, Davis EO, Colston MJ (1993) Cloning and sequencing of the gene which encodes the highly inducible acetamidase of *Mycobacterium smegmatis*. *J Gen Microbiol* 139: 575–583.
- Geiman DE, Kaushal D, Ko C, Tyagi S, Manabe YC, et al. (2004) Attenuation of late-stage disease in mice infected by the *Mycobacterium tuberculosis* mutant lacking the SigF alternate sigma factor and identification of SigF-dependent genes by microarray analysis. *Infect Immun* 72: 1733–1745.
- Kelley LA, MacCallum RM, Sternberg MJ (2000) Enhanced genome annotation using structural profiles in the program 3D-PSSM. *J Mol Biol* 299: 499–520.
- Boitel B, Ortiz-Lombardia M, Duran R, Pompeo F, Cole ST, et al. (2003) PknB kinase activity is regulated by phosphorylation in two Thr residues and dephosphorylation by PstP, the cognate phospho-Ser/Thr phosphatase, in *Mycobacterium tuberculosis*. *Mol Microbiol* 49: 1493–1508.
- Beaucher J, Rodrigue S, Jacques PE, Smith I, Brzezinski R, et al. (2002) Novel *Mycobacterium tuberculosis* anti-sigma factor antagonists control sigmaF activity by distinct mechanisms. *Mol Microbiol* 45: 1527–1540.
- Song T, Dove SL, Lee KH, Husson RN (2003) RshA, an anti-sigma factor that regulates the activity of the mycobacterial stress response sigma factor SigH. *Mol Microbiol* 50: 949–959.
- Greenstein AE, Echols N, Lombana TN, King DS, Alber T (2007) Allosteric activation by dimerization of the PknD receptor Ser/Thr protein kinase from *Mycobacterium tuberculosis*. *J Biol Chem*. E-pub 22 January 2007.
- Chen CC, Lewis RJ, Harris R, Yudkin MD, Delumeau O (2003) A supramolecular complex in the environmental stress signalling pathway of *Bacillus subtilis*. *Mol Microbiol* 49: 1657–1669.
- Chen CC, Yudkin MD, Delumeau O (2004) Phosphorylation and RsbX-dependent dephosphorylation of RsbR in the RsbR-RsbS complex of *Bacillus subtilis*. *J Bacteriol* 186: 6830–6836.
- Kim TJ, Gaidenko TA, Price CW (2004) A multicomponent protein complex mediates environmental stress signaling in *Bacillus subtilis*. *J Mol Biol* 341: 135–150.
- Greenstein AE, Grundner C, Echols N, Gay LM, Lombana TN, et al. (2005) Structure/function studies of Ser/Thr and Tyr protein phosphorylation in *Mycobacterium tuberculosis*. *J Mol Microbiol Biotechnol* 9: 167–181.
- Molle V, Brown AK, Besra GS, Cozzone AJ, Kremer L (2006) The condensing activities of the mycobacterium tuberculosis type II fatty acid synthase are differentially regulated by phosphorylation. *J Biol Chem* 281: 30094–30103.
- Molle V, Kremer L, Girard-Blanc C, Besra GS, Cozzone AJ, et al. (2003) An FHA phosphoprotein recognition domain mediates protein EmrB phosphorylation by PknH, a Ser/Thr protein kinase from *Mycobacterium tuberculosis*. *Biochemistry* 42: 15300–15309.
- Molle V, Soulat D, Jault JM, Grangeasse C, Cozzone AJ, et al. (2004) Two FHA domains on an ABC transporter, Rv1747, mediate its phosphorylation by PknF, a Ser/Thr protein kinase from *Mycobacterium tuberculosis*. *FEMS Microbiol Lett* 234: 215–223.
- Perez J, Garcia R, Bach H, de Waard JH, Jacobs WR Jr, et al. (2006) *Mycobacterium tuberculosis* transporter MmpL7 is a potential substrate for kinase PknD. *Biochem Biophys Res Commun* 348: 6–12.
- Cox JS, Chen B, McNeil M, Jacobs WR Jr. (1999) Complex lipid determines tissue-specific replication of *Mycobacterium tuberculosis* in mice. *Nature* 402: 79–83.
- Rodriguez GM, Voskuil MI, Gold B, Schoolnik GK, Smith I (2002) ideR, An essential gene in *Mycobacterium tuberculosis*: Role of IdeR in iron-dependent gene expression, iron metabolism, and oxidative stress response. *Infect Immun* 70: 3371–3381.
- Busso D, Delagoutte-Busso B, Moras D (2005) Construction of a set Gateway-based destination vectors for high-throughput cloning and expression screening in *Escherichia coli*. *Anal Biochem* 343: 313–321.
- Hartley JL, Temple GF, Brasch MA (2000) DNA cloning using in vitro site-specific recombination. *Genome Res* 10: 1788–1795.