

Lawrence Berkeley National Laboratory

Recent Work

Title

SHELL MODEL CALCULATION OF Po212

Permalink

<https://escholarship.org/uc/item/7w9299fm>

Authors

Glendenning, Norman K.
Harada, Kichinosuke.

Publication Date

1964-04-01

0 0 0 9 2 3 0 2 4 9 6

UCRL 11361
Abstract

UNIVERSITY OF
CALIFORNIA

*Radiation
Laboratory*

FOR REFERENCE

NOT TO BE TAKEN FROM THIS ROOM

BERKELEY, CALIFORNIA

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Shell Model Calculation of Po^{212} . * Norman K. Glendenning and Kichinosuke Harada[†],
Lawrence Radiation Laboratory, Berkeley.--So far there are a few shell model calculations¹⁻³ on Po^{212} , but unfortunately their results are more or less different from each other, owing to the use of different residual forces. A more realistic shell model calculation based on Glendenning's method is done for the lower lying levels and the high-spin isomer of Po^{212} . In this calculation we use the residual p-p, n-n and p-n interactions which are so determined as to reproduce the experimental Po^{210} , Pb^{210} and Bi^{210} spectra, and adopt different size parameters for proton- and neutron-oscillator potentials. The results will be discussed and compared with those of the previous works.

* Work performed under the auspices of the U. S. Atomic Energy Commission.

[†] On leave of absence from Japan Atomic Energy Research Institute.

¹ I. M. Band, L. A. Sliv and Yu, I. Kharitonov, Nuclear Physics 35, 136 (1962).

² N. K. Glendenning, Phys. Rev. 127, 923 (1962).

³ K. Harada, JAERI-Memo No. 1183 (1963).