

UC San Diego

UC San Diego Previously Published Works

Title

Asian Cohort for Alzheimer Disease (ACAD) Pilot Study: Vietnamese Americans.

Permalink

<https://escholarship.org/uc/item/80r819ns>

Journal

Alzheimer Disease and Associated Disorders, 38(3)

Authors

Peavy, Guerry

Võ, Namkhue

Revta, Carolyn

et al.

Publication Date

2024-07-01

DOI

10.1097/WAD.0000000000000631

Copyright Information

This work is made available under the terms of a Creative Commons Attribution-NonCommercial-NoDerivatives License, available at

<https://creativecommons.org/licenses/by-nc-nd/4.0/>

Peer reviewed

Asian Cohort for Alzheimer Disease (ACAD) Pilot Study

Vietnamese Americans

Guerry M. Peavy, PhD,*† Namkhuê Võ, BS,* Carolyn Revta, MPH,*
 Anna T. Lu, BA,* Jody-Lynn Lupo, MS,* Percival Nam, BS,‡
 Khải H. Nguyễn, MD, MHS,§ Li-San Wang, PhD,|| and
 Howard H. Feldman, MD*†

Introduction: The objective of this pilot study was to establish the feasibility of recruiting older Vietnamese Americans for research addressing genetic and nongenetic risk factors for Alzheimer disease (AD).

Methods: Twenty-six Vietnamese Americans were recruited from communities in San Diego. A Community Advisory Board provided cultural and linguistic advice. Bilingual/bicultural staff measured neuropsychological, neuropsychiatric, lifestyle, and medical/neurological functioning remotely. Saliva samples allowed DNA extraction. A consensus team reviewed clinical data to determine a diagnosis of normal control (NC), mild cognitive impairment (MCI), or dementia. Exploratory analyses addressed AD risk by measuring subjective cognitive complaints (SCC), depression, and vascular risk factors (VRFs).

Results: Twenty-five participants completed the study (mean age = 73.8 y). Eighty percent chose to communicate in Vietnamese. Referrals came primarily from word of mouth within Vietnamese communities. Diagnoses included 18 NC, 3 MCI, and 4 dementia. Participants reporting SCC acknowledged more depressive symptoms and had greater objective cognitive difficulty than those without SCC. Eighty-eight percent of participants reported at least 1 VRF.

Discussion: This pilot study supports the feasibility of conducting community-based research in older Vietnamese Americans. Challenges included developing linguistically and culturally appropriate

cognitive and neuropsychiatric assessment tools. Exploratory analyses addressing nongenetic AD risk factors suggest topics for future study.

Key Words: Vietnamese Americans, genetic diversity, Alzheimer, recruitment, subjective cognitive complaints, vascular risk factors

(*Alzheimer Dis Assoc Disord* 2024;38:277–284)

Addressing the underrepresentation of older Asian Americans in Alzheimer disease (AD) research is critical given the significant number of Asian Americans over the age of 65 (~2.5 million)¹ and the importance of age as a risk factor for AD.² In addition, Asian Americans, Native Hawaiians, and Pacific Islanders are often combined into one category that includes >40 ethnic subgroups.³ Disaggregating the larger Asian population into subgroups is essential, given significant demographic, cultural, linguistic, and socioeconomic differences. Increasing knowledge concerning healthcare and health disparities will depend on the “unmasking” of these differences, including the pursuit of information through studies of risk factors for AD and related dementias (ADRD).

The Asian Cohort for Alzheimer Disease (ACAD) investigates genetic and lifestyle factors that contribute to AD risk in Chinese, Korean, and Vietnamese populations across multiple sites in the United States and Canada. In this report, we present the results of a pilot study to determine the feasibility of conducting a more extensive investigation of genetic and lifestyle risk factors in Vietnamese Americans.

Vietnamese Americans number almost 2.4 million individuals either born in Vietnam or reporting Vietnamese ancestry. The number of Vietnamese Americans over the age of 65 is estimated at 23%,⁴ with 19% foreign-born. Their recruitment into AD research is critical to define phenotypes,⁵ identify differences in AD risk factors,⁶ and provide specific information about normal aging. There are, however, multiple factors that limit our ability to address the health needs of older Vietnamese Americans through research. These include resistance to research participation resulting from limited health literacy, stigma associated with cognitive decline and dementia, and mistrust in interactions with research personnel. In addition, there are very few neuropsychological and neuropsychiatric instruments that are culturally adapted, appropriately translated, and validated for Vietnamese cohorts.

Stress from past traumatic experiences and limited acculturation may reduce the degree to which Vietnamese

Received for publication April 28, 2024; accepted July 8, 2024.

From the *Department of Neurosciences, Alzheimer’s Disease Cooperative Study; †Department of Neurosciences, Alzheimer’s Disease Research Center; ‡Medical Student Y3, UC San Diego School of Medicine; §Division of Geriatrics, Gerontology, and Palliative Care, UC San Diego School of Medicine, San Diego, CA; and ||Department of Pathology and Laboratory Medicine, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA.

The study was supported by a supplement from the University of Pennsylvania Department of Pathology and Laboratory Medicine and National Institute on Aging Grants R56 AG069130 and U19 AG079774. Bio-samples collected in this study are deposited at the National Centralized Repository for Alzheimer’s Diseases and Related Dementias (NCRAD), which receives government support under a cooperative agreement grant U24 AG021886. Finally, the study was supported in part by the National Institute on Aging grant P30-AG062429 to the Shiley-Marcos Alzheimer’s Disease Research Center.

The authors declare no conflicts of interest.

Reprints: Guerry M. Peavy, PhD, Department of Neurosciences, Alzheimer’s Disease Cooperative Study, University of California San Diego, 9500 Gilman Drive, MC 0949, La Jolla, CA 92093-0949 (e-mail: gpeavy@health.ucsd.edu).

Copyright © 2024 The Author(s). Published by Wolters Kluwer Health, Inc. This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

DOI: 10.1097/WAD.0000000000000631

immigrants seek personal interactions, leaving them without a sense of belonging. In a study by Sorkin et al,⁷ Vietnamese Americans were more likely to report a need for mental health services than non-Hispanic Whites. Tran et al⁸ found that during the first 10 years of acculturation in the United States, Vietnamese Americans (ages 18 to 73 y) experienced more symptoms of depression than those with longer US residence and that for many, these symptoms were related to “persistent psychological distress, economic challenges, language barriers, and social isolation.” Compared with 5 groups of older Asian American immigrants, Vietnamese participants reported the second greatest incidence of depression, possibly associated with the perception of a cultural disconnect with their adult children⁹ and/or the subjective perception of cognitive decline.^{10,11}

Considering Vietnamese Americans of all ages, it is estimated that 35% are proficient in English.¹² For older adults, even fewer are English proficient, creating a potential barrier to their participation in ADRD research. In addition, differences in educational level and geographically determined language customs may influence the examiner's choice of words when conversing with Vietnamese Americans. Language barriers also impact the selection of cognitive assessment instruments necessary for accurate diagnoses in ADRD research. Dick et al¹³ developed the Cross-Cultural Neuropsychological Test Battery (CCNB) to measure cognition in 5 ethnic groups, including Vietnamese. Similarly, Kempler et al¹⁴ developed the Common Objects Memory Test (COMT) to assess learning and memory in these same 5 groups. They found that scores were influenced by age but relatively unaffected by education and ethnic group.

Subjective cognitive complaints (SCC) have been identified as a potential AD risk factor.¹⁵ Meyer et al¹¹ found that of 600 Vietnamese older adults living in communities in Da Nang, Vietnam, 64% complained of memory decline, with 39% reporting memory loss that interfered with activities of daily living (ADLs). These complaints were associated with both objective cognitive impairment and depressive symptoms. In another study, Lee et al¹⁰ found that SCCs reported by Vietnamese Americans were associated with depressive symptoms, but the study did not include objective cognitive measures.

The role of comorbid brain pathologies in aging and their association with cognitive impairment in older adults have been widely studied and found to relate to midlife vascular risk factors (VRFs).¹⁶ Boyle et al¹⁷ emphasized the added contribution of cerebrovascular pathologies to cognitive decline associated with aging and their interplay with neurodegenerative markers of AD pathophysiology, including amyloid- β . Nguyen et al¹⁸ reported a high prevalence of hypertension and a high rate of cerebrovascular mortality in Vietnamese. The co-occurrence of cerebrovascular disease and AD is common. A potentially causal relationship between the 2 as well as shared genetic risk¹⁹ may be relevant. Current data are typically reported only for Asian aggregates or not reported,²⁰ highlighting the need for better VRF characterization in Vietnamese Americans.

ACAD is the first large cohort recruited to examine genetic and lifestyle factors associated with risk for AD among older Asian Americans and Canadians. To date, 8 ACAD sites have recruited Chinese, Korean, or Vietnamese community participants, with coordination by the University of Pennsylvania (Principal investigator: L.-S.W.).²¹

ACAD sites have collected cognitive, neuropsychiatric, medical, and lifestyle information, as well as biosamples of blood or saliva for genetic analysis. A consensus team within each site has provided a diagnosis of normal control (NC), mild cognitive impairment (MCI), or dementia related to AD or another neurodegenerative disorder. In addition, a randomized external consensus review by 2 neurologists has been performed to ensure harmonization of diagnoses across all sites. A larger NIA-funded study to expand the sample size and pursue additional Asian American/Canadian subgroups is underway.

Here, we describe the ACAD pilot study focused on a cohort of older Vietnamese Americans enrolled at the University of California San Diego (UCSD) site. The overall goal was to establish feasibility by:

- (1) Developing strategies to engage members of the Vietnamese community and build trust in ACAD study personnel and research methods.
- (2) Recruiting and enrolling participants who agreed to procedures, including cognitive assessment and biosample collection for genetic testing.
- (3) Identifying and implementing cognitive and neuropsychiatric assessment tools with appropriate translations and cultural adaptations.
- (4) Exploring potential risk factors for cognitive decline, including subjective cognitive complaints, depressive symptomatology, and VRFs.

METHODS

Participants

Consenting participants who self-reported Vietnamese ancestry and were over the age of 60 were recruited from the community. Exclusion criteria included a non-neurodegenerative central nervous system disease, primary psychiatric disorder, or known major medical illness that could cause cognitive dysfunction. Participants could choose to communicate and complete study assessments in either English or Vietnamese.

Procedures

Participants were enrolled following the consent process that included a written informed consent approved by the UCSD Office of IRB Administration. Individuals directed questions about the consent to a well-informed bilingual member of the study team. If a participant was considered to lack capacity due to cognitive impairment, a legally authorized representative signed the consent with assent from the participant. All study procedures were conducted remotely using a customized electronic tablet programmed for ease of use by older adults. The tablet was delivered to the participant in his/her home or facility with instructions for a one-button connection with the research examiner who was available to discuss with the participant any questions or problems about the technology. Remote administration using the tablet allowed the participant to avoid potential exposure to illness during the COVID pandemic, participate in the research in a familiar environment, and avoid the burden of travel to a testing site.

Developing translations from English into Vietnamese (eg, outreach materials) was completed using a multistep approach to ensure translated materials were reliable, linguistically correct, and culturally appropriate. Bilingual individuals involved in the translation process included research team members, Community Advisory Board

(CAB) members, and professional translators. This multi-step process included an initial translation from English to Vietnamese. These were subsequently reviewed and verified to ensure that the translation done by the initial translator captured the meaning of the original/English instruments and was grammatically correct at an appropriate literacy level and culturally relevant. Finally, a certified service reviewed and certified the translations, a requirement of the local site Institutional Review Board.

The ACAD Outreach workgroup designed outreach materials describing the objectives and procedures of the study. Bicultural/bilingual research personnel, UCSD students with Vietnamese ancestry, and a CAB of trusted community leaders familiar with local Vietnamese communities contributed information concerning sociocultural customs and language, including existing stigmatized concepts (eg, dementia). The UCSD study team and community partners distributed the English and Vietnamese outreach materials to engage older adults in geographically targeted Vietnamese communities in San Diego County. Three community partners (ie, San Diego Vietnamese Association, Asian Pacific Health Foundation, UDW Home Care Providers Union) assisted in efforts to support research participation.

Bicultural/bilingual members of the UCSD research team provided potential participants with a description of study procedures to assess interest, prescreened individuals to ensure eligibility, and facilitated participant involvement to build trust. For physician medical examinations, a staff member or student fluent in Vietnamese provided translations.

The study did not specifically investigate potential safety issues. However, if the study team developed concerns about suicidality, unsafe driving, or elder abuse during the evaluation, specific steps outlined in the consent for such circumstances were followed. These included mandatory reporting or direction to appropriate resources for further assessment and management.

Data Collection Packets

There were 3 Data Collection Packets (DCPs). Part A included questions concerning demographics and diet (Mediterranean Diet Assessment Scale (MEDAS)),^{22,23} as well as the Clinical Dementia Rating Scale (CDR).²⁴ Part B included questions about potentially enriching physical, cultural, and cognitive activities (eg, reading, music lessons)

throughout the life span (Rush Early Life Enrichment Inventory).²⁵

Part C consisted of neuropsychological, psychiatric, functional, and medical assessments. Neuropsychological tests were chosen, in part, from previously validated cross-cultural instruments, with others adopted to test skills (eg, memory) important for ADRD diagnostic considerations. Neuropsychological tests included the Cognitive Abilities Screening Instrument (CASI)^{13,26} and the Common Objects Memory Test (COMT).¹⁴ Additional tests included the Clock Drawing Test²⁷ and category fluency (ie, animals, vegetables). Cognitively normal and mildly impaired participants^{28–30} completed the 15-item Short Form of the Geriatric Depression Scale (GDS).³¹ The Functional Assessment Questionnaire (FAQ)^{32,33} measured independence in instrumental ADLs. The clinical evaluation included medical history, health status, current medications, and a remotely conducted neurological examination. At consensus conferences following evaluations, team members used all data (ie, cognitive, behavioral, functional, medical) and informant reports (when available) to determine the diagnostic category (ie, NC, MCI, or dementia).

Statistical Analyses

Quantitative data included descriptive statistics as well as independent sample *t* tests and Levene test for equality of variance to determine differences between groups on continuous variables and differences in variance, respectively. Standardized effect sizes (Cohen *d*) were calculated to describe the magnitude of differences between groups.^{34,35} IBM SPSS, version 28.0.1.1 was used for analyses.

RESULTS

Twenty-six Vietnamese Americans enrolled in the study between July 2021 and April 2022. Data from 1 participant who withdrew consent was removed from the database and destroyed. The remaining 25 participants completed the study with adherence to all procedures. Of the 25 completers, 14 were female, and 11 male. Age ranged from 61 to 91 years, and education, from 3 to 20 years (Table 1). All participants were born in Vietnam and arrived in the United States between 1964 and 2014, with age of arrival ranging from 19 to 64 years. Eighty percent chose to complete study procedures in Vietnamese.

Although multiple strategies were used for recruitment³⁶ (Fig. 1), word of mouth within the

TABLE 1. Demographic/Clinical Variables for Total Cohort and Cohort Divided by Consensus Diagnosis (NC, MCI, and Dementia)

	Total sample (N = 25) [mean (SD)]	Consensus diagnosis [mean (SD)]		
		NC (n = 18)	MCI (n = 3)	Dementia (n = 4)
Age at testing (y)	73.8 (9.2)	70.7 (7.3)	75.3 (9.3)	87.0 (4.5)
Education (y)	12.6 (4.3)	13.1 (3.5)	17.3 (2.3)	7.3 (4.0)
Geriatric Depression Scale /15	3.9 (3.4)	4.2 (3.7)	2.3 (2.5)	NA
FAQ/30	—	1.4 (3.3)	1.7 (1.5)	29.3 (1.5)
Duration of illness	—	NA	2.0 (1.0)	11.3 (8.7)
Age of arrival in the United States	40.3 (14.4)	38.5 (12.8)	30.0 (17.3)	56.3 (8.4)
Years in the United States	33.5 (11.5)	32.2 (11.7)	45.3 (12.0)	30.8 (3.9)
Female (%)	56	61	0	75
Born in Vietnam (%)	100	100	100	100
Year of arrival in the United States (range)	1964–2014	1967–2014	1964–1989	1985–1994
Tested in Vietnamese (%)	80	83	33	100

FAQ indicates Functional Assessment Questionnaire; MCI, mild cognitive impairment; NC, normal control.




FIGURE 1. Recruitment and retention strategies for the Asian Cohort for Alzheimer Disease (ACAD) Vietnamese pilot study.

Vietnamese community accounted for most of the referrals eligible for enrollment. These included family, friends, neighbors, other ACAD participants, and members of the CAB. For many of the enrollees, a combination of referral sources led to their participation, but the CAB members were cited as at least one of the sources that influenced 11 individuals (44%) to participate.

Following the consensus diagnosis for 25 study completers, 18 received a diagnosis of NC, 3 MCI, and 4 dementia. Three of the 4 with dementia were severely cognitively impaired and unable to perform neuropsychological tests. The diagnosis was based on information provided by the informant. All 4 participants with dementia were further classified as having AD. Two of the 3 participants diagnosed with MCI were classified as single domain amnesic, and one, as executive type. Those with dementia were older than the NC participants ($t = -4.3$; $P < 0.001$) and less educated than those in both the NC ($t = 2.9$; $P = 0.004$) and MCI ($t = 3.8$; $P = 0.012$) groups. Three participants either refused or were unable to provide a saliva sample for DNA extraction.

Exploratory analyses focused on 3 factors associated with AD risk: subjective cognitive complaints (SCC), depressed mood, and vascular symptoms. There were no significant differences in age and education between the NC participants with and without SCC (Table 2). Based on medium to large effect sizes (0.59 to 0.79), those with SCC compared with those without SCC reported greater GDS and FAQ scores and had greater difficulty on a simple figure copy task and learning and recall of visually presented common objects.

The mean total GDS score fell within the normal range²⁸ and did not significantly differentiate NCs from MCIs. Females acknowledged more symptoms of depression than males ($t = -2.4$; $P = 0.03$) and showed significantly greater variability in total GDS scores. With a median age split of < 70 versus ≥ 70 years, the younger group (mean age = 65.3 y) reported more symptoms of depression ($t = 2.1$; $P = 0.059$; effect size = 0.88) and showed significantly greater variability ($F = 13.5$; $P = 0.002$) than the older group (mean age = 78.0 y). There was no difference in overall score on the GDS between those reporting less English proficiency and those reporting more; however, those who acknowledged a highly endorsed item on the GDS, a preference to stay home rather than going out and doing new things (48%), rated

their English skills less proficient than those who did not ($t = -2.5$; $P = 0.024$). For those who reported being afraid that something bad was going to happen to them (48%), the total GDS score was higher ($t = 2.2$; $P = 0.063$; effect size = 0.94) and more variable ($F = 14.0$; $P = 0.001$), and age was younger ($t = 2.7$; $P = 0.014$).

Eighty-eight percent (88%) of the 25 participants reported at least 1 VRF (ie, hypertension, hypercholesterolemia, diabetes mellitus). Five reported all 3 VRFs. Fifty-seven percent of the participants with MCI or dementia (4 of 7) reported 2 or 3 VRFs; this was true for only 33% (6 of 18) of NCs. When comparing subgroups on neuropsychological tests, we restricted the comparison of individuals with fewer (0 or 1) versus more (2 or 3) VRFs to NC participants to avoid the diagnostic confound due to circularity and found that, based on effect size, those with fewer VRFs performed better on visuospatial tasks (ie, CASI pentagons, clock drawing) than those with more (Table 3).

DISCUSSION

This UCSD ACAD pilot study demonstrated the feasibility of bringing Vietnamese Americans into scientific research designed to address genetic and lifestyle risk factors for AD. The study achieved a high participant completion rate by engaging and recruiting older adults from the Vietnamese community, a population that is significantly underrepresented in dementia research.

To succeed in this pilot study, we faced challenges and unknowns around recruitment success, the utility of the instruments used to measure variables across a range of severity, and the added task presented by COVID-19 to conduct clinical evaluations using remote technology. The pilot study incorporated multiple engagement and recruitment strategies, with word of mouth within the Vietnamese community being the most effective in reaching recruitment goals. Building trust was significantly enhanced by the inclusion of a Vietnamese advisory board and individuals from local Asian community organizations. Bicultural/bilingual research staff involvement in the local Vietnamese community, as well as the UCSD academic environment, allowed completion of screening and research procedures that emphasized sensitivity to cultural preferences and values of older Vietnamese adults within a research setting. Using a simplified remote technology platform with

TABLE 2. Demographic and Clinical Variables for NC Participants (N = 18) Divided by SCC

	NC without SCC (n = 7) [mean (SD)]	NC with SCC (n = 11) [mean (SD)]	Effect size (Cohen <i>d</i>)
Age (y)	69.9 (8.6)	71.2 (6.7)	−0.18
Education (y)	13.7 (2.4)	12.6 (4.1)	−0.31
Geriatric Depression Scale/15	2.9 (1.6)	5.0 (4.4)*	−0.60+
FAQ/30	0.1 (0.4)	2.2 (4.1)	−0.63+
Neuropsychological tests			
CASI total/100	91.8 (5.5)	91.5 (4.8)	0.05
CASI pentagons/10	9.9 (0.4)	9.3 (1.0)*	0.70+
Clock to command/15	13.6 (1.3)	12.5 (3.2)	0.43
Common Objects Memory Test			
Learning Trial 1 # recalled/10	6.9 (1.2)	6.6 (1.4)	−0.79+
Learning Trial 2 # recalled/10	8.1 (1.6)	7.2 (1.5)	−0.35
Learning Trial 3 # recalled/10	8.7 (1.1)	8.0 (1.1)	−0.34
Sum of Learning Trials/30	23.7 (2.9)	21.8 (3.4)	0.60+
Short Delay Free Recall/10	8.6 (1.0)	7.7 (1.4)	0.66+
Short Delay Recognition/20	20.0 (0.0)	20.0 (0.0)	—
Animal Fluency, # correct	11.3 (3.1)	10.9 (3.5)	0.11
Vegetable Fluency, # correct	9.6 (4.0)	12.7 (4.2)	−0.77+

*Significant difference in variance.

+Medium to large effect size.

CASI indicates Cognitive Abilities Screening Instrument; FAQ, Functional Assessment Questionnaire; NC, normal control; SCC, subjective cognitive complaint.

translations by bilingual staff enabled completion of well-received assessments. Eighty percent of participants in the pilot study chose to communicate in Vietnamese, a finding somewhat expected given that all participants were foreign-born, older Vietnamese Americans. Providing this choice for language was an essential factor for study recruitment and retention as well as trust-building.

Language had a significant impact on other aspects of the study as well. The selection and adaptation of linguistically and culturally appropriate translations of instructions and test items into Vietnamese resulted in the successful completion of these instruments in all but the 3 participants with severe dementia. The availability of this information, as well as the multidisciplinary, bicultural nature of the consensus team, strengthened the diagnostic process. However, the results underscored the need to

validate cognitive measures for assessment of participants with more advanced dementia with instruments that might include existing versions of the Severe Cognitive Impairment Profile (SCIP)³⁷ or the Severe Impairment Battery (SIB).³⁸ Finally, regardless of cognitive status, the modest number of validated test instruments available for use in the older Vietnamese American population will require significant instrument development to benefit future ADRD studies.

This pilot study has several limitations. While recruitment of a small sample size succeeded, a much larger sample enrollment may present challenges that necessitate additional approaches to recruitment. The pilot sample size supports only preliminary exploratory analyses of non-genetic AD risk factors. The findings will only be confirmed with the larger sample planned in the forthcoming NIH-funded U19 study titled “Asian Cohort for Alzheimer

TABLE 3. Normal Control Participants (N = 18) Divided by Number of Vascular Risk Factors (Hypertension, Diabetes, Hypercholesterolemia)

	No. vascular risk factors [mean (SD)]		Effect size (Cohen <i>d</i>)
	0 or 1 (n = 12)	2 or 3 (n = 6)	
Age (y)	68.4 (5.9)	75.2 (8.2)	−1.01
Education (y)	13.0 (4.0)	13.2 (2.2)	−0.05
Geriatric Depression Scale/15	4.3 (3.6)	4.0 (4.0)	0.07
Functional Assessment Scale/30	1.5 (4.0)	1.2 (1.6)	0.10
Neuropsychological tests			
CASI total/100	92.3 (5.2)	90.3 (4.6)	0.41
CASI pentagons/10	9.7 (0.7)	9.2 (1.2)	0.59
Clock to command/15	13.7 (1.1)	11.3 (4.0)*	0.96
Common Objects Memory Test			
Sum of Learning Trials/30	22.7 (2.9)	22.3 (4.2)	0.10
Short Delay Free Recall/10	7.9 (1.2)	8.3 (1.6)	−0.31
Animal Fluency, # correct	10.8 (3.4)	11.5 (3.3)	−0.20
Vegetable Fluency, # correct	11.3 (4.5)	12.0 (4.1)	−0.17
Female (%)	72.7	27.3	
Male (%)	57.1	42.9	

*Significant difference in variance.

CASI indicates Cognitive Abilities Screening Instrument.

Disease” (ACAD). The results, however, encourage that studies such as ACAD can be undertaken and can apply to a broad range of ages and years of education of foreign-born Vietnamese Americans. The finding that sufficient saliva samples for DNA analysis were not obtained from 12% of participants underscores some of the technical difficulties obtaining sufficient saliva in older participants in a remote home setting, as well as potential hesitation to consent to provide DNA. Also, consistent with other studies, the majority of participants in the cohort received a diagnosis of NC, highlighting the anticipated need to recruit larger numbers of cognitively impaired individuals to provide sufficient AD cases for genetic analysis with attention to age stratification. Finally, remote neuropsychological testing using an electronic tablet for video conferencing can limit the willingness of older adults to enroll in the study due to discomfort with technology. Remote testing can also limit confidence in the research data due to unproven reliability and validity of the instruments administered. However, the tablet we employed was customized for ease of use by older adults and was supported by the research staff, so technical problems were addressed before the assessment. Several survey studies^{39–41} have found that participants consider testing using remote audio-visual technologies to actually be user-friendly. In addition, recent studies have found remote neuropsychological tests that include measures of memory to be reliable and valid when compared with face-to-face assessments in a variety of research samples including: (1) older adults,^{41–46} (2) individuals with low education,⁴⁷ (3) rural/underserved populations,^{41,45} and (4) diverse groups.^{46,48}

Our preliminary exploratory analyses evaluated 3 potential AD risk factors, subjective cognitive complaints, depression, and vascular disorders. Our finding of 61% of normal controls reporting SCC, is consistent with findings from previous studies in which 64% of Vietnamese reported SCC,¹¹ and Vietnamese Americans reported more cognitive complaints compared with other groups (eg, African Americans, Chinese Americans).¹⁰ In addition, our findings that participants with SCC compared with those without reported more depressive symptoms and were more impaired on aspects of objective cognitive testing were consistent with results from the study by Meyer et al.¹¹

Depressive symptoms in the Vietnamese American population have been attributed to a variety of factors including economic challenges, language barriers, and social isolation.⁸ For older adults, a sense of cultural disconnect with adult children⁹ and the subjective perception of cognitive decline^{10,11} may also be factors. English proficiency did not make a difference in the report of depression symptomatology. However, those acknowledging a highly endorsed specific item on the GDS concerning a preference to stay home rated their English skills less proficient than those who did not. Comparing means, women reported more symptoms of depression than men, and younger individuals (below 70 y) reported more symptoms of depression than older participants. However, while GDS scores for the majority of participants fell within the normal range, the results showing differences in groups divided by sex and by age were likely driven by the scores of 4 younger women whose scores suggested clinically relevant depression. These results, the high likelihood that Vietnamese Americans have suffered from trauma, acculturation stress, and discrimination, and findings from previous studies^{7–11,49} support the need to address depression and other mental

health issues in Vietnamese Americans regardless of demographic characteristics.

Our finding that 88% of participants reported at least 1 VRF (ie, hypertension, high cholesterol, diabetes), and 40%, more than one, is consistent with observations reported by Nguyen et al¹⁸ that Vietnamese Americans have a significant VRF burden and also a high rate of cerebrovascular mortality. While the number of VRFs in our study increased with age, treatment in midlife compared with later life should be considered for impact on dementia risk.¹⁶ Other VRFs including access to care and smoking were not surveyed in this study but are important for investigation in future studies.^{18,50}

CONCLUSIONS

In general, the pilot study identified successful strategies for developing relationships with community partners, building trust with older adult Vietnamese American participants for research recruitment and retention, providing effective, linguistically and culturally appropriate language translations, carrying out technology-friendly remote assessments, and providing sufficient information for determining a diagnosis by a team of patient-facing raters and experienced dementia professionals. These support the feasibility of a much larger study to determine genetic and nongenetic risk factors for AD in a population whose underrepresentation in research is a significant concern. Funding has been approved for a multisite study (NIA ACAD U19: PI: L.-S.W.)²¹ designed to unmask differences in largely unexplored genetic profiles among Vietnamese Americans and other Asian subgroups (ie, Korean and Chinese Americans and Canadians) and in comparison to populations overrepresented in research (eg, Caucasians). The upcoming study will provide much larger sample sizes to aggregate and disaggregate North American Asian American and Canadian populations.

ACKNOWLEDGMENTS

The authors are extremely grateful to the research participants who volunteered their time and effort to make this study possible. They also thank the members of our local Vietnamese Community Advisory Board and 3 community partners, the San Diego Vietnamese Association, the Asian Pacific Health Foundation (APHF), and the UDW Home Care Providers Union for their support concerning linguistic cultural issues and help with participant recruitment.

REFERENCES

1. Wu B, Qi X. Addressing health disparities among older Asian American populations: research, data, and policy. *Public Policy Aging Rep.* 2022;32:105–111.
2. Larson EB, Kukull WA, Katzman RL. Cognitive impairment: dementia and Alzheimer's disease. *Annu Rev Public Health.* 1992;13:431–449.
3. Kanaya AM, Hsing AW, Panapasa SV, et al. Knowledge gaps, challenges, and opportunities in health and prevention research for Asian Americans, Native Hawaiians, and Pacific Islanders: a report from the 2021 National Institutes of Health Workshop. *Ann Intern Med.* 2022;175:574–589.
4. Harjanto L, Batalova J. Vietnamese Immigrants in the United States. Migration Information Source; 2023. Accessed October 11, 2023. <https://www.migrationpolicy.org>
5. Greene AS, Shen X, Noble S, et al. Brain-phenotype models fail for individuals who defy sample stereotypes. *Nature.* 2022; 609:109–118.

6. Campos M, Edland SD, Peavy GM. An exploratory study of APOE-ε4 genotype and risk of Alzheimer's disease in Mexican Hispanics. *J Am Geriatr Soc.* 2013;61:1038.
7. Sorkin D, Tan AL, Hays RD, et al. Self-reported health status of Vietnamese and non-Hispanic white older adults in California. *J Am Geriatr Soc.* 2008;56:1543–1548.
8. Tran TV, Manalo V, Nguyen VT. Nonlinear relationship between length of residence and depression in a community-based sample of Vietnamese Americans. *Int J Soc Psychiatry.* 2007;53:85–94.
9. Mui AC, Kang S-Y. Acculturation stress and depression among Asian immigrant elders. *Soc Work.* 2006;51:243–255.
10. Lee M, Nam J-H, Yi E, et al. Differences in subjective memory impairment, depressive symptoms, sleep, and physical activity in African American and Asian American elderly. *Brain Sci.* 2021;11:1155.
11. Meyer OL, Leggett A, Liu S, et al. Prevalence and correlates of subjective memory complaints in Vietnamese adults. *Int Psychogeriatr.* 2018;30:1039–1048.
12. Budiman A Vietnamese in the US Fact Sheet. Pew Research Center; 2021. Accessed April 29, 2021. <https://www.pewresearch.org/social-trends/fact-sheet/asian-americans-vietnamese-in-the-u-s-fact-sheet/>
13. Dick MB, Teng EL, Kempler D, et al. The Cross-Cultural Neuropsychological Test Battery (CCNB): effects of age, education, ethnicity, and cognitive status on performance. Ferraro FR, ed. *Studies on Neuropsychology, Development, and Cognition: Minority and Cross-Cultural Aspects of Neuropsychological Assessment.* Swets & Zeitlinger Publishers; 2002: 17–41.
14. Kempler D, Teng EL, Taussig M, et al. The common objects memory test (COMT): a simple test with cross-cultural applicability. *J Int Neuropsychol Soc.* 2010;16:537–545.
15. Mitchell AJ, Beaumont H, Ferguson D, et al. Risk of dementia and mild cognitive impairment in older people with subjective memory complaints: meta-analysis. *Acta Psychiatr Scand.* 2014; 130:439–451.
16. DeBette S, Seshadri S, Beiser A, et al. Midlife vascular risk factor exposure accelerates structural brain aging and cognitive decline. *Neurology.* 2011;77:461–468.
17. Boyle PA, Wang T, Yu L, et al. To what degree is late life cognitive decline driven by age-related neuropathologies? *Brain.* 2021;144:2166–2175.
18. Nguyen RT, Meyer O, Chu J, et al. Social determinants of health, cardiovascular risk factors, and atherosclerotic cardiovascular disease in individuals of Vietnamese origin. *Am J Cardiol.* 2023;189:11–21.
19. Laing KK, Simoes S, Baena-Caldas GP, et al. Cerebrovascular disease promotes tau pathology in Alzheimer's disease. *Brain Commun.* 2020;2:fcaa132.
20. Doãn LN, Takata Y, Sakuma KK, et al. Trends in clinical research including Asian American, Native Hawaiian, and Pacific Islander participants funded by the US National Institutes of Health, 1992 to 2018. *JAMA Netw Open.* 2019;2:e197432.
21. Ho PC, Yu WH, Tee BL, et al. Asian cohort for Alzheimer's Disease (ACAD) pilot study on genetic and non-genetic risk factors for Alzheimer's disease among Asian Americans and Canadians. *Alzheimers Dement.* 2024;20:2058–2071.
22. Schröder H, Fitó M, Estruch R, et al. A short screener is valid for assessing Mediterranean diet adherence among older Spanish men and women. *J Nutr.* 2011;141:1140–1145.
23. Martínez-González MA, García-Arellano A, Toledo E, et al. A 14-item Mediterranean diet assessment tool and obesity indexes among high-risk subjects: the PREDIMED trial; 2012.
24. Morris JC. The Clinical Dementia Rating (CDR): current version and scoring rules. *Neurology.* 1993;43:2412–2414.
25. Landau SM, Marks SM, Mormino EC, et al. Association of lifetime cognitive engagement and low β-amyloid deposition. *Arch Neurol.* 2012;69:623–629.
26. Teng EL, Hasegawa K, Homma A, et al. The Cognitive Abilities Screening Instrument (CASI): a practical test for cross-cultural epidemiological studies of dementia. *Int Psychogeriatr.* 1994;6:45–58; discussion 62.
27. Freedman M. *Clock Drawing: A Neuropsychological Analysis.* Oxford University Press; 1994.
28. Shin C, Park MH, Lee S-H, et al. Usefulness of the 15-item Geriatric Depression Scale (GDS-15) for classifying minor and major depressive disorders among community-dwelling elders. *J Affect Disord.* 2019;259:370–375.
29. Wancata J, Alexandrowicz R, Marquart B, et al. The criterion validity of the Geriatric Depression Scale: a systematic review. *Acta Psychiatr Scand.* 2006;114:398–410.
30. Balsamo M, Cataldi F, Carlucci L, et al. Assessment of late-life depression via self-report measures: a review. *Clin Interv Aging.* 2018;13:2021–2044.
31. Sheikh JJ, Yesavage JA. Geriatric Depression Scale (GDS): recent findings and development of a shorter version. Brink TL, ed. *Clinical Gerontology: A Guide to Assessment and Intervention.* The Haworth Press; 1986: 165–173.
32. Pfeffer RI, Kurosaki TT, Harrah CH, et al. Measurement of functional activities in older adults in the community. *J Gerontol.* 1982;37:323–329.
33. Loewenstein DA, Amigo E, Duara R, et al. A new scale for the assessment of functional status in Alzheimer's disease and related disorders. *J Gerontol.* 1989;44:P114–P121.
34. Sullivan GM, Feinn R. Using effect size—or why the *P* value is not enough. *J Grad Med Educ.* 2012;4:279–282.
35. Cohen J. *Statistical Power Analysis for the Behavioral Sciences.* Academic press; 2013.
36. Lu AT, Park VT, Peavy GM, et al. Successful Recruitment of Vietnamese Americans into the Asian Cohort for Alzheimer's Disease (ACAD) Pilot Study at UC San Diego. *Alzheimers Dement.* 2022;18:e067487.
37. Peavy GM, Salmon DP, Rice VA, et al. Neuropsychological assessment of severely demented elderly: the severe cognitive impairment profile. *Arch Neurol.* 1996;53:367–372.
38. Panisset M, Roudier M, Saxton J, et al. Severe impairment battery: a neuropsychological test for severely demented patients. *Arch Neurol.* 1994;51:41–45.
39. Parsons MW, Gardner MM, Sherman JC, et al. Feasibility and acceptance of direct-to-home tele-neuropsychology services during the COVID-19 pandemic. *J Int Neuropsychol Soc.* 2022;28:210–215.
40. Jacobs DM, Peavy GM, Banks SJ, et al. A survey of smartphone and interactive video technology use by participants in Alzheimer's disease research: Implications for remote cognitive assessment. *Alzheimers Dement (Amst).* 2021;13:e12188.
41. Zeghari R, Guerchouche R, Tran-Duc M, et al. Feasibility study of an internet-based platform for tele-neuropsychological assessment of elderly in remote areas. *Diagnostics.* 2022; 12:925.
42. Takakura Y, Otsuki M, Takagi R, et al. A validation study for wide-range remote assessment of cognitive functions in the healthy older Japanese population: a pilot randomised cross-over trial. *BMC Geriatr.* 2023;23:575.
43. Zadik L, Perlman S, Barak O, et al. Evaluation of Montreal Cognitive Assessment (MoCA) administered via videoconference. *J Am Med Dir Assoc.* 2023;24:1942.e3–1947.e3.
44. Smith V, Younes K, Poston KL, et al. Reliability of remote National Alzheimer's Coordinating Center Uniform Data Set data. *Alzheimers Dement (Amst).* 2023;15:e12498.
45. Tsiakiri A, Koutzmpis V, Megagianni S, et al. Remote neuropsychological evaluation of older adults. *Appl Neuro-psychol Adult.* 2022;20:1–8.
46. Marra DE, Hamlet KM, Bauer RM, et al. Validity of teleneuropsychology for older adults in response to COVID-19: a systematic and critical review. *Clin Neuropsychol.* 2020;34: 1411–1452.
47. Carotenuto A, Traini E, Fasanaro AM, et al. Tele-neuropsychological assessment of Alzheimer's disease. *J Pers Med.* 2021;11:688.
48. Messler AC, Kane KD, Serrano Y. Tele-neuropsychology in culturally and linguistically diverse populations within the US and US territories: a scoping review. *Clin Neuropsychol.* 2023; 7:1–23.

49. Chau V, Bowie JV, Juon H-S. The association of perceived discrimination and depressive symptoms among Chinese, Korean, and Vietnamese Americans. *Cultur Divers Ethnic Minor Psychol*. 2018;24:389.
50. Korinek K, Young Y, Teerawichitchainan B, et al. Is war hard on the heart? Gender, wartime stress and late life cardiovascular conditions in a population of Vietnamese older adults. *Soc Sci Med*. 2020;265:113380.