

UCSF

UC San Francisco Previously Published Works

Title

In vitro-expanded antigen-specific regulatory T cells suppress autoimmune diabetes

Permalink

<https://escholarship.org/uc/item/8124j43x>

Journal

Journal of Experimental Medicine, 199(11)

ISSN

0022-1007

Authors

Tang, Qizhi Z
Bluestone, Jeffrey A

Publication Date

2004-06-01

DOI

10.1084/jem.20040139

Peer reviewed

In Vitro–expanded Antigen-specific Regulatory T Cells Suppress Autoimmune Diabetes

Qizhi Tang,¹ Kammi J. Henriksen,¹ Mingying Bi,¹ Erik B. Finger,¹ Greg Szot,¹ Jianqin Ye,¹ Emma L. Masteller,¹ Hugh McDevitt,² Mark Bonyhadi,³ and Jeffrey A. Bluestone¹

¹UCSF Diabetes Center, Department of Medicine, University of California San Francisco, San Francisco, CA 94143

²Department of Microbiology and Immunology, Stanford University School of Medicine, Stanford, CA 94305

³Xcyte Therapies, Inc., Seattle, WA 98104

Abstract

The low number of CD4⁺ CD25⁺ regulatory T cells (T_{regs}), their anergic phenotype, and diverse antigen specificity present major challenges to harnessing this potent tolerogenic population to treat autoimmunity and transplant rejection. In this study, we describe a robust method to expand antigen-specific T_{regs} from autoimmune-prone nonobese diabetic mice. Purified CD4⁺ CD25⁺ T_{regs} were expanded up to 200-fold in less than 2 wk in vitro using a combination of anti-CD3, anti-CD28, and interleukin 2. The expanded T_{regs} express a classical cell surface phenotype and function both in vitro and in vivo to suppress effector T cell functions. Most significantly, small numbers of antigen-specific T_{regs} can reverse diabetes after disease onset, suggesting a novel approach to cellular immunotherapy for autoimmunity.

Key words: autoimmunity • tolerance • CD4⁺CD25⁺ T cells • NOD mice • immunoregulation

Introduction

It has become increasingly clear that the balance of pathogenic and immune regulatory pathways underlies disease progression in many autoimmune settings. The loss of regulatory pathways such as CTLA-4, TGF- β , and FoxP3 leads to lethal autoimmunity (1–7). This is best exemplified in type 1 diabetes (T1D) in nonobese diabetic (NOD) mice and humans, where genetic or biologic loss of function of these immunoregulatory pathways exacerbates disease development (8–10). Increasingly, these pathways have pointed to a novel CD4⁺ regulatory T cell (T_{reg}) lymphocyte subset as the central controller of autoimmunity in a variety of experimental animal models as well as an intrinsic regulator of spontaneous autoimmunity (for review see references 11–15). Although the most widely used markers for T_{regs} are the expression of CD4 and CD25, other molecules such as CD62L, CTLA-4, glucocorticoid-induced TNF receptor (GITR), and FoxP3 have emerged as additional markers of this unique T cell lineage (5–7, 16–20). Importantly, T_{reg} therapy can effectively delay and cure mice of a variety of immunological diseases including diabetes, colitis, gastritis, and graft-versus-host disease (8, 21–24). Several studies

have suggested that the T_{regs} are antigen specific, relying on TCR engagement to fully acquire suppressive activity in vivo. The regulatory cells appear to function preferentially at the site of inflammation to effect proliferation and/or cytokine production by the pathogenic T cells (17, 25, 26).

Recent studies have suggested that T_{regs} function via the production of immunosuppressive cytokines, particularly TGF- β and IL-10 (21, 27–29), whereas other studies indicate that suppressive function requires cell–cell contact and cannot be attributed to soluble inhibitors (30–36). Barthlott et al. (37) and Stockinger et al. (38) suggested that T_{regs} function to “take up space,” thus blocking the pathogenic cells from filling up their appropriate niche. The T_{reg} population is reduced in autoimmune-prone animals and patients (8, 39). It appears that T_{regs} may be defective in NOD mice (8, 39). For instance, T_{regs} constitute only ~5% of the circulating CD4⁺ T cells in NOD mice, significantly lower than that observed in other strains (8). Moreover, a large number of T_{regs} (1:1 T_{reg}/T_{eff} ratio) are required to suppress ongoing disease in this model and other autoimmune models (8, 16,

Address correspondence to Jeffrey A. Bluestone, UCSF Diabetes Center, University of California San Francisco, Box 0540, 513 Parnassus Avenue, San Francisco, CA 94143. Phone: (415) 514-1683; Fax: (415) 564-5813; email: jbluest@diabetes.ucsf.edu

Abbreviations used in this paper: APC, allophycocyanin; CFSE, carboxyfluorescein diacetate succinimidyl ester; Ct, threshold cycle(s); GITR, glucocorticoid-induced TNF receptor; GAD, glutamic acid decarboxylase; NOD, nonobese diabetic; T1D, type 1 diabetes; T_{eff}, T effector cell; Tg, transgenic; T_{reg}, regulatory T cell.

17). Finally, recent studies have suggested that it might be impossible to reverse ongoing autoimmune diabetes due to the autoreactive T cells becoming resistant to suppression during the active phase of the disease. However, the studies were limited to *in vitro* analyses presumably due to limited cell numbers (40). In spite of this complexity, the potential for T_{regs} to actively regulate autoimmunity and induce long-term tolerance has great potential applications both for understanding immune homeostasis and as a strategy for inducing long-lived tolerance.

It has been reported that T_{regs} preferentially respond to dendritic cells to proliferate *in vitro* and *in vivo*, but the *in vitro* T_{reg} expansion induced by dendritic cells was still very limited (41). In fact, taking advantage of T_{regs} has been complicated by the difficulty in expanding and characterizing this minor T cell subset. In this study, we developed a robust technique for expanding antigen-specific T_{regs} from autoimmune NOD mice. The expanded T_{regs} retained all the quintessential characteristics of this subset including expression of CD25, CD62L, FoxP3, and GITR. The ability of expanded NOD T_{regs} to suppress diabetes in prediabetic and diabetic mice *in vivo* was significantly enhanced using the autoantigen-specific T cells when compared with polyclonal T_{regs} . Antigen-specific T_{regs} effectively suppressed the development of diabetes in T_{reg} -deficient CD28^{-/-} mice, blocked syngeneic islet graft rejection in chronically diabetic animals, and in contrast to previous reports (40), T_{regs} are shown to reverse diabetes in mice with new onset disease.

Materials and Methods

Mice. NOD mice (Taconic), BALB/c mice (Charles River Laboratories), BDC2.5 TCR transgenic (Tg) mice, glutamic acid decarboxylase (GAD)286 TCR Tg mice (42), NOD.CD28^{-/-} mice, NOD.RAG^{-/-} mice, and NOD.TCR- α ^{-/-} mice were housed and bred under specific pathogen-free conditions at the University of California San Francisco Animal Barrier Facility.

Antibodies and Other Reagents. FITC-labeled mAbs against CD4 (GK1.5) and GITR (DTA-1; reference 19) were purified from hybridoma culture supernatant and conjugated in our lab. R-PE-conjugated anti-CD25 (7D4) mAbs were purchased from Southern Biotechnology Associates, Inc. Allophycocyanin (APC)-labeled mAbs against CD4 (RM4-5) and CD62L were purchased from BD Biosciences or eBioscience. The p31/I-A^{g7}mIlgG2a was generated in our lab (43). Carboxyfluorescein diacetate succinimidyl ester (CFSE) was purchased from Molecular Probes.

Cell Sorting and Flow Cytometry. CD4⁺ T cells were enriched from pooled LNs and spleens by negative selection using an AutoMACS (Miltenyi Biotec). The cells were then stained with anti-CD4-FITC, anti-CD25-PE, and anti-CD62L-APC, and the T_{regs} and CD4⁺ CD62L⁺ CD25⁻ T effector cells (T_{eff}) were sorted on a Mo-Flo cytometerTM (DakoCytomation) based on the expression of CD4, CD25, and CD62L to >98% purity. Flow cytometric analyses were performed on a FACScaliburTM flow cytometer with CELLQuestTM software (Becton Dickinson).

In Vitro Expansion of T Cells. FACS[®]-purified T cells were stimulated with anti-CD3 and anti-CD28 coupled to 4.5- μ m paramagnetic beads (provided by Xcyte Therapeutics Inc.) supplemented with 2,000 IU/ml rIL-2 (Chiron Corp.) in complete medium, which consisted of 10% heat-inactivated fetal bovine se-

rum (Biosource International), nonessential amino acids, 0.5 mM sodium pyruvate, 5 mM HEPES, 1 mM glutamax I (all from Invitrogen), and 55 μ M β -mercaptoethanol (Sigma-Aldrich) in DMEM base. The cultures were monitored daily and maintained at $0.7\text{--}1 \times 10^6$ /ml by diluting with IL-2-supplemented complete medium for 8–12 d. At the end of the culture, the anti-CD3 and anti-CD28 beads were removed using AutoMACS, and the cells were routinely assayed for CD4, CD62L, and CD25 expression by flow cytometry and for suppressive activity *in vitro*. It should be pointed out that conventional anti-CD3 plus anti-CD28-coated 6- μ m polystyrene beads can be adapted for use in this procedure.

In Vitro Suppression Assays. Graded numbers of expanded or fresh sorted T_{regs} were added to 50,000 CD4⁺ T cells stimulated with 50,000 irradiated splenic APC (2,000 rads) and 1 μ g/ml anti-CD3 in a U-bottomed 96-well plate. CD4⁺ T cell cultures without T_{regs} were stimulated in the same manner as positive controls. For some experiments, CD4⁺ T cells from DO11.10 TCR Tg mice were used as responders and the cocultures were stimulated with anti-CD3 as described above or with 0.1 μ g/ml OVA peptide. The cultures were maintained at 37°C for a total of 64 or 72 h and pulsed with 1 μ Ci/well [³H]thymidine for the last 14 or 8 h, respectively. For some experiments, the responder CD4⁺ cells were labeled with 2.5 μ M CFSE before the suppression assay, and the level of proliferation was assessed by determining the dilution of CFSE using flow cytometry 72 h after the initiation of the culture.

Real Time PCR Analysis. Total RNA was extracted using Trizol reagent (Invitrogen) or RNeasy (QIAGEN) from expanded T cells. cDNA was synthesized from 50 ng–2.5 μ g of each RNA sample using SuperScript II RNase H reverse transcriptase and oligo dT as primer (Invitrogen), and 0.6–31.25 ng of the cDNA was used in each quantitative real time PCR reaction. The same amount of RNA and cDNA was used for each T_{reg} and CD25⁻ sample pair. Primers and probes for FoxP3, CTLA-4, neuropilin, PD-1, TRAIL, and HPRT were purchased as reagent kits from Applied Biosystems. Primer sequences for SOCS-2 were 5'-GCGTCTGGCGAAAGCCCT (forward) and 5'-CTTCATTAACAGTCATACTTCC (reverse), and they were ordered from Integrated DNA Technologies Inc. The probe sequence for SOCS-2 was 5'-FAM-CGCGAGCTCAGTCAAACAGGATGGT-TAMRA-3', which was ordered from Applied Biosystems. The real time PCR was performed on an ABI prism 7700 using Taqman Universal PCR master mix (Applied Biosystems) in duplicates and the average threshold cycles (Ct) of the duplicates were used to calculate the fold change between expanded T_{regs} and CD4⁺ CD62L⁺ CD25⁻ cells. Ct for HPRT was used to normalize the samples. Expression ratios between T_{regs} and CD25⁻ cells were calculated using the following formula: Expression Ratio ($T_{\text{reg}}/CD25^{-}$)_{Gene} \times = 2^n , n = (CD25⁻ Ct_{Gene} \times -CD25⁻ Ct_{HPRT}) - (T_{reg} Ct_{Gene} \times - T_{reg} Ct_{HPRT}).

Western Blot. 5×10^5 of each cell type were lysed in sample buffer (6.25 mM Tris, pH 6.8, 12.5% glycerol, 2% SDS, 30 ng/ml bromophenolblue), sonicated, and passed through 28-gauge needles. The lysates were clarified by centrifugation and boiled for 5 min before separating on a 10% SDS PAGE gel. The samples were transferred to PVDF membrane after electrophoresis and incubated with rabbit anti-FoxP3 antisera (provided by S. Ziegler, Benaroya Research Institute at Virginia Mason, Seattle, WA) followed by horseradish peroxidase-conjugated anti-rabbit Ig. The blot was developed with SuperSignal[®] Chemiluminescent Substrate (Pierce, Rockford, IL) and visualized on a Kodak Image Station 440CF (Eastman Kodak Co.) and quantified using Kodak Digital Science 1D Image Analysis software 3.0.

Cytokine ELISA. The level of IL-2, IFN- γ , IL-10, and TGF- β in the culture supernatant was determined by ELISA using antibody pairs purchased from BD Biosciences. For TGF- β ELISA, the culture supernatant was first treated with acid to lower the pH to 2.0 to denature latency-associated peptide to allow the detection of active TGF- β . The supernatant was neutralized to pH 7.0 before ELISA.

Adoptive Transfer. Expanded T cells were labeled with 2.5 μ M CFSE, and $1-3 \times 10^6$ cells were transferred via retro-orbital injection. The recipient mice were killed on day 7 after cell transfer and the dilution of CFSE in splenic, peripheral LN, and pancreatic LN cell preparations was determined by flow cytometry. For adoptive transfer of diabetes to NOD.RAG $^{-/-}$ recipients, CD4 $^+$ CD62L $^+$ CD25 $^-$ cells from BDC2.5 TCR Tg mice were purified by cell sorting using a DakoCytomation Mo-Flo and $0.5-1 \times 10^6$ cells were transferred to each recipient mouse. For some experiments, the purified T $_{\text{eff}}$ s were activated with anti-CD3 and anti-CD28 for 8–10 d before transfer. To transfer diabetes with polyclonal T $_{\text{eff}}$ s, 25×10^6 pooled spleen and LN cells from diabetic NOD mice were injected into each recipient. When indicated, expanded T $_{\text{regs}}$ were depleted of anti-CD3 and anti-CD28 beads and washed extensively before mixing with effector cells for injection. The expanded T $_{\text{regs}}$ were similarly processed before transferring into NOD.CD28 $^{-/-}$, chronically diabetic NOD syngeneic islet transplant recipients, and NOD mice with new onset diabetes. Nonfasting blood glucose levels in recipient mice were monitored using an Accu-Check glucometer (Roche Diagnostics Corp.)

Murine Pancreatic Islet Isolation. Murine islets were isolated using a modified previously published protocol (44). In brief, a 3-ml collagenase P (Roche Molecular Biochemicals) solution (0.75 mg/ml) was injected into the pancreatic duct of 4-wk-old NOD mice. The distended pancreases were removed and incubated at 37°C for 17 min. The liberated free islets were purified

by centrifugation on Eurocollin-Ficoll gradients that comprised four different densities (1.108, 1.096, 1.069, and 1.037). After centrifugation, the islet-containing layers between densities 1.069 and 1.096 were collected and washed. Islets were then handpicked for transplantation.

Murine Islet Transplantation. Naturally diabetic NOD mice were used as recipients. The mice were diabetic for at least 2 wk before transplantation. The recipient mice were maintained with subcutaneous insulin pellets (Lin-Shin Canada, Inc.). 1 d before transplantation, the insulin pellets were removed and hyperglycemia was confirmed on the day of transplantation. 500 isolated and handpicked islets were transplanted beneath the left renal capsule of each recipient. Nonfasting blood glucose levels were determined in all animals daily after transplantation. Return to normoglycemia within 24 h after transplant was indicative of successful surgery. Rejection of the islets grafts was considered to have occurred when nonfasting blood glucose concentration exceeded 250 mg/dl for 3 consecutive days.

Results

Expansion of T $_{\text{regs}}$ from Autoantigen-specific TCR Tg NOD Mice. Previous studies have shown that the number and function of T $_{\text{regs}}$ in NOD mice decrease over time correlating with clinical disease onset between 16 and 24 wk of age (9, 40). These observations support the use of T $_{\text{regs}}$ to prevent or treat diabetes even after disease onset. However, the ability to use these cells therapeutically is severely limited by the small numbers of cells resident in the circulation or lymphoid organs (<5% of CD4 $^+$ T cells in NOD mice and <2% of CD4 $^+$ T cells in humans with T1D; references 8 and 39). Moreover, a large number of cells are required for therapeutic efficacy due to an inability at present to se-




Figure 1. In vitro expansion of T $_{\text{regs}}$. (A) Representative flow cytometry plots of CD25 and CD62L expression on CD4 cells from NOD (left), BDC2.5 (middle), and GAD286 (right) mice. FACS $^{\text{R}}$ -purified T $_{\text{regs}}$ (●) and CD4 $^+$ CD62L $^+$ CD25 $^-$ cells (○) from NOD (B), BDC2.5 TCR Tg (C), or GAD286 TCR Tg (D) mice were stimulated in vitro with anti-CD3- and anti-CD28-coated beads along with IL-2. (E) T cells from BDC2.5 TCR Tg mice were expanded as described above with p31-linked IA $^{\text{g7}}$ -mIgG2a immobilized on latex beads. All cultures were quantitated by viable cell counting.

lect the cells based on antigen specificity. Therefore, we developed a technique for rapid and efficient expansion of autoantigen-specific T_{regs} based on observations that these cells, present in TCR Tg mice, can be driven into cell cycle with coimmobilized anti-CD3 and anti-CD28 antibodies plus exogenous IL-2. As shown in Fig. 1, A and B, FACS[®]-purified NOD T_{regs} cultured with anti-CD3/anti-CD28-coated beads in the presence of IL-2 expanded 150–225-fold in 11 d (Fig. 1 B). In general, the $CD4^+ CD25^-$ T cells expanded more vigorously (ranging from 300–800-fold in multiple experiments). A purity of $>98\%$ $CD4^+ CD25^+ CD62L^+$ T cells was essential to enable successful T_{reg} expansion as a small contamination of either $CD25^- CD4^+$ or $CD8^+$ T cells significantly impacted the ability to specifically expand the T_{regs} (unpublished data). It should be noted that the T_{reg} expansion is dependent on the high level of IL-2 (2,000 IU/ml). No T_{reg} expansion was observed when 200 IU/ml IL-2 was used (unpublished data).

Previous studies have shown that $CD4^+ CD25^+ T_{\text{regs}}$ isolated from young NOD mice suppressed the ability of T_{effs} from diabetic NOD mice to transfer disease in immunodeficient NOD mice (8, 16, 17). However, the process was highly inefficient and the suppressive effects of T_{regs} in this setting required a 0.5:1 or 1:1 ratio of $T_{\text{reg}}/T_{\text{eff}}$. This is likely due to the low precursor frequency of antigen-specific T_{regs} . Thus, we examined whether T_{regs} from two different antigen-specific TCR Tg mice (Fig. 1 A) could be expanded in vitro using the same methodology as with the polyclonal NOD T_{regs} . BDC2.5 TCR Tg mice express a TCR specific for an islet antigen expressed in the granules of β cells, whereas the GAD286 TCR Tg recognizes a peptide derived from the islet antigen GAD. T_{regs} were purified from BDC2.5 and GAD286 mice and expanded using

the anti-CD3/anti-CD28 plus IL-2 cocktail (Fig. 1, C and D). The BDC2.5 cells expressed the Tg TCR $\alpha\beta$ based on efficient staining with an MHC peptide tetramer previously shown to react with this TCR (43), and the expanded GAD286 T_{regs} expressed the Tg TCR- β chain (unpublished data). The $CD4^+ CD62L^+ CD25^-$ and T_{regs} from BDC2.5 TCR Tg mice can also be expanded using immobilized MHC peptide dimers (Fig. 1 E). These results suggest that a population of $CD4^+ CD25^+ CD62L^+$ exists in both wild-type and TCR Tg mice that can be expanded using this protocol.

Next, we examined the phenotype of the expanded T_{regs} by flow cytometry, Western blot, and real time PCR. As can be seen in Fig. 2 A, the expanded T_{regs} maintained high levels of expression of CD25 as compared with expanded $CD25^-$ T cells, whereas the expression of CD62L remained high in both cell types. In addition, quantitative PCR showed that all of the T_{regs} expressed high levels of SOCS2, PD-1, and CTLA-4 as compared with similarly expanded $CD25^-$ T cells. Moreover, the recently identified markers neuropilin and TRAIL (20, 45) were also highly expressed on the expanded T_{regs} (Fig. 2 B). A high level of cell surface GITR expression was observed on the expanded T_{regs} . However, this previously identified T_{reg} marker was also induced on the expanded $CD25^-$ T cells (19, 20, and unpublished data). It should be noted that the quantitative PCR studies were performed on five separate expanded T_{reg} populations (including both polyclonal and BDC2.5 TCR Tg T_{regs}) and the relative expression of the T_{reg} -specific genes was highly reproducible. Finally, we examined the recently identified lineage/differentiation marker for T_{regs} , FoxP3 (Fig. 2, B and C). As noted by both real time PCR and Western blot analyses, the ex-




Figure 2. Phenotype of in vitro-expanded T_{regs} . (A) Expression of CD25 and CD62L on expanded T_{regs} and $CD4^+ CD62L^+ CD25^-$ cells was determined by flow cytometry on day 8 after the culture initiation. Results are representative of more than 20 independent experiments. (B) Levels of mRNA for the indicated genes in expanded NOD (filled symbols) or BDC2.5 TCR Tg T cells (open symbols) were determined by real time PCR analysis on day 10 after the initiation of the cultures. The relative expression ratio ($T_{\text{reg}}/T_{\text{CD25}^-}$) for each pair of cultures was calculated from Ct values as described in Materials and Methods. The dashed line represents the ratio of 1 (i.e., identical level of gene expression in T_{reg} and $CD4^+ CD62L^+ CD25^-$ cultures). (C) Western blot analysis of FoxP3

protein expression in fresh and expanded T cells. The level of tubulin expression was included as a loading control. Results are representative of three independent experiments. (D) Cytokine secretion by expanded BDC2.5 T cells 48 h after restimulation with antigenic peptide and splenic APC. Results are representative of two independent experiments.

expanded T_{regs} expressed levels of FoxP3 similar to those observed in fresh T_{regs} and significantly higher than those in fresh or expanded $CD25^{-}$ T cells. The RNA expression (10-fold) and protein amounts (20-fold) were consistent with previous studies of fresh T_{regs} , although there was clearly some increase in FoxP3 in $CD25^{-}$ T_{effs} , suggesting that the culture conditions may induce some T_{regs} within the $CD25^{-}$ subset, or FoxP3 is expressed at a low level in activated T_{effs} .

We also examined the ability of the expanded T_{regs} to secrete cytokines. Unlike activated $CD25^{-}$ T cells, the T_{regs} did not produce IL-2 or IFN- γ , but rather expressed the immunosuppressive cytokines IL-10 and TGF- β (Fig. 2 D). Thus, the extensive activation and proliferation of the T_{regs} does not alter the phenotype of the T_{regs} , which remained distinct from the $CD25^{-}$ T cell subset.

Functional Activity of In Vitro-expanded T_{regs} . Previous studies have shown that T_{regs} can effectively suppress proliferative responses of $CD25^{-}$ T cells stimulated with anti-CD3 and splenic APC. The expanded NOD T_{regs} efficiently suppressed proliferative responses (Fig. 3 A) and cytokine production including IL-2 and IFN- γ (unpublished data). In fact, in multiple experiments, the expanded T_{regs} suppressed significantly better than fresh NOD T_{regs} . The suppression was routinely observed at $T_{\text{reg}}/T_{\text{eff}}$ ratios of <1:10. Similar results were observed using the expanded T_{regs} from the TCR Tg mice, as the expanded BDC2.5 T_{regs} were effective in suppressing the proliferative response of BDC2.5 (unpublished data) as well as polyclonal NOD T cells (Fig. 3 B). Although the expanded T_{regs} expressed significant levels of IL-10 and TGF- β , suppressor activity was unaffected by the addition of anti-IL-10, anti-TGF- β , or a combination of both antibodies to the in vitro cultures (unpublished data). These results are consistent with numerous models of T_{reg} suppression where cell-cell contact is the primary means of immunosuppression in the in vitro setting (30–36).

To further assess the antigen specificity of the expanded T_{regs} and determine whether the expanded T_{regs} were constitutively suppressive, expanded T_{regs} from normal BALB/c mice were examined for their ability to suppress T cells from the OVA-specific DO11.10 TCR Tg mouse. T_{regs} and DO11.10 Tg T_{effs} were cocultured in the presence of OVA antigen (to activate only the T_{effs}) or anti-CD3 (to

activate both the T_{eff} and T_{regs}). Expanded BALB/c T_{regs} did not inhibit the proliferative response of the DO11.10 T cells stimulated by the OVA peptide. However, the anti-CD3 response was fully inhibited at low $T_{\text{reg}}/T_{\text{eff}}$ ratios. These results suggest that the expanded T_{regs} lack constitutive suppressive activity but require TCR-specific activation for effective suppression. This result also ruled out the trivial possibility that the cells were inhibiting the cultures by consuming available IL-2 through the high level of CD25 expression.

In Vivo Survival and Activation of Expanded T_{regs} . Effective suppression of immune responses in vivo by T_{regs} requires that the cells migrate to appropriate sites, respond to antigen, and survive long-term. We have observed recently that blockade of the CD28/B7 pathway resulted in rapid loss of T_{regs} in vivo and subsequent loss of critical immune regulation (8, 46). Thus, we examined the ability of expanded T_{regs} to survive and proliferate in vivo. Expanded T_{regs} were labeled with CFSE and transferred into normal nonlymphopenic syngeneic mice. At 30 d after transfer, the mice were killed and examined for the number of CFSE $^{+}$ cells as an indication of cell survival. As seen in Fig. 4 A, a significant number of CFSE $^{+}$ cells were recovered from mice transferred with expanded T_{regs} . The number of CFSE $^{+}$ T_{regs} was equal to that observed with fresh T_{regs} transferred in the same manner (Fig. 4 A). In fact, Thy1.1-marked expanded T_{regs} were observed at least 50 d after transfer (unpublished data).

Next, we analyzed the ability of the adoptively transferred T_{regs} to respond to antigen and proliferate in vivo. Expanded T_{regs} from NOD, BDC2.5, and GAD286 mice were labeled with CFSE and transferred into normal nonlymphopenic NOD recipients. At 7 d after transfer, the mice were killed and examined for the dilution of CFSE to assess in vivo proliferation. As seen in Fig. 4 B, top, a small but significant number of T_{regs} proliferated, as indicated by CFSE dilution. However, there was no selective proliferation of the NOD T_{regs} in the pancreatic LNs (Fig. 4 B, pancLN), suggesting that a detectable number of islet autoantigen-specific cells did not exist within the NOD T_{reg} repertoire. In contrast to the NOD T_{regs} , T_{regs} from BDC2.5 Tg mice proliferated extensively and selectively in the pancreatic LNs, dividing at least three to four times during the 7-d period (Fig. 4 B, bottom). Interestingly, the




Figure 3. In vitro suppression by expanded T_{regs} . (A) Fresh and expanded T_{regs} were compared for their ability to suppress the proliferation of $CD4^{+}$ responder T cells stimulated with anti-CD3 and T cell-depleted splenocytes. (B) Suppression by T_{regs} expanded from BDC2.5 TCR Tg or GAD286 TCR Tg mice was assayed as described in A. (C) Suppressive activity of BALB/c-expanded T_{regs} on $CD4^{+}$ responder T cells from DO11.10 TCR Tg mice stimulated with anti-CD3 or an OVA peptide. Results are representative of three independent experiments.




Figure 4. In vivo survival and activation of expanded T_{reg} s. (A) Freshly isolated and expanded BALB/c T_{reg} s were labeled with CFSE and then injected into normal BALB/c mice (10^6 /mouse). All recipient mice were killed on day 30 after injection and the numbers of CFSE⁺ cells in the peripheral LN and spleen (not shown) were determined by flow cytometry. The data are presented as the number of CD4⁺ CFSE⁺ cells/ 10^6 endogenous CD4⁺ T cells. Each circle represents the value from one mouse and the black bar represents the mean of the group. (B) Expanded T_{reg} s from NOD.Thy1.2 (top) and GAD286 TCR Tg Thy1.2 (middle) were labeled with CFSE and 3×10^6 were transferred to normal 8–12-wk-old NOD.Thy1.1 recipients. Expanded BDC2.5 TCR Tg Thy1.1 were labeled with CFSE and 3×10^6 were transferred to normal 8–12-wk-old NOD.Thy1.2 recipients. The presence of transferred cells and their activation status in spleens, LNs, and pancreatic LNs were determined by flow cytometry on day 7 after cell transfer. The dot plots shown are gated on the Thy1.2 for NOD and GAD286 cells and Thy1.1 for BDC2.5 cells. The percentages of cells with CFSE dilution are shown on the plots. Results are representative of at least three recipient mice in two separate experiments.

expression of the CD62L molecule was down-regulated on the surface of expanded T_{reg} s. This is surprising because the cells had undergone multiple proliferative cycles in vitro before transfer and had maintained high levels of CD62L expression. In contrast to the BDC2.5 T_{reg} s, the GAD286 T_{reg} s did not proliferate in vivo (Fig. 4 B, middle). The results of previous studies suggest that T cells in these two TCR Tg mice differ significantly in their thymic development. The BDC2.5 Tg mice do not negatively select the islet-specific T cells in the thymus, but rather develop a small, reproducible number of T_{reg} s. These cells have been

shown to block disease manifested by potential pathogenic CD4⁺ CD25⁻ T cells resident in these animals (47). By comparison, the majority of GAD286 TCR Tg T cells are deleted in the NOD thymus by negative selection. In fact, the minor population of cells that escape use alternative TCR- α chains. Thus, although the peripheral GAD286 TCR Tg cells respond to GAD peptide in vitro, the reactivity is weak and, in contrast to the BDC2.5, they are unable to induce diabetes upon adoptive transfer, suggesting the “absence” of an autoreactive repertoire (42). These results support the conclusion that the two Tg mice are resistant to diabetes for distinct reasons. The GAD286 TCR Tg mice are protected from the development of diabetes due to the potent central tolerance mechanism of clonal deletion and receptor editing. By comparison, the BDC2.5 TCR Tg have circulating autoreactive T_{reg} s that home to peripheral target tissues where they are activated and expand after encountering autoantigen, resulting in immune suppression and homeostasis.

In Vitro-expanded T_{reg} s Suppress Adoptive Transfer of Diabetes In Vivo. Next, we examined the ability of the expanded BDC2.5 T_{reg} s to suppress diabetes after in vivo cotransfer of activated BDC2.5 T cells into NOD.RAG mice. The T_{reg} s were effective in blocking the transfer of diabetes, functioning at as low as a 1:9 ratio of T_{reg}/T_{eff} (Fig. 5 A), whereas the GAD286 T_{reg} s did not protect even at a T_{reg}/T_{eff} ratio of 1:1 (Fig. 5 B). In fact, the expanded BDC2.5 T_{reg} s suppressed polyclonal T cell-mediated disease. As few as 2×10^6 expanded BDC2.5 T_{reg} s blocked the ability of 25×10^6 diabetogenic NOD spleen and LN cells to transfer disease (Fig. 5 C). The expanded antigen-specific T_{reg} s from the BDC2.5 mice were far more efficient than expanded polyclonal NOD T_{reg} s in preventing the onset of diabetes. The same number (2×10^6) as well as 5×10^6 expanded NOD T_{reg} s did not confer any protection under the same conditions (Fig. 5 C). In fact, even the transfer of four times as many expanded NOD T_{reg} s (8×10^6) only slightly delayed diabetes onset and prevented diabetes in only one diabetogenic cell recipient (Fig. 5 C). This result is consistent with previous findings suggesting that a high ratio of polyclonal T_{reg} s to T_{eff} s is necessary to efficiently suppress disease transfer in this setting (8, 16, 17). Importantly, these data suggest that in vitro activity of the T_{reg} s does not predict in vivo function in this disease setting.

Expanded T_{reg} s Prevent Diabetes In Vivo in a Nonlymphopenic Setting. Although there are multiple models demonstrating the immunoregulatory activity of T_{reg} s, many of the systems are based on adoptive transfer models that take advantage of lymphopenic mice to enhance T_{reg} proliferation (8, 21–24). Questions have been raised whether disease suppression observed in lymphopenic settings after T_{reg} transfer is due to active regulation or a side effect of competition for “space” (37, 38). Therefore, we examined the ability of the expanded T_{reg} s to prevent diabetes in a nonlymphopenic animal model. Previous studies have shown that CD28^{-/-} NOD mice have normal numbers of T cells and Th1 responses. In fact, these mice develop exacerbated




Figure 5. Prevention of diabetes transfer by expanded T_{regs} . (A) Activated diabetogenic BDC2.5 $CD4^+ CD62L^+ CD25^-$ cells (3.5×10^5) were cotransferred with BDC2.5-expanded T_{regs} to 8-wk-old NOD.RAG $^{-/-}$ recipients at the indicated ratio. The blood glucose for individual recipient mouse was monitored and plotted to access diabetes. $n = 3$ for no T_{regs} and 1:9 groups; $n = 4$ for 1:1 and 1:3 groups. Results are representative of three independent experiments. (B) Diabetes was induced in 6-wk-old NOD.RAG $^{-/-}$ mice in the same manner as described in A, except that the number of transferred expanded T_{regs} from GAD286 TCR Tg mice and BDC2.5 TCR Tg mice equaled the number of transferred T_{regs} ($n = 3$ mice/group). (C) Diabetes was induced in 5–8-wk-old NOD.RAG $^{-/-}$ or NOD.TCR- $\alpha^{-/-}$ recipients by injection of 25×10^6 pooled spleen and LN cells from diabetic donors ($n = 8$). Some recipient mice were coinjected with expanded T_{regs} from NOD (2×10^6 , $n = 3$; 5×10^6 , $n = 3$; 8×10^6 , $n = 4$) or BDC2.5 TCR Tg mice (2×10^6 , $n = 4$). Results represent two independent experiments.

autoimmunity due to a deficiency in Th2 and T_{regs} , which were shown to be exquisitely CD28 dependent (8, 21–24).

Thus, we examined whether wild-type expanded BDC2.5 T_{regs} transferred into CD28 $^{-/-}$ NOD mice could delay or prevent onset of disease. 5×10^5 T_{regs} were transferred into 5-wk-old CD28 $^{-/-}$ NOD mice and monitored for diabetes (Fig. 6). The transfer of expanded BDC2.5 T_{regs} prevented the development of diabetes in 100% of mice followed for as long as 20 wk after transfer. In contrast, the transfer of similar numbers of expanded NOD T_{regs} had no effect on disease incidence (unpublished data). These results suggest that the antigen-specific expanded T_{regs} functioned *in vivo* in the face of a fully functional pathogenic T cell response.

Expanded T_{regs} Reverse Diabetes *In Vivo*. The ultimate utility of T_{reg} therapy depends on an ability to treat individuals with ongoing disease. Thus, we examined the regulatory effects of expanded BDC2.5 T_{regs} in NOD mice that had been diabetic for at least 2 wk to ensure total endogenous islet cell destruction. Expanded BDC2.5 T_{regs} were transferred into diabetic NOD mice in conjunction with 500 syngeneic NOD islet transplant. Mice reverted to normoglycemia within 24–48 h after transplantation. However, unlike the control mice that rejected the transplanted islets within 2 wk, the transfer of 2×10^6 BDC2.5-expanded T_{regs} blocked rejection of the syngeneic islets, consistent with an ability of the suppressor cells to block ongoing autoimmunity in this setting. The transfer of 5×10^6 polyclonal NOD T_{regs} had no effect in this model. More significantly, the adoptive transfer of expanded BDC2.5 T_{regs} reversed diabetes in new-onset diabetic NOD mice (Fig. 7 B). Previous studies have shown that NOD mice diagnosed within the first week of hyperglycemia retain sufficient insulin-producing β cell activity, such that effective immunosuppression introduced at that time can reverse diabetes. To test the efficacy of expanded BDC2.5 T_{regs} in this setting, 10^7 T_{regs} were transferred into NOD mice diagnosed with disease based on recently elevated blood glucose levels (>300 mg/dL). The transferred T_{regs} reversed diabetes in 60% of the mice. Thus, the expanded T_{regs} were




Figure 6. Prevention of autoimmune diabetes in NOD.CD28 $^{-/-}$ mice with BDC2.5-expanded T_{regs} . 5-wk-old prediabetic NOD.CD28 $^{-/-}$ mice were injected with 5×10^5 BDC2.5-expanded T_{regs} ($n = 3$) or left untreated ($n = 4$). The development of diabetes was monitored and blood glucose levels of individual mice were plotted. Results are representative of at least five independent experiments.




Figure 7. Reversal of diabetes with expanded T_{regs} . (A) NOD mice with chronic diabetes were transplanted with syngeneic islets under the kidney capsule. On the day of transplantation, some recipient mice received 5×10^6 NOD-expanded T_{regs} ($n = 4$) or 2×10^6 BDC2.5-expanded T_{regs} ($n = 5$), and the remaining mice ($n = 3$) were left untreated. Blood glucose level was monitored. All islet recipients normalized blood glucose within the first day after transplantation. Results are representative of two independent experiments. (B) NOD mice with new onset diabetes (blood glucose > 300 mg/dL, $n = 7$) were injected with 10^7 BDC2.5-expanded T_{regs} and blood glucose was monitored. Two consecutive readings of blood glucose of < 250 mg/dL was considered remission of diabetes.

extremely effective in blocking and reversing diabetes in an ongoing autoimmune setting.

Discussion

The past few years have seen an increased interest in and understanding of the role of T_{regs} in immune homeostasis. As an example, we have recently shown that anti-CD3 therapy in new-onset diabetes leads to the production and expansion of TGF- β -dependent T_{regs} that reverse diabetes and promote long-term tolerance (9). Moreover, Edinger et al. (24) have shown that the adoptive transfer of T_{regs} in mice can block graft-versus-host disease without affecting graft-versus-leukemia responses. These studies and others have led investigators to conclude that these cells might be involved in human autoimmune diseases. The results have also prompted investigators to consider this cell type for immunotherapy. However, successful application of adoptive cellular immune therapy with these cells will depend on a large, reliable source of well-characterized T_{regs} .

In this study, we describe a robust and effective method for expanding T_{regs} while retaining their phenotype and suppressive activities. We demonstrate that expanded anti-

gen-specific T_{regs} prevent the development of diabetes and even restore an immune regulatory state that reverses diabetes and allows the mice to maintain long-term immune homeostasis. The expanded antigen-specific T_{regs} survived long-term in vivo, were less dependent on CD28 costimulation (unpublished data), but required antigen exposure for functional activity. To our knowledge, this is the first example of T_{reg} activity in a lymphocyte-sufficient diabetic animal. These results are especially important in light of a recent publication by Gregori et al. (40), suggesting that T_{regs} do not effectively suppress T_{eff} in the setting of diabetes. The difference between the two studies may reflect either of the different assay systems: in vivo versus in vitro or that the expanded T_{regs} might be more efficient. In this regard, we have noted that the expanded T_{regs} do indeed survive better in vivo than the fresh T_{regs} .

Among the more interesting and perhaps unexpected observations in this study was the differential dependency of antigen specificity for in vivo versus in vitro T_{reg} functions. Studies in multiple models have shown that polyclonal T_{regs} are effective in blocking autoimmunity (8, 21–24). In fact, in most settings (with a few notable exceptions such as allogeneic organ transplantation; reference 28), the ability to functionally suppress in vitro has been highly predictive of in vivo efficacy and presumed to be antigen non-specific. Yet, in this study, the islet autoantigen-specific BDC2.5 T_{regs} were significantly more efficient than polyclonal NOD T_{regs} in regulating autoimmune responses in vivo. This discriminating activity was not predicted by the in vitro studies that demonstrated equal efficacy in blocking anti-CD3 responses among the various expanded T_{regs} . There are several potential explanations for this observation. First, it is possible that the expansion method causes selective depletion of the autoantigen-specific T_{regs} in the polyclonal NOD populations. This seems unlikely as the BDC2.5 T_{regs} grew in vitro equally well when compared with the polyclonal NOD T_{regs} . Moreover, cursory analysis of the TCR usage in the expanded T_{regs} showed no preferential changes in the repertoire of the cells. Second, it is possible that the different expanded polyclonal T_{reg} populations homed or functioned differently in vivo. However, analysis of a panel of cell surface markers, intracellular proteins, and soluble cytokines suggested that the various cell populations all retained the essential properties of T_{regs} including the high expression of CD25 and FoxP3. Most likely, the differences observed reflected the model in which we have tested the T_{regs} . In many models, T_{regs} are cotransferred with effector cells into lymphoid cell-deficient animals, in which the T_{regs} home to the site of inflammation and inhibit T_{eff} response indirectly by competing for the “niche.” In fact, in one model of inflammatory bowel disease (37), CD25 $^-$ T cells are as effective as CD25 $^+$ T_{regs} with little evidence for antigen specificity. Similarly, there has been little evidence supporting an essential role for antigen-specific cells for the treatment of graft-versus-host disease. In sharp contrast, T_{reg} -mediated immune regulation is routinely found to be antigen specific in the non-

lymphopenic setting (28, 48–50). Thus, the greater functional activity of the antigen-specific T_{regs} in the NOD models described herein is most likely due to the fully functional immune system in these animals and the requirement for effective antigen-mediated reactivation of T_{regs} at the inflammatory site.

It is important to note that in the absence of antigen-specific activation, the T_{regs} have no effect on regulating disease. This is most evident in the adoptive transfer studies using the T_{regs} isolated from Tg mice on the NOD background expressing a TCR specific for peptide epitope 286–300 (p286) of GAD65. Although lymphocytes from these TCR Tg mice proliferated and produced cytokines when stimulated in vitro with GAD65 peptide 286–300 (42), the response was weak and the T cells escaping deletion expressed alternative TCR- α chains that were unable to transfer diabetes. In fact, the only model where the GAD Tg T cells were functional was in an adoptive transfer system where the p286-tetramer⁺ CD4⁺ T cells from TCR Tg mice delayed diabetes induced in NOD.SCID mice by diabetic NOD spleen cells independent of CD25 expression, not unlike other systems where the transferred T cells suppress independent of CD25 expression or antigen-specific mechanisms (42).

Finally, the observation that the T_{regs} were able to reverse diabetes has important implications for clinical therapy. We imagine a scenario in a number of autoimmune settings where T_{regs} are isolated from patients either during remission (as would be the case for systemic lupus erythematosus or multiple sclerosis) or soon after disease onset (as would be the case for T1D). The cells would be expanded and reintroduced at the time of maximal disease activity to moderate the inflammatory response. In some cases this could be combined with rapamycin, anti-CD3, or other drugs that cause deletion of the pathogenic cells without affecting the T_{regs} (unpublished data). Together, these therapies could both reduce the short-term pathogenic responses while reinstating a homeostatic balance for long-term tolerance induction. Findings in this study suggest that the efficacy of T_{reg} -based immune therapy is critically dependent on the antigen specificity of the T_{regs} , at least in the autoimmune diabetes setting. Therefore, it is important to develop a procedure to selectively and reproducibly expand antigen-specific T_{regs} from polyclonal populations for therapeutic use. Many islet-specific T cell antigens have been identified to contribute to diabetes development in both mice and humans and MHC multimer coupled with these antigenic epitopes has been developed (43, 51–53). It is possible that these MHC multimer reagents can be adapted for expanding islet-specific T_{regs} . Current efforts are underway to identify antigen-specific T_{regs} in the polyclonal population and develop a protocol to selectively expand islet antigen-specific T_{regs} from normal NOD mice using immobilized antigenic peptide-linked MHC multimers. In addition, we are developing similar techniques for purifying and expanding human T_{regs} for clinical use in these settings.

We thank Paul Wegfarht for expert assistance with the mice, Shuwei Jiang and Cliff McArthur for cell sorting, and Dr. Abul Abbas and all Bluestone Lab members for critical discussion. The authors would also like to thank Dr. Steve Ziegler for providing anti-FoxP3 antibodies for this study.

This work was supported by Juvenile Diabetes Research Foundation Center grant number 4-1999-841 and National Institutes of Health grant number R37 AI46643.

Submitted: 22 January 2004

Accepted: 7 April 2004

References

1. Waterhouse, P., J.M. Penninger, E. Timms, A. Wakeham, A. Shahinian, K.P. Lee, C.B. Thompson, H. Griesser, and T.W. Mak. 1995. Lymphoproliferative disorders with early lethality in mice deficient in Ctl4-4. *Science*. 270:985–988.
2. Tivol, E.A., F. Borriello, A.N. Schweitzer, W.P. Lynch, J.A. Bluestone, and A.H. Sharpe. 1995. Loss of CTLA-4 leads to massive lymphoproliferation and fatal multiorgan tissue destruction, revealing a critical negative regulatory role of CTLA-4. *Immunity*. 3:541–547.
3. Kulkarni, A.B., C.G. Huh, D. Becker, A. Geiser, M. Lyght, K.C. Flanders, A.B. Roberts, M.B. Sporn, J.M. Ward, and S. Karlsson. 1993. Transforming growth factor beta 1 null mutation in mice causes excessive inflammatory response and early death. *Proc. Natl. Acad. Sci. USA*. 90:770–774.
4. Shull, M.M., I. Ormsby, A.B. Kier, S. Pawlowski, R.J. Diebold, M. Yin, R. Allen, C. Sidman, G. Proetzel, D. Calvin, et al. 1992. Targeted disruption of the mouse transforming growth factor- β 1 gene results in multifocal inflammatory disease. *Nature*. 359:693–699.
5. Hori, S., T. Nomura, and S. Sakaguchi. 2003. Control of regulatory T cell development by the transcription factor Foxp3. *Science*. 299:1057–1061.
6. Khattry, R., T. Cox, S.A. Yasayko, and F. Ramsdell. 2003. An essential role for Scurfin in CD4⁺ CD25⁺ T regulatory cells. *Nat. Immunol.* 4:337–342.
7. Fontenot, J.D., M.A. Gavin, and A.Y. Rudensky. 2003. Foxp3 programs the development and function of CD4⁺ CD25⁺ regulatory T cells. *Nat. Immunol.* 4:330–336.
8. Salomon, B., D.J. Lenschow, L. Rhee, N. Ashourian, B. Singh, A. Sharpe, and J.A. Bluestone. 2000. B7/CD28 costimulation is essential for the homeostasis of the CD4⁺ CD25⁺ immunoregulatory T cells that control autoimmune diabetes. *Immunity*. 12:431–440.
9. Belghith, M., J.A. Bluestone, S. Barriot, J. Megret, J.F. Bach, and L. Chatenoud. 2003. TGF- β -dependent mechanisms mediate restoration of self-tolerance induced by antibodies to CD3 in overt autoimmune diabetes. *Nat. Med.* 9:1202–1208.
10. Ueda, H., J.M. Howson, L. Esposito, J. Heward, H. Snook, G. Chamberlain, D.B. Rainbow, K.M. Hunter, A.N. Smith, G. Di Genova, et al. 2003. Association of the T-cell regulatory gene CTLA4 with susceptibility to autoimmune disease. *Nature*. 423:506–511.
11. Sakaguchi, S., N. Sakaguchi, J. Shimizu, S. Yamazaki, T. Sakihama, M. Itoh, Y. Kuniyasu, T. Nomura, M. Toda, and T. Takahashi. 2001. Immunologic tolerance maintained by CD25⁺ CD4⁺ regulatory T cells: their common role in controlling autoimmunity, tumor immunity, and transplantation tolerance. *Immunol. Rev.* 182:18–32.

12. Chatenoud, L., B. Salomon, and J.A. Bluestone. 2001. Suppressor T cells—they're back and critical for regulation of autoimmunity! *Immunol. Rev.* 182:149–163.
13. Wood, K.J., and S. Sakaguchi. 2003. Regulatory T cells in transplantation tolerance. *Nat. Rev. Immunol.* 3:199–210.
14. Singh, B., S. Read, C. Asseman, V. Malmstrom, C. Mottet, L.A. Stephens, R. Stepankova, H. Tlaskalova, and F. Powrie. 2001. Control of intestinal inflammation by regulatory T cells. *Immunol. Rev.* 182:190–200.
15. Curotto de Lafaille, M.A., and J.J. Lafaille. 2002. CD4(+) regulatory T cells in autoimmunity and allergy. *Curr. Opin. Immunol.* 14:771–778.
16. Herbelin, A., J.M. Gombert, F. Lepault, J.F. Bach, and L. Chatenoud. 1998. Mature mainstream TCR alpha beta⁺ CD4⁺ thymocytes expressing I-selectin mediate “active tolerance” in the nonobese diabetic mouse. *J. Immunol.* 161:2620–2628.
17. Lepault, F., and M.C. Gagnerault. 2000. Characterization of peripheral regulatory CD4⁺ T cells that prevent diabetes onset in nonobese diabetic mice. *J. Immunol.* 164:240–247.
18. Takahashi, T., T. Tagami, S. Yamazaki, T. Uede, J. Shimizu, N. Sakaguchi, T.W. Mak, and S. Sakaguchi. 2000. Immunologic self-tolerance maintained by CD25⁺ CD4⁺ regulatory T cells constitutively expressing cytotoxic T lymphocyte-associated antigen 4. *J. Exp. Med.* 192:303–310.
19. Shimizu, J., S. Yamazaki, T. Takahashi, Y. Ishida, and S. Sakaguchi. 2002. Stimulation of CD25(+)CD4(+) regulatory T cells through GITR breaks immunological self-tolerance. *Nat. Immunol.* 3:135–142.
20. McHugh, R.S., M.J. Whitters, C.A. Piccirillo, D.A. Young, E.M. Shevach, M. Collins, and M.C. Byrne. 2002. CD4(+) CD25(+) immunoregulatory T cells: gene expression analysis reveals a functional role for the glucocorticoid-induced TNF receptor. *Immunity.* 16:311–323.
21. Read, S., V. Malmstrom, and F. Powrie. 2000. Cytotoxic T lymphocyte-associated antigen 4 plays an essential role in the function of CD25⁺ CD4⁺ regulatory cells that control intestinal inflammation. *J. Exp. Med.* 192:295–302.
22. Taylor, P.A., C.J. Lees, and B.R. Blazar. 2002. The infusion of ex vivo activated and expanded CD4(+)CD25(+) immune regulatory cells inhibits graft-versus-host disease lethality. *Blood.* 99:3493–3499.
23. Hoffmann, P., J. Ermann, M. Edinger, C.G. Fathman, and S. Strober. 2002. Donor-type CD4⁺ CD25⁺ regulatory T cells suppress lethal acute graft-versus-host disease after allogeneic bone marrow transplantation. *J. Exp. Med.* 196:389–399.
24. Edinger, M., P. Hoffmann, J. Ermann, K. Drago, C.G. Fathman, S. Strober, and R.S. Negrin. 2003. CD4⁺ CD25⁺ regulatory T cells preserve graft-versus-tumor activity while inhibiting graft-versus-host disease after bone marrow transplantation. *Nat. Med.* 9:1144–1150.
25. Lohr, J., B. Knoechel, S. Jiang, A.H. Sharpe, and A.K. Abbas. 2003. The inhibitory function of B7 costimulators in T cell responses to foreign and self-antigens. *Nat. Immunol.* 4:664–669.
26. Graca, L., S.P. Cobbold, and H. Waldmann. 2002. Identification of regulatory T cells in tolerated allografts. *J. Exp. Med.* 195:1641–1646.
27. Asseman, C., S. Mauze, M.W. Leach, R.L. Coffman, and F. Powrie. 1999. An essential role for interleukin 10 in the function of regulatory T cells that inhibit intestinal inflammation. *J. Exp. Med.* 190:995–1004.
28. Kingsley, C.I., M. Karim, A.R. Bushell, and K.J. Wood. 2002. CD25⁺ CD4⁺ regulatory T cells prevent graft rejection: CTLA-4- and IL-10-dependent immunoregulation of alloresponses. *J. Immunol.* 168:1080–1086.
29. Belkaid, Y., C.A. Piccirillo, S. Mendez, E.M. Shevach, and D.L. Sacks. 2002. CD4⁺ CD25⁺ regulatory T cells control *Leishmania major* persistence and immunity. *Nature.* 420:502–507.
30. Thornton, A.M., and E.M. Shevach. 1998. CD4⁺ CD25⁺ immunoregulatory T cells suppress polyclonal T cell activation in vitro by inhibiting interleukin 2 production. *J. Exp. Med.* 188:287–296.
31. Piccirillo, C.A., J.J. Letterio, A.M. Thornton, R.S. McHugh, M. Mamura, H. Mizuhara, and E.M. Shevach. 2002. CD4⁺ CD25⁺ regulatory T cells can mediate suppressor function in the absence of transforming growth factor β 1 production and responsiveness. *J. Exp. Med.* 196:237–246.
32. Dieckmann, D., H. Plattner, S. Berchtold, T. Berger, and G. Schuler. 2001. Ex vivo isolation and characterization of CD4⁺ CD25⁺ T cells with regulatory properties from human blood. *J. Exp. Med.* 193:1303–1310.
33. Apostolou, I., A. Sarukhan, L. Klein, and H. von Boehmer. 2002. Origin of regulatory T cells with known specificity for antigen. *Nat. Immunol.* 3:756–763.
34. Levings, M.K., R. Sangregorio, and M.G. Roncarolo. 2001. Human CD25⁺ CD4⁺ T regulatory cells suppress naive and memory T cell proliferation and can be expanded in vitro without loss of function. *J. Exp. Med.* 193:1295–1302.
35. Jonuleit, H., E. Schmitt, M. Stassen, A. Tuettenberg, J. Knop, and A.H. Enk. 2001. Identification and functional characterization of human CD4⁺ CD25⁺ T cells with regulatory properties isolated from peripheral blood. *J. Exp. Med.* 193:1285–1294.
36. Levings, M.K., R. Sangregorio, C. Sartirana, A.L. Moschin, M. Battaglia, P.C. Orban, and M.G. Roncarolo. 2002. Human CD25⁺ CD4⁺ T suppressor cell clones produce transforming growth factor β , but not interleukin 10, and are distinct from type 1 T regulatory cells. *J. Exp. Med.* 196:1335–1346.
37. Barthlott, T., G. Kassiotis, and B. Stockinger. 2003. T cell regulation as a side effect of homeostasis and competition. *J. Exp. Med.* 197:451–460.
38. Stockinger, B., T. Barthlott, and G. Kassiotis. 2001. T cell regulation: a special job or everyone's responsibility? *Nat. Immunol.* 2:757–758.
39. Kukreja, A., G. Cost, J. Marker, C. Zhang, Z. Sun, K. Lin-Su, S. Ten, M. Sanz, M. Exley, B. Wilson, et al. 2002. Multiple immuno-regulatory defects in type-1 diabetes. *J. Clin. Invest.* 109:131–140.
40. Gregori, S., N. Giarratana, S. Smiroldo, and L. Adorini. 2003. Dynamics of pathogenic and suppressor T cells in autoimmune diabetes development. *J. Immunol.* 171:4040–4047.
41. Yamazaki, S., T. Iyoda, K. Tarbell, K. Olson, K. Velinzon, K. Inaba, and R.M. Steinman. 2003. Direct expansion of functional CD25⁺ CD4⁺ regulatory T cells by antigen-processing dendritic cells. *J. Exp. Med.* 198:235–247.
42. Tarbell, K.V., M. Lee, E. Ranheim, C.C. Chao, M. Sanna, S.K. Kim, P. Dickie, L. Teyton, M. Davis, and H. McDevitt. 2002. CD4⁺ T cells from glutamic acid decarboxylase (GAD)65-specific T cell receptor transgenic mice are not diabetogenic and can delay diabetes transfer. *J. Exp. Med.* 196:481–492.
43. Masteller, E.L., M.R. Warner, W. Ferlin, V. Judkowski, D. Wilson, N. Glaichenhaus, and J.A. Bluestone. 2003. Peptide-MHC class II dimers as therapeutics to modulate antigen-specific T cell responses in autoimmune diabetes. *J. Immunol.* 171:5587–5595.
44. Lenschow, D.J., Y. Zeng, K.S. Hathcock, L.A. Zuckerman,

- G. Freeman, J.R. Thistlethwaite, G.S. Gray, R.J. Hodes, and J.A. Bluestone. 1995. Inhibition of transplant rejection following treatment with anti-B7-2 and anti-B7-1 antibodies. *Transplantation*. 60:1171–1178.
45. Gavin, M.A., S.R. Clarke, E. Negrou, A. Gallegos, and A. Rudensky. 2002. Homeostasis and anergy of CD4(+)CD25(+) suppressor T cells in vivo. *Nat. Immunol.* 3:33–41.
46. Tang, Q., K.J. Henriksen, E.K. Boden, A.J. Tooley, J. Ye, S.K. Subudhi, X.X. Zheng, T.B. Strom, and J.A. Bluestone. 2003. Cutting edge: CD28 controls peripheral homeostasis of CD4⁺ CD25⁺ regulatory T cells. *J. Immunol.* 171:3348–3352.
47. Kanagawa, O., A. Militech, and B.A. Vaupel. 2002. Regulation of diabetes development by regulatory T cells in pancreatic islet antigen-specific TCR transgenic nonobese diabetic mice. *J. Immunol.* 168:6159–6164.
48. Qin, S., S.P. Cobbold, H. Pope, J. Elliott, D. Kioussis, J. Davies, and H. Waldmann. 1993. “Infectious” transplantation tolerance. *Science*. 259:974–977.
49. Chen, Z.K., S.P. Cobbold, H. Waldmann, and S. Metcalfe. 1996. Amplification of natural regulatory immune mechanisms for transplantation tolerance. *Transplantation*. 62:1200–1206.
50. van Maurik, A., M. Herber, K.J. Wood, and N.D. Jones. 2002. Cutting edge: CD4⁺ CD25⁺ alloantigen-specific immunoregulatory cells that can prevent CD8⁺ T cell-mediated graft rejection: implications for anti-CD154 immunotherapy. *J. Immunol.* 169:5401–5404.
51. Reijonen, H., W.W. Kwok, and G.T. Nepom. 2003. Detection of CD4⁺ autoreactive T cells in T1D using HLA class II tetramers. *Ann. NY Acad. Sci.* 1005:82–87.
52. Eisenbarth, G.S., and G.T. Nepom. 2002. Class II peptide multimers: promise for type 1A diabetes? *Nat. Immunol.* 3:344–345.
53. Maus, M.V., J.L. Riley, W.W. Kwok, G.T. Nepom, and C.H. June. 2003. HLA tetramer-based artificial antigen-presenting cells for stimulation of CD4⁺ T cells. *Clin. Immunol.* 106:16–22.