

Lawrence Berkeley National Laboratory

Recent Work

Title

COLLISIONAL BREAKUP OF HIGH-ENERGY H₂⁺ IONS

Permalink

<https://escholarship.org/uc/item/8136t12q>

Authors

Berkner, Klaus H.

Kaplan, Selig N.

Pyle, Robert V.

et al.

Publication Date

1965-11-18

University of California
Ernest O. Lawrence
Radiation Laboratory

COLLISIONAL BREAKUP OF HIGH-ENERGY H_2^+ IONS

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

Berkeley, California

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Submitted to the Physical Review

UCRL-11217

UNIVERSITY OF CALIFORNIA
Lawrence Radiation Laboratory
Berkeley, California

AEC Contract No. W-7405-eng-48

COLLISIONAL BREAKUP OF HIGH-ENERGY H_2^+ IONS

Klaus H. Berkner, Selig N. Kaplan, Robert V. Pyle,
and J. Warren Stearns

November 18, 1965

Collisional Breakup of High-Energy H_2^+ Ions[†]

Klaus H. Berkner,* Selig N. Kaplan, Robert V. Pyle,[‡]
and J. Warren Stearns

Lawrence Radiation Laboratory
University of California
Berkeley, California

November 18, 1965

ABSTRACT

The gas collisional cross sections for breakup of 20.9-MeV H_2^+ ions into H_0+H^+ and H^++H^++e have been measured in H_2 , He, N_2 , and Ar. Total dissociation cross sections in the same gases and in the energy range 10 to 65 MeV have been deduced from radial-attenuation measurements on the internal beam of the Berkeley 88-inch cyclotron. These cross sections have also been calculated for each of the H_2^+ molecular-ion vibrational levels and for an average over the vibrational levels, with an approach proposed by Salpeter. The measured and calculated average cross sections are in good agreement.

INTRODUCTION

The gas collisional cross sections for breakup of H_2^+ ions have been studied fairly extensively in the past decade, particularly below about 250 keV. In attempts to explain the rather large differences among the earlier measurements, experimental attention has been focused on the effects of possibly differing, vibrational-population distributions. It is now generally agreed that measured dissociation cross sections are essentially independent of the type of ion source from which the H_2^+ beams are obtained. On the other hand, variations attributable to special vibrational distributions have been observed with H_2^+ beams produced by gas collisional breakup of H_3^+ ions^{1, 2} (up to 40% greater than from an ion source), or by ionization of H_2 gas with electrons having energies within a few electron volts of the H_2 ionization threshold^{3, 4} (cross section variations of about a factor of four).

No theoretical estimates of dissociation cross sections for energies below approximately 1 MeV have been published. In the high-energy range, Salpeter's approximate calculations, made a number of years ago for the H_2^+ ground-vibrational-state ion,⁵ gave quite good agreement with total cross-section values obtained by Effat.⁶ (Effat measured the radial attenuation of a cyclotron's internal beam in the energy range from 9 to 18 MeV in N_2 and Ar gases.) Recently Peek, Weihofen, and Green have reported more exact calculations for specific electronic transitions in a molecular-hydrogen target,⁷ and have obtained a total dissociation cross section

using a closure approximation. The approaches of Salpeter and of Peek et al. give results that are in very good agreement with each other.

In this paper we report measurements of the total and partial gas collisional cross sections in H_2 , He, N_2 , and Ar at a sufficiently high energy, 20.9 MeV, that the Salpeter model should apply in all cases. The ion source was a high-voltage PIG type that is assumed to produce ions with an initial vibrational distribution calculable by use of the Franck-Condon principle. We have used Salpeter's approach to calculate breakup cross sections for all of the bound vibrational levels of the H_2^+ molecule.

Several measurements with gas-cell targets have been reported for energies above 1 MeV (approximately the energy for which the model should begin to give reasonable results). Barnett,⁸ Sweetman and Riviere,⁹ and Pivovar, Tubaev, and Novikov¹⁰ have reported H_2 , He, N_2 , and Ar data at maximum energies of 2.25, 3 and, 1.2 MeV, respectively. Goldring et al.¹¹ made measurements in H_2 and N_2 at 3 MeV, and Röpke and Spehl¹² went up to 3.5 MeV in Ar. Their results are included for comparison with our measurements.


As an alternative to gas-cell measurements, it is possible to deduce total cross sections by observing the variation with radius of the internal beam of a cyclotron,⁶ provided certain assumptions are made about the dependence of the cross section on the H_2^+ energy and about the operating parameters of the cyclotron. We mention briefly some internal-cyclotron-beam measurements in the Berkeley 88-inch cyclotron for H_2^+ ions in the energy range 10 to 65 MeV, but our main emphasis is on the gas-cell measurements in the Berkeley heavy-ion linear accelerator (Hilac) beam; no assumptions about energy dependence and cyclotron parameters enter these calculations.

II. THEORY

The gas collisional breakup of H_2^+ can result in dissociation (denoted by the cross section σ_1)



or ionization (denoted by σ_2)



Electron capture is not important for H_2^+ energies above a few hundred keV. The contributions to σ_1 from the $1s\sigma_g - 2p\sigma_u$ electronic transition has been investigated by Peek⁷ in the first Born approximation for collisions with H; collisions with H_2 are being investigated by Peek, Weihofen, and Green.⁷ We know of no theoretical calculations for collisions with other gases; however, a method for estimating these cross sections at incident speeds that are large compared to the orbital speed of the bound electron in the H_2^+ ion has been prescribed by Salpeter:⁵ The breakup of H_2^+ may be achieved either by electronic excitation to an unstable state or by excitation of nuclear vibrations. Salpeter reduced the problem at high-impact energies to a study of collisions in which (a) a free electron of the same velocity as the H_2^+ ion experiences a change of momentum, K , sufficient either to excite an unstable electronic state of the H_2^+ ion ($K^2/2m_e > 12.5$ eV) or to remove the bound electron ($K^2/2m_e > 30$ eV); and (b) a free proton of the same velocity as the H_2^+ ion experiences a change of momentum, K , sufficient to dissociate H_2^+ ions ($K^2/2M_p > 2.65$ eV). (The numerical values are for the ground vibrational state). He calculated the cross sections for the necessary momentum transfers for collisions with atoms of H, N, O, and Ar, using Mott and Massey's results¹³ for large-angle scattering in the first Born approximation.

Although the Salpeter calculations are for H_2^+ ions in the ground vibrational state ($v=0$), the H_2^+ ions used in our experiment were undoubtedly distributed over various vibrational states. To determine the effects of such a distribution we have used the Salpeter prescription to calculate σ_1 and σ_2 for all vibrational states ($v=0$ to 18) of an H_2^+ ion in the lowest rotational state. The effect of a distribution over rotational states has not been estimated.

The breakup resulting from electronic excitation of H_2^+ was determined as follows: For a particular internuclear separation R , we calculated the probability that a free electron of the same velocity as the H_2^+ ion undergoes a change of momentum K sufficient to excite the $2p\sigma_u$ state (leading to the total breakup cross section)

$$\frac{K^2}{2m_e} \geq U_1(R) \equiv |V(2p\sigma_u) - V(1s\sigma_g)|_R \quad (3)$$

or the two-proton state (ionization)

$$\frac{K^2}{2m_e} \geq U_2(R) \equiv |V(H^+ + H^+) - V(1s\sigma_g)|_R \quad (4)$$

The quantities U_1 and U_2 are indicated in Fig. 1.

Momentum transfer to the electron can be achieved by either elastic or inelastic scattering from the target. The elastic scattering contribution to the cross section for each of these processes was obtained by integrating the differential scattering cross sections given by Motz, Olsen, and Koch¹⁴ from

$$\theta_{\min}(R) = \left(\frac{2U(R)}{m_e v^2} \right)^{1/2} \quad (5)$$

to π . Thus we have

$$\sigma_{2, el}^{(R)} = 2\pi \int_{\theta_{\min}(U_2)}^{\pi} I(\theta) \sin\theta \, d\theta \quad (6)$$

and

$$\sigma_{T, el}^{(R)} = 2\pi \int_{\theta_{\min}(U_1)}^{\pi} I(\theta) \sin\theta \, d\theta, \quad (7)$$

where $I(\theta)$ is the differential elastic-scattering cross section and σ_T is the total breakup cross section, $\sigma_T \equiv \sigma_1 + \sigma_2$, due to electronic excitation of H_2^+ .

Two different methods were used to calculate the contribution from inelastic scattering. One was to use the differential scattering cross sections of Marton and Schiff¹⁵ (based on first Born approximation with Hartree-Fock wave functions) for small momentum transfer, and those of Morse¹⁶ (first Born approximation with Thomas-Fermi screening) for large momentum transfer; a graphical interpolation was used for the intermediate region. Since the validity of our method is restricted to high impact energies, Eq. (5) was also used to relate the energy transfer to the scattering angle for inelastic collisions. Alternatively, Gryzinski's¹⁷ "semi-classical" result for transferring energy greater than U [Eq. (10), Ref. 17] was used. The value of U used in evaluating this equation was the greater of $U_1(U_2)$ or the lowest excitation energy of the target molecule under consideration.

The contribution from momentum transfer to either of the two protons, producing dissociation by vibrational excitation ($K^2/2M \geq$ the binding energy of vibrational state v), was calculated from the approximate formula used by Salpeter:

$$\sigma_p = 4\pi(Z + Z^2)e^4 [E_{in}E(v)]^{-1}, \quad (8)$$

where Z is the atomic number of the target, E_{in} is the H_2^+ energy, and $E(v)$ is the binding energy of the vibrational state v . For the lower lying levels, the contributions to the breakup cross sections from vibrational excitation are small. However, the implication by this formula-tion that cross sections for excitation to nearby vibrational levels are quite large thereby suggests the possibility of dissociation through cascade processes. The effects of cascading are not included in our analysis.

The cross section for the dissociation or ionization of a partic-ular vibrational state v was then obtained by averaging $\sigma(R)$ over all inter-nuclear positions described by the vibrational wave functions of Cohen, Hiskes, and Riddell.¹⁸

$$\sigma_2(v) = \sigma_{2,el}(v) + \sigma_{2,incl}(v) = \int \psi_v^2(R) [\sigma_{2,el}(R) + \sigma_{2,incl}(R)] R^2 dR \quad (9)$$

$$\sigma_T(v) = \sigma_{T,el}(v) + \sigma_{T,incl}(v) + \sigma_p = \int \psi_v^2(R) [\sigma_{T,el}(R) + \sigma_{T,incl}(R)] R^2 dR + \sigma_p(v) \quad (10)$$

$$\sigma_1(v) = \sigma_{1,el}(v) + \sigma_{1,incl}(v) + \sigma_p = \sigma_T(v) - \sigma_2(v). \quad (11)$$

Although we have integrated over the internuclear separations [Eqs. (9) and (10)] to obtain the results given in this article, consider-able computational effort can be saved by considering ψ_v to be local-ized at the classical outer turning points in the vibrational potential well. We find that with this approximation we tend to overestimate σ_2 by about 5 to 15% and σ_T by as much as 40%.

III. THE EXPERIMENTAL PROCEDURE

The experimental arrangement is shown in Fig. 2. A momentum-analyzed beam of H_2^+ , (nominally 20-MeV) from the Berkeley Hilac passed through the second 24-cm long gas cell (shown at the bottom of the figure) through a pair of magnets, and into plastic scintillator-photomultiplier counters. The signals from the counters were pulse-height analyzed to demonstrate that the proper particles were being counted; discriminators were set so as to make contributions from low-energy scattered particles negligible, and the pulses were scaled. The target gases were hydrogen, helium, nitrogen, and argon at pressures ranging from 1 to 100 millitorr. The quarter-inch-diameter beam was easily contained in the two-inch-diameter plastic scintillator counters. The method of taking data was to set a pressure, use the neutral hydrogen beam as a monitor, and count the H^+ and H_2^+ beams in the other counter by varying the magnetic field in the analyzing magnet. In a given run enough counts were obtained to give a maximum of 3% statistical uncertainty in a measurement, and measurements made at different times and with different apparatus over a period of one year gave consistent results.

Although the targets were fairly thin for beams of this energy, it was necessary to know the cross sections for ionizing neutral hydrogen atoms in interpreting the data. These cross sections were obtained by breaking up the H_2^+ atoms in the first gas target, sweeping the ions out of the way with immediately following magnet and determining ionization cross sections of the hydrogen atom with the second gas cell.

Background corrections were necessary for both the H^0 and H^+ count rates. The H^0 background was due to collisional breakup of H_2^+

ions in the low-pressure but fairly long drift tubes between the targets and magnets, and was always repeatable. The background in the H^+ count rate was also due in part to H_2^+ breakup on gas in the drift tubes, but to a greater degree was from grazing collisions with the collimators of the gas cell. Variation of the size of the contribution from the collimator collisions, due to beam steering from the accelerator, resulted in somewhat larger uncertainties in the σ_2 cross section than in σ_1 . During the experiment gas pressures in the targets were monitored with Schulz-Phelps ionization gauges that were cross calibrated with McLeod gauges and an oil manometer. The Gaede effect¹⁹ was determined by repeating the pressure calibrations with the mercury of the McLead gauge cooled to $0^\circ C$.¹⁹ The maximum effect was 5% for the pressures used in this experiment. From the internal consistency of the pressure calibrations, including day-to-day fluctuations, we set a maximum uncertainty of $\pm 10\%$ in target thickness. After combining these various effects we estimate that the standard error in σ_1 is $\pm 15\%$ and in σ_2 is $\pm 20\%$.

Radial beam-attenuation measurements were also carried out in the Berkeley 88-inch cyclotron (with the cooperation of Hermann Grunder) for 10- to 65-MeV H_2^+ ions in hydrogen, helium, nitrogen, and argon. As the results of careful measurements of the beam phase with respect to the rf as a function of radius were available,²⁰ the energy gain per turn could be determined precisely. Attenuation measurements were made at a number of pressures for each gas; with the assumption that the cross sections decrease as $1/E$, it was possible to make simple corrections for the background gas of unknown composition inside the cyclotron. However, there was an uncertainty in the pressure measurements because

the ion gauges were at the vacuum wall rather than in the cyclotron gap and, in addition, only total current to the probe was measured, so that contributions from low-energy "spurium" particles may have been present. If we neglect the latter factor, we might conclude that the cross section values obtained in this way should be good to about $\pm 30\%$, although an accurate estimate of the uncertainty is impossible. The total attenuation cross sections obtained with this technique agreed with the gas-cell results to within the 30% uncertainty estimate.

IV. RESULTS

The results of the 20.9-MeV H_2^+ and 10.4-MeV H^0 cross section measurements are summarized in Table I. Values of the total cross sections from this table are plotted in Figs. 3 through 6. Also shown are the results of a number of other experiments at energies above 100 keV and the average values of σ_T in the energy range 10 to 65 MeV that were deduced from our internal-cyclotron-beam measurements and Effat's similar measurements at 9 to 18 MeV. It must be emphasized that we have no real way to estimate the uncertainties of our cyclotron values so the data are of only casual interest.

In Tables II through V are the calculated dissociation cross sections for all 19 vibrational levels. The results are broken down to show the contributions from each of the terms in Eqs. (9) and (11). The inelastic and total cross sections are tabulated for both a Born approximation (B) and a semiclassical, Gryzinski calculation (G) of the inelastic contribution. The tabulated cross sections for \bar{v} were obtained by averaging over a vibrational-level population distribution calculated²¹ according to the Franck-Condon principle. For H_2 and N_2 the Born approximation cross sections given are simply twice the calculated atomic

cross sections. For the Gryzinski inelastic scattering probabilities, however, we interpreted the "minimum excitation energy U ", used in his Eqs. (7) and (10) of Ref. 17, to be the lowest electronic excitation energy of the molecule.²² The molecular targets could also have been treated as two atoms, as in the Born calculations. In this case, the only large difference is in $\sigma_{1,inel}$, the value of which is about 40% lower for two atoms than for a molecule of N_2 , and 15% higher for H_2 .

The solid lines of Figs. 3 through 6 are $1/E$ extrapolations of the Born-approximation results for representative values of v and \bar{v} . The $1/E$ dependence of the cross sections was confirmed by calculations at lower energies. Plotted as a broken line on each figure is the total cross section according to the Gryzinski calculation. For illustrative purposes the curves are extended to lower energies than the Born approximation warrants.

V. DISCUSSION

Our values for measured high-energy breakup cross sections are in good agreement with calculated values for averages over a Franck-Condon distribution of level populations,²³ when first-Born-approximation values are used for both the elastic and inelastic electron-scattering probabilities. Inelastic-scattering contributions from the Gryzinski semiclassical approach give apparently excessive values for the molecular gases.

A successful model should also predict the relative values of σ_1 and σ_2 . Examples of the ratio σ_2/σ_1 for a few vibrational levels are given in Table VI. The trend with Z , and the ratios for $v=0$ and \bar{v}

calculated on the basis of the Born approximation throughout, are in rough agreement with our measurements and with the results of the other experiments referenced in this paper. Unfortunately, neither the experimental results nor our knowledge of the vibrational distribution are precise enough to permit a good comparison.

The excellent agreement of the average cross section in H_2 calculated by Peek et al.⁷ (see Fig. 3) with that calculated in the Salpeter approximation, and the good agreement between our measured and calculated cross sections in H_2 , He, N_2 , and Ar inspired considerable confidence in the usefulness of Salpeter's approach. As mentioned earlier, the calculated values of the vibrational excitation terms (σ_p) do not make a significant contribution to the cross sections because they are relatively small for the lower lying vibrational states (the averaged cross sections correspond roughly to those for $v = 4$ or 5 , which are two of the most heavily weighted levels for Franck-Condon transitions). However, since σ_p varies inversely with threshold energy [Eq. (8)], and since the binding energy of the highest vibrational levels is quite small, this term becomes extremely large and, in fact, dominates the cross sections at the highest levels.

The only high-energy experiment that we are acquainted with that bears on collisional dissociation of the highly excited vibrational levels was reported by Riviere and Sweetman.²⁴ They measured the fraction of the total H_2^+ beam that was in the uppermost vibrational levels by electric field dissociation. After passing the beam through a gas target sufficiently thick to dissociate a large fraction of the incoming beam, they found that the fraction of the emerging beam in the uppermost levels had dropped only slightly. The small decrease does

not necessarily imply a failure in the model, since the form of Eq. (8) and the philosophy in which it was used would also imply very large cross sections for excitation to neighboring vibrational levels-- suggesting a mechanism for repopulation of the higher levels by an upward cascade. This characteristic of the cross section, however, remains to be demonstrated.

ACKNOWLEDGMENTS

We are grateful to Dr. C. M. Van Atta for supporting and encouraging this research. One of us (SNK) thanks Professors Burton J. Moyer and A. C. Helmholtz for the support and interest that enabled him to participate. We also thank Dr. John G. Conway, Dr. John R. Hiskes, and Dr. James M. Peek for several helpful discussions. Vincent J. Honey helped with the experimental work.

FOOTNOTES AND REFERENCES

† This work was performed under the auspices of the U. S. Atomic Energy Commission.

* Present address (until Sept. 1, 1966): UK AEA Culham Laboratory, Culham, Abingdon, Berks., England.

‡ Present address (until June 2, 1966): Physico-Technical Institute of the Academy of Sciences of the UkSSR, Khar'kov, USSR.

1. A. C. Riviere and D. R. Sweetman, Proc. Phys. Soc. (London) 78, 1215 (1961).
2. C. F. Barnett and J. A. Ray, "The Dissociation Cross Section of H_2^+ in Hydrogen Gas," in Atomic Collision Processes, M. R. C. McDowell, Ed. (North-Holland Publishing Co., Amsterdam, 1964), p. 743.
3. J. Wm. McGowan and Larkin Kerwin, Can. J. Phys. 42, 972 (1964).
4. N. N. Tunitskii, P. M. Smirnova, and M. V. Tknomirnov, Dokl. Akad. Nauk SSSR 101, 1083 (1955).
5. E. E. Salpeter, Proc. Phys. Soc. (London) A63, 1297 (1950).
6. K. E. A. Effat, Proc. Phys. Soc. (London) A65, 29 (1952).
7. James M. Peek, W. H. Weihofen, and Thomas A. Green, Paper NB3, Fourth International Conference on Physics of Electronic and Atomic Collisions, Quebec, Aug. 1965; and James M. Peek, Sandia Laboratory, Albuquerque, N. M. (private communication). See also Phys. Rev. 140, A11 (1965).
8. C. F. Barnett, Proc. of Second United Nations International Conference on the Peaceful Uses of Atomic Energy (United Nations, Geneva, 1958), 32, 398.

9. D. R. Sweetman, Proc. Roy. Soc. (London) A256, 416 (1960), and A. C. Riviere and D. R. Sweetman, private communication.
10. L. I. Pivovarov, V. M. Tubaev, and M. T. Novikov, Sov. Phys. JETP 13, 23 (1961).
11. G. Goldring, D. Kedem, U. Smilansky, and Z. Vager, Nucl. Instr. Methods 23, 231 (1963).
12. H. Röpke and H. Spehl, Nucl. Instr. Methods 17, 169 (1962).
13. N. F. Mott and H. S. W. Massey, The Theory of Atomic Collisions (Oxford, Clarendon Press, 1949).
14. J. W. Motz, H. Olsen, and H. W. Kock, Rev. Mod. Phys. 36, 881 (1964).
15. L. Marton and L. I. Schiff, J. Appl. Phys. 12, 759 (1941).
16. P. M. Morse, Phys. Zeitschrift 33, 443 (1932).
17. M. Gryzinski, Phys. Rev. 138, A336 (1965).
18. S. Cohen, J. R. Hiskes, and R. J. Riddell, Jr., Vibrational States of the Hydrogen Molecular Ion, Lawrence Radiation Laboratory Report UCRL-8871, 1959 (unpublished).
19. See, for example, Ch. Meinke and G. Reich, Vacuum 13, 579 (1963), or Vakuum Technik 12, 79 (1963).
20. These measurements were made concurrently by Alper Garren and Lloyd Smith for another purpose. We are indebted to them for their results and interpretation.
21. J. R. Hiskes, Nucl. Fusion 2, 38 (1962).
22. For example, the six 2p electrons of N_2 were considered to have an excitation threshold of 6.1 eV and the four 2s electrons 10.3 eV.

23. We are not suggesting that the experimental H_2^+ beam has a vibrational population distribution given by the Franck-Condon principle, even if it were created with one in the ion source. However, this distribution seems to be a plausible one to try.
24. A. C. Riviere and D. R. Sweetman, The Dissociation of Molecular Ions by Strong Electric Fields, in Proceedings of the Fifth Int. Conf. on Ionization Phenomena in Gases, Munich 1961 (North-Holland Publishing Company, Amsterdam, 1962), p. 1236.

Table I. Cross sections per target molecule in units of 10^{-18} cm².
 The absolute uncertainties in σ_1 and σ_2 are estimated to be $\pm 15\%$ and $\pm 20\%$, respectively. The uncertainty in the ratio σ_2/σ_1 is about 20%.

	20.9-MeV H_2^+				10.4-MeV H^0
	σ_1	σ_2	σ_T	σ_2/σ_1	σ_{01}
H ₂	1.53	1.69	3.22	1.10	2.24
He	1.06	1.77	2.83	1.67	1.59
N ₂	10.9	20.3	31.2	1.88	19.9
Ar	12.3	35.4	47.7	2.87	35.8

Table II. Calculated cross sections for 20-MeV H_2^+ ions in hydrogen, in units of 10^{-18} cm²/molecule. ^a

v	$\sigma_{1,el}$	$\sigma_{1,incl}$		σ_p	σ_1		$\sigma_{2,el}$	$\sigma_{2,incl}$		σ_2	
		B	G		B	G		B	G	B	G
0	0.24	1.07	6.42	0.02	1.33	6.68	0.51	0.63	1.68	1.14	2.19
1	0.26	1.31	6.60	0.02	1.59	6.88	0.54	0.68	1.84	1.22	2.38
2	0.25	1.42	6.34	0.02	1.70	6.61	0.52	0.66	1.83	1.18	2.35
3	0.26	1.58	6.21	0.03	1.87	6.50	0.52	0.66	1.87	1.18	2.39
4	0.28	1.82	6.06	0.03	2.13	6.37	0.54	0.71	2.02	1.24	2.56
5	0.29	2.02	6.42	0.04	2.35	6.75	0.55	0.72	2.10	1.27	2.65
6	0.30	2.24	6.41	0.04	2.58	6.75	0.56	0.74	2.20	1.30	2.75
7	0.31	2.47	6.41	0.05	2.83	6.76	0.56	0.76	2.30	1.33	2.86
8	0.32	2.72	6.23	0.06	3.10	6.60	0.57	0.79	2.41	1.36	2.98
9	0.31	2.95	6.22	0.07	3.33	6.60	0.56	0.79	2.44	1.35	3.00
10	0.32	3.42	6.28	0.09	3.82	6.68	0.59	0.84	2.67	1.46	3.26
11	0.33	3.64	6.19	0.11	4.07	6.63	0.58	0.84	2.68	1.41	3.26
12	0.34	4.18	6.32	0.15	4.68	6.82	0.60	0.89	2.90	1.49	3.50
13	0.34	4.61	6.19	0.20	5.16	6.74	0.60	0.90	3.00	1.51	3.61
14	0.35	5.40	6.05	0.31	6.06	6.71	0.62	0.94	3.25	1.55	3.86
15	0.36	6.17	6.03	0.51	7.04	6.90	0.63	0.98	3.39	1.60	4.01
16	0.36	7.42	5.93	1.05	8.84	7.34	0.62	1.02	3.58	1.66	4.22
17	0.36	9.12	5.62	3.15	12.63	9.14	0.65	1.07	3.92	1.72	4.57
18	0.35	10.3	5.20	29.1	39.8	34.7	0.67	1.15	4.42	1.82	5.10
Av	0.28	1.96	6.37	0.05	2.29	6.70	0.54	0.71	2.06	1.25	2.60

^aThe elastic scattering contributions $\sigma_{1,el}$ and $\sigma_{2,el}$ are calculated in the Born approximation. The inelastic contributions are calculated for both the Born and Gryzinski approximations. Total values of σ_1 and σ_2 are given in double entry indicating that Born (B) or Gryzinski (G) inelastic cross sections are used. The Gryzinski calculations are based on molecular values for the excitation and ionization energies. All other entries are twice the cross sections for a hydrogen atom.

Table III. Calculated cross sections for 20-MeV H_2^+ ions in helium, in units of 10^{-18} cm²/atom. See Table II footnote.

v	$\sigma_{1,el}$	$\sigma_{1,incl}$		σ_p	σ_1		$\sigma_{2,el}$	$\sigma_{2,incl}$		σ_2	
		B	G		B	G		B	G	B	G
0	0.12	0.60	1.48	0.029	0.75	1.63	0.55	0.68	2.08	1.23	2.63
1	0.13	0.68	1.46	0.032	0.84	1.62	0.58	0.73	2.29	1.31	2.87
2	0.13	0.69	1.26	0.036	0.86	1.43	0.55	0.71	2.28	1.26	2.83
3	0.13	0.73	1.13	0.041	0.90	1.30	0.55	0.71	2.35	1.25	2.89
4	0.14	0.81	1.06	0.047	0.99	1.24	0.57	0.75	2.53	1.32	3.10
5	0.14	0.87	0.94	0.054	1.06	1.13	0.57	0.76	2.65	1.33	3.22
6	0.14	0.93	0.82	0.062	1.13	1.02	0.58	0.78	2.77	1.35	3.35
7	0.14	1.00	0.70	0.073	1.21	0.92	0.58	0.79	2.89	1.37	3.47
8	0.15	1.08	0.58	0.087	1.32	0.82	0.58	0.82	3.00	1.40	3.58
9	0.14	1.14	0.52	0.105	1.38	0.76	0.57	0.80	2.98	1.38	3.55
10	0.16	1.29	0.40	0.130	1.58	0.69	0.59	0.86	3.19	1.44	3.77
11	0.14	1.35	0.40	0.166	1.66	0.71	0.58	0.84	3.12	1.43	3.70
12	0.15	1.52	0.35	0.220	1.89	0.72	0.60	0.89	3.27	1.49	3.88
13	0.15	1.66	0.30	0.307	2.12	0.75	0.60	0.89	3.29	1.49	3.88
14	0.15	1.92	0.24	0.461	2.53	0.85	0.61	0.92	3.35	1.53	3.96
15	0.15	2.15	0.18	0.772	3.07	1.10	0.62	0.94	3.42	1.56	4.04
16	0.15	2.51	0.12	1.58	4.24	1.86	0.62	0.97	3.48	1.60	4.10
17	0.15	2.96	0.07	4.73	7.84	4.95	0.63	1.00	3.52	1.63	4.14
18	0.14	3.26	0.02	43.7	47.1	43.9	0.64	1.05	3.57	1.69	4.21
Av	0.14	0.85	1.02	0.079	1.07	1.24	0.57	0.75	2.55	1.32	3.12

Table IV. Calculated cross sections for 20-MeV H_2^+ ions in nitrogen, in units of $10^{-18} \text{ cm}^2/\text{molecule}$. See Table II footnote.

v	$\sigma_{1, \text{el}}$	$\sigma_{1, \text{inel}}$		σ_p	σ_1		$\sigma_{2, \text{el}}$	$\sigma_{2, \text{inel}}$		σ_2	
		B	G		B	G		B	G	B	G
0	4.9	3.8	36.0	0.5	9.3	41.4	14.0	5.5	8.4	19.5	22.4
1	5.3	4.3	48.9	0.6	10.2	54.8	14.9	5.9	9.2	20.8	23.2
2	5.2	4.4	62.0	0.7	10.3	68.0	14.2	5.7	9.1	19.9	23.3
3	5.2	4.6	70.9	0.8	10.5	76.7	14.1	5.7	9.3	19.7	23.4
4	5.5	5.1	74.5	0.9	11.6	80.9	14.7	6.0	10.1	20.6	24.7
5	5.7	5.5	78.6	1.0	12.2	85.3	14.8	6.1	10.5	20.9	25.3
6	5.8	6.0	81.4	1.2	12.8	88.3	14.9	6.1	11.0	21.2	25.9
7	5.9	6.4	83.9	1.4	13.6	91.2	15.1	6.3	11.4	21.4	26.6
8	6.0	6.9	86.4	1.6	14.5	94.1	15.4	6.4	12.0	21.8	27.4
9	5.9	7.2	86.4	2.0	15.1	94.3	15.0	6.3	12.2	21.3	27.2
10	6.3	8.2	92.6	2.4	16.9	101	15.8	6.7	13.3	22.5	29.1
11	6.2	8.6	90.2	3.1	17.8	100	15.4	6.5	13.4	22.0	28.8
12	6.4	9.7	94.4	4.1	20.2	105	16.0	6.9	14.5	22.9	30.5
13	6.4	10.5	94.8	5.7	22.7	107	16.0	6.9	15.0	22.9	31.0
14	6.5	12.2	96.1	8.6	27.4	111	16.2	7.1	16.2	23.3	32.4
15	6.6	13.5	98.0	14.4	34.5	119	16.5	7.2	16.9	23.7	33.4
16	6.6	15.0	96.8	29.5	51.2	133	16.7	7.4	17.8	24.2	34.6
17	6.7	16.1	99.2	88.3	111	194	16.8	7.6	19.5	24.4	36.3
18	6.4	17.1	98.7	816	840	920	16.4	7.9	22.1	25.3	39.4
Av	5.5	5.4	71.3	1.5	12.4	78.3	14.7	6.0	10.3	20.7	25.9

Table V. Calculated cross sections for 20-MeV H_2^+ ions in argon, in units of $10^{-18} \text{ cm}^2/\text{atom}$. See Table II footnote.

v	$\sigma_{1, \text{el}}$	$\sigma_{1, \text{inel}}$		σ_p	σ_1		$\sigma_{2, \text{el}}$	$\sigma_{2, \text{inel}}$		σ_2	
		B	G		B	G		B	G	B	G
0	11.0	3.8	16.6	1.7	16.4	29.2	31.6	5.8	7.4	37.4	38.9
1	10.3	4.2	16.4	1.8	16.4	28.5	33.4	6.2	7.9	39.6	41.4
2	11.6	4.3	15.2	2.1	18.0	28.9	32.0	6.0	7.8	38.0	39.8
3	11.7	4.6	14.8	2.3	18.7	28.9	31.7	6.0	7.8	37.7	39.6
4	12.4	5.0	15.1	2.7	20.1	30.2	33.1	6.3	8.3	39.4	41.5
5	12.6	5.4	14.9	3.1	21.1	30.6	33.4	6.4	8.6	39.8	42.0
6	13.0	5.8	14.8	3.5	22.3	31.3	33.8	6.5	8.9	40.2	42.7
7	13.2	6.2	14.7	4.2	23.6	32.1	34.1	6.6	9.2	40.7	43.3
8	13.5	6.8	14.7	4.9	25.2	33.2	34.7	6.7	9.5	41.4	44.3
9	13.3	7.1	13.9	6.0	26.4	33.3	33.8	6.6	9.6	40.4	43.4
10	14.2	8.1	14.3	7.4	29.6	35.9	35.6	7.0	10.3	42.6	46.0
11	13.8	8.4	13.6	9.5	31.7	37.1	34.7	6.8	10.3	41.6	45.0
12	14.5	9.5	13.8	12.5	36.5	40.8	36.1	7.1	11.0	43.6	47.2
13	14.4	10.2	13.4	17.5	42.0	45.2	36.1	7.2	11.3	43.3	47.4
14	14.6	11.2	12.8	26.3	52.1	53.7	36.6	7.3	12.1	43.9	48.7
15	14.7	11.4	12.6	44.0	70.1	71.3	37.2	7.5	12.5	44.7	49.7
16	14.8	11.7	12.2	90.1	117	117	37.8	7.7	12.5	45.4	50.9
17	14.7	12.4	11.2	270	287	296	38.2	7.9	14.1	46.1	52.3
18	14.4	12.7	9.8	2490	2520	2520	39.1	8.2	15.6	47.3	54.8
Av	12.2	5.3	15.0	4.5	27.0	31.8	33.1	6.3	8.5	39.4	41.6

Table VI. The ratio σ_2/σ_1 for several vibrational levels with the inelastic contributions calculated from both the Born and the Gryzinski formulae.

	v	H ₂	He	N ₂	Ar
Born	0	0.86	1.64	2.10	2.28
	5	0.54	1.25	1.71	1.88
	10	0.38	0.91	1.33	1.44
	18	0.05	0.04	0.03	0.02
	\bar{v}_B	0.55	1.23	1.67	1.79
Gryzinski	0	0.33	1.61	0.54	1.33
	5	0.39	2.85	0.30	1.37
	10	0.49	5.46	0.29	1.28
	18	0.15	0.10	0.04	0.02
	\bar{v}_G	0.39	2.52	0.33	1.31
Exptl.		1.10	1.67	1.88	2.87
		± 0.22	± 0.35	± 0.38	± 0.57

FIGURE CAPTIONS

Fig. 1. Potential-energy diagram.

Fig. 2. Experimental arrangement.

Fig. 3. Total breakup cross section, $\sigma_T = \sigma_1 + \sigma_2$, in hydrogen. The solid lines are the Born approximation results of Table II plus similar calculations at other energies. These results are inversely proportional to the H_2^+ kinetic energy. The broken line labeled \bar{v}_G is for an average over vibrational levels, when the Gryzinski inelastic cross sections are used. The broken line labeled \bar{v}_{PWG} is the averaged cross section obtained by Peek et al. in a closure approximation. The square at 20.9 MeV is our gas-cell measurement, and the cross-hatched area indicates the value obtained from our internal-cyclotron-beam measurement, with its uncertainty of $\pm 30\%$. Other experimental results are 1. J. Guidini, Ionization Phenomena in Gases, Munich 1961 (North-Holland Publishing Company, Amsterdam, 1962), p. 1228; 2. Ref. 9; 3. Ref. 10; 4. Ref. 8; and 5. Ref. 11.

Fig. 4. The total breakup cross section, $\sigma_T = \sigma_1 + \sigma_2$, in helium. See also Fig. 3 caption.

Fig. 5. The total breakup cross section, $\sigma_T = \sigma_1 + \sigma_2$, in nitrogen. The cross-hatched area (Effat) shows the results of Ref. 6. See also Fig. 3 caption.

Fig. 6. The total breakup cross section, $\sigma_T = \sigma_1 + \sigma_2$, in argon. The experimental curve labeled 6 is from Ref. 12. The cross-hatched area (Effat) shows the results of Ref. 6. See also Fig. 3 caption.




Fig. 1

MUB-8874





Fig. 2

MU-32832-A



MUB-8875

Fig. 3




Fig. 4

MUB-8876




Fig. 5

MUB-8877




Fig 6

MUB-8878

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

