UCLA Proceedings of UCLA Health

Title

Cognitive Consequences of Curative Treatment for Hodgkin's Lymphoma

Permalink https://escholarship.org/uc/item/8183v6xq

Journal Proceedings of UCLA Health, 22(1)

Authors Hakakian, Benjamin Ashouri, Shahryar

Publication Date 2017-12-04

Cognitive Consequences of Curative Treatment for Hodgkin's Lymphoma

Benjamin Hakakian and Shahryar Ashouri, MD

A 26-year-old male graduate student in engineering, presented with enlarged lymph nodes in his right neck area, along with drenching night sweats, in October of 2016. Aside from this new issue, he was in good health with no co-morbidities. A core biopsy of his cervical lymph node was highly suspicious for classical Hodgkin lymphoma. The patient was referred to us for oncologic evaluation. Staging for Hodgkin's lymphoma was started and he was also underwent full excisional biopsy of his lymph node, for definitive diagnosis. A PET/CT scan revealed multiple masses in the lymph nodes of the neck, chest, and anterior mediastinum (3.1 x 11.5 x 12.9 cm) and a SUV of 15.1. The excisional biopsy of the right neck lymph node confirmed diagnosis of nodular sclerosis Hodgkin lymphoma. A bone marrow biopsy revealed no involvement of the bone marrow. His final stage was Stage IIB, with bulky disease.

Next, baseline echocardiogram and pulmonary function tests (PFT's) were done, in addition to being referred to fertility specialist for sperm banking. The patient was then started on combination chemotherapy, with doxorubicin, bleomycin, vinblastine, and dacarbazine regimen (ABVD). After his second cycle, the anterior mediastinal mass decreased to 2.2 x 6.3 x 6.4 cm with a SUV of 2.2. Echocardiogram and PFT's, were repeated after every 2 cycles and remained normal with a stable ejection fraction (EF) and diffusing capacity of the lungs for carbon monoxide (DLCO). Despite this, he developed mild dyspnea on exertion, therefore, given his good response on PET/CT, his bleomycin was discontinued after cycle #2. Upon completion of his sixth cycle, his anterior mediastinal mass decreased to 1.2 x 4.4 x 5.4 cm, with a SUV of 1.7. There were also no remnants of activity in other lymph nodes. His symptoms of dyspnea resolved, and he returned to his baseline activities within 2 months of completion of his chemotherapy. In addition, he was able to finish his master degree in engineering, with honors.

A growing number of studies have reported that chemotherapy and adjuvant therapies may impair cognitive functions, with some cognitive changes persisting in a subset of cancer survivors.^{1,2} The effect of adjuvant chemotherapy treatment is linked to decreased function in multiple facets of cognition, mainly including areas like attention, concentration, language, motor skills, multitasking, verbal memory, and visual memory.³⁻⁵ The effect of chemotherapy on areas of cognition is known colloquially as chemo-fog or chemo-brain.⁶ Researchers have used a battery of neurological and psychological tests on cancer patients to assess cognitive functioning. The most

common and persistent cognitive deficiencies reported were in the domain of frontal lobe function. These deficiencies include information processing speed, verbal memory, and working memory.^{1,7,8} Common chemotherapy regimens used to treat Hodgkins lymphoma involve ABVD (Adriamycin, Bleomycin, Vinblastine, and Dacarbazine), CHOP (Cyclophosphamide/ Hydroxydaunorubicin, Oncovin/Vincristine, and Prednisone or Prednisolone), or a combination of the monoclonal antibody Rituximab (R) with either ABVD or CHOP. Baudino et al. has shown that cancer patients with lymphoma treated with either ABVD, CHOP, R-ABVD, or R-CHOP have shown decreases in cognitive function compared to controls, specifically in areas involving frontal lobe function.⁸ These patients also showed significantly lower bilateral rate of glucose metabolism in prefrontal cortices, cerebellum, medial cortices, and limbic brain areas. There was a negative correlation between metabolism in these areas and number of chemotherapy treatment cycles, while there was no significant difference between chemotherapy treated patients and controls in regard to depression, anxiety, and distress.

Chemo-brain has varied duration, with some patients being affected in the short-term and others having symptoms that persist for years after treatment.^{1,4,9} Shagen et al. reported patients with complete recovery from cognitive deficiencies four years post-treatment.¹⁰ The model supported by Baudino et al. suggests cognitive decline due to chemotherapy tends to be transient, but long lasting. They posit that the cognitive deficits last for at least 6 months post treatment. Other studies also show that structural and functional changes accompany treatment with chemotherapy, with structural changes occurring both in grey and white matter.^{11,12} White matter hyperintensities (WMH) can become apparent as soon as 2 months after chemotherapy treatment, and usually last between 6 months to a year.¹³ WMH that last longer than a year seem to be permanent. A prospective study done by McDonald et al. found a similar trend with grey matter volume decline as the aforementioned trend.¹⁴ Patients treated with chemotherapy had a decrease in grey matter density in bilateral frontal, temporal, and cerebellar regions, as well as in the right thalamus as soon as one month post-treatment. Partial recovery was seen in some brain areas by end of year one, as well as what the study termed persistent gray matter volume decrease. Recovery in gray matter volume was not found for patients who were not at least partially recovering by the first year post-treatment.¹⁴ These studies imply is that the one year mark post-treatment might be a useful prognostic tool to gauge whether cognitive recovery

will occur long term. de Ruiter et al. investigated effects of adjuvant chemotherapy treatment on breast cancer survivors 9 years post-treatment to assess the long-term effects of treatment.¹⁵ Even after 9 years, these patients had hypoactivations in the bilateral posterior parietal cortex during memory encoding and visuospatial planning tests, as well as a reduction in grey matter volume in the left posterior parietal cortex. They also showed a loss in white matter volume in regions adjacent to the parietal cortex, possibly explaining persisting cognitive deficits.

These studies raise questions about optimal ways of treating cancer patients, whether through choice of chemotherapy regimens that have minimized incidence of cognitive deficiencies, alternative treatment options, or preventative interventions. Because the exact mechanism for pathogenesis of chemo-brain is not fully understood, and causes of chemobrain can vary from person to person, this is a difficult question. Possible mechanisms include vascular injury, oxidative damage, inflammation, direct injury to neurons, autoimmune responses, or chemotherapy-induced anemia.^{16,17} Interventions that help alleviate symptoms from these mechanisms have been tried as potential treatment. These include supplementing or increasing antioxidant intake as it can decrease vascular injury causing free radicals, supplementing Vitamin E to help against age-based cognitive decline, and performing aerobic exercise as it can increase blood flow to the brain and mitigate cognitive deficits.16,18,19

Pharmacological approaches have had minor success in minimizing risk for cognitive issues as a result of adjuvant chemotherapy treatment. One option was the use of D-Methylphenidate (d-MPH), a compound that has been clinically shown to help with attention-deficit/hyperactivity disorder and narcolepsy.²⁰ Unfortunately, two separate randomized controlled trials have shown that d-MPH did not improve cognitive function in breast cancer survivors compared to controls, while it did help with chemotherapy induced fatigue.^{21,22} Another pharmacological approach was the use of the psychostimulant Modafinil, which improved cognitive performance in breast cancer survivors by enhancing speed of memory, quality of episodic memory, and mean continuity of attention.²³

At this time, more research is needed to find a more definitive treatment option to prevent or ameliorate cognitive dysfunction associated with chemotherapy treatment. The literature supports the notion that chemo-brain resulting from adjuvant chemotherapy treatment is real and requires particular care in order to help patients overcome cancer without experiencing long-term cognitive side effects. Fortunately, it has not yet affected our patient.

REFERENCES

- Ahles TA, Saykin AJ, Furstenberg CT, Cole B, Mott LA, Skalla K, Whedon MB, Bivens S, Mitchell T, Greenberg ER, Silberfarb PM. Neuropsychologic impact of standard-dose systemic chemotherapy in longterm survivors of breast cancer and lymphoma. *J Clin Oncol.* 2002 Jan 15;20(2):485-93. PubMed PMID: 11786578.
- Castellon SA, Ganz PA, Bower JE, Petersen L, Abraham L, Greendale GA. Neurocognitive performance in breast cancer survivors exposed to adjuvant chemotherapy and tamoxifen. J Clin Exp Neuropsychol. 2004 Oct;26(7):955-69. PubMed PMID: 15742545.
- Anderson-Hanley C, Sherman ML, Riggs R, Agocha VB, Compas BE. Neuropsychological effects of treatments for adults with cancer: a meta-analysis and review of the literature. *J Int Neuropsychol Soc.* 2003 Nov;9(7):967-82. PubMed PMID: 14738279.
- 4. **Tannock IF, Ahles TA, Ganz PA, Van Dam FS**. Cognitive impairment associated with chemotherapy for cancer: report of a workshop. *J Clin Oncol.* 2004 Jun 1;22(11):2233-9. Review. PubMed PMID: 15169812.
- Meyers CA. Neurocognitive dysfunction in cancer patients. *Oncology (Williston Park)*. 2000 Jan;14(1):75-9; discussion 79, 81-2, 85. PubMed PMID: 10680150.
- Olin JJ. Cognitive function after systemic therapy for breast cancer. *Oncology (Williston Park)*. 2001 May;15(5):613-8; discussion 618, 621-4. Review. PubMed PMID: 11396356.
- Raffa RB, Duong PV, Finney J, Garber DA, Lam LM, Mathew SS, Patel NN, Plaskett KC, Shah M, Jen Weng HF. Is 'chemo-fog'/'chemo-brain' caused by cancer chemotherapy? *J Clin Pharm Ther.* 2006 Apr;31(2):129-38. Review. PubMed PMID: 16635046.
- Baudino B, D'agata F, Caroppo P, Castellano G, Cauda S, Manfredi M, Geda E, Castelli L, Mortara P, Orsi L, Cauda F, Sacco K, Ardito RB, Pinessi L, Geminiani G, Torta R, Bisi G. The chemotherapy long-term effect on cognitive functions and brain metabolism in lymphoma patients. *Q J Nucl Med Mol Imaging*. 2012 Dec;56(6):559-68. Epub 2012 Nov 21. PubMed PMID: 23172518.
- 9. Ahles TA. Do systemic cancer treatments affect cognitive function? *Lancet Oncol.* 2004 May;5(5):270-1. PubMed PMID: 15120662.
- 10. Schagen SB, Muller MJ, Boogerd W, Rosenbrand RM, van Rhijn D, Rodenhuis S, van Dam FS. Late effects of adjuvant chemotherapy on cognitive function: a follow-up study in breast cancer patients. *Ann Oncol.* 2002 Sep;13(9):1387-97. PubMed PMID: 12196364.
- Deprez S, Amant F, Yigit R, Porke K, Verhoeven J, Van den Stock J, Smeets A, Christiaens MR, Leemans A, Van Hecke W, Vandenberghe J, Vandenbulcke M, Sunaert S. Chemotherapy-induced structural changes in cerebral white matter and its correlation with impaired cognitive functioning in breast cancer patients. *Hum Brain Mapp.* 2011 Mar;32(3):480-93. doi: 10.1002/hbm.21033. PubMed PMID: 20725909.

- Brown MS, Simon JH, Stemmer SM, Stears JC, Scherzinger A, Cagnoni PJ, Jones RB. MR and proton spectroscopy of white matter disease induced by high-dose chemotherapy with bone marrow transplant in advanced breast carcinoma. *AJNR Am J Neuroradiol*. 1995 Nov-Dec;16(10):2013-20. PubMed PMID: 8585489.
- 13. Brown MS, Stemmer SM, Simon JH, Stears JC, Jones RB, Cagnoni PJ, Sheeder JL. White matter disease induced by high-dose chemotherapy: longitudinal study with MR imaging and proton spectroscopy. *AJNR Am J Neuroradiol.* 1998 Feb;19(2):217-21. PubMed PMID: 9504468.
- McDonald BC, Conroy SK, Ahles TA, West JD, Saykin AJ. Gray matter reduction associated with systemic chemotherapy for breast cancer: a prospective MRI study. *Breast Cancer Res Treat*. 2010 Oct;123(3):819-28. doi: 10.1007/s10549-010-1088-4. Epub 2010 Aug 6. PubMed PMID: 20690040; PubMed Central PMCID: PMC3661415.
- 15. de Ruiter MB, Reneman L, Boogerd W, Veltman DJ, Caan M, Douaud G, Lavini C, Linn SC, Boven E, van Dam FS, Schagen SB. Late effects of high-dose adjuvant chemotherapy on white and gray matter in breast cancer survivors: converging results from multimodal magnetic resonance imaging. *Hum Brain Mapp*. 2012 Dec;33(12):2971-83. doi: 10.1002/hbm.21422. Epub 2011 Sep 23. PubMed PMID: 22095746.
- Barton D, Loprinzi C. Novel approaches to preventing chemotherapy-induced cognitive dysfunction in breast cancer: the art of the possible. *Clin Breast Cancer*. 2002 Dec;3 Suppl 3:S121-7. Review. PubMed PMID: 12533273.
- Minisini A, Atalay G, Bottomley A, Puglisi F, Piccart M, Biganzoli L. What is the effect of systemic anticancer treatment on cognitive function? *Lancet Oncol.* 2004 May;5(5):273-82. Review. PubMed PMID: 15120664.
- Perkins AJ, Hendrie HC, Callahan CM, Gao S, Unverzagt FW, Xu Y, Hall KS, Hui SL. Association of antioxidants with memory in a multiethnic elderly sample using the Third National Health and Nutrition Examination Survey. *Am J Epidemiol*. 1999 Jul 1;150(1):37-44. PubMed PMID: 10400551.
- 19. Hassmén P, Ceci R, Bäckman L. Exercise for older women: a training method and its influences on physical and cognitive performance. *Eur J Appl Physiol Occup Physiol.* 1992;64(5):460-6. PubMed PMID: 1612088.
- Jensen PS. Clinical considerations for the diagnosis and treatment of ADHD in the managed care setting. *Am J Manag Care*. 2009 May;15(5 Suppl):S129-40. Review. PubMed PMID: 19601688.
- 21. Mar Fan HG, Clemons M, Xu W, Chemerynsky I, Breunis H, Braganza S, Tannock IF. A randomised, placebo-controlled, double-blind trial of the effects of dmethylphenidate on fatigue and cognitive dysfunction in women undergoing adjuvant chemotherapy for breast cancer. Support Care Cancer. 2008 Jun;16(6):577-83. Epub 2007 Oct 31. PubMed PMID: 17972110.

- Lower EE, Fleishman S, Cooper A, Zeldis J, Faleck H, Yu Z, Manning D. Efficacy of dexmethylphenidate for the treatment of fatigue after cancer chemotherapy: a randomized clinical trial. *J Pain Symptom Manage*. 2009 Nov;38(5):650-62. doi: 10.1016/j.jpainsymman.2009.03.011. PubMed PMID: 19896571.
- Kohli S, Fisher SG, Tra Y, Adams MJ, Mapstone ME, Wesnes KA, Roscoe JA, Morrow GR. The effect of modafinil on cognitive function in breast cancer survivors. *Cancer*. 2009 Jun 15;115(12):2605-16. doi: 10.1002/cncr.24287. PubMed PMID: 19309747; PubMed Central PMCID: PMC2796482.

Submitted December 4, 2017