

UC Davis

UC Davis Previously Published Works

Title

A class of two-dimensional AKLT models with a gap

Permalink

<https://escholarship.org/uc/item/8240v1qf>

Authors

Nachtergaele, Bruno

Young, Amanda

Lucia, Angelo

et al.

Publication Date

2023-12-13

DOI

10.1090/conm/741/14917

Peer reviewed

A CLASS OF TWO-DIMENSIONAL AKLT MODELS WITH A GAP

HOUSSAM ABDUL-RAHMAN, MARIUS LEMM, ANGELO LUCIA, BRUNO NACHTERGAELE,
AND AMANDA YOUNG

ABSTRACT. The AKLT spin chain is the prototypical example of a frustration-free quantum spin system with a spectral gap above its ground state. Affleck, Kennedy, Lieb, and Tasaki also conjectured that the two-dimensional version of their model on the hexagonal lattice exhibits a spectral gap. In this paper, we introduce a family of variants of the two-dimensional AKLT model depending on a positive integer n , which is defined by decorating the edges of the hexagonal lattice with one-dimensional AKLT spin chains of length n . We prove that these decorated models are gapped for all $n \geq 3$.

1. INTRODUCTION

A central question concerning a quantum spin system is whether it is gapped or gapless. (We say a system is gapped if its Hamiltonian exhibits a uniform spectral gap above the ground state. Otherwise, it is gapless.) The existence of a spectral gap is known to have wide-ranging consequences for the system's low energy physics. For instance, the ground states of gapped Hamiltonians display exponential clustering [21, 35] and, in one dimension, they are known to satisfy various notions of bounded complexity [3, 4, 20, 29]. Of particular interest are the spin liquid states conjectured to describe a number of interesting two- and three-dimensional systems [26, 45, 13]. Moreover, with the advent of Hastings' spectral flow [19] (also called quasi-adiabatic evolution), it has become possible to explore gapped ground state phases in considerable detail. Different gapped phases are separated from each other by quantum phase transitions, which are accompanied by a closing of the spectral gap [6, 12]. Accordingly, numerous recent works are concerned with the stability of the spectral gap under finite-range perturbations assuming local topological order [12, 31, 32, 36]. From these considerations, it would be desirable to have a multitude of gapped Hamiltonians that one can use as starting points for further analysis. However, proving the existence of a spectral gap is a non-trivial mathematical task and there exist only limited tools [15, 16, 22, 27, 30] and only a few special models in which a spectral gap has been rigorously established [2, 5, 7, 9, 10, 11, 27, 30], particularly in dimensions ≥ 2 . We also mention in passing that deciding whether a general Hamiltonian is gapped or not is known to be undecidable in general, even for reasonable (i.e., translation-invariant and local) Hamiltonians [8, 14]. See also [33].

The foundational work in the field was done by Affleck, Kennedy, Lieb and Tasaki (AKLT in the following) in 1988 [1, 2]. Motivated by a famous conjecture of Haldane [17, 18] that predicts a spectral gap for the one-dimensional integer-spin Heisenberg antiferromagnet, AKLT provided two main contributions that proved seminal in the years to come:

Date: October 31, 2019.


- (1) They defined what is now called the one-dimensional AKLT chain: a spin-1, isotropic (i.e., $SU(2)$ -invariant) antiferromagnet on a one-dimensional chain. They found that it has a unique ground state in the thermodynamic limit and rigorously established a spectral gap.
- (2) They defined analogous spin- $z/2$ AKLT models on any z -regular bipartite graph. They focused on the hexagonal lattice case (so spin-3/2) and derived exponential decay of correlations in the infinite-volume ground state. These facts led AKLT to conjecture that the hexagonal model is also gapped.

We make two further remarks about the AKLT model: (a) The ground state of the hexagonal AKLT model was proved to have a unique [24] thermodynamic limit when the limit is taken with boundary conditions in a certain natural class. It is also known that the ground state in finite-volume with periodic boundary conditions (a finite honeycomb lattice wrapped on a torus) is unique [25]. Kennedy, Lieb, and Tasaki also proved exponential decay of spin-spin correlations, which is significant because it shows that the AKLT antiferromagnet does not exhibit Néel order, in contrast to its spin-3/2 Heisenberg analog, and that there very likely is a spectral gap above the ground state. (b) Historically, the AKLT chain provided the first example of a Hamiltonian whose ground states are matrix product states. This notion has been vastly generalized, starting with [15], to what are now called tensor network states, and has developed into a central tenet of modern many-body physics [37, 42].

The conjecture of AKLT that the hexagonal AKLT model (or any other AKLT model in dimension ≥ 2) is gapped remains open to this day. This is insofar remarkable as all of the AKLT models share a key feature that makes the spectral gap problem in principle more amenable: They are frustration-free, meaning that the global ground state is also locally energy-minimizing.

In the present work, we introduce a novel family of AKLT models on ‘decorated’ hexagonal lattices depending on an integer parameter n , and prove that these models are gapped for sufficiently large values of n . We call these models the *edge-decorated AKLT models* (or *decorated AKLT models* for short). The positive integer n will be called the *decoration number*, and we explain its role in the next paragraph.

The decorated AKLT model is defined by replacing each edge of the hexagonal lattice with a copy of the one-dimensional AKLT chain of length n . Notice that this means that there are two types of vertices in the system: vertices of the hexagonal lattice which have degree 3 and spin 3/2, and “internal vertices” of the decorated edges which have degree 2 and spin 1; see Figure 1. The heuristic behind this construction is that the decorated AKLT model incorporates features of the one-dimensional AKLT chain, which is known to be gapped from the work of AKLT. While the decorated model is a bit contrived, it is not unreasonable to expect its ground state(s) to belong to the same gapped phase as those of the original AKLT model on the hexagonal lattice. It seems likely that the same features that generated interest in two-dimensional AKLT models [38, 40, 43] are also present for the decorated AKLT model. In particular, we mention [44], where it was shown that the valence-bond ground states of similarly decorated AKLT models can serve as a universal resource for quantum computation. Going beyond AKLT-type models, an $SU(3)$ spin liquid with \mathbb{Z}_3 topological order has recently been proposed in [28]. It too is expected to be gapped.

FIGURE 1. The decorated hexagonal lattice for $n = 2$.

The strategy that we use to derive the spectral gap of the decorated model follows two main steps. We begin by taking the square of the Hamiltonian, as usual. Step 1 is to employ an inequality due to Fannes, Nachtergaele, and Werner [15] which relates the anticommutator between interaction terms to the angle between ground state projections via a duality argument. This inequality reduces the claim to a sufficiently strong bound on the angle between two ground state projections that overlap along one decorated edge (so mainly along a one-dimensional AKLT chain of length n). Step 2 is to establish the desired angle bound by a computation with quasi-one-dimensional matrix product states. With an eye toward possible future applications, we generalize the last computation (of the angle between ground state projections overlapping on a chain) to other models with matrix product ground states.

2. AKLT MODELS ON DECORATED TWO-DIMENSIONAL LATTICES

For concreteness, we will first discuss in detail an AKLT model on a honeycomb lattice with additional spins on the edges. It will then be straightforward to consider generalizations to which the same arguments apply.

Let Γ be the hexagonal lattice and $n \geq 1$. The standard AKLT model on Γ [24] has a spin-3/2 degree of freedom at each vertex. For the ‘decorated’ models we introduce here, we add n spin 1’s along each edge of Γ and call the resulting ‘lattice’ $\Gamma^{(n)}$. On this graph with both degree 2 and degree 3 vertices (see Figure 1) we define the AKLT Hamiltonian as usual with nearest neighbor interactions given by the orthogonal projection $P^{(z(e)/2)}$ onto the space of total spin $z(e)/2$, where for any edge e , $z(e)$ is the sum of the degrees of its two vertices. For the AKLT model on $\Gamma^{(1)}$ all the interaction terms are $P^{(5/2)}$. For $n \geq 2$, the model also has interactions $P^{(2)}$ between neighboring spin 1’s. This class of models is a special case of the general class of AKLT-type models studied in [25]. There, it is shown that they are frustration-free and also that the ground state is non-degenerate if the model is considered with periodic boundary conditions. The frustration-freeness is easily proved in the same way as for the original AKLT models by using the Valence Bond Solid construction of a non-zero vector in the kernel of the manifestly non-negative Hamiltonian.

Let Λ be a suitable finite subset of Γ considered with periodic boundary conditions and denote by $\Lambda^{(n)}$ its decoration as above. Let $\mathcal{E}_{\Lambda^{(n)}}$ denote the set of edges of the decorated

graph and consider the Hamiltonian

$$(2.1) \quad H_{\Lambda^{(n)}} = \sum_{e \in \mathcal{E}_{\Lambda^{(n)}}} P_e^{(z(e)/2)}.$$

We claim that for n large enough there is $\gamma_n > 0$ such that for all nice Λ the gap of $H_{\Lambda^{(n)}}$ is lower bounded by γ_n . In this context, ‘nice’ Λ , means that one can consider the decorated graph $\Lambda^{(n)}$ as a union of overlapping subgraphs isomorphic to the H-shaped graph shown in Figure 2. For concreteness, we will explicitly treat the case of periodic boundary conditions (a finite rectangle cut out of the hexagonal lattice and wrapped around a torus). Other shapes can be considered without any significant change in the arguments.

To prove the claim we consider a comparable model defined as follows. For each vertex v in Λ and the three edges meeting in v , we consider the subsystem consisting of the spin $3/2$ at v and the $3n$ spin 1 ’s residing on the three edges. Let Y_v denote the corresponding set of $3n + 1$ vertices in $\Lambda^{(n)}$ and define h_v to be the AKLT Hamiltonian on Y_v . Then

$$(2.2) \quad H_{\Lambda^{(n)}} \leq \sum_{v \in \Lambda} h_v \leq 2H_{\Lambda^{(n)}}.$$

To simplify things further, define P_v to be the orthogonal projection onto $\text{ran } h_v$. It is a straightforward calculation to check that $\ker P_v = \ker h_v$ is 8-dimensional for $n = 1$, and hence for all larger values of n as well. We will estimate the gap of

$$(2.3) \quad \tilde{H}_{\Lambda^{(n)}} = \sum_{v \in \Lambda} P_v,$$

which is also comparable to $H_{\Lambda^{(n)}}$:

$$(2.4) \quad \frac{1}{2}\gamma_Y \tilde{H}_{\Lambda^{(n)}} \leq H_{\Lambda^{(n)}} \leq \|h_Y\| \tilde{H}_{\Lambda^{(n)}},$$

with $\gamma_Y > 0$. This inequality implies that $\tilde{H}_{\Lambda^{(n)}}$ is also frustration-free since $\ker(\tilde{H}_{\Lambda^{(n)}}) = \ker(H_{\Lambda^{(n)}}) \neq \{0\}$. Therefore, it suffices to study the gap of $\tilde{H}_{\Lambda^{(n)}}$.

We will obtain a lower bound for the gap of $\tilde{H}_{\Lambda^{(n)}}$, by finding a constant $\gamma > 0$ satisfying

$$(2.5) \quad (\tilde{H}_{\Lambda^{(n)}})^2 = \tilde{H}_{\Lambda^{(n)}} + \sum_{\{v,w\} \subset \Lambda, v \neq w} (P_v P_w + P_w P_v) \geq \gamma \tilde{H}_{\Lambda^{(n)}}.$$

If v and w are *not* nearest neighbors, P_v and P_w commute and $P_v P_w + P_w P_v \geq 0$. Therefore, in the second term we can drop all contributions from such pairs. For the nearest neighbor pairs (v, w) , instead of the combinatorial style argument in [27], which requires good estimates of a specific finite-volume gap, we apply the following inequality for a pair orthogonal projections E and F (for a proof see [15, Lemma 6.3]):

$$(2.6) \quad EF + FE \geq -\|EF - E \wedge F\|(E + F).$$

Here, $E \wedge F$ is the orthogonal projection onto $\text{ran } E \cap \text{ran } F$. We need this for $E = P_v, F = P_w$, with v, w nearest neighbors in Λ .

The norm in (2.6) remains unchanged if we replace E and F by $\mathbb{1} - E$ and $\mathbb{1} - F$, which, in our application, are the corresponding ground state projections. All nearest neighbor

pairs are equivalent in this consideration and we denote $\mathbb{1} - P_v = P_{\succ}$, $\mathbb{1} - P_w = P_{\prec}$, and $P_{\succ} \wedge P_{\prec} = P_{\succ\prec}$. Define

$$(2.7) \quad \varepsilon_n = \|P_{\succ}P_{\prec} - P_{\succ\prec}\|.$$

Since every $v \in \Lambda$ has 3 nearest neighbors, we have shown

$$(2.8) \quad (\tilde{H}_{\Lambda^{(n)}})^2 \geq (1 - 3\varepsilon_n)\tilde{H}_{\Lambda^{(n)}}.$$

Therefore,

$$(2.9) \quad \text{gap}(H_{\Lambda^{(n)}}) \geq \frac{1}{2}\gamma_Y(1 - 3\varepsilon_n).$$

It remains to show that $\varepsilon_n < 1/3$.

Proposition 2.1. *Let*

$$A_n = \frac{4}{3^n \left(1 - \frac{8(1+3^{-2n-1})}{3^n(1-3^{-2n})}\right)}.$$

Then, for all $n \geq 3$, the quantity ε_n defined in (2.7) satisfies

$$(2.10) \quad \varepsilon_n \leq A_n + A_n^2 \left(1 + \frac{8(1+3^{-2n-1})^2}{3^n(1-3^{-2n})^2}\right) < 1/3.$$

The proof of this proposition is contained in the next two sections. As a consequence, we can state the following theorem.


Theorem 2.2. *The spectral gap above the ground state of the AKLT model on the edge-decorated honeycomb lattice with $n \geq 3$ has a strictly positive lower bound uniformly for all finite volumes with periodic boundary conditions.*

3. GROUND STATE PROJECTIONS FOR SUBGRAPHS OVERLAPPING ON A CHAIN

To prove Proposition 2.1 we will formulate the quantity ε_n in terms of the ground state projections of quasi-one-dimensional Matrix Product States (MPS). A rather straightforward generalization of the arguments in [15] to MPS systems with matrices that may vary from site to site and are not necessarily square, will then yield the desired estimate. With an eye toward possible further generalizations and applications, we will estimate ε_n in a slightly more general setting, which we now introduce. The role of the finite subgraph $\succ\prec$ of the decorated honeycomb lattice will be played a finite graph G with a certain structure and we will make a number of assumptions on the ground states of a frustration free Hamiltonian on G . In Section 4 we will show that these assumptions are satisfied for the AKLT model on the decorated honeycomb lattice.

3.1. Assumptions on the tensor network states for the local patch G . Consider a finite graph $G = (\mathcal{V}, \mathcal{E})$ of the form $G_L - C_n - G_R$, meaning there are finite graphs G_L and G_R , $C_n = [v_1, v_n]$ is a chain of n vertices, and there exist $v_L \in G_L, v_R \in G_R$ such that \mathcal{V} is the disjoint union of the vertices of G_L, C_n and G_R and \mathcal{E} consists of the edges of G_L, C_n and G_R together with (v_L, v_1) and (v_n, v_R) , see Figure 2. We consider a frustration-free Hamiltonian on G of the following form:

$$(3.1) \quad H_G = H_{G_L} + H_{C_n} + H_{G_R} + h_{v_L, v_1} + h_{v_n, v_R},$$

FIGURE 2. The graph G for the decorated AKLT model.

where H_{G_L} , H_{G_R} , and H_{C_n} satisfy the following conditions. First, we assume H_{C_n} has ground states given by a translation invariant MPS with a primitive transfer matrix \mathbb{E} . Let D denote the bond dimension of this MPS and pick an orthonormal basis $\{|i\rangle \mid 1 \leq i \leq d\}$ for the physical degree of freedom at each of the n sites of C_n . In this case the transfer matrix \mathbb{E} , in isometric form, is given in terms of d $D \times D$ matrices V_i :

$$(3.2) \quad \mathbb{E}(B) = \sum_{i=1}^d V_i^* B V_i, \quad B \in M_D, \quad \text{with } \mathbb{E}(\mathbb{1}) = \mathbb{1}.$$

The primitivity assumption implies that there exists a non-singular density matrix $\rho \in M_D$ satisfying $\mathbb{E}^t(\rho) = \rho$, and constants $C \geq 0$ and $\lambda \in [0, 1)$, such that

$$(3.3) \quad a(n) := \|\mathbb{E}^n - |\mathbb{1}\rangle\langle\rho|\| \leq C\lambda^n.$$

For the spin-1 AKLT chain one has this estimate with $C = 1$ and $\lambda = 1/3$, and both constants are sharp in that case.

We now turn to the assumptions we make on H_{G_L} and H_{G_R} . For $\sharp \in \{L, R\}$, let \mathcal{H}_\sharp be the Hilbert space associated with the system on G_\sharp . We assume that the ground states of H_{G_\sharp} are given by a tensor T^\sharp as follows. We will consider T^L as a set of $\dim \mathcal{H}_L$ $D \times D_L$ matrices labeled by an orthonormal basis $\{|r\rangle_L\}$ of \mathcal{H}_L and T^R as a set of $\dim \mathcal{H}_R$ $D_R \times D$ matrices labeled by an orthonormal basis $\{|r\rangle_R\}$ of \mathcal{H}_R . The physical Hilbert space for the system on G is $\mathcal{H}_G = \mathcal{H}_{G_L} \otimes \mathcal{H}_{C_n} \otimes \mathcal{H}_{G_R}$, and the auxiliary space, which parametrizes the ground states, is $\mathcal{K}_G = \mathbb{C}^{D_L} \otimes \mathbb{C}^{D_R}$, which we identify with $\mathcal{L}(\mathbb{C}^{D_R}, \mathbb{C}^{D_L})$ and equip with the standard inner product $\langle \cdot, \cdot \rangle_{\mathcal{K}_G}$. The map $\Gamma_G : \mathcal{K}_G \rightarrow \mathcal{H}_G$ is then given by

$$(3.4) \quad \Gamma_G(B) = \sum_{l, i_1, \dots, i_n, r} \text{Tr}[B T_r^R V_{i_n} \cdots V_{i_1} T_l^L] |l\rangle_L \otimes |i_1, \dots, i_n\rangle \otimes |r\rangle_R, \quad B \in \mathcal{K}_G.$$

We assume that H_G is frustration free, which means that all terms in (3.1) are non-negative and $\ker H_G \neq \{0\}$ and, in addition, we assume $\ker H_G = \text{ran } \Gamma_G$.

We also introduce the transfer matrices associated with G_L and G_R , $\mathbb{E}_L : M_D \rightarrow M_{D_L}$ and $\mathbb{E}_R : M_{D_R} \rightarrow M_D$, as follows:

$$(3.5) \quad \mathbb{E}_L(B) = \sum_l (T_l^L)^* B T_l^L, \quad \mathbb{E}_R(B) = \sum_r (T_r^R)^* B T_r^R,$$

and define

$$(3.6) \quad Q_L = \mathbb{E}_L(\mathbb{1}), \quad Q_R = \mathbb{E}_R^t(\rho)$$

where $\mathbb{E}_R^t(B) = \sum_r T_r^R B (T_r^R)^*$. We assume that Q_L and Q_R are non-singular.

For $\Lambda \in \{G_L, C_n, G_R, G_L - C_n, C_n - G_R\}$, considered as subsets of G , let \mathcal{H}_Λ and \mathcal{K}_Λ denote the corresponding physical and auxiliary Hilbert spaces, respectively, and define the corresponding maps $\Gamma_\Lambda : \mathcal{K}_\Lambda \rightarrow \mathcal{H}_\Lambda$ in the obvious way. These maps are of the same form as Γ_G in (3.4); if one or both parts described by G_L or G_R are absent, the absent degrees of freedom associated with G_\sharp correspond to taking $\mathcal{H}_{G_\sharp} = \mathbb{C}$, $H_{G_\sharp} = 0$, and $T^\sharp \equiv 1$.

We assume that the maps Γ_Λ are injective, meaning

$$(3.7) \quad \dim \text{ran } \Gamma_\Lambda = \dim \mathcal{K}_\Lambda,$$

and $\text{ran } \Gamma_\Lambda = \ker H_\Lambda$.

3.2. General estimate of ε_n . Let \mathcal{G}_Λ denote $\ker H_\Lambda$, and P_Λ the orthogonal projection onto \mathcal{G}_Λ . Our next goal is to estimate $\varepsilon_n = \|P_{G_L - C_n} P_{C_n - G_R} - P_G\|$. It is easy to see that ε_n is explicitly given by the following expression:

$$(3.8) \quad \varepsilon_n = \sup \left\{ \frac{|\langle \phi, \psi \rangle|}{\|\phi\| \|\psi\|} \mid \phi \in \mathcal{G}_{G_L - C_n} \otimes \mathcal{H}_{G_R}, \psi \in \mathcal{H}_{G_L} \otimes \mathcal{G}_{C_n - G_R}, \phi, \psi \perp \mathcal{G}_G, \phi, \psi \neq 0 \right\}.$$

We will derive an estimate of the type of inner products that appear in (3.8), but first recall some basic properties of MPS.

Since ρ is non-singular and positive, it defines an inner product on M_D by

$$(3.9) \quad \langle A, B \rangle_\rho = \text{Tr} \rho A^* B \quad \text{for all } A, B \in M_D,$$

and let $\|\cdot\|_\rho$ denote the corresponding norm. We will also let ρ_{\min} denote the smallest eigenvalue of ρ , which is positive by assumption. It follows that the norm $\|\cdot\|_\rho$ is equivalent to the Hilbert-Schmidt norm on M_D , which is given by $\|\cdot\|_2 = \sqrt{\text{Tr} A^* A}$. Explicitly:

$$(3.10) \quad \|A\|_2 \leq \frac{1}{\sqrt{\rho_{\min}}} \|A\|_\rho, \quad A \in M_D.$$

The map $\Gamma_{C_n} : M_D \rightarrow \mathcal{H}_{C_n}$ is explicitly given by

$$(3.11) \quad \Gamma_{C_n}(B) = \sum_{i_1, \dots, i_n} \text{Tr}[B V_{i_n} \cdots V_{i_1}] |i_1, \dots, i_n\rangle, \quad B \in M_D.$$

Lemma 3.1 ([15, Lemma 5.2]). *For any $B, C \in M_D$,*

$$(3.12) \quad |\langle \Gamma_{C_n}(B), \Gamma_{C_n}(C) \rangle - \langle B, C \rangle_\rho| \leq a(n) \text{Tr} \rho^{-1} \|B\|_\rho \|C\|_\rho.$$

Proof. Using (3.11) we can express the inner product as follows:

$$\begin{aligned}
\langle \Gamma_{C_n}(B), \Gamma_{C_n}(C) \rangle &= \sum_{i_1, \dots, i_n} \overline{\text{Tr}[BV_{i_n} \cdots V_{i_1}]} \text{Tr}[CV_{i_n} \cdots V_{i_1}] \\
(3.13) \qquad \qquad \qquad &= \sum_{i_1, \dots, i_n} \text{Tr}[V_{i_1}^* \cdots V_{i_n}^* B^*] \text{Tr}[CV_{i_n} \cdots V_{i_1}].
\end{aligned}$$

By expanding the traces using any orthonormal basis $\{|1\rangle, \dots, |D\rangle\}$ for \mathbb{C}^D , we obtain

$$\begin{aligned}
\langle \Gamma_{C_n}(B), \Gamma_{C_n}(C) \rangle &= \sum_{\alpha, \beta=1}^D \sum_{i_1, \dots, i_n} \langle \alpha | V_{i_1}^* \cdots V_{i_n}^* B^* | \alpha \rangle \langle \beta | CV_{i_n} \cdots V_{i_1} | \beta \rangle \\
(3.14) \qquad \qquad \qquad &= \sum_{\alpha, \beta=1}^D \langle \alpha | \hat{\mathbb{E}}^n(B^* | \alpha \rangle \langle \beta | C) | \beta \rangle.
\end{aligned}$$

Now observe that

$$(3.15) \qquad \qquad \langle B, C \rangle_\rho = \sum_{\alpha, \beta=1}^D \langle \alpha | \mathbb{1} \langle \rho | (B^* | \alpha \rangle \langle \beta | C) | \beta \rangle.$$

Combining these two expressions and using (3.3), we obtain

$$\begin{aligned}
|\langle \Gamma_{C_n}(B), \Gamma_{C_n}(C) \rangle - \langle B, C \rangle_\rho| &\leq \sum_{\alpha, \beta=1}^D |\langle \alpha | (\mathbb{E}^n - \mathbb{1} \langle \rho |) (B^* | \alpha \rangle \langle \beta | C) | \beta \rangle| \\
&\leq a(n) \left(\sum_{\alpha=1}^D \|B^* | \alpha \rangle\| \right) \left(\sum_{\beta=1}^D \|C^* | \beta \rangle\| \right)
\end{aligned}$$

Now, pick for the orthonormal basis one that diagonalizes ρ , such that $\rho | \alpha \rangle = \rho_\alpha | \alpha \rangle$. Then

$$\left(\sum_{\alpha=1}^D \|B^* | \alpha \rangle\| \right)^2 = \left(\sum_{\alpha=1}^D \|B^* | \alpha \rangle\| \rho_\alpha^{1/2} \rho_\alpha^{-1/2} \right)^2 \leq \sum_{\alpha=1}^D \rho_\alpha \langle \alpha | BB^* | \alpha \rangle \sum_{\alpha=1}^D \rho_\alpha^{-1} = \|B\|_\rho^2 \text{Tr} \rho^{-1}.$$

Together with the analogous estimate for the second factor, this proves the lemma. \square

Note that one has the bound

$$\text{Tr} \rho^{-1} \leq \frac{D}{\rho_{\min}},$$

which is often saturated in models with symmetry. It will be convenient to define

$$(3.16) \qquad \qquad b(n) = a(n) \text{Tr} \rho^{-1}$$

The following is an immediate corollary of Lemma 3.1, and shows that Γ_{C_n} is injective for sufficiently large n .

Corollary 3.2. *For any $B \in M_D$, the bound*

$$(3.17) \qquad \qquad \|B\|_\rho \sqrt{1 - b(n)} \leq \|\Gamma_n(B)\| \leq \|B\|_\rho \sqrt{1 + b(n)}$$

holds for n sufficiently large so that $b(n) \leq 1$.

Proof. The bound

$$(3.18) \quad \left| \|\Gamma_{C_n}(B)\|^2 - \|B\|_\rho^2 \right| \leq b(n) \|B\|_\rho^2$$

follows immediately from (3.12). If $B = 0$, there is nothing to prove. Otherwise, this bound can be re-written as

$$(3.19) \quad -b(n) \leq \frac{\|\Gamma_{C_n}(B)\|^2}{\|B\|_\rho^2} - 1 \leq b(n)$$

from which the above claim readily follows. \square

Inner products of vectors of the form $\Gamma_G(B)$, $B \in \mathcal{K}_G$ can be estimated with a straightforward generalization of Lemma 3.1. To formulate the result, for each $\Lambda \in \{G, G_L - C_n, C_n - G_R\}$ we define an inner product on \mathcal{K}_Λ , denoted by $\langle \cdot, \cdot \rangle_\Lambda$, via

$$(3.20) \quad \langle B, C \rangle_G = \text{Tr}(Q_R B^* Q_L C)$$

$$(3.21) \quad \langle B, C \rangle_{G_L - C_n} = \text{Tr}(\rho B^* Q_L C)$$

$$(3.22) \quad \langle B, C \rangle_{C_n - G_R} = \text{Tr}(Q_R B^* C)$$

That these are inner products follows from the positive-definiteness of Q_L and Q_R . With respect to these inner products we obtain the following analog of Lemma 3.1.

Lemma 3.3. *Let $\Lambda \in \{G, G_L - C_n, C_n - G_R\}$. Then for any $B, C \in \mathcal{K}_\Lambda$,*

$$(3.23) \quad |\langle \Gamma_\Lambda(B), \Gamma_\Lambda(C) \rangle - \langle B, C \rangle_\Lambda| \leq a(n) D^2 C_\Lambda \|B\| \|C\|,$$

where

$$(3.24) \quad C_G = \|\mathbb{E}_L\| \|\mathbb{E}_R\|, \quad C_{G_L - C_n} = \|\mathbb{E}_L\|, \quad \text{and} \quad C_{C_n - G_R} = \|\mathbb{E}_R\|.$$

Proof. We prove the bound in the case of $\Lambda = G$. All other cases follow from similar arguments. Let $B, C \in \mathcal{K}_G$, and $\{|1\rangle, \dots, |D\rangle\}$ be an orthonormal bases of \mathbb{C}^D . Then, calculating similar to (3.13) and (3.14), we find

$$(3.25) \quad \begin{aligned} \langle \Gamma_G(B), \Gamma_G(C) \rangle &= \sum_{\substack{\ell, r \\ i_1, \dots, i_n}} \text{Tr}[V_{i_1}^* \dots V_{i_n}^* (T_r^R)^* B^* (T_\ell^L)^*] \text{Tr}[T_\ell^L C T_r^R V_{i_n} \dots V_{i_1}] \\ &= \sum_{\alpha, \beta=1}^D \langle \alpha | \mathbb{E}^n \circ \mathbb{E}_R [B^* \mathbb{E}_L (|\alpha\rangle\langle\beta|) C] | \beta \rangle, \end{aligned}$$

where we have also use cyclicity of the trace in the first equality. Now consider $\langle B, C \rangle_G$. It can easily be shown, e.g. by simplifying the RHS, that

$$(3.26) \quad \langle B, C \rangle_G = \sum_{\alpha, \beta=1}^D \langle \alpha | \mathbb{1} \rangle \langle \rho | \circ \mathbb{E}_R [B^* \mathbb{E}_L (|\alpha\rangle\langle\beta|) C] | \beta \rangle.$$

By substituting these (3.25) and (3.26) into (3.23) and then using (3.3), we estimate as follows:

$$\begin{aligned}
|\langle \Gamma_G(B), \Gamma_G(C) \rangle - \langle B, C \rangle_G| &\leq \sum_{\alpha, \beta=1}^D |\langle \alpha | (\mathbb{E}^n - |\mathbb{1}\rangle\langle \rho|) \circ \mathbb{E}_R [B^* \mathbb{E}_L(|\alpha\rangle\langle \beta|) C] | \beta \rangle| \\
&\leq a(n) \sum_{\alpha, \beta=1}^D \|\mathbb{E}_R [B^* \mathbb{E}_L(|\alpha\rangle\langle \beta|) C]\| \\
(3.27) \qquad \qquad \qquad &\leq a(n) D^2 \|\mathbb{E}_L\| \|\mathbb{E}_R\| \|B\| \|C\|.
\end{aligned}$$

This completes the claim. \square

Note that the bound in Lemma 3.3 is expressed in terms of the operator norms $\|B\|$ and $\|C\|$. This is just a common norm of reference. The natural norm to use is the one induced by the inner product that appears on the left of (3.23), as in done in Lemma 3.1. Since these norms are all equivalent to the operator norm, the estimates from (3.23) can be converted to the ‘natural’ norm by multiplying by an appropriate constant as follows: Let q_L (resp. q_r) be the minimal eigenvalue of Q_L (resp. Q_R). Then,

$$(3.28) \quad \|B\| \leq \frac{1}{\sqrt{q_L q_R}} \|B\|_G, \quad \|B\| \leq \frac{1}{\sqrt{\rho_{\min} q_L}} \|B\|_{G_L - C_n}, \quad \|B\| \leq \frac{1}{\sqrt{q_R}} \|B\|_{C_n - G_R}.$$

We can thus obtain a corollary to Lemma 3.3 similar to Corollary 3.2.

Corollary 3.4. *Let $\Lambda \in \{G, G_L - C_n, C_n - G_R\}$. Then for any $B \in \mathcal{K}_\Lambda$,*

$$(3.29) \quad \|B\|_\Lambda \sqrt{1 - b_\Lambda(n)} \leq \|\Gamma_\Lambda(B)\| \leq \|B\|_\Lambda \sqrt{1 + b_\Lambda(n)}$$

holds for n sufficiently large so that $b_\Lambda(n) \leq 1$, where

$$(3.30) \quad b_G(n) = \frac{a(n) D^2}{q_L q_R} \|\mathbb{E}_L\| \|\mathbb{E}_R\|, \quad b_{G_L - C_n}(n) = \frac{a(n) D^2}{\rho_{\min} q_L} \|\mathbb{E}_L\|, \quad b_{C_n - G_R}(n) = \frac{a(n) D^2}{q_R} \|\mathbb{E}_R\|.$$

In order to simplify notation, we will write $b_L(n)$ for $b_{G_L - C_n}(n)$ and $b_R(n)$ for $b_{C_n - G_R}(n)$.

Ultimately, we will want a bound for the inner product in (3.8) in terms of the norms of the vectors ϕ and ψ defined below. Lemma 3.3 can also be used to show that this is once again straightforward at the cost of another prefactor in the bound.

Since $\Gamma_{G_L - C_n}$ and $\Gamma_{C_n - G_R}$ are assumed to be injective, there exist D_R $D_L \times D$ matrices $B_\phi(r)$, and D_L $D \times D_R$ matrices $B_\psi(l)$, uniquely determined by ϕ and ψ , such that

$$(3.31) \quad \phi = \sum_{l, i_1, \dots, i_n, r} \text{Tr}[B_\phi(r) V_{i_n} \cdots V_{i_1} T_l^L] |l\rangle_L \otimes |i_1, \dots, i_n\rangle \otimes |r\rangle_R$$

$$(3.32) \quad \psi = \sum_{l, i_1, \dots, i_n, r} \text{Tr}[B_\psi(l) T_r^R V_{i_n} \cdots V_{i_1}] |l\rangle_L \otimes |i_1, \dots, i_n\rangle \otimes |r\rangle_R.$$

These expressions are simply expansions of

$$(3.33) \quad \phi = \sum_r \Gamma_{G_L-C_n}(B_\phi(r)) \otimes |r\rangle_R$$

$$(3.34) \quad \psi = \sum_l |l\rangle_L \otimes \Gamma_{C_n-G_R}(B_\psi(l)).$$

It will be convenient to define $C_\phi, C_\psi \in \mathcal{K}_G$ as follows:

$$(3.35) \quad C_\phi = \sum_r B_\phi(r) \rho(T_r^R)^*$$

$$(3.36) \quad D_\psi = \sum_l (T_l^L)^* B_\psi(l).$$

Next, we consider inner products of the form $\langle \phi, \psi \rangle$, with $\phi \in \mathcal{G}_{G_L-C_n} \otimes \mathcal{H}_{G_R}$ and $\psi \in \mathcal{H}_{G_L} \otimes \mathcal{G}_{C_n-G_R}$.

Lemma 3.5. *Suppose that n is large enough so that $b(n) < 1$. Then, for all $\phi \in \mathcal{G}_{G_L-C_n} \otimes \mathcal{H}_{G_R}$ and $\psi \in \mathcal{H}_{G_L} \otimes \mathcal{G}_{C_n-G_R}$, we have*

$$(3.37) \quad |\langle \phi, \psi \rangle - \langle C_\phi, D_\psi \rangle_{\mathcal{K}_G}| \leq \frac{b(n)}{\sqrt{1-b_{LR}(n)}} \|\phi\| \|\psi\|,$$

where

$$(3.38) \quad b_{LR}(n) = b_L(n) + b_R(n) - b_L(n)b_R(n),$$

and $b_L(n)$ and $b_R(n)$ are defined in (3.30).

Proof. Using the expansions (3.31) and (3.32) we find

$$(3.39) \quad \begin{aligned} \langle \phi, \psi \rangle &= \sum_{l, i_1, \dots, i_n, r} \overline{\text{Tr}[B_\phi(r) V_{i_n} \cdots V_{i_1} T_l^L]} \text{Tr}[B_\psi(l) T_r^R V_{i_n} \cdots V_{i_1}] \\ &= \sum_{l, r} \langle \Gamma_{C_n}(T_l^L B_\phi(r)), \Gamma_{C_n}(B_\psi(l) T_r^R) \rangle \end{aligned}$$

Similarly, we observe that $\langle C_\phi, D_\psi \rangle_{\mathcal{K}_G}$ can be expressed as a sum of inner products:

$$\begin{aligned} \langle C_\phi, D_\psi \rangle_{\mathcal{K}_G} &= \text{Tr} \left(\sum_r T_r^R \rho B_\phi(r)^* \right) \left(\sum_l (T_l^L)^* B_\psi(l) \right) \\ &= \sum_{l, r} \text{Tr} \rho B_\phi(r)^* (T_l^L)^* B_\psi(l) T_r^R \\ &= \sum_{l, r} \langle T_l^L B_\phi(r), B_\psi(l) T_r^R \rangle_\rho. \end{aligned}$$

Now we apply Lemma 3.1 term by term to obtain, using Cauchy-Schwarz:

$$\begin{aligned}
|\langle \varphi, \psi \rangle - \langle C_\phi, D_\psi \rangle_{\mathcal{K}_G}| &\leq \sum_{l,r} |\langle \Gamma_{C_n}(T_l^L B_\phi(r)), \Gamma_{C_n}(B_\psi(l) T_r^R) \rangle - \langle T_l^L B_\phi(r), B_\psi(l) T_r^R \rangle_\rho| \\
&\leq b(n) \sum_{l,r} \|T_l^L B_\phi(r)\|_\rho \cdot \|B_\psi(l) T_r^R\|_\rho \\
(3.40) \quad &\leq b(n) \sqrt{\sum_{l,r} \|T_l^L B_\phi(r)\|_\rho^2} \cdot \sqrt{\sum_{l,r} \|B_\psi(l) T_r^R\|_\rho^2}.
\end{aligned}$$

The quantity under the first square root can be bounded in terms of $\|\phi\|$ as follows:

$$\begin{aligned}
\sum_{l,r} \|T_l^L B_\phi(r)\|_\rho^2 &= \sum_{l,r} \text{Tr} \rho B_\phi(r)^* (T_l^L)^* T_l^L B_\phi(r) \\
&= \sum_r \text{Tr} \rho B_\phi(r)^* Q_L B_\phi(r) = \sum_r \|B_\phi(r)\|_{G_L - C_n}^2 \\
&\leq \frac{1}{1 - b_L(n)} \sum_r \|\Gamma_{G_L - C_n}(B_\phi(r))\|^2 \\
&= \frac{1}{1 - b_L(n)} \|\phi\|^2,
\end{aligned}$$

where we have used the definition of Q_L , Corollary 3.4 and (3.33). The quantity under the second square root is similarly estimated in terms of $\|\psi\|$:

$$(3.41) \quad \sum_{l,r} \|B_\psi(l) T_r^R\|_\rho^2 \leq \frac{1}{1 - b_R(n)} \|\psi\|^2.$$

Inserting these into (3.40) yields

$$(3.42) \quad |\langle \varphi, \psi \rangle - \langle C_\phi, D_\psi \rangle_{\mathcal{K}_G}| \leq \frac{b(n)}{\sqrt{(1 - b_L(n))(1 - b_R(n))}} \|\phi\| \|\psi\|.$$

□

Now, we are ready to estimate the quantity of interest in (3.8), which is an inner product of the form considered in Lemma 3.5 with the additional information that ϕ and ψ are both orthogonal to \mathcal{G}_G .

Proposition 3.6. *Under the assumptions stated in Section 3.1 and with the notations introduced there, we have the following estimate for the quantity ε_n defined in (3.8):*

$$(3.43) \quad \varepsilon_n \leq \frac{b(n)}{\sqrt{1 - b_{LR}(n)}} + \left(\frac{b(n)}{\sqrt{1 - b_{LR}(n)}} \right)^2 (1 + b_G(n)),$$

with

$$(3.44) \quad b(n) = a(n) \text{Tr} \rho^{-1}, \quad a(n) = \|\mathbb{E}^n - |\mathbf{1}\rangle\langle \rho|\|,$$

and $b_{LR}(n)$ and $b_G(n)$ are defined in (3.30) and (3.38).

Proof. Any $\xi \in \mathcal{G}_G$ belongs to both $\mathcal{G}_{G_L - C_n} \otimes \mathcal{H}_{G_R}$ and $\mathcal{H}_{G_L} \otimes \mathcal{G}_{C_n - G_R}$. Therefore, there are unique matrices $B_\xi(r)$ and $B_\xi(l)$ and corresponding expressions (3.31) and (3.32) for ξ . Since $\xi \in \mathcal{G}_G$, there also exists $X \in \mathcal{K}_G$ such that $\xi = \Gamma_G(X)$. By injectivity it follows that

$$(3.45) \quad B_\xi^L(l) = T_l^L X, \quad B_\xi^R(r) = X T_r^R.$$

Inserting the first relation above into the expression for D_ψ and the second into C_ϕ we find the following special form of these matrices for a ground state ξ :

$$(3.46) \quad C_\xi = \sum_r B_\phi(r) \rho(T_r^R)^* = \sum_r X T_r^R \rho(T_r^R)^* = X Q_R$$

$$(3.47) \quad D_\xi = \sum_l (T_l^L)^* B_\psi(l) = \sum_l (T_l^L)^* T_l^L X = Q_L X.$$

We use this to extract information from the orthogonality of ϕ and ψ to \mathcal{G}_G . Using $\langle \phi, \xi \rangle = \langle \xi, \psi \rangle = 0$, from Lemma 3.5 we have that for all $X \in \mathbb{C}^{D_R \times D_L}$:

$$(3.48) \quad |\langle C_\phi, Q_L X \rangle_{\mathcal{K}_G}| \leq \frac{b(n)}{\sqrt{1 - b_{LR}(n)}} \|\phi\| \|\xi\|$$

$$(3.49) \quad |\langle X Q_R, D_\psi \rangle_{\mathcal{K}_G}| \leq \frac{b(n)}{\sqrt{1 - b_{LR}(n)}} \|\xi\| \|\psi\|.$$

Applying Corollary 3.4 gives

$$\|\xi\|^2 = \|\Gamma_G(X)\|^2 \leq (1 + b_G(n)) \|X\|_G^2.$$

Using this with (3.48) and (3.49) yields

$$(3.50) \quad |\langle C_\phi, Q_L X \rangle_{\mathcal{K}_G}| \leq \delta(n) \|\phi\| \|X\|_G$$

$$(3.51) \quad |\langle X Q_R, D_\psi \rangle_{\mathcal{K}_G}| \leq \delta(n) \|\psi\| \|X\|_G,$$

where

$$\delta(n) = \frac{b(n)}{\sqrt{1 - b_{LR}(n)}} \sqrt{1 + b_G(n)}.$$

The LHS of these inequalities can be expressed in terms of the inner product $\langle \cdot, \cdot \rangle_G$ as follows:

$$\langle C_\phi, Q_L X \rangle_{\mathcal{K}_G} = \text{Tr} C_\phi^* Q_L X = \text{Tr} Q_R Q_R^{-1} C_\phi^* Q_L X = \langle C_\phi Q_R^{-1}, X \rangle_G$$

$$\langle X Q_R, D_\psi \rangle_{\mathcal{K}_G} = \text{Tr} Q_R X^* D_\psi = \text{Tr} Q_R X^* Q_L Q_L^{-1} D_\psi = \langle X, Q_L^{-1} D_\psi \rangle_G.$$

The estimates (3.50) now become: for all X

$$(3.52) \quad |\langle C_\phi Q_R^{-1}, X \rangle_G| \leq \delta(n) \|\phi\| \|X\|_G$$

$$(3.53) \quad |\langle X, Q_L^{-1} D_\psi \rangle_G| \leq \delta(n) \|\psi\| \|X\|_G,$$

which imply

$$(3.54) \quad \|C_\phi Q_R^{-1}\|_G \leq \delta(n) \|\phi\|$$

$$(3.55) \quad \|Q_L^{-1} D_\psi\|_G \leq \delta(n) \|\psi\|.$$

Noting the identity

$$\langle C_\phi, D_\psi \rangle_{\mathcal{K}_G} = \text{Tr} Q_R (Q_R^{-1} C_\phi^*) Q_L (Q_L^{-1} D_\psi) = \langle C_\phi Q_R^{-1}, Q_L^{-1} D_\psi \rangle_G,$$

we have

$$|\langle C_\phi, D_\psi \rangle_{\mathcal{K}_G}| \leq \|C_\phi Q_R^{-1}\|_G \|Q_L^{-1} D_\psi\|_G \leq \delta(n)^2 \|\phi\| \|\psi\|.$$




FIGURE 3. The VBS picture for the decorated AKLT model.

Moreover, it follows from (3.37) that

$$(3.56) \quad |\langle \phi, \psi \rangle| \leq |\langle C_\phi, D_\psi \rangle \kappa_G| + \frac{b(n)}{\sqrt{1 - b_{LR}(n)}} \|\phi\| \|\psi\|.$$

Combining the last two inequalities we obtain the final estimate

$$(3.57) \quad |\langle \phi, \psi \rangle| \leq \left[\frac{b(n)}{\sqrt{1 - b_{LR}(n)}} + \left(\frac{b(n)}{\sqrt{1 - b_{LR}(n)}} \right)^2 (1 + b_G(n)) \right] \|\phi\| \|\psi\|.$$

□

In the next section we verify the assumptions stated in this section for the AKLT model on the decorated honeycomb lattice with $n \geq 2$ and apply Proposition 3.6 to show that for this model we have $\varepsilon_n < 1/3$, for all $n \geq 3$.

4. GAP OF THE DECORATED AKLT MODEL

In this section we prove Proposition 2.1 and Theorem 2.2 by applying the results of Section 3 to the decorated AKLT model discussed in Section 2. In this case, the graph G is given by $Y_v \cup Y_w$ for two adjacent sites v and w in Γ . We decompose G as $G = G_L - C_n - G_R$, where $C_n = Y_v \cap Y_w$, $G_L = Y_v \setminus C_n$ and $G_R = Y_w \setminus C_n$. The VBS (Valence Bond Solid) or PEPS (Product of Entangled Pairs) ground states on G are depicted in Figure 3. The corresponding Hilbert spaces are given by

$$\mathcal{H}_{G_L} = \mathcal{H}_{G_R} = (\mathbb{C}^3 \otimes \mathbb{C}^3)^{\otimes n} \otimes \mathbb{C}^4, \quad \text{and} \quad \mathcal{H}_{C_n} = (\mathbb{C}^3)^{\otimes n}.$$

For $s \in \{1, 3/2\}$, we use $\mathcal{B}_s = \{|s\rangle, |s-1\rangle, \dots, |-s\rangle\}$ to denote an orthonormal basis of \mathbb{C}^{2s+1} consisting of eigenvectors of the third component of spin associated to the spin- s irreducible representation of $SU(2)$. We begin by discussing the MPS Γ_{C_n} and its associated transfer operator \mathbb{E} . We then define the operator \mathbb{E}_L associated with G_L and prove that $\Gamma_{G_L - C_n}$ is injective for $n \geq 2$, after which we prove the main results.

On C_n we have the one-dimensional AKLT spin chain, which has the bond dimension $D = 2$. For this model, every physical spin-1 vertex is identified with the symmetric subspace of the virtual space $\mathbb{C}^2 \otimes \mathbb{C}^2$. We will denote by S^X, S^Y, S^Z the usual spin-1/2 operators, and by S^\pm the corresponding lowering and raising operators. To differentiate between the physical and virtual spins, we will use $|\uparrow\rangle, |\downarrow\rangle \in \mathbb{C}^2$ to denote the standard orthonormal eigenbasis of S^Z rather than $|\pm 1/2\rangle$. The intertwiner $P_{\text{sym}}^{(1)} : \mathbb{C}^2 \otimes \mathbb{C}^2 \rightarrow \mathbb{C}^3$ that maps between the virtual and physical space of a site v is given by:

$$P_{\text{sym}}^{(1)} = |1\rangle\langle\uparrow\uparrow| + |0\rangle\langle\psi^+| + |-1\rangle\langle\downarrow\downarrow|,$$

where $|\psi^+\rangle = \frac{1}{\sqrt{2}}(|\uparrow\downarrow\rangle + |\downarrow\uparrow\rangle)$.

Recall that the symmetric subspace of $\mathbb{C}^2 \otimes \mathbb{C}^2$ can be encoded into the MPS matrices

$$P_1^{(1)} = |\uparrow\rangle\langle\uparrow|, \quad P_0^{(1)} = \sqrt{2}S^X, \quad P_{-1}^{(1)} = |\downarrow\rangle\langle\downarrow|.$$

The ground states of the one-dimensional AKLT model can then be described as a valence-bond solid state obtained from projecting each (virtual) edge of the graph into the singlet states $|\psi^-\rangle = \frac{1}{\sqrt{2}}(|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle)$; this is represented by the MPS matrix

$$K = \frac{1}{\sqrt{2}}(|\uparrow\rangle\langle\downarrow| - |\downarrow\rangle\langle\uparrow|) = \sqrt{2}iS^Y.$$

With these matrices, and with a convenient choice of normalization, the ground state space of the one-dimensional AKLT matrix is given by

$$\Gamma_{C_n}(B) = \sum_{i_1, \dots, i_n \in \{\pm 1, 0\}} \text{Tr}[BV_{i_n} \dots V_{i_1}] |i_1 \dots i_n\rangle$$

where $V_i = \frac{2}{\sqrt{3}}KP_i^{(1)}$. Explicitly,

$$(4.1) \quad V_1 = -\sqrt{\frac{2}{3}}S^+, \quad V_0 = \frac{2}{\sqrt{3}}S^Z, \quad V_{-1} = \sqrt{\frac{2}{3}}S^-.$$

Given the form of Γ_{C_n} , the choice of multiplying on the *left* by K in the definition of V_i corresponds to projecting the edge to the *right* of the associated site into the singlet state. This convention will also be used to define the tensors T_ℓ^L and T_r^R . For more details on this and other MPS constructions, see [41, 39, 37].

Noting that $S^+BS^- + S^-BS^+ = 2(S^XBS^X + S^YBS^Y)$, the transfer operator $\mathbb{E} : M_2 \rightarrow M_2$ associated with Γ_{C_n} takes the form

$$(4.2) \quad \mathbb{E}(B) = \sum_{i \in \{\pm 1, 0\}} V_i^* B V_i = \frac{4}{3}(S^XBS^X + S^YBS^Y + S^ZBS^Z),$$

which can be easily diagonalized as

$$(4.3) \quad \mathbb{E} = |\mathbb{1}\rangle\langle\rho| - \frac{2}{3} \sum_{U \in \{X, Y, Z\}} |S^U\rangle\langle S^U|,$$

where $\rho = \mathbb{1}/2$ is the maximally mixed state. This allows to easily compute

$$\mathbb{E}^n = |\mathbb{1}\rangle\langle\rho| + 2 \frac{(-1)^n}{3^n} \sum_{U \in \{X, Y, Z\}} |S^U\rangle\langle S^U|,$$

from which (3.3) takes the explicit form

$$a(n) = \|\mathbb{E}^n - |\mathbb{1}\rangle\langle\rho|\| = 3^{-n},$$

and $b(n) = \text{Tr}(\rho^{-1})a(n) = 4 \cdot 3^{-n}$. By Corollary 3.2, this implies that Γ_{C_n} is injective when $n \geq 2$. It can easily be shown that it is not injective for $n = 1$.

We now consider G_L and G_R . For the decorated AKLT model, $D_L = D_R = 4$ and so $\mathcal{K}_G = M_4$. We first construct the operator \mathbb{E}_L associated with G_L , and use this to prove $\Gamma_{G_L-C_n}$ is injective of $n \geq 2$. The analogous operator \mathbb{E}_R for G_R and the injectivity of $\Gamma_{C_n-G_R}$ follow from similar calculations.

We first note that G_L can be written as $[u_1^1, u_n^1] \times [u_1^2, u_n^2] \times \{v\}$, where the sites u_k^i correspond to the $2n$ spin-1's, and v is the spin-3/2. By grouping the sites u_i^1 and u_i^2 into a single site (u_i^1, u_i^2) , we can recognize the ground states of H_{G_L} as a PEPS. We choose the product basis for \mathcal{H}_{G_L} given by

$$|i_1, j_1\rangle \otimes \cdots \otimes |i_n, j_n\rangle \otimes |k\rangle \quad i_1, \dots, i_n, j_1, \dots, j_n \in \{\pm 1, 0\}, \quad k \in \left\{ \pm \frac{3}{2}, \pm \frac{1}{2} \right\}.$$

For each element $|l\rangle_L = |i_1, j_1\rangle \otimes \cdots \otimes |i_n, j_n\rangle \otimes |k\rangle$ of the basis, the 2×4 matrix T_l^L is given by

$$T_l^L = W_k^L V_{i_n} \otimes V_{j_n} \cdots V_{i_1} \otimes V_{j_1},$$

where the V_i are as defined in (4.1), and the $W_k^L \in \mathcal{L}(\mathbb{C}^4, \mathbb{C}^2)$ are given by the PEPS representation of the AKLT on the hexagonal lattice, which we now define. Analogous to the spin-1 case, the virtual space of a spin-3/2 particle is the symmetric subspace of three spin-1/2 particles, and so the intertwiner $P_{\text{sym}}^{(3/2)} : \mathbb{C}^2 \otimes \mathbb{C}^2 \otimes \mathbb{C}^2 \rightarrow \mathbb{C}^4$ between the virtual and physical space is given by

$$P_{\text{sym}}^{(3/2)} = |3/2\rangle\langle\uparrow\uparrow\uparrow| + |1/2\rangle\langle\phi^+| + |-1/2\rangle\langle\phi^-| + |-3/2\rangle\langle\downarrow\downarrow\downarrow|,$$

where

$$(4.4) \quad |\phi^+\rangle = \frac{1}{\sqrt{3}}(|\uparrow\uparrow\downarrow\rangle + |\uparrow\downarrow\uparrow\rangle + |\downarrow\uparrow\uparrow\rangle) = \frac{1}{\sqrt{3}}|\downarrow\rangle|\uparrow\uparrow\rangle + \sqrt{\frac{2}{3}}|\uparrow\rangle|\psi^+\rangle,$$

$$(4.5) \quad |\phi^-\rangle = \frac{1}{\sqrt{3}}(|\uparrow\downarrow\downarrow\rangle + |\downarrow\uparrow\downarrow\rangle + |\downarrow\downarrow\uparrow\rangle) = \frac{1}{\sqrt{3}}|\uparrow\rangle|\downarrow\downarrow\rangle + \sqrt{\frac{2}{3}}|\downarrow\rangle|\psi^+\rangle.$$

By grouping two virtual edges to the left of v , see Figure 3, the virtual space can be represented by the MPS matrices $P_k^{(3/2)} \in M_{2 \times 4}$ defined by

$$\begin{aligned} P_{3/2}^{(3/2)} &= |\uparrow\rangle\langle\uparrow\uparrow|, & P_{1/2}^{(3/2)} &= \frac{1}{\sqrt{3}}|\downarrow\rangle\langle\uparrow\uparrow| + \sqrt{\frac{2}{3}}|\uparrow\rangle\langle\psi^+| \\ P_{-3/2}^{(3/2)} &= |\downarrow\rangle\langle\downarrow\downarrow|, & P_{-1/2}^{(3/2)} &= \frac{1}{\sqrt{3}}|\uparrow\rangle\langle\downarrow\downarrow| + \sqrt{\frac{2}{3}}|\downarrow\rangle\langle\psi^+|. \end{aligned}$$

Once again projecting edges on the *right* of v into a singlet state (and choosing a convenient normalization) we define $W_k^L = \sqrt{2}K P_k^{(3/2)}$. Explicitly,

$$(4.6) \quad W_{3/2}^L = -|\downarrow\rangle\langle\uparrow\uparrow|, \quad W_{1/2}^L = \frac{1}{\sqrt{3}}|\uparrow\rangle\langle\uparrow\uparrow| - \sqrt{\frac{2}{3}}|\downarrow\rangle\langle\psi^+|,$$

$$(4.7) \quad W_{-3/2}^L = |\uparrow\rangle\langle\downarrow\downarrow|, \quad W_{-1/2}^L = -\frac{1}{\sqrt{3}}|\downarrow\rangle\langle\downarrow\downarrow| + \sqrt{\frac{2}{3}}|\uparrow\rangle\langle\psi^+|,$$

which satisfies $\sum_{i=1}^4 W_i^L (W_i^L)^* = 2\mathbb{1}_{\mathbb{C}^2}$. For $B \in M_2$, define $\mathbb{E}^\gamma(B) = \sum_i (W_i^L)^* B W_i^L$. While this is a completely positive map from M_2 to M_4 , unlike the MPS case it is not unital, since

$$\mathbb{E}^\gamma(\mathbb{1}) = \sum_{i=1}^4 (W_i^L)^* W_i^L = \frac{4}{3}(|\uparrow\uparrow\rangle\langle\uparrow\uparrow| + |\downarrow\downarrow\rangle\langle\downarrow\downarrow| + |\psi^+\rangle\langle\psi^+|) = \mathbb{1} + \frac{4}{3}\mathbf{S} \cdot \mathbf{S},$$

where as usual $\mathbf{S} = (S^X, S^Y, S^Z)$ and $\mathbf{S} \cdot \mathbf{S} = S^X \otimes S^X + S^Y \otimes S^Y + S^Z \otimes S^Z$. By direct calculation, we see that

$$(4.8) \quad (\mathbb{E}^\gamma)^*(B) = c(B)\rho + \sum_{U \in \{X, Y, Z\}} c_U(B)S^U,$$

where

$$\begin{aligned} c(B) &= \frac{4}{3}(\langle\uparrow\uparrow|B|\uparrow\uparrow\rangle + \langle\downarrow\downarrow|B|\downarrow\downarrow\rangle + \langle\psi^+|B|\psi^+\rangle) \\ c_X(B) &= -\frac{2\sqrt{2}}{3}\text{Re}[\langle\psi^+|B|\uparrow\uparrow\rangle + \langle\downarrow\downarrow|B|\psi^+\rangle] \\ c_Y(B) &= -\frac{2\sqrt{2}}{3}\text{Im}[\langle\psi^+|B|\uparrow\uparrow\rangle + \langle\downarrow\downarrow|B|\psi^+\rangle] \\ c_Z(B) &= \frac{2}{3}(\langle\downarrow\downarrow|B|\downarrow\downarrow\rangle - \langle\uparrow\uparrow|B|\uparrow\uparrow\rangle). \end{aligned}$$

It can easily be checked that $(\mathbb{E}^\gamma)^t \circ \tau = (\mathbb{E}^\gamma)^t$ where $\tau : M_4 \rightarrow M_4$ is the transposition operator

$$\tau(A \otimes B) = B \otimes A$$

Combining this with (4.8) allows us to verify that

$$(\mathbb{E}^\gamma)^t(\rho \otimes \rho) = \rho, \quad (\mathbb{E}^\gamma)^t(S^U \otimes S^{U'}) = \delta_{U,U'} \frac{1}{3}\rho, \quad (\mathbb{E}^\gamma)^t(\rho \otimes S^U) = (\mathbb{E}^\gamma)^t(S^U \otimes \rho) = -\frac{1}{3}S^U,$$

or equivalently

$$(4.9) \quad \mathbb{E}^\gamma = |\mathbb{1} \otimes \mathbb{1}\rangle\langle\rho| + \frac{4}{3}|\mathbf{S} \cdot \mathbf{S}\rangle\langle\rho| - \frac{4}{3} \sum_{U \in \{X, Y, Z\}} (|S^U \otimes \mathbb{1}\rangle + |\mathbb{1} \otimes S^U\rangle) \langle S^U|.$$

To simplify notation we define

$$|\Omega^U\rangle = |S^U \otimes \mathbb{1}\rangle + |\mathbb{1} \otimes S^U\rangle \quad \forall U \in \{X, Y, Z\},$$

and notice that

$$\|\mathbf{S} \cdot \mathbf{S}\|_2 = \frac{\sqrt{3}}{2}, \quad \|\Omega^U\|_2 = \sqrt{2}, \quad \text{and} \quad \langle\mathbf{S} \cdot \mathbf{S}, \Omega^U\rangle_2 = 0$$

for all U . The transfer matrix for G_L , defined in (3.5), is then given by $\mathbb{E}_L = (\mathbb{E}^n \otimes \mathbb{E}^n) \circ \mathbb{E}^\gamma$, which can be simplified to

$$\begin{aligned} \mathbb{E}_L &= \left(|\mathbb{1}\rangle\langle\rho| + 2\frac{(-1)^n}{3^n} \sum_U |S^U\rangle\langle S^U| \right) \otimes \left(|\mathbb{1}\rangle\langle\rho| + 2\frac{(-1)^n}{3^n} \sum_U |S^U\rangle\langle S^U| \right) \mathbb{E}^\gamma \\ (4.10) \quad &= |\mathbb{1} \otimes \mathbb{1}\rangle\langle\rho| + 2\frac{(-1)^{n+1}}{3^{n+1}} \sum_U |\Omega^U\rangle\langle S^U| + \frac{4}{3^{2n+1}} |\mathbf{S} \cdot \mathbf{S}\rangle\langle\rho|. \end{aligned}$$

Using this decomposition to compute Q_L gives

$$(4.11) \quad Q_L = \mathbb{1} + \frac{4}{3^{2n+1}} \mathbf{S} \cdot \mathbf{S}, \implies \text{spec}(Q_L) = \left\{ 1 - \frac{1}{3^{2n}}, 1 + \frac{1}{3^{2n+1}} \right\}.$$

Therefore, $q_L = 1 - \frac{1}{3^{2n}}$, and moreover, since \mathbb{E}_L is a completely positive map,

$$\|\mathbb{E}_L\| = \|Q_L\| = 1 + \frac{1}{3^{2n+1}}.$$

Since Q_L is invertible, the theory of Section 3 applies and we can use the above relations to prove the following result.

Lemma 4.1. $\Gamma_{G_L-C_n}$ is injective for $n \geq 2$.

Proof. Let $B \in M_D$ and consider $\Gamma_{G_L-C_n}(B)$. Applying Corollary 3.4 gives

$$\|\Gamma_{G_L-C_n}(B)\|^2 \geq (1 - b_L(n)) \|B\|_{G_L-C_n}^2$$

with $b_L(n) = \frac{4a(n)\|\mathbb{E}_L\|}{\rho_{\min} q_L}$. By inserting the values of $a(n)$, $\|\mathbb{E}_L\|$, ρ_{\min} , and q_L into this expression, one finds that

$$1 - b_L(n) = 1 - \frac{8(1 + 3^{-2n-1})}{3^n(1 - 3^{-2n})}.$$

This quantity is strictly positive for any $n \geq 2$ from which it follows that $\Gamma_{G_L-C_n}$ is injective. \square

We now consider G_R . The operator \mathbb{E}_R is obtained using a similar construction as \mathbb{E}_L . As with G_L , we can once again group the spin-1 particles into pairs and to construct an orthonormal basis

$$|r\rangle = |i_1, j_1\rangle \otimes \cdots \otimes |i_n, j_n\rangle \otimes |k\rangle \quad i_1, \dots, i_n, j_1, \dots, j_n \in \{\pm 1, 0\}, k \in \left\{ \pm \frac{3}{2}, \pm \frac{1}{2} \right\},$$

for which the corresponding tensor is given by

$$(4.12) \quad T_r^R = V_{i_n} \otimes V_{j_n} \cdots V_{i_1} \otimes V_{j_1} W_k^R,$$

where V_1, V_0 and V_{-1} are as before, and $W_k^R = 2K \otimes K(P_k^{(3/2)})^*$. Explicitly,

$$(4.13) \quad W_{3/2}^R = |\downarrow\downarrow\rangle\langle\uparrow|, \quad W_{1/2}^R = \frac{1}{\sqrt{3}}|\downarrow\downarrow\rangle\langle\downarrow| - \sqrt{\frac{2}{3}}|\psi^+\rangle\langle\uparrow|,$$

$$(4.14) \quad W_{-3/2}^R = |\uparrow\uparrow\rangle\langle\downarrow|, \quad W_{-1/2}^R = \frac{1}{\sqrt{3}}|\uparrow\uparrow\rangle\langle\uparrow| - \sqrt{\frac{2}{3}}|\psi^+\rangle\langle\downarrow|.$$

Similar to the case of \mathbb{E}^L , we have $\mathbb{E}^R := \mathbb{E}^\sphericalangle \circ (\mathbb{E}^n \otimes \mathbb{E}^n)$ where $\mathbb{E}^\sphericalangle : M_4 \rightarrow M_2$ is defined by

$$(4.15) \quad \mathbb{E}^\sphericalangle(B) := \sum_i (W_i^R)^* B W_i^R = \sum_i W_i^L B (W_i^L)^* = (\mathbb{E}^\sphericalangle)^t(B).$$

The final equality above follows from recognizing

$$(4.16) \quad W_{3/2}^R = (W_{-3/2}^L)^*, \quad W_{1/2}^R = -(W_{-1/2}^L)^*$$

$$(4.17) \quad W_{-3/2}^R = -(W_{3/2}^L)^*, \quad W_{-1/2}^R = (W_{1/2}^L)^*.$$

It follows from the analogous arguments as used in Lemma 4.1 above that $\Gamma_{C_n-G_R}$ is also injective for $n \geq 2$. We can now prove Proposition 2.1, and Theorem 2.2.

Proof of Proposition 2.1 and Theorem 2.2. Since $\mathbb{E} = \mathbb{E}^t$, from (4.15) it follows that

$$\mathbb{E}_R = [(\mathbb{E}^n \otimes \mathbb{E}^n) \circ \mathbb{E}^\gamma]^t = (\mathbb{E}_L)^t.$$

Therefore, $\|\mathbb{E}_R\| = \|\mathbb{E}_L\|$ and

$$(4.18) \quad Q_R := (\mathbb{E}_R)^t(\rho) = \mathbb{E}_L(\rho) = \frac{1}{2}Q_L.$$

As a consequence,

$$q_R = \frac{1}{2}q_L = \rho_{\min}q_L.$$

Using (3.43) to estimate (3.8), we find

$$(4.19) \quad \varepsilon_n \leq \frac{4a(n)}{\sqrt{1 - b_{LR}(n)}} + \left(\frac{4a(n)}{\sqrt{1 - b_{LR}(n)}} \right)^2 (1 + b_G(n)).$$

From (3.30) and the values above, it is clear that $b_L(n) = b_R(n)$, and so

$$(4.20) \quad \begin{aligned} 1 - b_{LR}(n) &= (1 - b_L(n))(1 - b_R(n)) = \left(1 - \frac{8(1+3^{-2n-1})}{3^n(1-3^{-2n})}\right)^2, \\ 1 + b_G(n) &= \left(1 + \frac{8(1+3^{-2n-1})^2}{3^n(1-3^{-2n})^2}\right). \end{aligned}$$

This establishes Proposition 2.1.

Inserting these into (4.19), we find that $\varepsilon_n < \frac{1}{3}$ whenever $n \geq 3$. By (2.9) this implies that the decorated AKLT model has a positive spectral gap above the ground state energy for $n \geq 3$. This completes the proof of Theorem 2.2. \square

5. DISCUSSION

We proved an explicit positive lower bound for the spectral gap above the ground state of the AKLT model on the decorated honeycomb lattice for $n \geq 3$, where n is the number of vertices inserted on each edge of the honeycomb lattice. It is natural to ask whether the approach of this paper could be used to prove that the AKLT model on the honeycomb lattice itself ($n = 0$) is gapped too, which is expected. It is clear to us, however, that significant changes to the arguments would be necessary to achieve this. For example, a numerical calculation shows that $\varepsilon_1 \sim .478 > 1/3$. Therefore, our method does not work for $n = 1$. For the case $n = 2$, we do not have a good estimate of ε_2 , but it is conceivable that our approach could be extended to the case $n = 2$. For the model with $n = 3$, however, we proved a positive lower bound. By using a numerically calculated value for the gap for the small system on Y , which appears in (2.4), ($\gamma_Y \sim 0.2966$), and the rigorous estimate showing $\varepsilon_3 < 0.2683$ (Proposition 2.1), we found the following uniform lower bound for the gap: $\gamma > 0.0289$.

About generalizations to frustration-free models on other decorated lattices on the other hand, we can be rather optimistic. For example, we expect that similar arguments will work to study the spectral of AKLT models on decorated hypercubic lattices of any dimension. One could also try to apply our approach to some of the more exotic hybrid valence bond models discussed in [44].

For physical reasons, one wants the spectral gap to be robust under small perturbations of the interactions. It seems very likely that the AKLT models on the decorated honeycomb lattices (and likely also on the honeycomb lattice itself) satisfy the Local Topological Quantum Order condition introduced by Bravyi, Hastings, and Michalakis [12]. If so, the stability theorem of Michalakis and Zwolak [31] would apply to the AKLT models on decorated lattices and provide the desired robustness of the spectral gap.

ACKNOWLEDGMENTS

This work arose from discussions during the follow-up workshop on *Gapped Ground State Phases of Quantum Many-Body Systems* to the *2018 Arizona School of Analysis and Mathematical Physics*, organized by Robert Sims and two of the authors (H. A. and A. Y.) and supported by NSF Grant DMS-1800724. A. L. acknowledges support from the Walter Burke Institute for Theoretical Physics in the form of the Sherman Fairchild Fellowship as well as support from the Institute for Quantum Information and Matter (IQIM), an NSF Physics Frontiers Center (NSF Grant PHY-1733907). B. N. acknowledges support by the National Science Foundation under Grant DMS-1813149 and a CRM-Simons Professorship for a stay at the Centre de Recherches Mathématiques (Montréal) during Fall 2018, where part of this work was carried out.

REFERENCES

1. I. Affleck, T. Kennedy, E. H. Lieb, and H. Tasaki, *Rigorous results on valence-bond ground states in antiferromagnets*, Phys. Rev. Lett. **59** (1987), 799
2. I. Affleck, T. Kennedy, E. H. Lieb, and H. Tasaki, *Valence Bond Ground States in Isotropic Quantum Antiferromagnets*, Comm. Math. Phys. **115** (1988), no. 3, 477 – 528.
3. I. Arad, A. Kitaev, Z. Landau, and U. Vazirani, *An area law and sub-exponential algorithm for 1D systems*, arXiv:1301.1162
4. I. Arad, Z. Landau, U. Vazirani and T. Vidick, *Rigorous RG algorithms and area laws for low energy eigenstates in 1D*, Comm. Math. Phys. **356** (2018), no. 1, 65 – 105
5. S. Bachmann, E. Hamza, B. Nachtergaele and A. Young, *Product Vacua and Boundary State Models in d Dimensions*, J. Stat. Phys. **160** (2015), 636 – 658
6. S. Bachmann, S. Michalakis, B. Nachtergaele, and R. Sims, *Automorphic equivalence within gapped phases of quantum lattice systems*, Comm. Math. Phys. **309** (2012), no. 3, 835 – 871
7. S. Bachmann and B. Nachtergaele, *Product Vacua with Boundary States and the Classification of Gapped Phases*, Comm. Math. Phys. **329** (2014), 509 – 544
8. J. Bausch, T. Cubitt, A. Lucia, D. Perez-Garcia, *Undecidability of the spectral gap in one dimension*, arXiv:1810.01858
9. M. Bishop, *Spectral gaps for the Two-Species Product Vacua and Boundary States models on the d -dimensional lattice*, arXiv:1705.04755
10. M. Bishop, B. Nachtergaele, A. Young, *Spectral Gap and Edge Excitations of d -dimensional PVBS models on half-spaces* J. Stat. Phys. **162** (2016), 1485 – 1521
11. S. Bravyi and D. Gosset, *Gapped and gapless phases of frustration-free spin 1/2 chains* J. Math. Phys. **56** (2015), no. 6, 061902
12. S. Bravyi, M. Hastings, S. Michalakis, *Topological quantum order: stability under local perturbations*, J. Math. Phys. **51** (2010), 093512
13. B. Canals and C. Lacroix, *Pyrochlore Antiferromagnet: A Three-Dimensional Quantum Spin Liquid*, Phys. Rev. Lett. **80** (1998) 2933–2926
14. T.S. Cubitt, D. Perez-Garcia, and M.M. Wolf, *Undecidability of the spectral gap*, Nature **528** (2015), 207 – 211

15. M. Fannes, B. Nachtergaele, R.F. Werner, *Finitely Correlated States on Quantum Spin Chains*, Commun. Math. Phys. **144** (1992), 443 – 490
16. D. Gosset and E. Mozgunov, *Local gap threshold for frustration-free spin systems*, J. Math. Phys. **57** (2016), 091901
17. F.D.M. Haldane, *Continuum dynamics of the 1-d Heisenberg antiferromagnet: identification with the $O(3)$ nonlinear sigma model* Phys. Lett. **93** (1983), 464 – 468
18. F.D.M. Haldane, *Nonlinear field theory of large-spin Heisenberg antiferromagnets: semiclassically quantized solutions of the one-dimensional easy-axis Neel state* Phys. Rev. Lett. **50** (1983), 1153–1156
19. M. Hastings, *Lieb-Schultz-Mattis in higher dimensions* Phys. Rev. B **69** (2004), 104431
20. M. B. Hastings, *An area law for one-dimensional quantum systems*, J. Stat. Mech. Theory Exp. **2007** (2007), no. 8, P08024
21. M. Hastings, T. Koma, *Spectral gap and exponential decay of correlations*, Comm. Math. Phys. **265** (2006), no. 3, 781–804
22. M. J. Kastoryano and A. Lucia, *Divide and conquer method for proving gaps of frustration free Hamiltonians*, J. Stat. Mech. (2018) 033105
23. M. J. Kastoryano, A. Lucia, and D. Pérez-García, *Locality at the boundary implies gap in the bulk for 2D PEPS*, arXiv:1709.07691.
24. T. Kennedy, E.H. Lieb, and H. Tasaki, *A two-dimensional isotropic quantum antiferromagnet with unique disordered ground state*, J. Stat. Phys. **53** (1988), 383 – 415.
25. A.N. Kirillov and V.E. Korepin, *The resonating valence bond in quasicrystals*, Algebra and Analysis, **1** (2), 47 (1989) (in Russian), and arXiv:0909.2211 (in English).
26. S.A. Kivelson, D.S. Rokhsar, and J.P. Sethna, *Topology of the resonating valence-bond state: Solitons and high- T_c superconductivity*, Phys. Rev. B **35** (1987) 8865–8868
27. S. Knabe, *Energy gaps and elementary excitations for certain VBS-quantum antiferromagnets*, J. Stat. Phys. **52** (1988), no. 3-4, 627 – 638
28. I. Kurečić, L. Vanderstraeten, and N. Schuch, *A gapped $SU(3)$ spin liquid with \mathbb{Z}_3 topological order*, arXiv:1805.11628
29. Z. Landau, U. Vazirani and T. Vidick, *A polynomial-time algorithm for the ground state of 1D gapped local Hamiltonians*, Nature Physics **11** (2015), 566 – 569
30. M. Lemm and E. Mozgunov, *Spectral gaps of frustration-free spin systems with boundary*, arXiv:1801.08915
31. S. Michalakis, and J. Zwolak, *Stability of frustration-free Hamiltonians*, Comm. Math. Phys. **322** (2013), no. 2, 277 – 302
32. A. Moon and B. Nachtergaele, *Stability of gapped ground state phases of spins and fermions in one dimension*, J. Math. Phys. **59** (2018), no. 9, 091415
33. R. Movassagh, *Generic local Hamiltonians are gapless*, Phys. Rev. Lett. **119** (2017), 220504
34. B. Nachtergaele, *The spectral gap for some spin chains with discrete symmetry breaking*, Comm. Math. Phys. **175** (1996), 565–606
35. B. Nachtergaele, and R. Sims, *Lieb-Robinson bounds and the exponential clustering theorem*, Comm. Math. Phys. **265** (2006), no. 1, 119 – 130
36. B. Nachtergaele, R. Sims and A. Young, *Quasi-Localty Bounds for Quantum Lattice Systems. Part I. Lieb-Robinson Bounds, Quasi-Local Maps, and Spectral Flow Automorphisms*, arXiv:1810.02428
37. R. Orús, *A practical introduction to tensor networks: Matrix product states and projected entangled pair states*, Ann. Phys. **349** (2014), 117 – 158
38. R. Pan, D-L. Deng, and C.W. Clark, *Implementing Majorana fermions in a cold-atom honeycomb lattice with textured pairings*, Phys. Rev. A **98** (2018), 033604.
39. D. Pérez-García, F. Verstraete, M.M. Wolf, J.I. Cirac *Matrix Product State Representations*, Quantum Inf. Comput. **7**, (2007), 401–430
40. D. Poilblanc, N. Schuch, D. Pérez-García, and J.I. Cirac, *Topological and entanglement properties of resonating valence bond wave functions*, Phys. Rev. B **86** (2012), 014404.
41. U. Schollwöck, *The density-matrix renormalization group in the age of matrix product states*, Ann. Phys. **326**, (2011), 96–192.
42. N. Schuch, J.I. Cirac, and D. Perez-Garcia, *PEPS as ground states: degeneracy and topology*, Ann. Phys. **325** (2010), 2153

43. T-C. Wei, I. Affleck, and R. Raussendorf, *Affleck-Kennedy-Lieb-Tasaki state on a honeycomb lattice is a universal quantum computational resource*, Phys. Rev. Lett. **106** (2011), 070501.
44. T-C. Wei, P. Haghnegahdar, and R. Raussendorf, *Hybrid valence-bond states for universal quantum computation*, Phys. Rev. A **90** (2014), 042333
45. X.G. Wen, *Mean-field theory of spin-liquid states with finite energy gap and topological orders*, Phys. Rev. B **44** (1991) 2664–2672

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ARIZONA, TUCSON, AZ 85721, USA

Email address: `houssam@math.arizona.edu`

DEPARTMENT OF MATHEMATICS, HARVARD UNIVERSITY, CAMBRIDGE, MA 02138, USA

Email address: `mlemm@math.harvard.edu`

WALTER BURKE INSTITUTE FOR THEORETICAL PHYSICS AND INSTITUTE FOR QUANTUM INFORMATION & MATTER, CALIFORNIA INSTITUTE OF TECHNOLOGY, PASADENA, CA 91125, USA

Email address: `alucia@caltech.edu`

DEPARTMENT OF MATHEMATICS AND CENTER FOR QUANTUM MATHEMATICS AND PHYSICS, UNIVERSITY OF CALIFORNIA, DAVIS, DAVIS, CA 95616, USA

Email address: `bxn@math.ucdavis.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ARIZONA, TUCSON, AZ 85721, USA

Email address: `amyoung@math.arizona.edu`