

UC Merced

UC Merced Previously Published Works

Title

High quality draft genome sequence of the slightly halophilic bacterium *Halomonas zhanjiangensis* type strain JSM 078169T (DSM 21076T) from a sea urchin in southern China

Permalink

<https://escholarship.org/uc/item/8282561w>

Journal

Environmental Microbiome, 9(3)

ISSN

1944-3277

Authors

Zhou, Yu

Li, Rui

Gao, Xiao-Yang

et al.

Publication Date

2014-11-01

DOI

10.4056/sigs.5449586

Peer reviewed

High quality draft genome sequence of the slightly halophilic bacterium *Halomonas zhanjiangensis* type strain JSM 078169T (DSM 21076T) from a sea urchin in southern China

Yu Zhou^{1,2}, Rui Li², Xiao-Yang Gao^{3,4}, Alla Lapidus^{5,6}, James Han⁷, Matthew Haynes⁷, Elizabeth Lobos⁷, Marcel Huntemann⁷, Amrita Pati⁷, Natalia N. Ivanova⁷, Manfred Rohde⁸, Konstantinos Mavromatis⁷, Brian J. Tindall⁹, Victor Markowitz¹⁰, Tanja Woyke⁷, Hans-Peter Klenk^{9*}, Nikos C. Kyrpides^{7,11} and Wen-Jun Li^{1,3*}

¹ Key Laboratory of Microbial Diversity in Southwest China, Ministry of Education and the Laboratory for Conservation and Utilization of Bio-Resources, Yunnan Institute of Microbiology, Yunnan University, Kunming, China

² State Key Laboratory Breeding Base for Zhejiang Sustainable Plant Pest Control; Institute of Quality and Standard for Agro-products, Zhejiang Academy of Agricultural Sciences, Hangzhou, Zhejiang, China

³ Key Laboratory of Biogeography and Bioresource in Arid Land, Xinjiang Institute of Ecology and Geography, Chinese Academy of Sciences, Urumqi, China

⁴ University of Chinese Academy of Sciences, Beijing, China

⁵ Theodosius Dobzhansky Center for Genome Bioinformatics, St. Petersburg State University, St. Petersburg, Russia

⁶ Algorithmic Biology Lab, St. Petersburg Academic University, St. Petersburg, Russia

⁷ DOE Joint Genome Institute, Walnut Creek, California, USA

⁸ HZI – Helmholtz Centre for Infection Research, Braunschweig, Germany

⁹ Leibniz-Institute DSMZ - German Collection of Microorganisms and Cell Cultures, Braunschweig, Germany

¹⁰ Biological Data Management and Technology Center, Lawrence Berkeley National Laboratory, Berkeley, California, USA

¹¹ Department of Biological Sciences, King Abdulaziz University, Jeddah, Saudi Arabia

* Correspondence: W.J. Li (wjli@ynu.edu.cn) and H.P. Klenk (hpk@dsMZ.de)

Halomonas zhanjiangensis Chen et al. 2009 is a member of the genus *Halomonas*, family *Halomonadaceae*, class *Gammaproteobacteria*. Representatives of the genus *Halomonas* are a group of halophilic bacteria often isolated from salty environments. The type strain *H. zhanjiangensis* JSM 078169^T was isolated from a sea urchin (*Hemicentrotus pulcherrimus*) collected from the South China Sea. The genome of strain JSM 078169^T is the fourteenth sequenced genome in the genus *Halomonas* and the fifteenth in the family *Halomonadaceae*. The other thirteen genomes from the genus *Halomonas* are *H. halocynthiae*, *H. venusta*, *H. alkaliphila*, *H. lutea*, *H. anticariensis*, *H. jeotgali*, *H. titanicae*, *H. desiderata*, *H. smyrnensis*, *H. salifodinae*, *H. boliviensis*, *H. elongata* and *H. stevensii*. Here, we describe the features of strain JSM 078169^T, together with the complete genome sequence and annotation from a culture of DSM 21076^T. The 4,060,520 bp long draft genome consists of 17 scaffolds with the 3,659 protein-coding and 80 RNA genes and is a part of *Genomic Encyclopedia of Type Strains*, Phase I: the one thousand microbial genomes (KMG) project.

Keywords: strictly aerobic, motile Gram-negative, chemoorganotrophic, slightly halophilic, *Halomonadaceae*

Introduction

Strain JSM 078169^T (= DSM 21076 = KCTC 22279 = CCTCC AB 208031) is the type strain of the species *Halomonas zhanjiangensis* [1], one out of 84 species with a validly published name in the genus

Halomonas [2], family *Halomonadaceae* [3]. The family *Halomonadaceae* currently comprises thirteen genera (*Aidingimonas*, *Carnimonas*, *Chromohalobacter*, *Cobetia*, *Halomonas*, *Halotalea*, *Halo-*

vibrio, *Kushneria*, *Marinospirillum*, *Modicisali-bacter*, *Candidatus Portiera*, *Salinicola* and *Zy-mobacter*) with *Halomonas* being the largest genus in this family [3-6]. Members of the genus *Halomonas* have been isolated from various saline environments and showed halophilic characteristics [7-11]. Strain JSM 078169^T was originally isolated from a sea urchin (*Hemicentrotus pulcher-rimus*) that was collected from the South China Sea. The genus name was derived from the Greek words 'halos' meaning 'salt' and 'monas' meaning 'monad', yielding the Neo-Latin word 'halomonas' [2]; the species epithet was derived from Latin word 'zhanjiangensis', of Zhanjiang, a city in China near where the sample was collected [1]. Strain JSM 078169^T was found to assimilate several mono- and disaccharides and to produce numerous acid and alkaline phosphatases, leucine arylamidase, naphthol-ASBI-phosphohydrolase and valine arylamidase [1]. There are no PubMed records that document the use of these strain for any biotechnological studies; only comparative analyses performed for the description of later members of the genus *Halomonas* are recorded. However, the NamesforLife [12] database reports at least 70 patents in which *Halomonas* ssp. are referenced. Here we present a summary classification and a set of feature for *H. zhanjiangensis* JSM 078169^T, together with the description of the genomic sequencing and annotation of DSM 21076.

Classification and features

16S rRNA analysis

The original assembly of the genome did not contain longer stretches of 16S rRNA copies. Therefore, a 1,413 bp long fragment of the 16S rRNA gene was later patched into the genome sequence assembly. This almost full length version of the 16S rRNA sequence was compared using NCBI BLAST [13,14] under default settings (e.g., considering only the high-scoring segment pairs (HSPs) from the best 250 hits) with the most recent version of the Greengenes database [15] and the relative frequencies of taxa and unidentified clones (or strains) were calculated by BLAST scores. The most frequently occurring genus was *Halomonas* (74.8%), and the unidentified clones or isolates

represented 25.5% for the total BLAST results. Except for sequences of representatives of the genus *Halomonas*, no sequences from other genera were observed in the BLAST search. The highest degree of sequence similarity was reported with *H. alkantarctica* str. CRSS.

Figure 1 shows the phylogenetic neighborhood of *H. zhanjiangensis* JSM 078169^T in a tree based on 16S rRNA genes. The 1,413 bp long sequence fragment of the 16S rRNA gene differs by three nucleotides from the previously published 16S rRNA sequence (FJ429198). The tree provided a precise insight into the nomenclature and classification of members of the genus *Halomonas*. The phylogenetic analysis showed that strain *H. zhanjiangensis* JSM 078169^T was most closely related to *H. nanhaiensis* YIM M 13059^T with 98.3% sequence similarity.

Morphology and physiology

H. zhanjiangensis JSM 078169^T is a Gram-negative-staining, non-sporulating, strictly aerobic (Table 1), catalase-positive, oxidase-negative and slightly halophilic bacterium that reduces nitrate [1]. Cells of JSM 078169^T are short rods (0.4-0.7 μm × 0.6-1.0 μm) and motile with peritrichous flagella (not visible in Figure 2). Colonies are yellow-pigmented, flat and non-translucent with glistening surfaces and circular/slightly irregular margins, 2-3 mm in diameter after incubation on Marine Agar (MA) at 28 °C for 3-5 days. No diffusible pigments are produced. Growth occurs at 4-40 °C with an optimum growth at 25-30 °C, at pH range of 6.0-10.5 with an optimum pH of 7.5. The salinity range suitable for growth was 1.0-20.0% (w/v) total salts with an optimum between 3.0-5.0% (w/v) total salts. No growth occurs in the absence of NaCl or with NaCl as the sole salt. Strain JSM 078169^T grows on Marine Agar and the medium contained the following: 5.0 g peptone, 1.0 g yeast extract, 0.1 g ferric citrate, 19.45 g NaCl, 8.8 g MgCl₂, 3.24 g Na₂SO₄, 1.8 g CaCl₂, 0.55 g KCl, 0.16 g NaHCO₃, 0.08 g KBr, 0.034 g SrCl₂, 0.022 g H₃BO₃, 0.004 g sodium silicate, 0.0024 g sodium fluoride, 0.0016 g ammonium nitrate, 0.008 g disodium phosphate and 15 g agar.




Figure 1. Phylogenetic tree highlighting the position of *H. zhanjiangensis* relative to the closest related type strains of the other species within the family *Halomonadaceae*. All the 16S rRNA gene sequences of the type strains within the genus *Halomonas* were included and combined with the representative 16S rRNA gene sequences of the type species in other genera, according to the most recent release of the EzTaxon database. The tree was inferred from 1,381 aligned characters [16] under the neighbor-joining (NJ) [17], and maximum-likelihood (ML) [18] method with 1,000 randomly selected bootstrap replicates using MEGA version 5.2 [19]. The branches are scaled in terms of the expected number of substitutions per site. Numbers adjacent to the branches are support values from 1,000 NJ bootstrap (left) and from 1,000 ML bootstrap (right) replicates [20] if larger than 60%. Lineages with type strain genome sequencing projects registered in GOLD [21] are labeled with one asterisk, those also listed as ‘Complete and Published’ with two asterisks [22].




Figure 2 Scanning electron micrograph of *H. zhanjiangensis* JSM 078169^T

Table 1 Classification and general features of *H. zhanjiangensis* JSM 078169^T according the MIGS recommendations [23], (published by the Genomic Standards Consortium [24]), List of Prokaryotic names with Standing in Nomenclature [25] and the Names for Life database [12].

MIGS ID	Property	Term	Evidence code
MIGS-2	Domain	<i>Bacteria</i>	TAS [26]
	Phylum	<i>Proteobacteria</i>	TAS [27]
	Class	<i>Gammaproteobacteria</i>	TAS [28-30]
	Order	<i>Oceanospirillales</i>	TAS [29,31]
MIGS-3	Family	<i>Halomonadaceae</i>	TAS [3,32-35]
	Genus	<i>Halomonas</i>	TAS [2,36]
	Species	<i>Halomonas zhanjiangensis</i>	TAS [1]
	Type strain	JSM 078169	TAS [1]
MIGS-37.1	Cell shape	rod-shaped	TAS [1]
MIGS-37.2	Motility	motile	TAS [1]
MIGS-37.3	Sporulation	non-sporulating	TAS [1]
MIGS-37.12	Temperature range	4-40°C	TAS [1]
	Optimum temperature	25-30°C	TAS [1]
	Salinity	1-20% NaCl (w/v), optimum 3-5%	TAS [1]
	pH	6.0-10.5	TAS [1]
MIGS-37.5	Cell diameter	0.7-1.4 μm	TAS [1]
MIGS-37.6	Cell length	1.5-2.5 μm	TAS [1]
MIGS-37.9	Cell arrangement	singles, pairs	TAS [1]
MIGS-37.11	Energy metabolism	chemoorganotrophic	TAS [1]
MIGS-6	Habitat	aquatic, marine host	TAS [1]
MIGS-22	Oxygen requirement	aerobe	TAS [1]
MIGS-15	Biotic relationship	not reported	TAS [1]
MIGS-16	Host name	<i>Hemicentrotus pulcherrimus</i> (sea urchin)	TAS [1]
MIGS-4.5	Isolation site	Unspecified	TAS [1]
		Naozhou island, South China Sea near	
MIGS-4	Geographic location	Zhanjiang	TAS [1]
MIGS-4.1	Latitude	20.90	TAS [1]
MIGS-4.2	Longitude	110.59	TAS [1]
MIGS-4.3	Altitude	not reported	
MIGS-4.4	Depth	not reported	

Evidence codes-TAS: Traceable Author Statement (i.e., a direct report exists in the literature); NAS: Non-traceable Author Statement (i.e., not directly observed for the living, isolated sample, but based on a generally accepted property for the species, or anecdotal evidence). Evidence codes are from the Gene Ontology project [37].

Chemotaxonomy

The predominant respiratory quinone is Q-9 which is consistent to the other members of the genus *Halomonas* [1]. The predominant fatty acids are C_{18:1 ω7c} (48.9%), C_{16:0} (17.0%) and C_{12:0 3-OH} (10.7%). The profile of major fatty acids is also similar to the other representatives of the genus *Halomonas* [38-41].

Genome sequencing and annotation

Genome project history

This organism was selected for sequencing on the basis of its phylogenetic position [42,43]. Sequenc-

ing strain JSM 078169^T is part of *Genomic Encyclopedia of Type Strains*, Phase I: the one thousand microbial genomes (KMG) project [44], a follow-up of the GEBA project [45], which aims in increasing the sequencing coverage of key reference microbial genomes. The genome project is deposited in the Genomes OnLine Database [21] and the permanent draft genome sequence is deposited in GenBank. Sequencing, finishing and annotation were performed by the DOE Joint Genome Institute (JGI) using state of the art sequencing technology [46]. A summary of the project information is shown in Table 2.

Table 2 Genome sequencing project information

MIGS ID	Property	Term
MIGS-31.1	Sequencing quality	Level 2: High-Quality Draft
MIGS-28.1	Libraries used	One Illumina standard shotgun library
MIGS-29	Sequencing method	Illumina HiSeq 2000,
	Assembly method	Velvet v. 1.1.04; ALLPATHS v. r41043
	Gene calling method	Prodigal
MIGS-30	INSDC ID	ARIT00000000
MIGS-32	Genbank date of release	December 12, 2013
	GOLD ID	Gi11554
MIGS-1.1	NCBI project ID	178047
MIGS-1.2	Straininfo ID	845770
	Database: IMG	2517572236
MIGS-13	Source material identifier	DSM 21076
MIGS-38.2	Project relevance	Tree of Life, GEBA-KMG

Growth conditions and DNA isolation

H. zhanjiangensis JSM 078169^T, DSM 21076, was grown in DSMZ medium 1510 (modified medium 514 for *Halomonas* sp.) [47] at 28 °C. DNA was isolated from 0.5-1.0 g of cell paste using MasterPure Gram-positive DNA purification kit (Epicentre MGP04100) following the standard protocol as recommended by the manufacturer with modification st/DL for cell lysis as described by Wu *et al.* [45]. DNA is available through the DNA Bank Network [48].

Genome sequencing and assembly

The draft genome sequence was generated using the Illumina technology [49]. An Illumina Standard shotgun library was constructed and sequenced using the Illumina HiSeq 2000 platform which generated 15,593,002 reads totaling 2,339.0 Mbp. All general aspects of library construction and sequencing performed at the JGI can be found at [50]. All raw Illumina sequence data was passed through DUK, a filtering program developed at JGI, which removes known Illumina se-

quencing and library preparation artifacts [51]. Following steps were then performed for assembly: (1) filtered Illumina reads were assembled using Velvet [52], (2) 1–3 kbp simulated paired end reads were created from Velvet contigs using wgsim [53], (3) Illumina reads were assembled with simulated read pairs using Allpaths-LG [54]. Parameters for assembly steps were: 1) Velvet (velveth: 63 –shortPaired and velvetg: –very clean yes –export-Filtered yes –min contig lgth 500 –scaffolding no –cov cutoff 10) 2) wgsim (–e 0 –1 100 –2 100 –r 0 –R 0 –X 0) 3) Allpaths-LG (PrepareAllpathsInputs: PHRED 64=1 PLOIDY=1 FRAG COVERAGE=125 JUMP COVERAGE=25 LONG JUMP COV=50, RunAllpathsLG: THREADS=8 RUN=std shredpairs TARGETS=standard VAPI WARN ONLY=True OVERWRITE=True). The final draft assembly contained 18 contigs in 17 scaffolds. The total size of the genome is 4.1 Mbp and the final assembly is based on 501.3 Mbp of Illumina data, which provides an average 123.5 × coverage of the genome.

Genome annotation

Genes were identified using Prodigal [55] as part of the DOE-JGI genome annotation pipeline [56], following by a round of manual curation using the JGI GenePRIMP pipeline [57]. The predicted CDSs were translated and used to search the National Center for Biotechnology Information (NCBI) non-redundant database, UniProt, TIGR-Fam, Pfam, PRIAM, KEGG, COG, and InterPro database. These data sources were combined to assert a product description for each predicted protein. Additional gene prediction analysis and functional annotation was performed within the Integrated Microbial Genomes-Expert Review (IMG-ER) platform [58].

Genome properties

The assembly of the draft genome sequence consists of 17 scaffolds amounting to 4,060,520 bp, and the G+C content is 54.5% (Table 3 and Figure 3). Of the 3,739 genes predicted, 3,659 were protein-coding genes, and 80 RNAs. The majority of the protein-coding genes (87.1%) were assigned a putative function while the remaining ones were annotated as hypothetical proteins. The distribution of genes into COGs functional categories is presented in Table 4 and Figure 3.

Table 3. Genome statistics

Attribute	Value	% of total
Genome size (bp)	4,060,520	100.00%
DNA coding region (bp)	3,661,597	90.18%
DNA G+C content (bp)	2,212,212	54.48%
Number of scaffolds	17	
Extrachromosomal elements	unknown	
Total genes	3,739	100.00%
RNA genes	80	2.14%
rRNA operons	unknown	
tRNA genes	56	1.50%
Protein-coding genes	3,659	97.86%
Pseudo genes	0	0.00%
Genes with function prediction (proteins)	3,256	87.08%
Genes in paralog clusters	2,856	76.38%
Genes assigned to COGs	3,175	84.92%
Genes assigned Pfam domains	3,303	88.34%
Genes with signal peptides	313	8.37%
Genes with transmembrane helices	941	25.17%
CRISPR repeats	1	




Figure 3 The graphical map of the largest scaffold of the genome. From bottom to the top: Genes on forward strand (color by COG categories), Genes on reverse strand (color by COG categories), RNA genes (tRNA green, rRNA red, other RNAs black), GC content, GC skew (purple/olive).

Table 4 Number of genes associated with the general COG functional categories

Code	Value	% age	Description
J	180	5.03	Translation, ribosomal structure and biogenesis
A	1	0.03	RNA processing and modification
K	251	7.02	Transcription
L	143	4.00	Replication, recombination and repair
B	6	0.17	Chromatin structure and dynamics
D	38	1.06	Cell cycle control, cell division, chromosome partitioning
Y	0	0.00	Nuclear structure
V	36	1.01	Defense mechanisms
T	168	4.70	Signal transduction mechanisms
M	206	5.76	Cell wall/membrane/envelope biogenesis
N	93	2.60	Cell motility
Z	0	0.00	Cytoskeleton
W	0	0.00	Extracellular structures
U	80	2.24	Intracellular trafficking, secretion, and vesicular transport
O	137	3.83	Posttranslational modification, protein turnover, chaperones
C	211	5.90	Energy production and conversion
G	267	7.47	Carbohydrate transport and metabolism
E	342	9.56	Amino acid transport and metabolism
F	77	2.15	Nucleotide transport and metabolism
H	173	4.84	Coenzyme transport and metabolism
I	145	4.05	Lipid transport and metabolism
P	217	6.07	Inorganic ion transport and metabolism
Q	95	2.66	Secondary metabolites biosynthesis, transport and catabolism
R	404	11.30	General function prediction only
S	306	8.56	Function unknown
-	563	15.06	Not in COGs

Insights into the genome sequence

One complete genome sequence from a type strain of the family *Halomonas* - *H. elongata* [22] is available in GenBank, and four other permanent draft genomes of *H. anticariensis*, *H. lutea*, *H. jeotgali* and *H. halocynthiae* are available from IMG. The genome size of *H. zhanjiangensis* is smaller than those of *H. elongata*, *H. lutea* and *H. anticariensis* (4.06-5.02 Mbp), but much larger than those of *H. jeotgali* and *H. halocynthiae* (2.85-2.88 Mbp). Using the genome-to-genome distance calculator [59-61] version 2.0 revealed that all digital DNA-DNA hybridization (DDH) values are much lower than 70% using the program NCBI-BLAST, which demonstrated that *H. zhanjiangensis* is distinct from *H. elongata*, *H. anticariensis*, *H. lutea*, *H. jeotgali* and *H. halocynthiae* at the species level. Distance is 0.1845 between the type strain genomes of *H. zhanjiangensis* and *H. elongata*, which corresponds to a DDH value of $13.00 \pm 2.99\%$. The distances of *H. zhanjiangensis* from *H. anticariensis*, *H. lutea*, *H. jeotgali* and *H. halocynthiae* are 0.1842, 0.1837, 0.1835 and 0.1849, which correspond to

DDH values of $20.30 \pm 2.41\%$, $20.30 \pm 2.41\%$, $20.40 \pm 2.41\%$ and $20.20 \pm 2.41\%$, respectively.

A major feature of the previously sequenced genomes from this family is the presence of large numbers of proteins for the TRAP-type C4-dicarboxylate transport systems. A total of 267 genes in the genome of *H. zhanjiangensis* encode proteins for carbohydrate transport and metabolism, 68 genes are related to TRAP-type C4-dicarboxylate transport systems and encoded 22 large permease proteins, 24 periplasmic proteins and 22 small permease proteins. Genomic analysis of *H. elongata*, *H. anticariensis*, *H. lutea*, *H. jeotgali* and *H. halocynthiae* showed that they encode 58, 65, 61, 7 and 32 proteins related to TRAP-type C4-dicarboxylate transport system respectively. Proteins for TRAP-type C4-dicarboxylate transport systems constitute 1.86% as the total protein-coding sequences of the *H. zhanjiangensis* genome. In the genomes of *H. elongata*, *H. anticariensis*, *H. lutea*, *H. jeotgali* and *H. halocynthiae*, TRAP-type

C4-dicarboxylate transport system related proteins are accounted for 1.67%, 1.37%, 1.42%, 0.27% and 1.18% of the total protein-coding genes respectively. Therefore, *H. zhanjiangensis* has the highest percentage of TRAP-type C4-dicarboxylate transport system related encoding proteins in this group of bacteria to date.

Of the signal transduction mechanisms, Methyl-accepting Chemotaxis Proteins (MCPs) are transmembrane sensor proteins of bacteria. The MCPs allow bacteria to detect concentrations of molecules in the extracellular matrix so that they may smoothly swim or tumble accordingly [62,63]. Various environmental conditions give rise to diversity in bacterial signaling receptors, and consequently there are many genes encoding MCPs [64]. A number of MCPs (23) are present in *H. zhanjiangensis*, while *H. elongata*, *H. anticariensis*, *H. lutea*, and *H. jeotgali* have only 4, 21, 16, and 17 MCPs, respectively. MCPs are not found in the genome of *H. halocynthiae*. *H. zhanjiangensis* has the largest numbers of MCPs in this family. The analysis of bacterial genomes reveals that the family *Halomonadaceae* differs enormously in the num-

ber of MCPs from *E. coli*, and the number of MCPs in *Halomonadaceae* is about two times than that of *E. coli* strains.

Acknowledgements

The authors gratefully acknowledge the assistance of Susanne Schneider for growing *H. Zhanjiangensis* cultures and Evelyne-Marie Brambilla for DNA extraction and quality control (both at the DSMZ). This work was performed under the auspices of the US Department of Energy Office of Science, Biological and Environmental Research Program, and by the University of California, Lawrence Berkeley National Laboratory under contract No. DE-AC02-05CH11231, Lawrence Livermore National Laboratory under Contract No. DE-AC52-07NA27344 Genome analysis was supported by the National Basic Research Program of China (No. 2010CB833801). W-J Li was also supported by 'Hundred Talents Program' of the Chinese Academy of Sciences. A.L. was supported in part by Russian Ministry of Science Mega-grant no.11.G34.31.0068 (PI. Dr Stephen J O'Brien)

References

- Chen YG, Zhang YQ, Huang HY, Klenk HP, Tang SK, Huang K, Chen QH, Cui XL, Li WJ. *Halomonas zhanjiangensis* sp. nov., a halophilic bacterium isolated from a sea urchin. *Int J Syst Evol Microbiol* 2009; **59**:2888-2893. PubMed <http://dx.doi.org/10.1099/ijs.0.010173-0>
- Vreeland R, Litchfield CD, Martin EL, Elliot E. *Halomonas elongata*, a new genus and species of extremely salt-tolerant bacteria. *Int J Syst Evol Microbiol* 1980; **30**:485-495.
- Franzmann PD, Wehmeyer U, Stackebrandt E. *Halomonadaceae* fam. nov., a new family of the class *Proteobacteria* to accommodate the genera *Halomonas* and *Deleya*. *Syst Appl Microbiol* 1988; **11**:16-19. [http://dx.doi.org/10.1016/S0723-2020\(88\)80043-2](http://dx.doi.org/10.1016/S0723-2020(88)80043-2)
- Anzai Y, Kim H, Park JY, Wakabayashi H, Oyaizu H. Phylogenetic affiliation of the pseudomonads based on 16S rRNA sequence. *Int J Syst Evol Microbiol* 2000; **50**:1563-1589. PubMed <http://dx.doi.org/10.1099/00207713-50-4-1563>
- Arahal DR, Ludwig W, Schleifer KH, Ventosa A. Phylogeny of the family *Halomonadaceae* based on 23S and 16S rDNA sequence analyses. *Int J Syst Evol Microbiol* 2002; **52**:241-249. PubMed
- Mellado E, Moore ERB, Nieto JJ, Ventosa A. Phylogenetic inferences and taxonomic consequences of 16S ribosomal DNA sequence comparison of *Chromohalobacter marismortui*, *Volcaniella eurihalina*, and *Deleya salina* and reclassification of *V. eurihalina* as *Halomonas eurihalina* comb. nov. *Int J Syst Evol Microbiol* 1995; **45**:712-716. PubMed
- Cabrera A, Aguilera M, Fuentes S, Incerti C, Russell NJ, Ramos-Cormenzana A, Monteoliva-Sánchez M. *Halomonas indalinina* sp. nov., a moderately halophilic bacterium isolated from a solar saltern in Cabo de Gata, Almería, southern Spain. *Int J Syst Evol Microbiol* 2007; **57**:376-380. PubMed <http://dx.doi.org/10.1099/ijs.0.64702-0>
- Kim KK, Jin L, Yang HC, Lee ST. *Halomonas gomseomensis* sp. nov., *Halomonas janggokensis* sp. nov., *Halomonas salaria* sp. nov. and *Halomonas denitrificans* sp. nov., moderately halophilic bacteria isolated from saline water. *Int J Syst Evol Microbiol* 2007; **57**:675-681. PubMed <http://dx.doi.org/10.1099/ijs.0.64767-0>
- Soto-Ramírez N, Sánchez-Porro C, Rosas S, González W, Quiñones M, Ventosa A, Montalvo-Rodríguez R. *Halomonas avicenniae* sp. nov., isolated from the salty leaves of the black mangrove

- Avicennia germinans in Puerto Rico. *Int J Syst Evol Microbiol* 2007; **57**:900-905. PubMed <http://dx.doi.org/10.1099/ijs.0.64818-0>
10. Ventosa JJ, Nieto, Oren A. Biology of moderately halophilic aerobic bacteria. *Microbiol Mol Biol Rev* 1998; **62**:504-544. PubMed
 11. Wang YN, Cai H, Chi CQ, Lu AH, Lin XG, Jiang ZF, Wu XL. *Halomonas shengliensis* sp. nov., a moderately halophilic, denitrifying, crude-oil-utilizing bacterium. *Int J Syst Evol Microbiol* 2007; **57**:1222-1226. PubMed <http://dx.doi.org/10.1099/ijs.0.64973-0>
 12. Garrity GM. Names for Life Browser Tool takes expertise out of the database and puts it right in the browser. *Microbiol Today* 2010; **37**:9.
 13. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. *J Mol Biol* 1990; **215**:403-410. PubMed [http://dx.doi.org/10.1016/S0022-2836\(05\)80360-2](http://dx.doi.org/10.1016/S0022-2836(05)80360-2)
 14. Korf I, Yandell M, Bedell J. BLAST, O'Reilly, Sebastopol, 2003.
 15. DeSantis TZ, Hugenholtz P, Larsen N, Rojas M, Brodie EL, Keller K, Huber T, Dalevi D, Hu P, Andersen GL. Greengenes, a chimera-checked 16S rRNA gene database and workbench compatible with ARB. *Appl Environ Microbiol* 2006; **72**:5069-5072. PubMed <http://dx.doi.org/10.1128/AEM.03006-05>
 16. Lee C, Grasso C, Sharlow MF. Multiple sequence alignment using partial order graphs. *Bioinformatics* 2002; **18**:452-464. PubMed <http://dx.doi.org/10.1093/bioinformatics/18.3.452>
 17. Saitou N, Nei M. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* 1987; **4**:406-425. PubMed
 18. Strimmer K, von Haeseler A. Quartet puzzling: a quartet maximum-likelihood method for reconstructing tree topologies. *Mol Biol Evol* 1996; **13**:964-969. <http://dx.doi.org/10.1093/oxfordjournals.molbev.a025664>
 19. Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S. MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol* 2011; **28**:2731-2739. PubMed <http://dx.doi.org/10.1093/molbev/msr121>
 20. Pattengale ND, Alipour M, Bininda-Emonds OR, Moret BM, Stamatakis A. How many bootstrap replicates are necessary? [doi:10.1089/cmb.2009.0179]. [pmid:20377449]. *J Comput Biol* 2010; **17**:337-354.
 21. Pagani I, Liolios K, Jansson J, Chen IM, Smirnova T, Nosrat B, Markowitz VM, Kyrpides NC. The Genomes OnLine Database (GOLD) v.4: status of genomic and metagenomic projects and their associated metadata. *Nucleic Acids Res* 2012; **40**:D571-D579. PubMed <http://dx.doi.org/10.1093/nar/gkr1100>
 22. Schwibbert K, Marin-Sanguino A, Bagyan I, Heidrich G, Lentzen G, Seitz H, Rampp M, Schuster SC, Klenk HP, Pfeiffer F, et al. A blueprint of ectoine metabolism from the genome of the industrial producer *Halomonas elongata* DSM 2581T. *Environ Microbiol* 2011; **13**:1973-1994. PubMed <http://dx.doi.org/10.1111/j.1462-2920.2010.02336.x>
 23. Field D, Garrity G, Gray T, Morrison N, Selengut J, Sterk P, Tatusova T, Thomson N, Allen MJ, Angiuoli SV, et al. The minimum information about a genome sequence (MIGS) specification. *Nat Biotechnol* 2008; **26**:541-547. PubMed <http://dx.doi.org/10.1038/nbt1360>
 24. Field D, Amaral-Zettler L, Cochrane G, Cole JR, Dawyndt P, Garrity GM, Gilbert J, Glöckner FO, Hirschman L, Karsch-Mzrachi I, et al. *PLoS Biol* 2011; **9**:e1001088. PubMed <http://dx.doi.org/10.1371/journal.pbio.1001088>
 25. Euzéby JP. List of bacterial names with standing in nomenclature: A folder available on the Internet. *Int J Syst Bacteriol* 1997; **47**:590-592. PubMed <http://dx.doi.org/10.1099/00207713-47-2-590>
 26. Woese CR, Kandler O, Wheelis ML. Towards a natural system of organisms: proposal for the domains *Archaea*, *Bacteria*, and *Eucarya*. *Proc Natl Acad Sci USA* 1990; **87**:4576-4579. PubMed <http://dx.doi.org/10.1073/pnas.87.12.4576>
 27. Garrity GM, Lilburn T. Phylum XIV. *Proteobacteria* phyl. nov. In: Garrity GM, Krieg NR, Staley JT (eds), *Bergey's Manual of Systematic Bacteriology*, Second Edition, Springer, New York, 2005.
 28. Garrity GM, Bell JA, Lilburn T. Class III. *Gammaproteobacteria* class. nov. In: Brenner DJ, Krieg NR, Staley JT and Garrity GM (eds), *Bergey's Manual of Systematic Bacteriology*, Second Edition, Volume 2, Part C, Springer, New York, 2005, p. 1.
 29. Validation of publication of new names and new combinations previously effectively published outside the IJSEM. List no. 106. *Int J Syst Evol*

- Microbiol* 2005; **55**:2235-2238.
<http://dx.doi.org/10.1099/ijs.0.64108-0>
30. Williams KP, Kelly DP. Proposal for a new class within the phylum *Proteobacteria*, *Acidithiobacillia* classis nov., with the type order *Acidithiobacillales*, and emended description of the class *Gammaproteobacteria*. *Int J Syst Evol Microbiol* 2013; **63**:2901-2906. PubMed <http://dx.doi.org/10.1099/ijs.0.049270-0>
 31. Garrity GM, Bell JA, Lilburn T. Order VIII. *Oceanospirillales* ord. nov. In: Brenner DJ, Krieg NR, Staley JT, Garrity GM (eds), *Bergey's Manual of Systematic Bacteriology*, second edition, vol. 2 (*The Proteobacteria*), part B (*The Gammaproteobacteria*), Springer, New York, 2005, p. 270.
 32. Validation List no. 29. Validation of the publication of new names and new combinations previously effectively published outside the IJSB. *Int J Syst Bacteriol* 1989; **39**:205-206.
<http://dx.doi.org/10.1099/00207713-39-2-205>
 33. Dobson SJ, Franzmann PD. Unification of the genera *Deleya* (Baumann et al. 1983), *Halomonas* (Vreeland et al. 1980), and *Halovibrio* (Fendrich 1988) and the species *Paracoccus halodenitrificans* (Robinson and Gibbons 1952) into a single genus, *Halomonas*, and placement of the genus *Zymobacter* in the family *Halomonadaceae*. *Int J Syst Bacteriol* 1996; **46**:550-558. <http://dx.doi.org/10.1099/00207713-46-2-550>
 34. Ntougias S, Zervakis GI, Fasseas C. *Halotalea alkalilenta* gen. nov., sp. nov., a novel osmotolerant and alkalitolerant bacterium from alkaline olive mill wastes, and emended description of the family *Halomonadaceae* Franzmann et al. 1989, emend. Dobson and Franzmann 1996. *Int J Syst Evol Microbiol* 2007; **57**:1975-1983. PubMed <http://dx.doi.org/10.1099/ijs.0.65078-0>
 35. Ben Ali Gam Z, Abdelkafi S, Casalot L, Tholozan JL, Oueslati R, Labat M. *Modicisalibacter tunisiensis* gen. nov., sp. nov., an aerobic, moderately halophilic bacterium isolated from an oil-field-water injection sample, and emended description of the family *Halomonadaceae* Franzmann et al. 1989 emend Dobson and Franzmann 1996 emend. Ntougias et al. 2007. *Int J Syst Evol Microbiol* 2007; **57**:2307-2313. PubMed <http://dx.doi.org/10.1099/ijs.0.65088-0>
 36. Dobson SJ, Franzmann PD. Unification of the genera *Deleya* (Baumann et al. 1983), *Halomonas* (Vreeland et al. 1980), and *Halovibrio* (Fendrich 1988) and the species *Paracoccus halodenitrificans* (Robinson and Gibbons 1952) into a single genus, *Halomonas*, and placement of the genus *Zymobacter* in the family *Halomonadaceae*. *Int J Syst Bacteriol* 1996; **46**:550-558. <http://dx.doi.org/10.1099/00207713-46-2-550>
 37. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig JT, et al. Gene ontology: tool for the unification of biology. *Nat Genet* 2000; **25**:25-29. PubMed <http://dx.doi.org/10.1038/75556>
 38. Berendes F, Gottschalk G, Heine-Dobbernack E, Moore ERB, Tindall BJ. *Halomonas desiderata* sp. nov, a new alkaliphilic, halotolerant and denitrifying bacterium isolated from a municipal sewage works. *Syst Appl Microbiol* 1996; **19**:158-167. [http://dx.doi.org/10.1016/S0723-2020\(96\)80041-5](http://dx.doi.org/10.1016/S0723-2020(96)80041-5)
 39. Romano I, Gottschalk G, Heine-Dobbernack E, Moore ERB, Tindall BJ. Characterization of a haloalkaliphilic strictly aerobic bacterium, isolated from *Pantelleria island*. *Syst Appl Microbiol* 1996; **19**:326-333. [http://dx.doi.org/10.1016/S0723-2020\(96\)80059-2](http://dx.doi.org/10.1016/S0723-2020(96)80059-2)
 40. Heyrman J, Balcaen A, De Vos P, Swings J. *Halomonas muralis* sp. nov., isolated from microbial biofilms colonizing the walls and murals of the Saint-Catherine chapel (Castle Herberstein, Austria). *Int J Syst Evol Microbiol* 2002; **52**:2049-2054. PubMed <http://dx.doi.org/10.1099/ijs.0.02166-0>
 41. Jeon CO, Lim JM, Lee JR, Lee GS, Park DJ, Lee JC, Oh HW, Kim CJ. *Halomonas kribbensis* sp. nov., a novel moderately halophilic bacterium isolated from a solar saltern in Korea. *Int J Syst Evol Microbiol* 2007; **57**:2194-2198. PubMed <http://dx.doi.org/10.1099/ijs.0.65285-0>
 42. Klenk HP, Göker M. En route to a genome - based classification of *Archaea* and *Bacteria*? *Syst Appl Microbiol* 2010; **33**:175-182. PubMed <http://dx.doi.org/10.1016/j.syapm.2010.03.003>
 43. Göker M, Klenk HP. Phylogeny-driven target selection for genome-sequencing (and other) projects. *Stand Genomic Sci* 2013; **8**:360-374. PubMed <http://dx.doi.org/10.4056/sigs.3446951>
 44. Kyrpides NC, Woyke T, Eisen JA, Garrity G, Lilburn TG, Beck BJ, Whitman WB, Hugenholtz P, Klenk HP. Genomic Encyclopedia of Type Strains, Phase I: the one thousand microbial genomes (KMG-I) project. *Stand Genomic Sci* 2013; **9**:628-634. <http://dx.doi.org/10.4056/sigs.5068949>
 45. Wu D, Hugenholtz P, Mavromatis K, Pukall R, Dalin E, Ivanova NN, Kunin V, Goodwin L, Wu

- M, Tindall BJ, *et al.* A phylogeny-driven genomic encyclopaedia of *Bacteria* and *Archaea*. *Nature* 2009; **462**:1056-1060. PubMed <http://dx.doi.org/10.1038/nature08656>
46. Mavromatis K, Land ML, Brettin TS, Quest DJ, Copeland A, Clum A, Goodwin L, Woyke T, Lapidus A, Klenk HP, *et al.* The fast changing landscape of sequencing technologies and their impact on microbial genome assemblies and annotation. *PLoS ONE* 2012; **7**:e48837. PubMed <http://dx.doi.org/10.1371/journal.pone.0048837>
47. List of growth media used at DSMZ. <http://www.dsmz.de/catalogues/catalogue-microorganisms/culture-technology/list-of-media-for-microorganisms.html>.
48. Gemeinholzer B, Dröge G, Zetzsche H, Haszprunar G, Klenk HP, Güntsch A, Berendsohn WG, Wägele JW. The DNA Bank Network: the start from a German initiative. *Biopreserv Biobank* 2011; **9**:51-55. <http://dx.doi.org/10.1089/bio.2010.0029>
49. Bennett S. Solexa Ltd. *Pharmacogenomics* 2004; **5**:433-438. PubMed <http://dx.doi.org/10.1517/14622416.5.4.433>
50. The DOE Joint Genome Institute. <http://www.jgi.doe.gov>
51. Mingkun L, Copeland A, Han J. DUK, unpublished, 2011.
52. Zerbino D, Birney E. Velvet: algorithms for de novo short read assembly using de Bruijn graphs. *Genome Res* 2008; **18**:821-829. PubMed <http://dx.doi.org/10.1101/gr.074492.107>
53. <https://github.com/lh3/wgsim>
54. Gnerre S, MacCallum I. High-quality draft assemblies of mammalian genomes from massively parallel sequence data. *Proc Natl Acad Sci USA* 2011; **108**:1513-1518. PubMed <http://dx.doi.org/10.1073/pnas.1017351108>
55. Hyatt D, Chen GL, LoCascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics* 2010; **11**:119. PubMed <http://dx.doi.org/10.1186/1471-2105-11-119>
56. Mavromatis K, Ivanova NN, Chen IM, Szeto E, Markowitz VM, Kyrpides NC. The DOE-JGI Standard operating procedure for the annotations of microbial genomes. *Stand Genomic Sci* 2009; **1**:63-67. PubMed <http://dx.doi.org/10.4056/sigs.632>
57. Pati A, Ivanova NN, Mikhailova N, Ovchinnikova G, Hooper SD, Lykidis A, Kyrpides NC. GenePRIMP: a gene prediction improvement pipeline for prokaryotic genomes. *Nat Methods* 2010; **7**:455-457. PubMed <http://dx.doi.org/10.1038/nmeth.1457>
58. Markowitz VM, Ivanova NN, Chen IMA, Chu K, Kyrpides NC. IMG ER: a system for microbial genome annotation expert review and curation. *Bioinformatics* 2009; **25**:2271-2278. PubMed <http://dx.doi.org/10.1093/bioinformatics/btp393>
59. Meier-Kolthoff JP, Auch AF, Klenk HP, Göker M. Genome sequence-based species delimitation with confidence intervals and improved distance functions. *BMC Bioinformatics* 2013; **14**:60. PubMed <http://dx.doi.org/10.1186/1471-2105-14-60>
60. Auch AF, von Jan M, Klenk HP, Göker M. Digital DNA-DNA hybridization for microbial species delineation by means of genome-to-genome sequence comparison. *Stand Genomic Sci* 2010; **2**:117-134. PubMed <http://dx.doi.org/10.4056/sigs.531120>
61. Auch AF, Klenk HP, Göker M. Standard operating procedure for calculating genome-to-genome distances based on high-scoring segment pairs. *Stand Genomic Sci* 2010; **2**:142-148. PubMed <http://dx.doi.org/10.4056/sigs.541628>
62. Derr P, Boder E, Goulian M. Changing the specificity of a bacterial chemoreceptor. *J Mol Biol* 2006; **355**:923-932. PubMed <http://dx.doi.org/10.1016/j.jmb.2005.11.025>
63. Alexander RP, Zhulin IB. Evolutionary genomics reveals conserved structural determinants of signaling and adaptation in microbial chemoreceptors. *Proc Natl Acad Sci USA* 2007; **104**:2885-2890. PubMed <http://dx.doi.org/10.1073/pnas.0609359104>
64. Harrison DM, Skidmore J, Armitage JP, Maddock JR. Localization and environmental regulation of MCP-like proteins in *Rhodobacter sphaeroides*. *Mol Microbiol* 1999; **31**:885-892. PubMed <http://dx.doi.org/10.1046/j.1365-2958.1999.01226.x>