

UC Davis

UC Davis Previously Published Works

Title

Complete genome sequence of the extremely halophilic Halanaerobium praevalens type strain (GSLT)

Permalink

<https://escholarship.org/uc/item/8314q0ww>

Journal

Environmental Microbiome, 4(3)

ISSN

2524-6372

Authors

Ivanova, Natalia
Sikorski, Johannes
Chertkov, Olga
[et al.](#)

Publication Date

2011-05-01

DOI

10.4056/sigs.1824509

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Complete genome sequence of the extremely halophilic *Halanaerobium praevalens* type strain (GSL^T)

Natalia Ivanova¹, Johannes Sikorski², Olga Chertkov^{1,3}, Matt Nolan¹, Susan Lucas¹, Nancy Hammon¹, Shweta Deshpande¹, Jan-Fang Cheng¹, Roxanne Tapia^{1,3}, Cliff Han^{1,3}, Lynne Goodwin^{1,3}, Sam Pitluck¹, Marcel Huntemann¹, Konstantinos Liolios¹, Ioanna Pagani¹, Konstantinos Mavromatis¹, Galina Ovchinnikova¹, Amrita Pati¹, Amy Chen⁴, Krishna Palaniappan⁴, Miriam Land^{1,5}, Loren Hauser^{1,5}, Evelyne-Marie Brambilla², K. Palani Kannan², Manfred Rohde⁶, Brian J. Tindall², Markus Göker², John C. Detter^{1,3}, Tanja Woyke¹, James Bristow¹, Jonathan A. Eisen^{1,7}, Victor Markowitz⁴, Philip Hugenholtz^{1,8}, Nikos C. Kyrpides¹, Hans-Peter Klenk², and Alla Lapidus^{1*}

¹ DOE Joint Genome Institute, Walnut Creek, California, USA

² DSMZ - German Collection of Microorganisms and Cell Cultures GmbH, Braunschweig, Germany

³ Los Alamos National Laboratory, Bioscience Division, Los Alamos, New Mexico, USA

⁴ Biological Data Management and Technology Center, Lawrence Berkeley National Laboratory, Berkeley, California, USA

⁵ Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA

⁶ HZI – Helmholtz Centre for Infection Research, Braunschweig, Germany

⁷ University of California Davis Genome Center, Davis, California, USA

⁸ Australian Centre for Ecogenomics, School of Chemistry and Molecular Biosciences, The University of Queensland, Brisbane, Australia

*Corresponding author: Alla Lapidus

Keywords: strictly anaerobic, non-motile, Gram-negative, straight rod-shaped, halophilic, moderate alkaliphile, mesophilic, chemoorganotroph, *Halanaerobiaceae*, GEBA

Halanaerobium praevalens Zeikus et al. 1984 is the type species of the genus *Halanaerobium*, which in turn is the type genus of the family *Halanaerobiaceae*. The species is of interest because it is able to reduce a variety of nitro-substituted aromatic compounds at a high rate, and because of its ability to degrade organic pollutants. The strain is also of interest because it functions as a hydrolytic bacterium, fermenting complex organic matter and producing intermediary metabolites for other trophic groups such as sulfate-reducing and methanogenic bacteria. It is further reported as being involved in carbon removal in the Great Salt Lake, its source of isolation. This is the first completed genome sequence of a representative of the genus *Halanaerobium* and the second genome sequence from a type strain of the family *Halanaerobiaceae*. The 2,309,262 bp long genome with its 2,110 protein-coding and 70 RNA genes is a part of the *Genomic Encyclopedia of Bacteria and Archaea* project.

Introduction

Strain GSL^T (= DSM 2228 =ATCC 33744) is the type strain of the species *Halanaerobium praevalens*, which is the type species of its genus *Halanaerobium* [1]. Originally described as *Haloanaerobium* [2], the name was later changed to *Halanaerobium* to conform with rule 61 of the Bacteriological Code [3]. The genus currently consists of nine validly named species [4]. The genus name is derived from the Latinized Greek word *hals*; *halos* meaning *salt*, the Latinized Greek word *an-* meaning *not*, the Latinized Greek word *aer* meaning *air* and the Latinized

Greek word *bios* meaning *life*, yielding the Neo-Latin word 'Halanaerobium' meaning 'salt organism which grows in the absence of air' [4]. The species epithet is derived from the Latin word 'praevalens' (very powerful, very strong, here prevalent) [4]. Strain GSL^T was isolated from the hypersaline surface sediments of Great Salt Lake, Utah, USA [2]. Further strains of *H. praevalens* have been isolated from canned salted Swedish fermented herrings referred to as Surströmming [5] and probably also from the Red Sea [6]. Other

members of the genus have been isolated also from high salt environments distributed worldwide [7-10]. The enzymatic activities of the fatty acid synthetase complex and the *D*-BAPA (N'-benzoyl-arginine-*p*-nitroanilide)-ase of *H. praevalens* have been studied in more detail [11,12]. Here we present a summary classification and a set of features for *H. praevalens* strain GSL^T, together with the description of the complete genome sequencing and annotation.

Classification and features

A representative genomic 16S rRNA sequence of *H. praevalens* was compared using NCBI BLAST under default values (e.g., considering only the best 250 hits) with the most recent release of the Greengenes database [13] and the relative frequencies, weighted by BLAST scores, of taxa and keywords (reduced to their stem [14]) were determined. The five most frequent genera were *Halanaerobium* (81.9%), *Halo-thermothrix* (7.8%), *Halanaerobacter* (2.7%), *Acetohalobium* (2.3%) and *Natroniella* (1.9%). Regarding hits to sequences from other members of the genus, the average identity within HSPs (high-scoring segment pairs) was 97.8%, whereas the average coverage by HSPs was 96.3%. The species yielding the highest score was *Halanaerobium saccharolyticum*. (Note that the Greengenes database uses the INSDC (= EMBL/NCBI/DDBJ) annotation, which is not an authoritative source for nomenclature or classification.) The five most frequent keywords within the labels of environmental samples which yielded hits were 'microbi' (9.4%), 'hypersalin' (9.1%), 'mat' (8.6%), 'len, miniprim, new, view, world' (8.5%) and 'food' (3.4%). The single most frequent keyword within the labels of environmental samples which yielded hits of a higher score than the highest scoring species was 'hypersalin, len, mat, microbi, miniprim, new, view, world' (12.5%). These key words are in line with the ecology and the niche from where strains of *H. praevalens* have been isolated.

Figure 1 shows the phylogenetic neighborhood of *H. praevalens* GSL^T in a 16S rRNA gene based tree. The sequences of the four 16S rRNA gene copies in the genome differ from each other by up to five nucleotides, and differ by up to five nucleotides from the previously published 16S rRNA gene sequence (AB022034).

The cells of strain GSL^T are straight and rod-shaped (1 × 2.4 µm) (Figure 2) when grown to the mid-log phase at 37°C on CS medium containing 12.5% NaCl and 0.5% glucose [2]. When grown at higher

NaCl concentrations (> 20%) the cells appear granulated and shorter in length [2]. Single colonies were reported as white to translucent in color, 0.5-2.0 mm in diameter, and glistened, when grown on agar plates containing CS medium, 12.5% NaCl, and 0.5% glucose [2]. *H. praevalens* cells stain Gram-negative [2] and electron microscopy in thin section revealed architectural features typical of Gram-negative bacteria [2]. However, the positive *D*-BAPA-ase reaction [11] confirms its phylogenetic affiliation to the endospore-forming firmicutes (Table 1). In this respect, *H. praevalens* is able to hydrolyze only the D- but not the L- isomer of N'-benzoyl-arginine-*p*-nitroanilide (BAPA) [11]. The activity of *D*-BAPA-ase was highest at low NaCl concentration (100 mM) and completely inhibited at NaCl concentration equal or higher than 1.0 M (~12%) [11]. Strain GSL^T is described to be non-motile, although many flagellar genes have been identified in the genome (see below). Other isolates of *H. praevalens* were described as motile [5,6], as were other members of the genus [7,9,10], suggesting strain GSL^T is atypical with regard to motility. The organism is a strictly anaerobic chemoorgano-troph [2]. It grows at NaCl concentrations between 2% and 30%, with optimal growth at approximately 13% [2]. The doubling time is 4 h at 12.5% NaCl and 7 h at 25% NaCl in complex CS medium [2]. The temperature range for growth ranges from 5°C to 60°C, with an optimum at 37°C [2]. The pH range for growth is between 6.0 and 9.0, with an optimum at pH between 7.0 and 7.4 [2]. Strain GSL^T is able to utilize carbohydrates (including pectin and *N*-acetylglucosamine), amino acids, yeast extract, and trypticase; the two latter serving as carbon and energy sources on complex medium [2]. The fermentation of glucose yielded butyrate, acetate, propionate, H₂, and CO₂ as major products [2]. Also, fructose, D-mannose and maltose are utilized and methionine is transformed to methylmercaptan [2]. Penicillin, tetracycline, cycloserine, chloramphenicol (each at 100 µg/ml culture) or sodium azide (500 µg/ml) completely inhibit the growth of *H. praevalens* [2]. Strain GSL^T was also able to degrade nitro-substituted aromatic compounds such as nitrobenzene, *o*-nitrophenol, *m*-nitrophenol, *p*-nitrophenol, 2,4-dinitrophenol, and 2,4-dinitroaniline [34]. The fatty acid synthetase of *H. praevalens* is only slightly inhibited at 17.5% and was the first reported to be active in the presence of high salt concentrations [12]. *H. praevalens* was reported to be involved in carbon sequestration in the Great Salt Lake [35], since it is present in the sediments of

this lake in high numbers ($\geq 10^8$ cells/ml) [2,36]. *H. praevalens* regulates its internal osmotic pressure by the accumulation of salts (Na^+ , K^+ , Cl^-) rather than by compatible solutes [36]. High concentrations of these salts were measured inside the cells, in sufficient concentration to be isotonic or hypertonic with the medium [37]. Thiosulfate reduction

and rhodanese-like enzyme (thiosulfate:cyanide sulfur-transferase) activities also tested positive in strain GSLT [8]. Early in 1987, Matheson *et al.* [38] established the primary structure of the ribosomal A-protein of the strain GSLT, which is the equivalent to the ribosomal protein L12 from *Escherichia coli*.




Figure 1. Phylogenetic tree highlighting the position of *H. praevalens* GSLT relative to the other type strains within the family *Halanaerobiaceae*. The tree was inferred from 1,460 aligned characters [15,16] of the 16S rRNA gene sequence under the maximum likelihood criterion [17] and rooted in accordance with the current taxonomy. The branches are scaled in terms of the expected number of substitutions per site. Numbers to the right of bifurcations are support values from 1,000 bootstrap replicates [18] if larger than 60%. Lineages with type strain genome sequencing projects registered in GOLD [19] are labeled with an asterisk, published genomes with two asterisks [20].




Figure 2. Scanning electron micrograph of *H. praevalens* GSLT

Table 1. Classification and general features of *H. praevalens* GSL^T according to the MIGS recommendations [21] and the NamesforLife database [22].

MIGS ID	Property	Term	Evidence code
		Domain <i>Bacteria</i>	TAS [23]
		Phylum <i>Firmicutes</i>	TAS [24,25]
		Class <i>Clostridia</i>	TAS [26,27]
	Current classification	Order <i>Halanaerobiales</i>	TAS [28-30]
		Family <i>Halanaerobiaceae</i>	TAS [1,31]
		Genus <i>Halanaerobium</i>	TAS [1-3,30]
		Species <i>Halanaerobium praevalens</i>	TAS [1,2]
		Type strain GSL	TAS [2]
	Gram stain	negative	TAS [2]
	Cell shape	straight rods	TAS [2]
	Motility	non-motile	TAS [2]
	Sporulation	none	TAS [2]
	Temperature range	above 5°C and below 60°C	TAS [2]
	Optimum temperature	37°C	TAS [2]
	Salinity	2%-30% NaCl, optimum at 13%	TAS [2]
MIGS-22	Oxygen requirement	strictly anaerobic	TAS [2]
	Carbon source	yeast extract, trypticase	TAS [2]
	Energy metabolism	chemoorganotroph	TAS [2]
MIGS-6	Habitat	saline environments	TAS [2]
MIGS-15	Biotic relationship	not reported	NAS
MIGS-14	Pathogenicity	not reported	NAS
	Biosafety level	1	TAS [32]
	Isolation	surface sediments of a saline lake	TAS [2]
MIGS-4	Geographic location	Great Salt Lake, Utah, USA	TAS [2]
MIGS-5	Sample collection time	between August 1979 and August 1980	TAS [2]
MIGS-4.1	Latitude	41.15	NAS
MIGS-4.2	Longitude	-112.67	NAS
MIGS-4.3	Depth	10 m bottom sediment	TAS [2]
MIGS-4.4	Altitude	1.755 m above sea level	NAS

Evidence codes - IDA: Inferred from Direct Assay (first time in publication); TAS: Traceable Author Statement (i.e., a direct report exists in the literature); NAS: Non-traceable Author Statement (i.e., not directly observed for the living, isolated sample, but based on a generally accepted property for the species, or anecdotal evidence). These evidence codes are from of the Gene Ontology project [33]. If the evidence code is IDA, the property was directly observed by one of the authors or an expert mentioned in the acknowledgements.

Chemotaxonomy

When grown on CS medium, at 5% NaCl, more lipids are produced than at 25% NaCl (3.74% and 2.54% of the dry weight of the organism, respectively) [2]. At 5% NaCl, the fractions of glycolipids and phospholipids are 46.9% and 44.5% of the total lipids, respectively. At 25%, the proportion changes in favor of phospholipids (49.1%), whereas glycolipids decrease (43.0%) [2]. The glycolipids consist of a single component diacylglycerol derivative, while the phospholipids consist mainly of cardiolipin (CL), phosphatidyl glycerol (PG),

and three minor unidentified constituents [2]. When grown on CS medium, at 5%, the major fatty acids are C_{14:0} (49.3%), C_{16:1} (31.3%) and C_{16:0} (11.4%). At 25%, these fractions change to 36.8%, 39%, and 22.7%, respectively. Similar though more detailed results on the fatty acid composition have been reported recently [6].

Genome sequencing and annotation

Genome project history

This organism was selected for sequencing on the basis of its phylogenetic position [39], and is part of the *Genomic Encyclopedia of Bacteria and Archaea* project [40]. The genome project is deposited

in the Genome On Line Database [19] and the complete genome sequence is deposited in GenBank. Sequencing, finishing and annotation were performed by the DOE Joint Genome Institute (JGI). A summary of the project information is shown in Table 2.

Table 2. Genome sequencing project information

MIGS ID	Property	Term
MIGS-31	Finishing quality	Finished
MIGS-28	Libraries used	Three genomic libraries: one 454 pyrosequence standard library, one 454 PE library (13 kb insert size), one Illumina library
MIGS-29	Sequencing platforms	Illumina GAii, 454 GS FLX Titanium
MIGS-31.2	Sequencing coverage	201.2 × Illumina; 174.2 × pyrosequence
MIGS-30	Assemblers	Newbler version 2.3, Velvet, phrap
MIGS-32	Gene calling method	Prodigal 1.4, GenePRIMP
	INSDC ID	CP002175
	Genbank Date of Release	October 21, 2010
	GOLD ID	Gc01415
	NCBI project ID	32591
	Database: IMG-GEBA	2503283011
MIGS-13	Source material identifier	DSM 2228
	Project relevance	Tree of Life, GEBA

Growth conditions and DNA isolation

H. praevalens GSLT, DSM 2228, was grown anaerobically in DSMZ medium 210 (*'Haloanaerobium'* medium) [41] at 30–37°C. DNA was isolated from 0.5–1 g of cell paste using MasterPure Gram-positive DNA purification kit (Epicentre MGP04100) following the standard protocol as recommended by the manufacturer, with modification st/DL for cell lysis as described in Wu *et al.* [40]. DNA is available through the DNA Bank Network [42].

Genome sequencing and assembly

The genome was sequenced using a combination of Illumina and 454 sequencing platforms. All general aspects of library construction and sequencing can be found at the JGI website [43]. Pyrosequencing reads were assembled using the Newbler assembler (Roche). The initial Newbler assembly, consisting of 85 contigs in 31 scaffolds, was converted into a phrap [44] assembly by making fake reads from the consensus, to collect the read pairs in the 454 paired end library. Illumina sequencing data (360 Mb) was assembled with Velvet [45] and the consensus sequences were shredded into 1.5 kb overlapped fake reads and assembled together with the 454 data. The 454

draft assembly was based on 401.6 Mb 454 draft data and all of the 454 paired end data. Newbler parameters are -consed -a 50 -l 350 -g -m -ml 20. The Phred/Phrap/Consed software package [44] was used for sequence assembly and quality assessment in the subsequent finishing process. After the shotgun stage, reads were assembled with parallel phrap (High Performance Software, LLC). Possible mis-assemblies were corrected with gapResolution [43], Dupfinisher [46], or sequencing cloned bridging PCR fragments with subcloning. Gaps between contigs were closed by editing in Consed, by PCR and by Bubble PCR primer walks (J.-F. Chang, unpublished). A total of 417 additional reactions and ten shatter libraries were necessary to close gaps and to raise the quality of the finished sequence. Illumina reads were also used to correct potential base errors and increase consensus quality using a software Polisher developed at JGI [47]. The error rate of the completed genome sequence is less than 1 in 100,000. Together, the combination of the Illumina and 454 sequencing platforms provided 375.4 × coverage of the genome. The final assembly contained 838,597 pyrosequence and 12,903,210 Illumina reads.

Genome annotation

Genes were identified using Prodigal [48] as part of the Oak Ridge National Laboratory genome annotation pipeline, followed by a round of manual curation using the JGI GenePRIMP pipeline [49]. The predicted CDSs were translated and used to search the National Center for Biotechnology Information (NCBI) non-redundant database, UniProt, TIGR-Fam, Pfam, PRIAM, KEGG, COG, and InterPro databases. Additional gene prediction analysis and functional annotation was performed within the Integrated Microbial Genomes - Expert Review (IMG-ER) platform [50].

Genome properties

The genome consists of a 2,309,262 bp long chromosome with a G+C content of 30.3% (Figure 3 and Table 3). Of the 2,180 genes predicted, 2,110 were protein-coding genes, and 70 RNAs; 42 pseudogenes were also identified. The majority of the protein-coding genes (77.7%) were assigned with a putative function while the remaining ones were annotated as hypothetical proteins. The distribution of genes into COGs functional categories is presented in Table 4.




Figure 3. Graphical circular map of the chromosome. From outside to the center: Genes on forward strand (color by COG categories), Genes on reverse strand (color by COG categories), RNA genes (tRNAs green, rRNAs red, other RNAs black), GC content, GC skew.

Table 3. Genome Statistics

Attribute	Value	% of Total
Genome size (bp)	2,309,262	100.00%
DNA coding region (bp)	2,059,925	89.20%
DNA G+C content (bp)	699,559	30.29%
Number of replicons	1	
Extrachromosomal elements	0	
Total genes	2,180	100.00%
RNA genes	70	3.21%
rRNA operons	4	
Protein-coding genes	2,110	96.79%
Pseudo genes	42	1.93%
Genes with function prediction	1,694	77.71%
Genes in paralog clusters	310	14.22%
Genes assigned to COGs	1,760	80.73%
Genes assigned Pfam domains	1,860	85.32%
Genes with signal peptides	446	20.46%
Genes with transmembrane helices	582	26.70%
CRISPR repeats	1	

Table 4. Number of genes associated with the general COG functional categories

Code	value	%age	Description
J	133	6.9	Translation, ribosomal structure and biogenesis
A	0	0.0	RNA processing and modification
K	124	6.4	Transcription
L	115	6.0	Replication, recombination and repair
B	1	0.0	Chromatin structure and dynamics
D	24	1.2	Cell cycle control, cell division, chromosome partitioning
Y	0	0.0	Nuclear structure
V	18	0.9	Defense mechanisms
T	126	6.5	Signal transduction mechanisms
M	106	5.5	Cell wall/membrane/envelope biogenesis
N	76	3.9	Cell motility
Z	0	0.0	Cytoskeleton
W	0	0.0	Extracellular structures
U	44	2.3	Intracellular trafficking, secretion, and vesicular transport
O	66	3.4	Posttranslational modification, protein turnover, chaperones
C	123	6.4	Energy production and conversion
G	150	7.8	Carbohydrate transport and metabolism
E	136	7.0	Amino acid transport and metabolism
F	74	3.8	Nucleotide transport and metabolism
H	80	4.1	Coenzyme transport and metabolism
I	45	2.3	Lipid transport and metabolism
P	121	6.3	Inorganic ion transport and metabolism
Q	18	0.9	Secondary metabolites biosynthesis, transport and catabolism
R	186	9.6	General function prediction only
S	168	8.7	Function unknown
-	420	19.3	Not in COGs

Acknowledgements

We would like to gratefully acknowledge the help of Olivier D. Ngatchou-Djao (HZI) for drafting the manuscript, and Helga Pomrenke (DSMZ) for growing *H. praevalens* cultures. This work was performed under the auspices of the US Department of Energy Office of Science, Biological and Environmental Research Program, and by the University of California, Lawrence Berkeley National Laboratory under contract No. DE-

AC02-05CH11231, Lawrence Livermore National Laboratory under Contract No. DE-AC52-07NA27344, and Los Alamos National Laboratory under contract No. DE-AC02-06NA25396, UT-Battelle and Oak Ridge National Laboratory under contract DE-AC05-00OR22725, as well as German Research Foundation (DFG) INST 599/1-2.

References

1. Validation of the publication of new names and new combinations previously effectively published outside the IJSB. List No. 16. *Int J Syst Bacteriol* 1984; **34**:503-504. [doi:10.1099/00207713-34-4-503](https://doi.org/10.1099/00207713-34-4-503)
2. Zeikus JG, Hegge PW, Thompson TE, Phelps TJ, Langworthy TA. Isolation and description of *Haloanaerobium praevalens* gen. nov. and sp. nov., an obligately anaerobic halophile common to Great Salt Lake sediments. *Curr Microbiol* 1983; **9**:225-233. [doi:10.1007/BF01567586](https://doi.org/10.1007/BF01567586)
3. Oren A. Change of the names *Haloanaerobiales*, *Haloanaerobiaceae* and *Haloanaerobium* to *Halanaerobiales*, *Halanaerobiaceae* and *Halanaerobium*, respectively, and further nomenclatural changes within the order *Halanaerobiales*. *Int J Syst Evol Microbiol* 2000; **50**:2229-2230. [PubMed](https://pubmed.ncbi.nlm.nih.gov/11111111/) [doi:10.1099/00207713-50-6-2229](https://doi.org/10.1099/00207713-50-6-2229)
4. Euzéby JP. List of bacterial names with standing in nomenclature: a folder available on the Internet. *Int J Syst Bacteriol* 1997; **47**:590-592. [PubMed](https://pubmed.ncbi.nlm.nih.gov/11111111/) [doi:10.1099/00207713-47-2-590](https://doi.org/10.1099/00207713-47-2-590)
5. Kobayashi T, Kimura B, Fujii T. Strictly anaerobic halophiles isolated from canned Swedish fermented herrings (Surströmming). *Int J Food Microbiol* 2000; **54**:81-89. [PubMed](https://pubmed.ncbi.nlm.nih.gov/11111111/) [doi:10.1016/S0168-1605\(99\)00172-5](https://doi.org/10.1016/S0168-1605(99)00172-5)
6. Eder W, Jahnke LL, Schmidt M, Huber R. Microbial diversity of the brine-seawater interface of the Kebrit Deep, Red Sea, studied via 16S rRNA gene sequences and cultivation methods. *Appl Environ Microbiol* 2001; **67**:3077-3085. [PubMed](https://pubmed.ncbi.nlm.nih.gov/11111111/) [doi:10.1128/AEM.67.7.3077-3085.2001](https://doi.org/10.1128/AEM.67.7.3077-3085.2001)
7. Tsai CR, Garcia JL, Patel BKC, Cayol JL, Baresi L, Mah RA. *Haloanaerobium alcaliphilum* sp. nov., an anaerobic moderate halophile from the sediments of Great Salt Lake, Utah. *Int J Syst Bacteriol* 1995; **45**:301-307. [PubMed](https://pubmed.ncbi.nlm.nih.gov/11111111/) [doi:10.1099/00207713-45-2-301](https://doi.org/10.1099/00207713-45-2-301)
8. Ravot G, Casalot L, Ollivier B, Loison G, Magot M. rdlA, a new gene encoding a rhodanese-like protein in *Halanaerobium congolense* and other thiosulfate-reducing anaerobes. *Res Microbiol* 2005; **156**:1031-1038. [PubMed](https://pubmed.ncbi.nlm.nih.gov/11111111/) [doi:10.1016/j.resmic.2005.05.009](https://doi.org/10.1016/j.resmic.2005.05.009)
9. Cayol JL, Ollivier B, Patel BKC, Ageron E, Grimon PAD, Prensier G, Garcia JL. *Haloanaerobium lacusroseus* sp. nov., an extremely halophilic fermentative bacterium from the sediments of a hypersaline Lake. *Int J Syst Bacteriol* 1995; **45**:790-797. [PubMed](https://pubmed.ncbi.nlm.nih.gov/11111111/) [doi:10.1099/00207713-45-4-790](https://doi.org/10.1099/00207713-45-4-790)
10. Bhupathiraju VK, McInerney MJ, Woese CR, Tanner RS. *Haloanaerobium kushneri* sp. nov., an obligately halophilic, anaerobic bacterium from an oil brine. *Int J Syst Bacteriol* 1999; **49**:953-960. [PubMed](https://pubmed.ncbi.nlm.nih.gov/11111111/) [doi:10.1099/00207713-49-3-953](https://doi.org/10.1099/00207713-49-3-953)
11. Oren A, Gofshtein-Gandman LV, Keynan A. Hydrolysis of N¹-benzoyl-D-arginine-p-nitroanilide by members of the *Haloanaerobiaceae*: additional evidence that *Haloanaerobium praevalens* is related to endospore-forming bacteria. *FEMS Microbiol Lett* 1989; **58**:5-10.
12. Oren A, Gurevich P. The fatty acid synthetase complex of *Haloanaerobium praevalens* is not inhibited by salt. *FEMS Microbiol Lett* 1993; **108**:287-290. [PubMed](https://pubmed.ncbi.nlm.nih.gov/11111111/) [doi:10.1111/j.1574-6968.1993.tb06117.x](https://doi.org/10.1111/j.1574-6968.1993.tb06117.x)
13. DeSantis TZ, Hugenholtz P, Larsen N, Rojas M, Brodie E, Keller K, Huber T, Dalevi D, Hu P, Andersen G. Greengenes, a chimera-checked 16S rRNA gene database and workbench compatible with ARB. *Appl Environ Microbiol* 2006; **72**:5069-5072. [PubMed](https://pubmed.ncbi.nlm.nih.gov/11111111/) [doi:10.1128/AEM.03006-05](https://doi.org/10.1128/AEM.03006-05)
14. Porter MF. An algorithm for suffix stripping. Program: electronic library and information systems. 1980; **14**:130-137.
15. Castresana J. Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. *Mol Biol Evol* 2000; **17**:540-552. [PubMed](https://pubmed.ncbi.nlm.nih.gov/11111111/)

16. Lee C, Grasso C, Sharlow MF. Multiple sequence alignment using partial order graphs. *Bioinformatics* 2002; **18**:452-464. [PubMed](#) [doi:10.1093/bioinformatics/18.3.452](https://doi.org/10.1093/bioinformatics/18.3.452)
17. Stamatakis A, Hoover P, Rougemont J. A rapid bootstrap algorithm for the RAxML Web servers. *Syst Biol* 2008; **57**:758-771. [PubMed](#) [doi:10.1080/10635150802429642](https://doi.org/10.1080/10635150802429642)
18. Pattengale ND, Alipour M, Bininda-Emonds ORP, Moret BME, Stamatakis A. How many bootstrap replicates are necessary? *Lect Notes Comput Sci* 2009; **5541**:184-200. [PubMed](#) [doi:10.1007/978-3-642-02008-7_13](https://doi.org/10.1007/978-3-642-02008-7_13)
19. Liolios K, Chen IM, Mavromatis K, Tavernarakis N, Hugenholtz P, Markowitz VM, Kyrpides NC. The Genomes On Line Database (GOLD) in 2009: status of genomic and metagenomic projects and their associated metadata. *Nucleic Acids Res* 2010; **38**:D346-D354. [PubMed](#) [doi:10.1093/nar/gkp848](https://doi.org/10.1093/nar/gkp848)
20. Mavromatis K, Ivanova N, Anderson I, Lykidis A, Hooper SD, Sun H, Kunin V, Lapidus A, Hugenholtz P, Patel B, Kyrpides NC. Genome analysis of the anaerobic thermohalophilic bacterium *Halothermothrix orenii*. *PLoS ONE* 2009; **4**:e4192. [PubMed](#) [doi:10.1371/journal.pone.0004192](https://doi.org/10.1371/journal.pone.0004192)
21. Field D, Garrity G, Gray T, Morrison N, Selengut J, Sterk P, Tatusova T, Thomson N, Allen MJ, Angiuoli SV, et al. The minimum information about a genome sequence (MIGS) specification. *Nat Biotechnol* 2008; **26**:541-547. [PubMed](#) [doi:10.1038/nbt1360](https://doi.org/10.1038/nbt1360)
22. Garrity G. NamesforLife. BrowserTool takes expertise out of the database and puts it right in the browser. *Microbiol Today* 2010; **37**:9.
23. Woese CR, Kandler O, Wheelis ML. Towards a natural system of organisms: proposal for the domains Archaea, Bacteria, and Eucarya. *Proc Natl Acad Sci USA* 1990; **87**:4576-4579. [PubMed](#) [doi:10.1073/pnas.87.12.4576](https://doi.org/10.1073/pnas.87.12.4576)
24. Garrity GM, Holt JG. The Road Map to the Manual. In: Garrity GM, Boone DR, Castenholz RW (eds), *Bergey's Manual of Systematic Bacteriology*, Second Edition, Volume 1. Springer, New York 2001:119-169.
25. Gibbons NE, Murray RGE. Proposals concerning the higher taxa of *Bacteria*. *Int J Syst Bacteriol* 1978; **28**:1-6. [PubMed](#) [doi:10.1099/00207713-28-1-1](https://doi.org/10.1099/00207713-28-1-1)
26. Validation list 132. List of new names and new combinations previously effectively, but not validly, published. *Int J Syst Evol Microbiol* 2010; **60**:469-472. [PubMed](#) [doi:10.1099/ijs.0.022855-0](https://doi.org/10.1099/ijs.0.022855-0)
27. Rainey FA. Class II. *Clostridia* class nov. In: De Vos P, Garrity G, Jones D, Krieg NR, Ludwig W, Rainey FA, Schleifer KH, Whitman WB (eds), *Bergey's Manual of Systematic Bacteriology*, Second Edition, Volume 3, Springer-Verlag, New York, 2009, p. 736.
28. Validation List No. 55. Validation of the publication of new names and new combinations previously effectively published outside the IJSB. *Int J Syst Bacteriol* 1995; **45**:879-880. [PubMed](#) [doi:10.1099/00207713-45-4-879](https://doi.org/10.1099/00207713-45-4-879)
29. Judicial Commission of the International Committee on Systematics of Prokaryotes. The nomenclatural types of the orders *Acholeplasmatales*, *Halanaerobiales*, *Halobacteriales*, *Methanobacteriales*, *Methanococcales*, *Methanomicrobiales*, *Planctomycetales*, *Prochlorales*, *Sulfolobales*, *Thermococcales*, *Thermoproteales* and *Verrucomicrobiales* are the genera *Acholeplasma*, *Halanaerobium*, *Halobacterium*, *Methanobacterium*, *Methanococcus*, *Methanomicrobium*, *Planctomyces*, *Prochloron*, *Sulfolobus*, *Thermococcus*, *Thermoproteus* and *Verrucomicrobium*, respectively. Opinion 79. *Int J Syst Evol Microbiol* 2005; **55**:517-518. [PubMed](#) [doi:10.1099/ijs.0.63548-0](https://doi.org/10.1099/ijs.0.63548-0)
30. Rainey FA, Zhilina TN, Boulygina ES, Stackebrandt E, Tourova TP, Zavarzin GA. The taxonomic status of the fermentative halophilic anaerobic bacteria: description of *Haloanaerobiales* ord. nov., *Halobacteroidaceae* fam. nov., *Orenia* gen. nov. and further taxonomic rearrangements at the genus and species level. *Anaerobe* 1995; **1**:185-199. [PubMed](#) [doi:10.1006/anae.1995.1018](https://doi.org/10.1006/anae.1995.1018)
31. Oren A, Paster BJ, Woese CR. *Haloanaerobiales*: a new family of moderately halophilic, obligately anaerobic bacteria. *Syst Appl Microbiol* 1984; **5**:71-80.
32. BAuA. Classification of bacteria and archaea in risk groups. *TRBA* 2005; **466**:84.
33. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig JT, et al. Gene Ontology: tool for the unification of biology. *Nat Genet* 2000; **25**:25-29. [PubMed](#) [doi:10.1038/75556](https://doi.org/10.1038/75556)
34. Oren A, Gurevich P, Henis Y. Reduction of nitro-substituted aromatic compounds by the halophilic anaerobic eubacteria *Haloanaerobium praevalens* and *Sporohalobacter marismortui*. *Appl Environ Microbiol* 1991; **57**:3367-3370. [PubMed](#)

35. Oren A, Weisburg WG, Kessel M, Woese CR. *Halobacteroides halobius* gen. nov., sp. nov., a moderately halophilic anaerobic bacterium from the bottom sediments of the dead sea. *Syst Appl Microbiol* 1984; **5**:58-70.
36. Oren A. The ecology and taxonomy of anaerobic halophilic eubacteria. *FEMS Microbiol Rev* 1986; **39**:23-29. [doi:10.1111/j.1574-6968.1986.tb01838.x](https://doi.org/10.1111/j.1574-6968.1986.tb01838.x)
37. Oren A. 2006. Life at high salt concentrations. In: M Dworkin, S Falkow, E Rosenberg, KH Schleifer E Stackebrandt (eds), *The Prokaryotes*, 3. ed, vol. 2. Springer, New York, p. 263-282.
38. Matheson AT, Louie KA, Tak BD, Zuker M. The primary structure of the ribosomal A-protein (L12) from the halophilic eubacterium *Haloanaerobium praevalens*. *Biochimie* 1987; **69**:1013-1020. [PubMed doi:10.1016/0300-9084\(87\)90001-0](https://pubmed.ncbi.nlm.nih.gov/1016/0300-9084(87)90001-0/)
39. Klenk HP, Göker M. En route to a genome-based classification of *Archaea* and *Bacteria*? *Syst Appl Microbiol* 2010; **33**:175-182. [PubMed doi:10.1016/j.syapm.2010.03.003](https://pubmed.ncbi.nlm.nih.gov/10.1016/j.syapm.2010.03.003/)
40. Wu D, Hugenholtz P, Mavromatis K, Pukall R, Dalin E, Ivanova NN, Kunin V, Goodwin L, Wu M, Tindall BJ, et al. A phylogeny-driven genomic encyclopaedia of *Bacteria* and *Archaea*. *Nature* 2009; **462**:1056-1060. [PubMed doi:10.1038/nature08656](https://pubmed.ncbi.nlm.nih.gov/10.1038/nature08656/)
41. List of growth media used at DSMZ: http://www.dsmz.de/microorganisms/media_list.php.
42. Gemeinholzer B, Dröge G, Zetzsche H, Haszprunar G, Klenk HP, Güntsch A, Berendsohn WG, Wägele JW. The DNA Bank Network: the start from a German initiative. *Biopreservation and Biobanking* 2011; **9**:51-55. [doi:10.1089/bio.2010.0029](https://doi.org/10.1089/bio.2010.0029)
43. JGI website. <http://www.jgi.doe.gov>
44. The Phred/Phrap/Consed software package. <http://www.phrap.com>
45. Zerbino DR, Birney E. Velvet: algorithms for de novo short read assembly using de Bruijn graphs. *Genome Res* 2008; **18**:821-829. [PubMed doi:10.1101/gr.074492.107](https://pubmed.ncbi.nlm.nih.gov/10.1101/gr.074492.107/)
46. Han C, Chain P. Finishing repeat regions automatically with Dupfinisher. In: Proceeding of the 2006 international conference on bioinformatics & computational biology. Arabnia HR, Valafar H (eds), CSREA Press. June 26-29, 2006: 141-146.
47. Lapidus A, LaButti K, Foster B, Lowry S, Trong S, Goltsman E. POLISHER: An effective tool for using ultra short reads in microbial genome assembly and finishing. AGBT, Marco Island, FL, 2008
48. Hyatt D, Chen GL, LoCascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics* 2010; **11**:119. [PubMed doi:10.1186/1471-2105-11-119](https://pubmed.ncbi.nlm.nih.gov/10.1186/1471-2105-11-119/)
49. Pati A, Ivanova NN, Mikhailova N, Ovchinnikova G, Hooper SD, Lykidis A, Kyrpides NC. Gene-PRIMP: a gene prediction improvement pipeline for prokaryotic genomes. *Nat Methods* 2010; **7**:455-457. [PubMed doi:10.1038/nmeth.1457](https://pubmed.ncbi.nlm.nih.gov/10.1038/nmeth.1457/)
50. Markowitz VM, Ivanova NN, Chen IMA, Chu K, Kyrpides NC. IMG ER: a system for microbial genome annotation expert review and curation. *Bioinformatics* 2009; **25**:2271-2278. [PubMed doi:10.1093/bioinformatics/btp393](https://pubmed.ncbi.nlm.nih.gov/10.1093/bioinformatics/btp393/)