

Lawrence Berkeley National Laboratory

LBL Publications

Title

Polarization of High-Energy Deuterons

Permalink

<https://escholarship.org/uc/item/83r8128p>

Authors

Chamberlain, Owen

Segrè, Emilio

Tripp, Robert

et al.

Publication Date

1954-06-01

UNIVERSITY OF
CALIFORNIA

*Radiation
Laboratory*

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

BERKELEY, CALIFORNIA

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Cy. 2

UCRL-2622

UNIVERSITY OF CALIFORNIA
Radiation Laboratory

Contract No. W-7405-eng-48

POLARIZATION OF HIGH-ENERGY DEUTERONS

Owen Chamberlain, Emilio Segrè, Robert Tripp, Clyde Wiegand,
and Thomas Ypsilantis

June 16, 1954

Berkeley, California

POLARIZATION OF HIGH-ENERGY DEUTERONS

Owen Chamberlain, Emilio Segrè, Robert Tripp, Clyde Wiegand,
and Thomas Ypsilantis

Radiation Laboratory, Department of Physics,
University of California, Berkeley, California

June 16, 1954

In analogy to experiments on proton polarization¹⁻⁴ we obtained a polarized deuteron beam by the same scattering method. The trajectories of 167-Mev deuterons as produced by the cyclotron are essentially the same as those of 312-Mev protons if the magnetic field of the steering magnet is properly adjusted and the deuteron beam enters the cave through the same channel as the proton beam.

In order to polarize and analyze the deuteron beam we have used carbon as scatterers A and B. After the second scatterer we have measured the intensity of the scattered beam as a function of ϕ . We have endeavored to limit ourselves to elastically scattered deuterons by the use of absorbers, as in the proton work.

It has been shown by Lakin and Wolfenstein⁵ that the most general intensity distribution produced by polarized deuterons is:

$$\begin{aligned}
 I(\theta, \phi) = & I_0(\theta) + A(\theta) \langle T_{20} \rangle + \\
 & \left(B(\theta) \langle T_{21} \rangle + C(\theta) \left| \langle T_{11} \rangle \right| \right) \sin \theta \cos \phi + \\
 & D(\theta) \langle T_{22} \rangle \sin^2 \theta \cos 2\phi. \quad (1)
 \end{aligned}$$

The incident deuterons travel in the Z direction and have been polarized by scattering in the x, y plane. I, A, B, C, D are polynomials in $\cos \theta$ and the T_{ik} 's are expectation values, for the beam before it undergoes the second scattering, of quantities such as

$$T_{21} = -\frac{\sqrt{3}}{2} \left[(S_x + iS_y) S_z + S_z (S_x + iS_y) \right]$$

$$T_{11} = -\frac{\sqrt{3}}{2} (S_x + iS_y)$$

$$T_{22} = \frac{\sqrt{3}}{2} (S_x + iS_y)^2$$

where \vec{S} is the spin operator for the deuteron.

Experimentally we have found $I(20^\circ, \phi)$ as in Fig. 1. An analysis of this scattering gives

$$I(20^\circ, \phi) = p + q \cos \phi + r \cos 2\phi \quad (2)$$

with $p = 50.3 \pm 2.2$, $q = 15.3 \pm 1.9$, and $r = -1.8 \pm 3.6$ in units of $10^{-27} \text{ cm}^2 / \text{steradian}$. The errors are based on counting statistics only, and are valid for the ratios $p:q:r$ but not necessarily for the absolute values of these parameters. Measurements at smaller values of θ give similar results with smaller values of q/p .

This equation agrees well with Eq. (1) and shows that the coefficient of $\cos 2\phi$ is very small or zero, at least in the cases investigated up to now.

A possible explanation which is consistent with the usual spin-orbit model has been pointed out by Dr. M. Ruderman. If it is assumed that the deuteron polarization originates in an $\vec{L} \cdot \vec{S}$ coupling which acts in the scattering process in addition to a central force, and if the magnitude of the spin-orbit potential is small compared to the central part, it then follows that the $\cos 2\phi$ term is negligible because $\langle T_{22} \rangle$ is very small.

It must be noted that the description of the polarization of a deuteron beam is substantially more complicated than the similar description for particles of spin $1/2$, and that our analysis gives only a small part of the relevant information.

¹ Oxley, Cartwright, and Rouvina, Phys. Rev. 93, 806 (1954).

² Chamberlain, Segrè, Tripp, Wiegand, and Ypsilantis, Phys. Rev. 93, 1430 (1954). Fig. 1 of the reference paper is applicable to the present work and the same notations are used throughout.

³ Marshall, Marshall, and de Carvalho, Phys. Rev. 93, 1431 (1954).

⁴ Dickson, and Salter, Nature 173, 946 (1954).

⁵ W. Lakin and L. Wolfenstein - Unpublished report. We thank the authors for having shown us the manuscript.

This work was done under the auspices of the U. S. Atomic Energy Commission.

Figure 1. Differential cross section versus azimuthal angle Φ for polarized deuterons scattered elastically by carbon.

$d\sigma/d\Omega$ IN 10^{-27} CM²/STERADIAN

