

# UC San Diego

## UC San Diego Previously Published Works

### Title

Greater change in arbuscular mycorrhizal fungal richness as a response to short-term rainfall exclusion across the North American monsoon season

### Permalink

<https://escholarship.org/uc/item/844168t4>

### Journal

Elementa: Science of the Anthropocene, 12(1)

### ISSN

2785-4558

### Authors

Dickey, Jonathan R

Fordyce, James A

### Publication Date

2024-02-22

### DOI

10.1525/elementa.2022.00116

Peer reviewed

## RESEARCH ARTICLE

# Greater change in arbuscular mycorrhizal fungal richness as a response to short-term rainfall exclusion across the North American monsoon season

Jonathan R. Dickey<sup>1,2,\*</sup>  and James A. Fordyce<sup>1</sup>

Arid ecosystems around the world are projected to experience reduced and more infrequent precipitation events. The effects of reduced rainfall are well understood for plants and aboveground interactions; however, the effect of rainfall on belowground microbial interactions remains understudied. Here, we illustrate the strength and direction of change that short-term manipulative rainfall exclusion treatments have on the composition and relative abundance of Arbuscular Mycorrhizal (AM) fungal communities throughout the monsoon season in the Sonoran Desert. Additionally, we use these rainfall exclusion treatments to further understand the impact of soil moisture on labile forms of nitrogen and phosphorous in a natural environment. Rainfall exclusion treatments were installed to remove 0% (control), 60% (moderate exclusion), and 80% (high exclusion) of rainfall over a single wet season. AM fungal community composition varied among rainfall treatments and across the monsoon season, where change was dependent on initial conditions within treatments. Phosphorous content was also a strong predictor of AM fungal composition and relative abundance. By the end of the season, the difference in AM fungal richness was influenced by rainfall exclusion, with the greatest positive change in richness in the moderate rainfall exclusion treatments. Lastly, AM fungal community turnover was predicted by rainfall exclusion treatments, where rainfall contributed to greater turnover in the control treatment in comparison to other treatments. This study further illustrates the complex association between soil abiotic factors, how they are influenced by environmental stress, and how in turn cause shifts in AM fungal communities.

**Keywords:** Arbuscular mycorrhizal fungi, Rainfall exclusion, Sonoran Desert, Community ecology, 18S rRNA amplicon sequencing

## Introduction

As climate change alters seasonal weather patterns, interactions among organisms can become decoupled through range shifts or changing phenology (Van Der Putten et al., 2010). Arid regions around the world are predicted to experience stark reductions in seasonal precipitation while global temperatures begin to rise (Diffenbaugh and Giorgi, 2012). The North American Monsoon (NAM) season, present in Southeastern Arizona and Western New Mexico, provides roughly 40% of the regions mean annual precipitation (Mitchell et al., 2002). This seasonal effect has been shifting spatiotemporally and becoming difficult to predict under future climate conditions (Comrie and Glenn, 1999; Demaria et al., 2019). Climate models have

illustrated the importance seasonal precipitation and temperature have on plants, animals, and, thus, the interactions among them (Bradie and Leung, 2017). Plants also have numerous belowground interactions with microorganisms, such as bacteria and fungi, whereby mutualistic associations provide vital access to nutrients, water, as well as resistance to pathogens (Newsham et al., 1995). For example, these associations promote plant productivity, affect plant defense strategies, and influence overall health (Van Der Heijden et al., 2008; Lau and Lennon, 2011). Yet, there remains a need to understand how changing seasonal precipitation regimes in dry, arid climates might affect plant–fungal interactions.

Arbuscular mycorrhizal (AM) fungi are a clade of plant host-associated microorganisms (*Glomeromycotina*) that form associations with over 70% of terrestrial plant species within the rhizosphere microhabitat (Soudzilovskaia et al., 2020). AM fungi are highly abundant in soils and have a wide distribution globally, extending from tropical forests to grasslands (Hayman, 1982). The communities of AM fungi are the key components of soil nutrient cycles,

<sup>1</sup>Department of Ecology and Evolutionary Biology, The University of Tennessee, Knoxville, TN, USA

<sup>2</sup>Current Address: School of Biological Sciences, University of California San Diego, La Jolla, CA, USA

\* Corresponding author:  
Email: [jrdickey@ucsd.edu](mailto:jrdickey@ucsd.edu)

exchanging phosphorous, nitrogen, and other trace minerals with plant-derived carbohydrates and lipids (Allen, 1991). At a regional scale, AM fungal communities and the type of symbiosis (e.g., parasitism, commensalism, mutualism) are often more strongly predicted by environmental factors than plant hosts or dispersal limitation (Johnson et al., 2010; Kivlin et al., 2014). These environmental factors could be one or more of the following: soil fertility (soil organic matter content; Anderson et al., 1984), soil moisture (Miller, 2000; Gao et al., 2016), soil pH (Green et al., 1976), soil texture (e.g., silt, sand, etc.; Johnson et al., 1992), and soil nutrients (Johnson et al., 2010). The changes in soil moisture or H<sub>2</sub>O saturation can also influence the labile nature of soil abiotic factors of the rhizosphere (Deepika and Kothamasi, 2015; Kanwal et al., 2015; Bobille et al., 2019). Therefore, it is important to investigate the relative importance perturbations in soil moisture have on AM fungal communities through both direct and indirect pathways.

Environmental stress can also have large direct consequences on AM fungal community composition and relative abundance (Hernandez et al., 2021). Stress can range from increased seasonal temperatures (Zhou et al., 2016) to changes in precipitation regimes (Waring and Hawkes, 2015). AM fungi have been shown to exist in a wider range of annual precipitation than other mycorrhizal fungi (Tedersoo et al., 2014), and thus, AM fungi have been regarded as key regulators of plant stress brought on by drought (Birhane et al., 2012). For example, in greenhouse experiments, seasonal drought was induced through water pulsing, where irregular water regimes have a positive effect on AM fungal colonization and persistence in comparison to treatments with a continuous supply of water (Birhane et al., 2012). AM fungi confer drought tolerance in plants by increasing osmotic adjustment and regulating stomatal conductance, reducing transpiration and leading to enhanced plant growth and vigor (Wu, 2017). In natural environments, AM fungal communities may be predisposed to tolerate drought conditions based on prior rainfall history (Kuehn et al., 1991). Deveautour et al. (2020) concluded that under reduced rainfall, estimated AM fungal alpha diversity is larger in comparison to the control treatment (ambient rainfall). On the other hand, plants that experienced complete rainfall removal, or higher environmental stress, showed lower alpha diversity estimates than the control and the reduced rainfall treatment. Thus, there lacks a consensus regarding the direction and magnitude of change in composition as soil moisture varies among ecosystem types and temporal scales.


In this study, we examine the relative importance of manipulative rainfall exclusion, local soil abiotic factors, and sampling date have on rhizospheric AM fungal communities across the NAM season of 2019. We first ask, what is the strength and direction of change in AM fungal communities (i.e., composition, richness, and turnover) in response to the manipulation of seasonal rainfall in the Sonoran Desert? And secondly, what is the relative influence soil abiotic factors and sampling date have on these communities? To answer these questions, experimental rainfall exclusion treatments were set up over the course

of the monsoon season (June 2019–August 2019) to remove 0% (ambient rainfall/control), 60% (moderate exclusion), and 80% (high exclusion) of seasonal rainfall. Soil abiotic factors and AM fungal communities were sampled at the beginning and end of the monsoon season. Due to the complex ecological communities found in the rhizosphere microhabitat, there are 2 predictive hypotheses that were derived from our questions. First, with reduced rainfall, AM fungal communities could show no observed net gain or loss in taxa across the monsoon season in response to rainfall exclusion, as AM fungi have been noted to have higher desiccation tolerance than other fungal clades, such as ectomycorrhizal fungi (Allen, 1991). Alternatively, AM fungal communities could differ in turnover among rainfall exclusion treatments with ambient seasonal rainfall contributing to greater community turnover than rhizosphere communities that undergo rainfall exclusion throughout the NAM season. This could be a result of the plant-host growing season contributing to AM fungal community succession (Santos-González et al., 2007; Dumbrell et al., 2011). Understanding the response of plant-host associated AM fungal communities to drier microhabitats could shed light on how AM fungi might continue to associate with hosts in more extreme, future climate conditions.

## Materials and methods

### Experimental design

Monsoon rainfall occurs in the Sonoran Desert ecoregion of the United States around late June–mid August. This rainfall often influences how organisms interact and can help maintain population-level persistence throughout the remaining portion of the year. This season can function as a natural experiment to manipulate seasonal rainfall intensity on a stable plant population and their interactions with arbuscular mycorrhizas. Monsoon rainfall exclusion shelters were established following guidelines outlined in Beier et al. (2012). Three treatments were established at the beginning of June 2019, a control treatment that received no rainfall exclusion ( $n = 22$ ), moderate rainfall exclusion ( $n = 22$ ), and high rainfall exclusion ( $n = 22$ ). Short-term rainfall exclusion shelters were constructed of 6-mil heavy-duty clear plastic sheeting with 60% (moderate exclusion), or 80% (high exclusion), or 100% (control treatment) of a 1-m<sup>2</sup> surface area removed (**Figure 1**). Rainfall exclusion shelters were supported by two 30-cm and two 64-cm wooden landscape stakes and four 1-m wooden dowels along each edge. This design raised one edge of the shelter higher than the other providing airflow and to allow rainfall water to run off plot to ensure soil moisture manipulation. Each sampled plant rhizosphere was centered in the middle of each exclusion shelter. *Lupinus neomexicanus* Greene and herbaceous legume were chosen as an experimental system because of an elongated flowering time allowing for consistent identification throughout the monsoon season. This allowed us to ensure that all plants sampled were within the same life stage and had similar aboveground biomass. This species is endemic to midrange elevations (1,600–2,000 m) in Southern Arizona (e.g., 31°53'50.3"N 109°16'38.5"W).



**Figure 1. Rainfall exclusion treatment design.** Rainfall exclusion treatment designs to manipulate monsoon rainfall on arbuscular mycorrhizal fungal communities of the rhizosphere with 60% (moderate) exclusion, 80% (high) exclusion, and no shelter (control treatment).

#### **Sampling soil abiotic factors**

The rhizosphere soil microhabitat was sampled twice, once prior to the monsoon season (Date A; June 2019) and once at the tail end of the monsoon season (Date B; August 2019). Baseline estimates for AM fungal community structure, soil moisture, and other soil abiotic factors are from the first sampling period prior to monsoon shelter establishment. Soil was dried in a laboratory oven for 48 h at 54°C. Soil moisture content was measured as the difference in soil weight between field wet soil and oven dried soil divided by the weight of oven dried soil (Standards Association of Australia, 1977). By the end of the monsoon season, the high rainfall exclusion treatment was on average  $1.66 \pm 0.29$  g water/g soil drier than the moderate rainfall exclusion treatment and  $2.64 \pm 0.17$  g water/g soil drier than the control treatment. To estimate soil abiotic factors (e.g., ammonium, nitrate, and phosphorous) in the rhizosphere environment, all rhizosphere samples were subsampled twice (2 g of rhizosphere soil per subsample) and extracted with potassium sulfate and then analyzed in triplicate ( $K_2SO_4$ ; Brookes et al., 1985; Vance et al., 1987; Wu et al., 1990). Since AM fungi play a pivotal role in nitrogen cycling, nitrogen was measured in 2 ways (ammonium and nitrate). Ammonium was measured using the indophenol-blue Berthelot method (Weatherburn, 1967; Verdouw et al., 1978), and a vanadium (III) chloride acid solution and Griess reagents (Mulvaney, 1996) were used to extract nitrate from the  $K_2SO_4$  samples. Phosphorous content for each rhizosphere sample was measured by a reaction between the  $K_2SO_4$  extractions, ammonium paramolybdate (AMP), concentrated sulfuric acid, and a malachite green solution (Lajtha et al., 1999).

#### **Sampling AM fungal communities**

AM fungal communities were sampled using sterilized, handheld soil cores and stored in Nasco Whirl-Paks (Nasco, Fort Atkinson, WI, USA). All field equipment was surface sterilized with 70% ethanol and soil particulates were removed in-between sampling. To collect AM fungal communities, the sterilized soil core was pushed approximately 5–7 cm into the rhizosphere microhabitat along the root base, where the plant stem meets the soil. All rhizosphere samples were stored on dry ice until temporary storage ( $-80^\circ\text{C}$ ) at the Southwest Research Station in Portal, AZ, USA. Samples were then shipped on dry ice to the University of Tennessee, Knoxville, and immediately stored at  $-80^\circ\text{C}$  upon arrival. Prior to DNA isolation, all rhizosphere soil samples were thawed for 1 h at room temperature. DNA isolation proceeded with 250 mg of rhizosphere soil per sample and the DNeasy PowerSoil HTP 96 Kit (MoBio Laboratories, Inc., Carlsbad, CA, USA). The final elution volume was modified, so that 50  $\mu\text{L}$  of Solution C6 was used. Additionally, the solutions sat on the DNA membrane for 5 min at room temperature prior to centrifuging the concentrated DNA down into a sterile 96 well collection plate.

AM fungal specific libraries of the 18S rRNA gene were amplified using a nested PCR protocol (Kivlin et al., 2021). The first reaction utilized the primer pair NS1/NS4, which amplified approximately a 1,200 bp product (White et al., 1990; Raja et al., 2017). The forward NS1 primer sequence was 5'-GTAGTCATATGCTTGCTC-3' and the reverse NS4 primer sequence was 5'-CTTCCGCAATTCCTTAAG-3'. Each reaction used 2.5  $\mu\text{L}$  of the DNA template, 5  $\mu\text{L}$  of both the forward and reverse primers, and 12.5  $\mu\text{L}$  of KAPA HiFi HotStart Ready Mix (KAPA Biosystems, Wilmington, MA, USA). This reaction ran at  $95^\circ\text{C}$  for 3 min,

followed by 30 cycles of 95°C for 30s, 40°C for 1 min, and 72°C for 1 min with a final elongation step of 72°C for 10 min. These reactions produced 25 µL of PCR product. The following reaction amplified approximately 550 bp with the mixed primer pair, NS31 and AML2, which contain the sequences 5'-TTGGAGGGCAAGTCTGGTGCC-3' and 5'-GAACCCAAACACTTTGGTTTC-3', respectively (Simon et al., 1992; Lee et al., 2008). These reactions required 2.5 µL of product from the first reactions, 5 µL of NS31 and AML2, and 12.5 µL of KAPA HiFi HotStart Ready Mix. They began with 95°C for 5 min followed by 40 cycles of 95°C for 45 s, 63.1°C for 1 min, and 72°C for 1.5 min with a final elongation step of 72°C for 10 min. This nested PCR produced 25 µL of AM fungal specific 18S rRNA gene product per sample. All 18S PCR products were then stored at -20°C until further use.

PCR products were combined with 20 µL of Agencourt AMPure XP magnetic beads, placed on a plate magnet, and rinsed twice with 200 µL of 80% ethanol for product purification. The remaining ethanol was allowed to evaporate before purified product was resuspended in 50 µL of Tris-HCl. Purified products were stored at -20°C until 18S rRNA communities per sample were multiplexed. Each sample was multiplexed with a forward and reverse index primer, which are specifically designed for cluster generation on an Illumina MiSeq flow cell (Illumina Corporation, San Diego, CA, USA). These primers were obtained from the i5/i7 Nextera XT Index Kit. For the indexing PCR, 5 µL of purified PCR product from the previous reaction is required, along with 5 µL of the uniquely combined forward and reverse Nextera index primers, 25 µL of KAPA HiFi HotStart Ready Mix, and 10 µL of sterile nuclease free water. These reactions ran for an initial cycle of 95°C for 3 min followed by an 8-cycle sequence of 95°C for 3 s, 55°C for 30 s, and 72°C for 30 s and held at 72°C for 5 min. These reactions yielded 50 µL of product that was purified with 56 µL of Agencourt AMPure XP magnetic beads, and 2 washes of 220 µL freshly prepared 80% ethanol. Cleaned and purified product was then resuspended in 25 µL of Tris-HCl. All DNA libraries were then pooled by equal concentrations. The pooled solution was diluted to 4 pM with Tris and loaded onto a V3 flow cell with a 20% PhiX spike. This flow cell was run on an Illumina MiSeq instrument at the University of Tennessee Genomics Core, Knoxville, TN, USA, set to a 2 × 300 paired end cycle.

### Bioinformatics

Demultiplexing was completed by the Illumina MiSeq instrument prior to all other processing. 18S rRNA gene amplicon sequencing data were processed with the open-source DADA2 pipeline version 1.8 (<https://github.com/benjjneb/dada2>) and R version 3.5.2. Amplification primers for reads corresponding to 18S rRNA genes were trimmed at the 5' end according to the length of the primer (NS31: 21 and AML2: 21). For 18S, all forward and reverse reads were truncated at 275 and 270 base pairs and filtered to remove reads below  $Q = 30$ . Error rates for the forward and reverse reads were generated by performing sample inference and error estimations repeatedly until congruence was met. Next, the derepFastq function

was used in the DADA2 package to dereplicate amplicons for every sample. Here, sequences with 99% similarity were grouped, and creating unique amplicon sequence variants (ASVs) and read abundance was totaled (Callahan et al., 2016). Forward and reverse sequence reads were concatenated since the amplicon length for the AM fungal specific target region within the 18S rRNA gene is too long, such that sufficient overlap in reads was negligible between pairs. Chimeras were detected across all samples and were tossed from the data set using the removeBimeraDenovo function. Across all rhizosphere samples, fungal ASVs with less than 12 reads were filtered out of the data set.

The decostand function in the vegan package v.2.5-6 (Oksanen et al., 2019) was used to proportionally standardize all AM fungal ASVs within sample (method = "total"). Next, all AM fungal ASVs were matched against the MaarjAM database, an AM fungal sequence repository for the NS31/AML2 primer pair, using BLAST, the Basic Local Alignment Search Tool (Öpik et al., 2010). The program Practical Alignment using Saté and TrAnsitivity (Mirarab et al., 2015) was used to conduct multiple sequence alignment with MAFFT software (Kato and Standley, 2013). Then, the aligned FASTA file was used to infer a maximum likelihood phylogenetic tree using FastTree-2 (Price et al., 2010). FastTree-2 created a generalized time-reversible CAT model of rate heterogeneity to estimate 18S rRNA phylogeny. Lastly, the treePL program was utilized to correct for differences in divergence time among branch lengths using a penalized likelihood approach (Smith and O'Meara, 2012). To understand how sensitive our inferences are to differences in sampling effort, we rarefied to the lowest number of reads among samples (1,768) for 1,000 iterations, creating 1,000 rarefied AM fungal communities per sample.

### Statistical analyses

To address the question, to what extent does AM fungal community composition respond to monsoon rainfall exclusion, sampling date, and soil abiotic factors, a distance-based redundancy analysis (dbRDA) was utilized (R package vegan, Oksanen et al., 2019). The quantitative Jaccard distance (also known as the Ružička distance) was used to test community dissimilarity in composition among rainfall exclusion treatments, sampling date, and soil abiotic factors. A distance matrix was calculated for each of the 1,000 rarefied communities and a mean distance matrix was calculated for incorporation into model selection. Forward stepwise model selection on the dbRDA analysis was completed using the function ordiR2-step in the vegan package (permutations = 1,000). The full scope incorporated into model selection had the following variables as main effects: rainfall exclusion treatment, sampling date, log of phosphorous, log of ammonium, and the square root of nitrate. Model selection on the dbRDA using the mean quantitative Jaccard distance from 1,000 rarefied communities determined rainfall exclusion treatment and log of phosphorous as main effects. For hypothesis testing, an analysis of variation (permutations = 9,999) was conducted on this dbRDA. A constrained


ordination was made from this model using the 2 most predictive axes and their associated species scores.

Hill numbers were used to examine the effective number of AM fungal ASVs ( ${}^qD$ ) associated with each treatment and date (Hill, 1973). The effective number of AM fungal ASVs ( $D$ ) is sensitive to the order of diversity ( $q$ ), and the importance of rarer ASVs is reduced when calculating diversity as the value of  $q$  increases. For instance,  $q = 0$  represents species richness, where rare and common ASVs are weighted equally. At  $q = 1$  (exponential Shannon's entropy), ASVs are weighted by their proportional abundance. Lastly,  $q = 2$  is the inverse of the Simpson's index, where rare ASVs are downweighted when calculating the effective number of AM fungal ASVs.

We fitted Bayesian linear models to assess  $\Delta$  AM fungal richness and community turnover associated with rainfall exclusion treatment at  $q = 0, 1,$  and  $2$ . This was accomplished using the brms package (Bürkner, 2017), which calls the probabilistic programming language Stan (Carpenter et al., 2017).  $\Delta$  AM fungal richness was calculated as the difference in the effective number of AM fungal ASVs across the monsoon season (Date A – Date B) for each rhizosphere sample. Community turnover was calculated as  $1 -$ the pairwise beta diversity for each rhizosphere, thus providing the proportional change in community composition between Date A and Date B. Each model used 4

Markov chain Monte Carlo (MCMC) chains, each with 2,000 iterations with a starting warmup period made up of 1,000 iterations and 1,000 post-warm up draws. The default priors provided by the brms package were used for all models. Chain convergence was confirmed using the Gelman–Rubin convergence statistic ( $R$  hat = 1 for all models). Effective sample size for all models was  $>3,000$ . Inferences were based on examining the 95% highest density interval of model parameters calculated using the hdi function in the HDInterval package (v. 0.2.2) (Meredith and Kruschke, 2020). Hypotheses regarding differences in  $\Delta$  AM fungal richness and community turnover among treatments were examined using the posterior joint probability distribution for the parameters using the hypothesis() function in brms, which provides the proportion of times a parameter is either greater than, or less than, another parameter across MCMC steps, in this case zero where zero represents no change in  $\Delta$  AM fungal richness and AM fungal community turnover.  $\Delta$  AM fungal richness and AM fungal community turnover were further analyzed using standardized abundance ASV data and rarefied data to confirm that Bayesian inference was robust against the differences in reads among samples.

To examine the phylogenetic community structure among rainfall exclusion treatments, we used a phylogenetically informed Hill number approach at  $q = 1$  for



**Figure 2. Heat map of arbuscular mycorrhizal (AM) fungi recovered through sequencing.** Heat map tree illustrating the AM fungal variants recovered from Illumina MiSeq run from this study. A number of AM fungal variants are represented here as presence/absence, where darker pink colors and larger circles show higher abundance.

alpha diversity, which is analogous to a phylo-Horn N-assembly overlap measure (Chao et al., 2010). This measure considers phylogenetic relationships by calculating the effective number of AM fungal ASVs (richness) given relative abundance and a phylogenetic tree. At  $q = 1$ , we used the `hill_div` function in the `hilldiv` package v.1.5.1 for proportional abundance data and all rarefied community data (Alberdi and Gilbert, 2019). We calculated the phylogenetically informed  $\Delta$  AM fungal richness across the monsoon season for each sample and modeled it as a fitted Bayesian linear model as described above. The same one-sided hypothesis was used to test significant patterns in phylogenetically informed  $\Delta$  AM fungal richness among treatments.

**Results**


**Summary**

After running the DADA2 customized pipeline as described above, a total of 34,011 ASVs were generated. Based on the best practices for sequencing and analyzing AM fungal communities, ASVs with less than 10–12 reads (for this study <12 was chosen as the threshold) were omitted from the data set. This resulted in a total of 5,012 ASVs that correspond to the 18S rRNA gene region. After ASVs were confirmed to be AM fungi in the MaarjAM database, 5,011 ASVs remained in the final data set. One ASV did not match any taxa from the MaarjAM database and was removed. All other reads were assigned to the subphylum *Glomeromycotina* (Soudzilovskaia et al.,

2020). **Figure 2** illustrates the relative abundance in reads assigned to the subphylum *Glomeromycotina* and the distribution of these reads among orders and families within this subphylum. AM fungal communities from Date A, prior to the onset of rainfall exclusion treatment, were utilized as a baseline understanding of AM fungal diversity and relative abundance present in the *L. neomexicanus* rhizosphere. Of the 4 orders within *Glomeromycotina*; *Glomerales*, *Diversisporales*, *Paraglomerales*, and *Archaeosporales* represented 80%, 10%, 6%, and 4% of reads recovered. The genus *Glomus* was the most abundant genus recovered from the rhizosphere microhabitat, representing approximately 67% of all total reads at Date A. Further, the second most abundant genus recovered was *Claroideoglomus* with approximately 12% of total reads sampled at Date A. Lastly, the third most abundant genus was *Paraglomus* (*Paraglomerales*) comprising approximately 7% of reads in *Glomeromycotina*. The most abundant AM fungal variant recovered was *Glomus Valera-Cervero15 BG21* representing 14% of all total reads recovered at Date A in this study.

**AM fungal community composition is influenced by rainfall exclusion**

AM fungal community composition was characterized by assigning taxonomy and documenting change in relative read abundance among treatments before and after the monsoon season (**Figure 3**). Across all treatments, *Diversisporaceae* decreased in relative abundance across the




**Figure 3. Relative abundance changes among rainfall exclusion treatments.** Stacked bar chart for the orders *Archaeosporales*, *Diversisporales*, *Paraglomerales*, and *Glomerales*. Here, each treatment is broken up by Date (A or B) to show changes within arbuscular mycorrhizal fungal families across the monsoon season. Here, 0% represents the control treatment, 60% the moderate rainfall exclusion treatment, and 80% the high rainfall exclusion treatment.

monsoon season with the largest observed difference in the high rainfall exclusion treatment. On the other hand, **Figure 3** illustrates an increase in *Paraglomeraceae* in response to monsoon rainfall, with the most extreme observed change in the control treatment. *Glomeraceae* was observed to be the most common AM fungal family recovered, and relative abundance decreased over time. *Claroideoglomerales* decreased in relative abundance among all treatments, but most notably in the control treatment.

We used the mean quantitative Jaccard distance matrix from 1,000 rarefactions to test the strength of rainfall exclusion and the log of phosphorous on AM fungal community composition and relative abundance. This model's constrained variables explained 3.55% of total variation in multivariate space. Additionally, axes 1 and 2 explained 47.14% and 34.04% of the 3.55% of total variation, respectively. **Figure 4** shows the relative change in direction and strength of AM fungal communities given rainfall

exclusion treatments relative to estimates in composition before treatments were installed. Points here represent the median centroid in multivariate space with  $\pm 1$  standard error for each treatment and sampling date. A permutational analysis of variance showed that rainfall exclusion treatments were a significant predictor for AM fungal community structure and composition ( $P < 0.05$ ). Based on this model, the log of phosphorous was also a significant predictor of AM fungal community composition in multivariate space ( $P < 0.001$ ), contributing to the observed changes in **Figure 4** from community estimates before treatments were installed. All measured soil abiotic factors were shown to not be confounded or interact with rainfall exclusion treatments (Supplemental File 1).

**Richness and community turnover from  $q = 0$  to  $q = 2$**   
The differences in alpha diversity between Date A and Date B, or  $\Delta$  AM fungal richness, were analyzed from diversity




**Figure 4. Principal coordinate analysis for arbuscular mycorrhizal (AM) fungal composition among rainfall treatments.** Principal coordinate analysis for AM fungal community composition from the dbRDA using the mean quantitative Jaccard dissimilarity matrix of 1,000 rarefied communities. Axes 1 and 2 explain 47.14% and 34.04% of the 3.55% of total variation explained by this model. Colors represent the median centroid of the control treatment (black), the moderate rainfall exclusion treatment (teal), and the high rainfall exclusion treatment (pink). Date A and Date B are represented by triangles and circles, respectively. Bars are  $\pm 1$  standard error.



order  $q = 0-2$  using linear models. From  $q = 0$  to  $q = 2$ , the greatest observed gain in the effective number of AM fungal ASVs was in the moderate rainfall exclusion treatment (Figure 5). On the contrary, the control treatment on average lost AM fungal ASVs throughout the monsoon season. As  $q$  increases, the 95% highest density intervals decrease around the median change in AM fungal richness across the monsoon season. The control treatment was observed to have an estimated median change of  $-15.13$ ,  $-4.85$ , and  $-2.96$  in AM fungal richness across the monsoon season for  $q = 0, 1, \text{ and } 2$ , respectively. We can infer

that rainfall had a negative effect on  $\Delta$  AM fungal richness of rhizospheres. This hypothesis was observed to be true in our posterior samples 77%, 89%, and 96% of the time for  $q = 0, 1, \text{ and } 2$  (Table 1). The median estimated  $\Delta$  AM fungal richness for the moderate rainfall exclusion treatment was 4.68, 6.10, and 3.10 at  $q = 0, 1, \text{ and } 2$ . For the moderate rainfall exclusion treatment, we can infer positive change in richness throughout the monsoon season for each value of  $q$ . For  $q = 0, 1, \text{ and } 2$ , this hypothesis was observed 76%, 97%, and 99% of the time of the posterior samples. In the high rainfall exclusion treatment, we can



**Figure 5. Delta richness of arbuscular mycorrhizal (AM) fungi across the monsoon season by rainfall exclusion treatment.**  $\Delta$  AM fungal richness and the 95% highest density interval around the median change in alpha diversity between the 2 time points. Bayesian inference was determined using linear models, where the difference in AM fungal alpha diversity was predicted by rainfall exclusion treatment. This was conducted for  $q = 0, 1, \text{ and } 2$ . Black represents the control treatment, teal the moderate (60%) rainfall exclusion treatment, and pink the high (80%) rainfall exclusion treatment. Percentages are the proportion of samples from the joint probability distribution, where parameters are greater or less than zero (zero indicating no change in richness). The percentage for control describes the change in diversity between time A and B, and the percentages for the moderate and high groups describe the change in diversity relative to the control group. Across all values of  $q$ , the moderate rainfall exclusion treatment, on average, had the largest change in AM fungal richness compared to other treatments.

**Table 1. Hypothesis testing on  $\Delta$  arbuscular mycorrhizal fungal richness for each rainfall exclusion treatment and value of  $q$** 

Value of $q$	Hypothesis	Lower 95% HDI	Median	Higher 95% HDI	Posterior Probability
0	Control < 0	-55.55	-15.13	24.95	0.77
1	Control < 0	-12.34	-4.85	3.27	0.89
2	Control < 0	-6.36	-2.96	0.41	0.96
0	Moderate > 0	-35.83	4.68	45.53	0.76
1	Moderate > 0	-1.44	6.1	14.02	0.97
2	Moderate > 0	-0.57	3.1	6.75	0.99
0	High > 0	-43.01	2.15	42.06	0.73
1	High > 0	-8.40	-0.42	7.62	0.78
2	High > 0	-4.66	-0.56	2.89	0.83

Hypotheses are listed per value of  $q$  with the estimated median and 95% highest density interval (HDI). Here, posterior probability represents the proportion of times the given hypothesis was supported throughout the Markov chain Monte Carlo chains.

infer that minimal, positive change in AM fungal richness was observed 73%, 78%, and 83% of the time of the posterior samples. The patterns here illustrate that rainfall exclusion had a positive effect on  $\Delta$  AM fungal richness and that as rare AM fungal taxa are removed from our analysis ( $q = 2$ ), the likelihood of this to be truer increased. These inferences in  $\Delta$  AM fungal richness are consistent with inferences generated from the 1,000 rarefied samples (Supplemental File 2: Figure S1).


Phylogenetic relatedness among rainfall exclusion treatments was measured as  $\Delta$  AM fungal richness across the monsoon season using the phylo-Horn N-assembly overlap measure ( $q = 1$  when informed by phylogeny). Median change in phylogenetic relatedness was -1.05, 1.21, and 0.41 for the control treatment, moderate rainfall exclusion treatment, and the high rainfall exclusion treatment, respectively. For the control treatment, we can infer negative change in phylogenetic relatedness 85% of the time throughout the monsoon season. The moderate and the high rainfall exclusion treatments differ in inference from the control treatment showing positive change in phylogenetic relatedness 94% and 83% of the time, respectively. A full summary of hypothesis testing for each treatment including the 95% highest density intervals and associated posterior probabilities can be found in Supplemental File 2: Table S1. Additionally, inferences per treatment for rarefied communities mirror that of inferences uncovered through this Bayesian analyses applied to proportional abundance data.

From  $q = 0$  to  $q = 2$ , AM fungal community turnover was measured as 1-pairwise beta diversity for each sample, which represents proportional change in AM fungal community composition across the monsoon season (between Date A and Date B). Across all values of  $q$ , the control treatment had the greatest AM fungal community turnover in comparison to other treatments (Figure 6). At  $q = 0, 1$ , and  $2$ , the median estimate for AM fungal turnover for the control treatment was estimated as 0.72, 0.57,

and 0.60 and was greater than zero 100% of the time for each value of  $q$  (Table 2). This indicates that there was community turnover in the control treatment due to rainfall. Median community turnover estimates for the moderate and high rainfall exclusion treatments were more similar to each other and differed from the control treatment. From  $q = 0$  to  $q = 2$ , rhizospheres that underwent moderate rainfall exclusion were shown to have lower estimates of turnover in comparison to the control treatment 96%, 93%, and 78% of the posterior samples were less than zero, respectively. This suggests that rainfall exclusion treatments influenced AM fungal community composition across the monsoon season (Figure 6). Further, rhizosphere microhabitats that experienced high rainfall exclusion across the monsoon season had the least AM fungal community turnover out of all treatments and differed from the control treatment where 99%, 99%, and 96% of the posterior samples were less than zero for  $q = 0, 1$ , and  $2$ . As rare species were down weighted from calculating AM fungal community turnover, the proportion of times our hypotheses were observed in the MCMC chains decreases. Figure 6 also illustrates that as the value of  $q$  increases from 0 to 2, so does the highest density intervals around the estimated values of AM fungal community turnover. Inferences made for AM fungal community turnover using proportional abundance data are in concordance with inferences made from 1,000 rarefied samples (Supplemental File 2; Figure S2). Lastly, median point estimates for rarefied AM fungal community turnover can be found in Supplemental File 2 as Table S2. Here, one can see that the 95% highest density intervals around the median are less than the highest density intervals seen around the median point estimates calculated using proportional abundance data.

## Discussion

In this study, we explicitly show the importance rainfall exclusion has on AM fungal richness and community



**Figure 6. Arbuscular mycorrhizal (AM) fungal community turnover among rainfall exclusion treatments.** AM fungal community turnover between time points and the 95% highest density interval around the median between the 2 time points. Bayesian inference was determined using linear models, where AM fungal community turnover was predicted by rainfall exclusion. This was conducted for  $q = 0, 1,$  and  $2$ . Here, the color black represents the control treatment, whereas teal and pink represent moderate (60%) and high (80%) rainfall exclusion treatments, respectively. Percentages are the proportion of samples from the joint probability distribution, where parameters are greater or less than zero (zero indicating no change in turnover). The percentage for control describes the change in diversity between time A and B, and the percentages for the moderate and high groups describe the change in diversity relative to the control group. Across all values of  $q$ , the control treatment, on average, had the largest observed AM fungal community turnover compared to other treatments.

turnover in the rhizosphere microhabitats *L. neomexicana*, an herbaceous plant found within the NAM region of the Sonoran Desert. Our results illustrate that variation in AM fungal community structure may be partially predicted by phosphorous and short-term rainfall exclusion. As soils get drier, AM fungal communities have been shown to have less seasonal community turnover and may be more influenced by phosphorous. Based on model selection, all other soil abiotic factors failed to explain significant variation in AM fungal community composition and richness. By the end of the monsoon season,  $\Delta$  AM fungal richness at  $q = 0, 1,$  and  $2$  responded to rainfall exclusion treatments; however, the response

differed among rainfall exclusion treatments, where the control treatment showed a loss in taxa due to rainfall and the moderate and high rainfall exclusion treatments showed increase in fungal ASVs. Rainfall exclusion also contributes to differences in phylogenetic relatedness among rhizosphere microhabitats in the Sonoran Desert, with rainfall contributing to a decrease in phylogenetic community composition. Lastly, AM fungal community turnover across the monsoon season was dependent on rainfall exclusion treatments, whereby the greatest turnover across all values of  $q$  was observed in the control treatment. Here, moderate rainfall exclusion could be a threshold that favors the coexistence of different AM

**Table 2. Hypothesis testing on pairwise arbuscular mycorrhizal fungal community turnover for each rainfall exclusion treatment and value of  $q$** 

Value of $q$	Hypothesis	Lower 95% HDI	Median	Higher 95% HDI	Posterior Probability
0	Control > 0	0.69	0.72	0.76	1
1	Control > 0	0.48	0.57	0.65	1
2	Control > 0	0.47	0.6	0.72	1
0	Moderate < 0	0.59	0.67	0.77	0.96
1	Moderate < 0	0.27	0.48	0.68	0.93
2	Moderate < 0	0.23	0.69	0.85	0.78
0	High < 0	0.57	0.66	0.75	0.99
1	High < 0	0.23	0.44	0.64	0.99
2	High < 0	0.14	0.44	0.75	0.96

For each hypothesis, the parameter estimates and the lower and upper bounds of the 95% highest density intervals are included. The probability for each hypothesis represents the proportion of times that the hypothesis was supported throughout the Markov chain Monte Carlo chains. HDI = highest density interval.

fungal taxa by the end of the monsoon season (Tilman, 1977). These results support our second prediction that AM fungal composition responds to short-term rainfall exclusion treatments, and more importantly, emphasizes the relationship between rainfall perturbations and soil abiotic factors like phosphorous content in the soil.

Our results for  $\Delta$  AM fungal richness across the monsoon season are consistent (Deveautour et al., 2020) as our moderate exclusion treatment showed an increase in AM fungal ASVs from  $q = 0$  to  $q = 2$  relative to the control treatment. Further, more extreme rainfall removal resulted in lesser changes in alpha diversity, which may be a result of taxa specific requirements for population-level persistence throughout decreased soil moisture and warmer summer temperatures (Tilman, 1977). In comparison to other ecological factors, precipitation and soil water availability more strongly influences AM fungal composition in semiarid ecosystems (Gao et al., 2016). In a study by Xiao et al. (2019), AM fungal diversity was influenced by phosphorous limitation, while total nitrogen affected the relative abundance of AM fungi. We show that in the context of naturally occurring AM fungal communities in the Sonoran Desert that phosphorous content in the rhizosphere microhabitat plays a substantial role in structuring the compositional spread of AM fungal communities and that nitrate content did not contribute to the variation in the effective number of AM fungal ASVs based on model selection. This suggests that as precipitation becomes more infrequent that AM fungal communities may be more dependent on specific, limiting soil abiotic factors in dry, arid climates.

The effects of sampling date on plant communities have been well studied and supported, providing some foundation to examine the temporal aspects of microbial communities (Smith and Huston, 1990; Guo and Brown, 1996). Temporal effects on the maintenance of AM fungal communities have been studied in a variety of ecological contexts ranging from volcanic ash fields (Higo et al.,

2015), mixed hardwood forests (Davison et al., 2012), to grasslands (Deveautour et al., 2020). There is mixed evidence showing that AM fungal community composition is temporally dynamic. For example, in mixed hardwood forests, AM fungal communities have been shown to be spatially distinct but temporally stable across the growing season, perhaps due to the heterogeneous plant community and a consistent AM fungal species pool over their 2-year sampling scheme (Davison et al., 2012). Yet, on the same time scale, Higo et al. (2015) state that differences in AM fungal composition are dependent on plant cover from the previous year, thus contributing to fungal community turnover year to year. Temporal effects have also been observed on much smaller time scales, such as over one season in a wet tropical forest (Waring and Hawkes, 2015). Our multivariate models show distinct clustering for the composition of AM fungal communities over the NAM season and by rainfall exclusion treatment. Additionally, we show that AM fungal community turnover responds to rainfall exclusion treatments from  $q = 0$  to  $q = 2$ . The effect seen here may illustrate the contribution available soil moisture has on the number of ASVs throughout the monsoon season and on the relative abundance of each ASV in a dry, arid ecosystem. Our results further highlight the interconnected relationships between regional climate and local soil abiotic factors that could hinder a generalizable pattern consensus of a spatio-temporal response of AM fungal community composition.

Here, we demonstrate that rainfall exclusion affects variation in AM fungal composition and the difference in the effective number of AM fungal ASVs, where moderate rainfall exclusion led to increased estimates in alpha diversity. Additionally, we show that AM fungal community turnover is dependent on rainfall exclusion when common and rare ASVs are weighted equally and also when rare ASVs are down weighted in estimating community turnover. The work outlined in this study leaves several forward directions for AM fungal ecology. Specifically,

what are the relative temporal scales needed to observe AM fungal responses to manipulative field experiments? How does temporal turnover in AM fungal communities differ between within year and among year exposure to environmental stress? AM fungi can be somewhat ubiquitous at local scales (Davison et al., 2015) and have been shown to be heterokaryotic (i.e., multinucleate with genetically distinct nuclei; Bruns et al., 2018). This could lead to lack of agreement in empirical biodiversity studies, as species richness estimates might be dependent on spatial scale, temporal scale, and the weight of rare ASVs. Therefore, while we did find that AM fungal communities respond to changes in seasonal monsoon precipitation, more studies across a wider range of ecological contexts could further uncover the direction and magnitude of changing climate conditions on plant–host AM fungal interactions.

#### Data accessibility statement

Sequence data are available in the NCBI Sequence Read Archive under BioProject ID: PRJNA1065923.

#### Supplemental files

The supplemental files for this article can be found as follows:

- Supplemental File 1.xlsx
- Supplemental File 2. docx

#### Acknowledgments

The authors would like to begin by giving thanks to the Collections Committee and the TENN Herbarium at the University of Tennessee, Knoxville (Department of Ecology and Evolutionary Biology) for recommendations regarding experimental design and funding field portions of this study. Secondly, they would like to thank the Arnold Lab at the University of Arizona, School of Plant Sciences for hosting during field collection. They additionally acknowledge Michele Lanan and the Southwest Research Station (American Museum of Natural History) for coordination and lab space while in the Chiricahua Mountains. They would like to thank the U.S. Department of Agriculture, Forest Service, and the Douglas Ranger District for permitting this study during 2019. They would also like to thank Stephanie Kivlin for providing access to lab space for 18S library preparation and soil assays during the coronavirus pandemic and for providing comments that substantially improved this work.

#### Funding

The Kivlin lab was funded by the NSF DEB 1936195, Texas Ecological Laboratory, and The University of Tennessee, Knoxville. This work was also supported by The National Science Foundation DEB 1638922 to JAF.

#### Competing interests

The authors declare that they have no competing or conflict of interest.

#### Author contributions

Contributed to conception and design: JRD, JAF.

Contributed to acquisition of data: JRD.

Contributed to analysis and interpretation of data: JRD, JAF.

Drafted and/or revised the manuscript: JRD, JAF.

Approved the final version for submission: JRD, JAF.

#### References

- Alberdi, A, Gilbert, MTP.** 2019. Hilldiv: An R package for the integral analysis of diversity based on hill numbers. *Biorxiv*: 545665. Cold Spring Harbor Laboratory.
- Allen, MF.** 1991. *The ecology of mycorrhizae*. Cambridge, UK: Cambridge University Press.
- Anderson, RC, Liberta, AE, Dickman, LA.** 1984. Interaction of vascular plants and vesicular-arbuscular mycorrhizal fungi across a soil moisture-nutrient gradient. *Oecologia* **64**(1): 111–117. DOI: <http://dx.doi.org/10.1007/BF00377552>.
- Beier, C, Beierkuhnlein, C, Wohlgemuth, T, Penuelas, J, Emmett, B, Körner, C, de Boeck, H, Christensen, JH, Leuzinger, S, Janssens, IA, Hansen, K.** 2012. Precipitation manipulation experiments—Challenges and recommendations for the future. *Ecology Letters* **15**(8): 899–911. DOI: <http://dx.doi.org/10.1111/j.1461-0248.2012.01793.x>.
- Birhane, E, Sterck, FJ, Fetene, M, Bongers, F, Kuyper, TW.** 2012. Arbuscular mycorrhizal fungi enhance photosynthesis, water use efficiency, and growth of frankincense seedlings under pulsed water availability conditions. *Oecologia* **169**(4): 895–904. DOI: <http://dx.doi.org/10.1007/s00442-012-2258-3>.
- Bobille, H, Fustec, J, Robins, RJ, Cukier, C, Limami, AM.** 2019. Effect of water availability on changes in root amino acids and associated rhizosphere on root exudation of amino acids in *Pisum sativum* L. *Phytochemistry* **161**: 75–85. DOI: <http://dx.doi.org/10.1016/j.phytochem.2019.01.015>.
- Bradie, J, Leung, B.** 2017. A quantitative synthesis of the importance of variables used in MaxEnt species distribution models. *Journal of Biogeography* **44**(6): 1344–1361. DOI: <http://dx.doi.org/10.1111/jbi.12894>.
- Brookes, PC, Landman, A, Pruden, G, Jenkinson, DS.** 1985. Chloroform fumigation and the release of soil nitrogen: A rapid direct extraction method to measure microbial biomass nitrogen in soil. *Soil Biology & Biochemistry* **17**(6): 837–842. DOI: [http://dx.doi.org/10.1016/0038-0717\(85\)90144-0](http://dx.doi.org/10.1016/0038-0717(85)90144-0).
- Bruns, TD, Corradi, N, Redecker, D, Taylor, JW, Öpik, M.** 2018. Glomeromycotina: What is a species and why should we care? *New Phytologist* **220**(4): 963–967. DOI: <http://dx.doi.org/10.1111/nph.14913>.
- Bürkner, PC.** 2017. brms: An R package for Bayesian multilevel models using Stan. *Journal of Statistical Software* **80**(1). DOI: <https://doi.org/10.18637/jss.v080.i01>.
- Callahan, BJ, McMurdie, PJ, Rosen, MJ, Han, AW, Johnson, AJA, Holmes, SP.** 2016. DADA2: High-resolution sample inference from Illumina amplicon

- data. *Nature Methods* **13**(7): 581–583. DOI: <http://dx.doi.org/10.1038/nmeth.3869>.
- Carpenter, B, Gelman, A, Hoffman, MD, Lee, D, Goodrich, B, Betancourt, M, Brubaker, M, Guo, J, Li, P, Riddell, A.** 2017. Stan: A probabilistic programming language. *Journal of Statistical Software* **76**(1): 1–32.
- Chao, A, Chiu, CH, Jost, L.** 2010. Phylogenetic diversity measures based on hill numbers. *Philosophical Transactions of the Royal Society Biological Sciences* **365**(1558): 3599–3609. DOI: <http://dx.doi.org/10.1098/rstb.2010.0272>.
- Comrie, AC, Glenn, EC.** 1999. Principal components-based regionalization of precipitation regimes across the southwest United States and northern Mexico, with an application to monsoon precipitation variability. *Climate Research* **10**(3): 201–215. DOI: <http://dx.doi.org/10.3354/cr010201>.
- Davison, J, Moora, M, Öpik, M, Adholeya, A, Ainsaar, L, Bâ, A, Burla, S, Diedhiou, AG, Hiiesalu, I, Jairus, T.** 2015. Global assessment of arbuscular mycorrhizal fungus diversity reveals very low endemism. *Science* **349**(6251): 970–973.
- Davison, J, Öpik, M, Zobel, M, Vasar, M, Metsis, M, Moora, M.** 2012. Communities of arbuscular mycorrhizal fungi detected in forest soil are spatially heterogeneous but do not vary throughout the growing season. *PLoS One* **7**(8). DOI: <http://dx.doi.org/10.1371/journal.pone.0041938>.
- Deepika, S, Kothamasi, D.** 2015. Soil moisture—A regulator of arbuscular mycorrhizal fungal community assembly and symbiotic phosphorus uptake. *Mycorrhiza* **25**(1): 67–75. DOI: <http://dx.doi.org/10.1007/s00572-014-0596-1>.
- Demaria, EMC, Hazenberg, P, Scott, RL, Meles, MB, Nichols, M, Goodrich, D.** 2019. Intensification of the North American monsoon rainfall as observed from a long-term high-density gauge network. *Sciences Geophysical Research Letters* **46**(12): 6837–6847. DOI: <http://dx.doi.org/10.1029/2019GL082461>.
- Deveautour, C, Power, SA, Barnett, KL, Ochoa-Hueso, R, Donn, S, Bennett, AE, Powell, JR.** 2020. Temporal dynamics of mycorrhizal fungal communities and co-associations with grassland plant communities following experimental manipulation of rainfall. *Journal of Ecology* **108**(2): 515–527. DOI: <http://dx.doi.org/10.1111/1365-2745.13267>.
- Diffenbaugh, NS, Giorgi, F.** 2012. Climate change hotspots in the CMIP5 global climate model ensemble. *Climate Change* **114**(3–4): 813–822. DOI: <http://dx.doi.org/10.1007/s10584-012-0570-x>.
- Dumbrell, AJ, Ashton, PD, Aziz, N, Feng, G, Nelson, M, Dytham, C, Fitter, AH, Helgason, T.** 2011. Distinct seasonal assemblages of arbuscular mycorrhizal fungi revealed by massively parallel pyrosequencing. *New Phytologist* **190**(3): 794–804. DOI: <http://dx.doi.org/10.1111/j.1469-8137.2010.03636.x>.
- Gao, C, Kim, YC, Zheng, Y, Yang, W, Chen, L, Ji, NN, Wan, SQ, Guo, LD.** 2016. Increased precipitation, rather than warming, exerts a strong influence on arbuscular mycorrhizal fungal community in a semi-arid steppe ecosystem. *Botany* **94**(6): 459–469. DOI: <http://dx.doi.org/10.1139/cjb-2015-0210>.
- Green, NE, Graham, SO, Schenck, NC.** 1976. The influence of pH on the germination of vesicular-arbuscular mycorrhizal spores. *Mycologia* **68**(4): 929–934.
- Guo, Q, Brown, JH.** 1996. Temporal fluctuations and experimental effects in desert plant communities. *Oecologia* **107**(4): 568–577. DOI: <http://dx.doi.org/10.1007/BF00333950>.
- Hayman, DS.** 1982. Influence of soils and fertility on activity and survival of vesicular-arbuscular mycorrhizal fungi. *Phytopathology* **72**(8): 1119–1125.
- Hernandez, DJ, David, AS, Menges, ES, Searcy, CA, Afkhami, ME.** 2021. Environmental stress destabilizes microbial networks. *ISME Journal* **15**(6): 1722–1734. DOI: <http://dx.doi.org/10.1038/s41396-020-00882-x>.
- Higo, M, Isobe, K, Kondo, T, Yamaguchi, M, Takeyama, S, Drijber, RA, Torigoe, Y.** 2015. Temporal variation of the molecular diversity of arbuscular mycorrhizal communities in three different winter cover crop rotational systems. *Biology and Fertility of Soils* **51**(1): 21–32. DOI: <http://dx.doi.org/10.1007/s00374-014-0945-4>.
- Hill, MO.** 1973. Diversity and evenness: A unifying notation and its consequences. *Ecology* **54**(2): 427–432. DOI: <http://dx.doi.org/10.2307/1934352>.
- Johnson, NC, Tilman, D, Wedin, D.** 1992. Plant and soil controls on mycorrhizal fungal communities. *Ecology* **73**(6): 2034–2042. DOI: <http://dx.doi.org/10.2307/1941453>.
- Johnson, NC, Wilson, GWT, Bowker, MA, Wilson, JA, Miller, RM.** 2010. Resource limitation is a driver of local adaptation in mycorrhizal symbioses. *Proceedings of the National Academy of Sciences of the United States of America* **107**(5): 2093–2098. DOI: <http://dx.doi.org/10.1073/pnas.0906710107>.
- Kanwal, S, Bano, A, Malik, RN.** 2015. Effects of arbuscular mycorrhizal fungi on metals uptake, physiological and biochemical response of *Medicago Sativa* L. with increasing Zn and Cd concentrations in soil. *American Journal of Plant Sciences* **06**(18): 2906–2923. DOI: <http://dx.doi.org/10.4236/ajps.2015.618287>.
- Katoh, K, Standley, DM.** 2013. MAFFT multiple sequence alignment software version 7: Improvements in performance and usability. *Molecular Biology and Evolution* **30**(4): 772–780. DOI: <http://dx.doi.org/10.1093/molbev/mst010>.
- Kivlin, SN, Harpe, VR, Turner, JH, Moore, JA, Moorhead, LC, Beals, KK, Hubert, MM, Papes, M, Schweitzer, JA.** 2021. Arbuscular mycorrhizal fungal response to fire and urbanization in the Great Smoky Mountains National Park. *Elementa: Science of the Anthropocene* **9**(1): 00037. DOI: <http://dx.doi.org/10.1525/elementa.2021.00037>.
- Kivlin, SN, Winston, GC, Goulden, ML, Treseder, KK.** 2014. Environmental filtering affects soil fungal community composition more than dispersal limitation

- at regional scales. *Fungal Ecology* **12**(C): 14–25. DOI: <http://dx.doi.org/10.1016/j.funeco.2014.04.004>.
- Kuehn, K-D, Weber, H-C, Dehne, H-W, Gworgwor, NA.** 1991. Distribution of vesicular-arbuscular mycorrhizal fungi on a fallow agriculture site, pt. 1: Dry habitat. *Angewandte Botanik* **65**: 169–186.
- Lajtha, K, Driscoll, CT, Jarrell, WM, Elliott, ET.** 1999. Soil P characterization and total element analysis, in Robertson, GP, Coleman, DC, Bledsoe, CS, Sollins, P eds., *Stand soil methods long-term ecological research*. Oxford, UK: Oxford University Press.
- Lau, JA, Lennon, JT.** 2011. Evolutionary ecology of plant–microbe interactions: Soil microbial structure alters selection on plant traits. *New Phytologist* **192**(1): 215–224. DOI: <http://dx.doi.org/10.1111/j.1469-8137.2011.03790.x>.
- Lee, J, Lee, S, Young, JPW.** 2008. Improved PCR primers for the detection and identification of arbuscular mycorrhizal fungi. *FEMS Microbiology Ecology* **65**(2): 339–349. DOI: <http://dx.doi.org/10.1111/j.1574-6941.2008.00531.x>.
- Meredith, M, Kruschke, J.** 2020. HDInterval: Highest (posterior) density intervals. R Package v.02 2.
- Miller, SP.** 2000. Arbuscular mycorrhizal colonization of semi-aquatic grasses along a wide hydrologic gradient. *New Phytologist* **145**(1): 145–155. DOI: <http://dx.doi.org/10.1046/j.1469-8137.2000.00566.x>.
- Mirarab, S, Nguyen, N, Guo, S, Wang, LS, Kim, J, Warnow, T.** 2015. PASTA: Ultra-large multiple sequence alignment for nucleotide and amino-acid sequences. *Journal of Computational Biology* **22**(5): 377–386. DOI: <http://dx.doi.org/10.1089/cmb.2014.0156>.
- Mitchell, DL, Ivanova, D, Rabin, R, Brown, TJ, Redmond, K.** 2002. Gulf of California sea surface temperatures and the North American monsoon: Mechanistic implications from observations. *Journal of Climate* **15**(17): 2261–2281. DOI: [http://dx.doi.org/10.1175/1520-0442\(2002\)015<2261:GOCSSST>2.0.CO;2](http://dx.doi.org/10.1175/1520-0442(2002)015<2261:GOCSSST>2.0.CO;2).
- Mulvaney, RL.** 1996. Nitrogen-inorganic forms, in Sparks, DL, Page, AL, Helmke, PA, Loeppert, RH, Soltanpoor, PN, Tabatabai, MA, Johnston, CT, Sumner, ME eds., *Methods of Soil Analysis. Part 3 Chemical Methods*. Madison, WI: SSSA: 1123–1184. (SSSA Book Series No. 5).
- Newsham, KK, Fitter, AH, Watkinson, AR.** 1995. Multifunctionality and biodiversity in arbuscular mycorrhizas. *Trends in Ecology & Evolution* **10**(10): 407–411. DOI: [http://dx.doi.org/10.1016/S0169-5347\(00\)89157-0](http://dx.doi.org/10.1016/S0169-5347(00)89157-0).
- Oksanen, AJ, Blanchet, FG, Friendly, M, Kindt, R, Legendre, P, Mcglinn, D, Minchin, PR, Hara, RBO, Simpson, GL, Solymos, P, Stevens, H, Wagner, HH.** 2019. Package “vegan.” Available at <https://cran.r-project.org>; <https://github.com/veg>. Accessed March 1, 2022.
- Öpik, M, Vanatoa, A, Vanatoa, E, Moora, M, Davison, J, Kalwij, JM, Reier, Ü, Zobel, M.** 2010. The online database MaarjAM reveals global and ecosystemic distribution patterns in arbuscular mycorrhizal fungi (Glomeromycota). *New Phytologist* **188**(1): 223–241. DOI: <http://dx.doi.org/10.1111/j.1469-8137.2010.03334.x>.
- Price, MN, Dehal, PS, Arkin, AP.** 2010. FastTree 2—Approximately maximum-likelihood trees for large alignments. *PLoS One* **5**(3). DOI: <http://dx.doi.org/10.1371/journal.pone.0009490>.
- Raja, HA, Miller, AN, Pearce, CJ, Oberlies, NH.** 2017. Fungal identification using molecular tools: A primer for the natural products research community. *Journal of Natural Products* **80**(3): 756–770. DOI: <http://dx.doi.org/10.1021/acs.jnatprod.6b01085>.
- Santos-González, JC, Finlay, RD, Tehler, A.** 2007. Seasonal dynamics of arbuscular mycorrhizal fungal communities in roots in a seminatural grassland. *Applied and Environmental Microbiology* **73**(17): 5613–5623. DOI: <http://dx.doi.org/10.1128/AEM.00262-07>.
- Simon, L, Lalonde, M, Bruns, TD.** 1992. Specific amplification of 18 S fungal ribosomal genes from vesicular-arbuscular mycorrhizal fungal communities. *Applied and Environmental Microbiology* **58**(1): 291–295.
- Smith, SA, O'Meara, BC.** 2012. treePL: Divergence time estimation using penalized likelihood for large phylogenies. *Bioinformatics* **28**(20): 2689–2690. DOI: <http://dx.doi.org/10.1093/bioinformatics/bts492>.
- Smith, T, Huston, M.** 1990. A theory of the spatial and temporal dynamics of plant communities. *Progress in Theoretical Vegetation Science* **83**(1–2): 49–69.
- Soudzilovskaia, NA, Vaessen, S, Barcelo, M, He, J, Rahimlou, S, Abarenkov, K, Brundrett, MC, Gomes, SIF, Merckx, V, Tedersoo, L.** 2020. Fungal-Root: Global online database of plant mycorrhizal associations. *New Phytologist* **227**(3): 955–966. DOI: <http://dx.doi.org/10.1111/nph.16569>.
- Standards Association of Australia.** 1977. *Methods of testing soils for engineering purposes*. North Sydney, NSW: The Association.
- Tedersoo, L, Bahram, M, Pölme, S, Kõljalg, U, Yorou, NS, Wijesundera, R, Ruiz, LV, Vasco-Palacios, AM, Thu, PQ, Suija, A, Smith, ME, Sharp, C, Saluveer, E, Saitta, A, Rosas, M, Riit, T, Ratkowsky, D, Pritsch, K, Pöldmaa, K, Piepenbring, M, Phosri, C, Peterson, M, Parts, K, Pärtel, K, Otsing, E, Nouhra, E, Njouonkou, AL, Nilsson, RH, Morgado, LN, Mayor, J, May, TW, Majuakim, L, Lodge, DJ, Lee, SS, Larsson, KH, Kohout, P, Hosaka, K, Hiiesalu, I, Henkel, TW, Harend, H, Guo, LD, Greslebin, A, Grelet, G, Geml, J, Gates, G, Dunstan, W, Dunk, C, Drenkhan, R, Dearnaley, J, De Kesel, A, Dang, T, Chen, X, Buegger, F, Brearley, FQ, Bonito, G, Anslan, S, Abell, S, Abarenkov, K.** 2014. Global diversity and geography of soil fungi. *Science* **346**(6213): 1256688. DOI: <http://dx.doi.org/10.1126/science.1256688>.
- Tilman, D.** 1977. Resource competition between plankton algae: An experimental and theoretical approach. *Ecology* **58**(2): 338–348.

- Vance, ED, Brookes, PC, Jenkinson, DS.** 1987. An extraction method for measuring soil microbial biomass C. *Soil Biology and Biochemistry* **19**(6): 703–707. DOI: [http://dx.doi.org/10.1016/0038-0717\(87\)90052-6](http://dx.doi.org/10.1016/0038-0717(87)90052-6).
- Van Der Heijden, MGA, Bardgett, RD, Van Straalen, NM.** 2008. The unseen majority: Soil microbes as drivers of plant diversity and productivity in terrestrial ecosystems. *Ecology Letters* **11**(3): 296–310. DOI: <http://dx.doi.org/10.1111/j.1461-0248.2007.01139.x>.
- Van Der Putten, WH, Macel, M, Visser, ME.** 2010. Predicting species distribution and abundance responses to climate change: Why it is essential to include biotic interactions across trophic levels. *Philosophical Transactions of the Royal Society Biological Sciences* **365**(1549): 2025–2034. DOI: <http://dx.doi.org/10.1098/rstb.2010.0037>.
- Verdouw, H, Van Echteld, CJA, Dekkers, EMJ.** 1978. Ammonia determination based on indophenol formation with sodium salicylate. *Water Research* **12**(6): 399–402. DOI: [http://dx.doi.org/10.1016/0043-1354\(78\)90107-0](http://dx.doi.org/10.1016/0043-1354(78)90107-0).
- Waring, BG, Hawkes, CV.** 2015. Short-term precipitation exclusion alters microbial responses to soil moisture in a wet tropical forest. *Microbial Ecology* **69**(4): 843–854. DOI: <http://dx.doi.org/10.1007/s00248-014-0436-z>.
- Weatherburn, MW.** 1967. Phenol-hypochlorite reaction for determination of ammonia. *Analytical Chemistry* **39**(8): 971–974. DOI: <http://dx.doi.org/10.1021/ac60252a045>.
- White, TJ, Bruns, T, Lee, S, Taylor, J.** 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics, in Innis, MA, Gelfand, DH, Sninsky, JJ, White, TJ eds., *PCR protocols: A guide to methods and applications*. San Diego, CA: Academic Press: 315–322.
- Wu, J, Joergensen, RG, Pommerening, B, Chaussod, R, Brookes, PC.** 1990. Measurement of soil microbial biomass C by fumigation-extraction-an automated procedure. *Soil Biology and Biochemistry* **22**(8): 1167–1169.
- Wu, QS.** 2017. *Arbuscular mycorrhizas and stress tolerance of plants*. Singapore: Springer. DOI: <http://dx.doi.org/10.1007/978-981-10-4115-0>.
- Xiao, D, Che, R, Liu, X, Tan, Y, Yang, R, Zhang, W, He, X, Xu, Z, Wang, K.** 2019. Arbuscular mycorrhizal fungi abundance was sensitive to nitrogen addition but diversity was sensitive to phosphorus addition in karst ecosystems. *Biology and Fertility of Soils* **55**(5): 457–469. DOI: <http://dx.doi.org/10.1007/s00374-019-01362-x>.
- Zhou, J, Deng, Y, Shen, L, Wen, C, Yan, Q, Ning, D, Qin, Y, Xue, K, Wu, L, He, Z, Voordeckers, JW, Nostrand, JD, Buzzard, V, Michaletz, ST, Enquist, BJ, Weiser, MD, Kaspary, M, Waide, R, Yang, Y, Brown, JH.** 2016. Temperature mediates continental-scale diversity of microbes in forest soils. *Nature Communications* **7**: 12083. DOI: <http://dx.doi.org/10.1038/ncomms12083>.

**How to cite this article:** Dickey, JR, Fordyce, JA. 2024. Greater change in arbuscular mycorrhizal fungal richness as a response to short-term rainfall exclusion across the North American monsoon season. *Elementa: Science of the Anthropocene* 12(1). DOI: <https://doi.org/10.1525/elementa.2022.00116>

**Domain Editor-in-Chief:** Steven Allison, University of California Irvine, Irvine, CA, USA

**Associate Editor:** Rebecca A. Lybrand, Department of Land, Air and Water Resources, University of California, Davis, CA, USA

**Knowledge Domain:** Ecology and Earth Systems

**Published:** February 22, 2024    **Accepted:** December 22, 2023    **Submitted:** September 12, 2022

**Copyright:** © 2024 The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC-BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. See <http://creativecommons.org/licenses/by/4.0/>.

