

UC Davis

Faculty

Title

Lectures on the Topological Recursion for Higgs Bundles and Quantum Curves

Permalink

<https://escholarship.org/uc/item/858382hb>

Authors

Dumitrescu, O
Mulase, M

Publication Date

2018

DOI

10.1142/9789813229099_0003

Peer reviewed

LECTURES ON THE TOPOLOGICAL RECURSION FOR HIGGS BUNDLES AND QUANTUM CURVES

OLIVIA DUMITRESCU AND MOTOHICO MULASE

ABSTRACT. The paper aims at giving an introduction to the notion of *quantum curves*. The main purpose is to describe the new discovery of the relation between the following two disparate subjects: one is the topological recursion, that has its origin in random matrix theory and has been effectively applied to many enumerative geometry problems; and the other is the quantization of Hitchin spectral curves associated with Higgs bundles. Our emphasis is on explaining the motivation and examples. Concrete examples of the direct relation between Hitchin spectral curves and enumeration problems are given. A general geometric framework of quantum curves is also discussed.

CONTENTS

1. Introduction	2
1.1. On the other side of the rainbow	2
1.2. Quantum curves, semi-classical limit, and the WKB analysis	11
1.3. The topological recursion as quantization	13
1.4. Non-Abelian Hodge correspondence and quantum curves	16
1.5. The Lax operator for Witten-Kontsevich KdV equation	19
1.6. All things considered	21
2. From Catalan numbers to the topological recursion	24
2.1. Counting graphs on a surface	25
2.2. The spectral curve of a Higgs bundle and its desingularization	29
2.3. The generating function, or the Laplace transform	31
2.4. The unstable geometries and the initial value of the topological recursion	33
2.5. Geometry of the topological recursion	34
2.6. The quantum curve for Catalan numbers	38
2.7. Counting lattice points on the moduli space $\mathcal{M}_{g,n}$	40
3. Quantization of spectral curves	45
3.1. Geometry of non-singular Hitchin spectral curves of rank 2	45
3.2. The generalized topological recursion for the Hitchin spectral curves	47
3.3. Quantization of Hitchin spectral curves	51
3.4. Classical differential equations	54
4. Difference operators as quantum curves	59
4.1. Simple and orbifold Hurwitz numbers	60
4.2. Gromov-Witten invariants of the projective line	63
References	66

2010 *Mathematics Subject Classification*. Primary: 14H15, 14N35, 81T45; Secondary: 14F10, 14J26, 33C05, 33C10, 33C15, 34M60, 53D37.

Key words and phrases. Topological quantum field theory; topological recursion; quantum curves; Hitchin spectral curves, Higgs bundles.

We're not going to tell you the story
the way it happened.
We're going to tell it
the way we remember it.

1. INTRODUCTION

Mathematicians often keep their childhood dream for a long time. When you saw a perfect rainbow as a child, you might have wondered what awaited you when you went through the arch. In a lucky situation, you might have seen the double, or even triple, rainbow arches spanning above the brightest one, with the reversing color patterns on the higher arches. Yet we see nothing underneath the brightest arch.

One of the purposes of these lectures is to offer you a vision: **on the other side of the rainbow, you see quantum invariants.** This statement describes only the tip of the iceberg. We believe something like the following is happening: Let C be a smooth projective curve over \mathbb{C} and (E, ϕ) a *meromorphic* Higgs bundle on C . Then the asymptotic expansion at an essential singularity of a solution (flat section) of the \hbar -connection on C , that is the image of the **non-abelian Hodge correspondence** applied to the Hitchin spectral curve of (E, ϕ) , carries the information of quantum invariants of a totally different geometric structure, which should be considered as the **mirror** to the Higgs bundle.

In this introduction, we are going to tell a story of an example to this mirror correspondence using the rainbow integral of Airy. The Hitchin spectral curve is a singular compactification of a parabola $x = y^2$ in a Hirzebruch surface. The corresponding quantum invariants, the ones we see under the rainbow, are the cotangent class intersection numbers of the moduli space $\overline{\mathcal{M}}_{g,n}$. These numbers then determine the coefficients of the tautological relations among the generators of the tautological rings of \mathcal{M}_g and $\overline{\mathcal{M}}_{g,n}$. The uniqueness of the asymptotic expansion relates the WKB analysis of the quantization of the parabola at the infinity to the intersection numbers on $\overline{\mathcal{M}}_{g,n}$, through a combinatorial estimate of the *volume* of the moduli space $\mathcal{M}_{g,n}$.

The story begins in 1838.

1.1. On the other side of the rainbow. Sir George Biddel Airy devised a simple formula, which he called the *rainbow integral*

$$(1.1) \quad Ai(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{ipx} e^{i\frac{p^3}{3}} dp$$

and now carries his name, in his attempt of explaining the rainbow phenomena in terms of wave optics [2].




FIGURE 1.1. Rainbow arches

The angle between the sun and the observer measured at the brightest arch is always about 42° . The higher arches also have definite angles, independent of the rainbow. Airy tried to explain these angles and the brightness of the rainbow arches by the peak points of the rainbow integral.




FIGURE 1.2. The Airy function

We note that (1.1) is an *oscillatory integral*, and determines a real analytic function in $x \in \mathbb{R}$. It is easy to see, by integration by parts and taking care of the boundary contributions in oscillatory integral, that

$$\begin{aligned} \frac{d^2}{dx^2} Ai(x) &= \frac{1}{2\pi} \int_{-\infty}^{\infty} (-p^2) e^{ipx} e^{i\frac{p^3}{3}} dp \\ &= \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{ipx} \left(i \frac{d}{dp} e^{i\frac{p^3}{3}} \right) dp \\ &= -\frac{1}{2\pi} \int_{-\infty}^{\infty} \left(i \frac{d}{dp} e^{ipx} \right) e^{i\frac{p^3}{3}} dp \\ &= \frac{1}{2\pi} x \int_{-\infty}^{\infty} e^{ipx} e^{i\frac{p^3}{3}} dp. \end{aligned}$$

Thus the Airy function satisfies a second order differential equation (known as the Airy differential equation)

$$(1.2) \quad \left(\frac{d^2}{dx^2} - x \right) Ai(x) = 0.$$

Now we consider $x \in \mathbb{C}$ as a complex variable. Since the coefficients of (1.2) are entire functions (since they are just 1 and x), any solution of this differential equation is automatically entire, and has a convergent power series expansion

$$(1.3) \quad Ai(x) = \sum_{n=0}^{\infty} a_n x^n$$

at the origin with the radius of convergence ∞ . Plugging (1.3) into (1.2), we obtain a recursion formula

$$(1.4) \quad a_{n+2} = \frac{1}{(n+2)(n+1)} a_{n-1}, \quad n \geq 0,$$

with the initial condition $a_{-1} = 0$. Thus we find

$$a_{3n} = a_0 \cdot \frac{\prod_{j=1}^n (3j-2)}{(3n)!}, \quad a_{3n+1} = a_1 \cdot \frac{\prod_{j=1}^n (3j-1)}{(3n+1)!}, \quad a_{3n+2} = 0.$$

Here, a_0 and a_1 are arbitrary constants, so that the Airy differential equation has a two-dimensional space of solutions. These coefficients do not seem to be particularly interesting. The oscillatory integral (1.1) tells us that as $x \rightarrow +\infty$ on the real axis, the Airy function defined by the rainbow integral vanishes, because $e^{ipx+ip^3/3}$ oscillates so much that the integral cancels. More precisely, $Ai(x)$ satisfies a limiting formula

$$(1.5) \quad \lim_{x \rightarrow +\infty} \frac{Ai(x)}{\frac{1}{2\sqrt{\pi}} \cdot \frac{1}{\sqrt[4]{x}} \exp\left(-\frac{2}{3}x^{3/2}\right)} = 1.$$

Hence it exponentially decays, as $x \rightarrow +\infty$, $x \in \mathbb{R}$. Among the Taylor series solutions (1.3), there is *only one* solution that satisfies this exponential decay property, which is given by the following initial condition for (1.4):

$$a_0 = \frac{1}{3^{2/3}\Gamma(\frac{2}{3})}, \quad a_1 = -\frac{1}{3^{1/3}\Gamma(\frac{1}{3})}.$$

The exponential decay on the positive real axis explains why we do not see any rainbows under the brightest arch. Then what do we really see underneath the rainbow? Or on the *other side* of the rainbow?

The differential equation (1.2) tells us that the Airy function has an essential singularity at $x = \infty$. Otherwise, the solution would be a polynomial in x , but (1.4) does not terminate at a finite n . How do we analyze the behavior of a holomorphic function at its essential singularity? And what kind of information does it tell us?

Definition 1.1 (Asymptotic expansion). Let $f(z)$ be a holomorphic function defined on an open domain Ω of the complex plane \mathbb{C} having the origin 0 on its boundary. A formal power series

$$\sum_{n=0}^{\infty} a_n z^n$$

is an *asymptotic expansion* of $f(z)$ on Ω at $z = 0$ if

$$(1.6) \quad \lim_{\substack{z \rightarrow 0 \\ z \in \Omega}} \frac{1}{z^{m+1}} \left(f(z) - \sum_{n=0}^m a_n z^n \right) = a_{m+1}$$

holds for every $m \geq 0$.

The asymptotic expansion is a domain specific notion. For example, $f(z) = e^{-1/z}$ is holomorphic on $\mathbb{C}^* = \mathbb{C} \setminus \{0\}$, but it does not have any asymptotic expansion on the whole \mathbb{C}^* . However, it has an asymptotic expansion

$$e^{-1/z} \sim 0$$

on

$$\Omega = \left\{ z \in \mathbb{C}^* \mid |\operatorname{Arg}(z)| < \frac{\pi}{2} - \epsilon \right\}, \quad \epsilon > 0.$$

If there is an asymptotic expansion of f on a domain Ω , then it is unique, and if $\Omega' \subset \Omega$ with $0 \in \partial\Omega'$, then obviously the same asymptotic expansion holds on Ω' .

The Taylor expansion (1.3) shows that $Ai(x)$ is real valued on the real axis, and from (1.5), we see that the value is positive for $x > 0$. Therefore, $\log Ai(x)$ is a holomorphic function on $Re(x) > 0$.

Theorem 1.2 (Asymptotic expansion of the Airy function). *Define*

$$(1.7) \quad S_0(x) = -\frac{2}{3}x^{\frac{3}{2}}, \quad S_1(x) = -\frac{1}{4}\log x - \log(2\sqrt{\pi}).$$

Then $\log Ai(x) - S_0(x) - S_1(x)$ has the following asymptotic expansion on $Re(x) > 0$.

$$(1.8) \quad \log Ai(x) - S_0(x) - S_1(x) = \sum_{m=2}^{\infty} S_m(x),$$

$$(1.9) \quad S_m(x) := x^{-\frac{3}{2}(m-1)} \cdot \frac{1}{2^{m-1}} \sum_{\substack{g \geq 0, n > 0 \\ 2g-2+n=m-1}} \frac{(-1)^n}{n!} \sum_{\substack{d_1 + \dots + d_n \\ = 3g-3+n}} \langle \tau_{d_1} \cdots \tau_{d_n} \rangle_{g,n} \prod_{i=1}^n (2d_i - 1)!!$$

for $m \geq 2$. The key coefficients are defined by

$$(1.10) \quad \langle \tau_{d_1} \cdots \tau_{d_n} \rangle_{g,n} := \int_{\overline{\mathcal{M}}_{g,n}} \psi_1^{d_1} \cdots \psi_n^{d_n},$$

where, $\overline{\mathcal{M}}_{g,n}$ is the moduli space of stable curves of genus g with n smooth marked points. Let $[C, (p_1, \dots, p_n)] \in \overline{\mathcal{M}}_{g,n}$ be a point of the moduli space. We can construct a line bundle \mathbb{L}_i on the smooth Deligne-Mumford stack $\overline{\mathcal{M}}_{g,n}$ by attaching the cotangent line $T_{p_i}^*C$ at the point $[C, (p_1, \dots, p_n)]$ of the moduli space. The symbol

$$\psi_i = c_1(\mathbb{L}_i) \in H^2(\overline{\mathcal{M}}_{g,n}, \mathbb{Q})$$

denotes its first Chern class. Since $\overline{\mathcal{M}}_{g,n}$ has dimension $3g - 3 + n$, the integral (1.10) is automatically 0 unless $d_1 + \dots + d_n = 3g - 3 + n$.

Surprisingly, on the other side of the rainbow, i.e., when $x > 0$, we see the intersection numbers (1.10)!

Remark 1.3. The relation between the Airy function and the intersection numbers was discovered by Kontsevich [55]. He replaces the variables x and p in (1.1) by *Hermitian matrices*. It is a general property that a large class of Hermitian matrix integrals satisfy an integrable system of KdV and KP type (see, for example, [62]). Because of the cubic polynomial in the integrand, the matrix Airy function of [55] counts trivalent ribbon graphs through the Feynman diagram expansion, which represent open dense subsets of $\overline{\mathcal{M}}_{g,n}$. By identifying the intersection numbers and the Euclidean volume of these open subsets, Kontsevich proves the Witten conjecture [84]. Our formulas (1.8) and (1.9) are a consequence of the results reported in [15, 27, 64, 67]. We will explain the relation more concisely in these lectures.

Remark 1.4. The numerical value of the asymptotic expansion (1.8) is given by

$$(1.11) \quad \begin{aligned} \log Ai(x) = & -\frac{2}{3}x^{\frac{3}{2}} - \frac{1}{4}\log x - \log(2\sqrt{\pi}) \\ & - \frac{5}{48}x^{-\frac{3}{2}} + \frac{5}{64}x^{-3} - \frac{1105}{9216}x^{-\frac{9}{2}} + \frac{565}{2048}x^{-6} - \frac{82825}{98304}x^{-\frac{15}{2}} + \frac{19675}{6144}x^{-9} \\ & - \frac{1282031525}{88080384}x^{-\frac{21}{2}} + \frac{80727925}{1048576}x^{-12} - \frac{1683480621875}{3623878656}x^{-\frac{27}{2}} + \dots \end{aligned}$$

This follows from the WKB analysis of the Airy differential equation, which will be explained in this introduction.

Remark 1.5. Although the asymptotic expansion is not *equal* to the holomorphic function itself, we use the equality sign in these lectures to avoid further cumbersome notations.

The Airy differential equation appears in many different places, showing the feature of a *universal object* in the WKB analysis. It reflects the fact that the intersection numbers (1.10) are the most fundamental objects in Gromov-Witten theory. In contrast to the Airy differential equation, the gamma function is a universal object in the context of *difference equations*. We recall that

$$\Gamma(z + 1) = z\Gamma(z),$$

and its asymptotic expansion for $Re(z) > 0$ is given by

$$(1.12) \quad \log \Gamma(z) = z \log z - z - \frac{1}{2} \log z + \frac{1}{2} \log(2\pi) + \sum_{m=1}^{\infty} \frac{B_{2m}}{2m(2m-1)} z^{-(2m-1)},$$

where B_{2m} is the $(2m)$ -th Bernoulli number defined by the generating function

$$\frac{x}{e^x - 1} = \sum_{n=0}^{\infty} \frac{B_n}{n!} x^n.$$

This is called **Stirling's formula**, and its main part gives the well-known approximation of the factorial:

$$n! \sim \sqrt{2\pi n} \frac{n^n}{e^n}.$$

The asymptotic expansion of the gamma function is deeply related to the moduli theory of algebraic curves. For example, from Harer and Zagier [45] we learn that the orbifold Euler characteristic of the moduli space of smooth algebraic curves is given by the formula

$$(1.13) \quad \chi(\mathcal{M}_{g,n}) = (-1)^{n-1} \frac{(2g-3+n)!}{(2g-2)!n!} \zeta(1-2g),$$

and the special value of the Riemann zeta function is the Bernoulli number

$$\zeta(1-2g) = -\frac{B_{2g}}{2g}.$$

The expression (1.13) is valid for $g = 0, n \geq 3$ if we use the gamma function for $(2g-2)!$. Stirling's formula (1.12) is much simpler than the expansion of $\log Ai(x)$. As the prime factorization of one of the coefficients

$$\frac{1683480621875}{3623878656} = \frac{5^5 \cdot 13 \cdot 17 \cdot 2437619}{2^{27} \cdot 3^3}$$

shows, we do not expect any simple closed formula for the coefficients of (1.11), like Bernoulli numbers. Amazingly, still there is a close relation between these two asymptotic expansions (1.11) and (1.12) through the work on *tautological relations* of Chow classes on the moduli space \mathcal{M}_g by Mumford [69], followed by recent exciting developments on the Faber-Zagier conjecture [36, 49, 76]. In Theorem 2.7, we will see yet another close relationship between the Euler characteristic of $\mathcal{M}_{g,n}$ and the intersection numbers on $\overline{\mathcal{M}}_{g,n}$, through two special values of the *same* function.

The asymptotic expansion of the Airy function $Ai(x)$ itself for $Re(x) > 0$ has actually a rather simple expression:

$$(1.14) \quad \begin{aligned} Ai(x) &= \frac{e^{-\frac{2}{3}x^{\frac{3}{2}}}}{2\sqrt{\pi}x^{\frac{1}{4}}} \sum_{m=0}^{\infty} \frac{\left(-\frac{3}{4}\right)^m \Gamma\left(m + \frac{5}{6}\right) \Gamma\left(m + \frac{1}{6}\right)}{2\pi m!} x^{-\frac{3}{2}m} \\ &= \frac{e^{-\frac{2}{3}x^{\frac{3}{2}}}}{2\sqrt{\pi}x^{\frac{1}{4}}} \sum_{m=0}^{\infty} (-1)^m \left(\frac{1}{576}\right)^m \frac{(6m)!}{(2m)!(3m)!} x^{-\frac{3}{2}m}. \end{aligned}$$

The expansion in terms of the gamma function values of the first line of (1.14) naturally arises from a hypergeometric function. The first line is equal to the second line because

$$\Gamma(z)\Gamma(1-z) = \frac{\pi}{\sin \pi z},$$

and the induction on m . Since the $m = 0$ term in the summation is 1, we can apply the formal logarithm expansion

$$\log(1-X) = -\sum_{j=1}^{\infty} \frac{1}{j} X^j$$

to (1.14) with

$$X = -\sum_{m=1}^{\infty} (-1)^m \left(\frac{1}{576}\right)^m \frac{(6m)!}{(2m)!(3m)!} x^{-\frac{3}{2}m},$$

and obtain

$$(1.15) \quad \begin{aligned} \log Ai(x) &= -\frac{2}{3}x^{\frac{3}{2}} - \frac{1}{4} \log x - \log(2\sqrt{\pi}) \\ &\quad - \frac{1}{576} \frac{6!}{2!3!} x^{-\frac{3}{2}} + \left(\frac{1}{576}\right)^2 \left(\frac{12!}{4!6!} - \frac{1}{2} \left(\frac{6!}{2!3!}\right)^2\right) x^{-3} \\ &\quad - \left(\frac{1}{576}\right)^3 \left(\frac{18!}{6!9!} - \frac{12!}{4!6!} \cdot \frac{6!}{2!3!} + \frac{1}{3} \left(\frac{6!}{2!3!}\right)^3\right) x^{-\frac{9}{2}} + \dots \end{aligned}$$

In general, for $m \geq 1$, we have

$$(1.16) \quad S_{m+1}(x) = (-1)^m x^{-\frac{3}{2}m} \left(\frac{1}{576}\right)^m \sum_{\lambda \vdash m} (-1)^{\ell(\lambda)-1} \frac{(\ell(\lambda)-1)!}{|\text{Aut}(\lambda)|} \prod_{i=1}^{\ell(\lambda)} \frac{(6\lambda_i)!}{(2\lambda_i)!(3\lambda_i)!},$$

where λ is a partition of m , $\ell(\lambda)$ its length, and $\text{Aut}(\lambda)$ is the group of permutations of the parts of λ of equal length.

Comparing (1.9) and (1.16), we establish concrete relations among the intersection numbers.

Theorem 1.6 (Rainbow formula). *The cotangent class intersection numbers of $\overline{\mathcal{M}}_{g,n}$ satisfy the following relation for every $m \geq 1$:*

$$(1.17) \quad \begin{aligned} &\sum_{\substack{g \geq 0, n > 0 \\ 2g-2+n=m}} \frac{1}{n!} \sum_{\substack{d_1 + \dots + d_n \\ = 3g-3+n}} \langle \tau_{d_1} \cdots \tau_{d_n} \rangle_{g,n} \prod_{i=1}^n (2d_i - 1)!! \\ &= \left(\frac{1}{288}\right)^m \sum_{\lambda \vdash m} (-1)^{\ell(\lambda)-1} \frac{(\ell(\lambda)-1)!}{|\text{Aut}(\lambda)|} \prod_{i=1}^{\ell(\lambda)} \frac{(6\lambda_i)!}{(2\lambda_i)!(3\lambda_i)!}. \end{aligned}$$

Here, we use the fact that $(-1)^n = (-1)^m$ if $2g - 2 + n = m$. For example, for $m = 2$, we have

$$\frac{1}{6}\langle\tau_0^3\tau_1\rangle_{0,4} + \frac{1}{2}\langle\tau_1^2\rangle_{1,2} + 3\langle\tau_0\tau_2\rangle_{1,2} = \left(\frac{1}{288}\right)^2 \left(\frac{12!}{4!6!} - \frac{1}{2}\left(\frac{6!}{2!3!}\right)^2\right) = \frac{5}{64}.$$

This can be verified by evaluating

$$\langle\tau_0^3\tau_1\rangle_{0,4} = 1, \quad \langle\tau_1^2\rangle_{1,2} = \langle\tau_0\tau_2\rangle_{1,2} = \frac{1}{24}.$$

Remark 1.7. The main purpose of these lectures is to relate the topological recursion of [34] and quantization of Hitchin spectral curves. The left-hand side of (1.17) represents the topological recursion in this example, since the intersection numbers can be computed through this mechanism, as explained below. Actually, this is an important example that leads to the universal structure of the topological recursion. The right-hand side is the asymptotic expansion of a function that is coming from the geometry of the Hitchin spectral curve of a Higgs bundle.

The structure of the cohomology ring (or more fundamental Chow ring) of the moduli space $\overline{\mathcal{M}}_{g,n}$, and its open part $\mathcal{M}_{g,n}$ consisting of smooth curves, attracted much attention since the publication of the inspiring paper by Mumford [69] mentioned above. Let us focus on a simple situation

$$\pi : \mathcal{M}_{g,1} \longrightarrow \mathcal{M}_g,$$

which *forgets* the marked point on a smooth curve. By gluing the canonical line bundle of the fiber of each point on the base \mathcal{M}_g , which is the curve represented by the point on the moduli, we obtain the relative dualizing sheaf ω on $\mathcal{M}_{g,1}$. Its first Chern class, considered as a divisor on $\mathcal{M}_{g,1}$ and an element of the Chow group $A^1(\mathcal{M}_{g,1})$, is denoted by ψ . In the notation of (1.10), this is the same as ψ_1 . Mumford defines *tautological* classes

$$\kappa_a := \pi_*(\psi^{a+1}) \in A^a(\mathcal{M}_g).$$

One of the statements of the Faber-Zagier conjecture of [36], now a theorem due to Ionel [49], says the following.

Conjecture 1.8 (A part of the Faber-Zagier Conjecture [36]). *Define rational numbers $a_j \in \mathbb{Q}$ by*

$$(1.18) \quad \sum_{j=1}^{\infty} a_j t^j = -\log \left(\sum_{m=0}^{\infty} \frac{(6m)!}{(2m)!(3m)!} t^m \right).$$

Then the coefficient of t^ℓ of the expression

$$\exp \left(\sum_{j=1}^{\infty} a_j \kappa_j t^j \right) \in (\mathbb{Q}[\kappa_1, \kappa_2, \dots])[t]$$

for each $\ell \geq 1$ gives the unique codimension ℓ tautological relation among the κ -classes on the moduli space $\mathcal{M}_{3\ell-1}$.

We see from (1.17), these coefficients a_j 's are given by the intersection numbers (1.10), by a change of the variables

$$t = -\frac{1}{576} x^{-\frac{3}{2}}.$$

Indeed, we have

$$(1.19) \quad a_j = -288^j \sum_{\substack{g \geq 0, n > 0 \\ 2g-2+n=j}} \frac{1}{n!} \sum_{\substack{d_1 + \dots + d_n \\ = 3g-3+n}} \langle \tau_{d_1} \cdots \tau_{d_n} \rangle_{g,n} \prod_{i=1}^n (2d_i - 1)!! .$$

These tautological relations are generalized for the moduli spaces $\overline{\mathcal{M}}_{g,n}$, and are proved in [76]. Amazingly, still the rainbow integral (1.1) plays the essential role in determining the tautological relations in this generalized work.

Remark 1.9. As we have seen above, the asymptotic expansions of the gamma function and the Airy function carry information of quantum invariants, in particular, certain topological information of $\mathcal{M}_{g,n}$ and $\overline{\mathcal{M}}_{g,n}$. We note that these quantum invariants are stored in the **insignificant** part of the asymptotic expansions.

Here come questions.

Question 1.10. *The Airy function is a one-variable function. It cannot be a generating function of all the intersection numbers (1.10). Then how do we obtain all intersection numbers from the Airy function, or the Airy differential equation, alone?*

Question 1.11. *The relations between the Airy function, the gamma function, and intersection numbers are all great. But then what does this relation have anything to do with Higgs bundles?*

Question 1.12. *As we remarked, the information of the quantum invariants is stored in the insignificant part of the asymptotic expansion. In the Airy example, they correspond to $S_m(x)$ of (1.9) for $m \geq 2$. Then what does the main part of the asymptotic behavior of the function, i.e., those functions in (1.7), determine?*

As we have remarked earlier, Kontsevich [55] utilized the matrix integral technique to answer Question 1.10. The key idea is to replace the variables in (1.1) by Hermitian matrices, and then use the asymptotic expansion on the result. Through the Feynman diagram expansion, he was able to obtain a generating function of all the intersection numbers.

What we explain in these lectures is the concept of **topological recursion** of [34]. Without going into matrix integrals, we can directly obtain (a totally different set of) generating functions of the intersection numbers from the Airy function. Here, the Airy differential equation is identified as a **quantum curve**, and application of the **semi-classical limit** and the topological recursion enable us to calculate generating functions of the intersection numbers.

But before going into detail, let us briefly answer Question 1.11 below. The point is that the geometry of the Airy function is a special example of Higgs bundles.

For the example of the Airy differential equation, the topological recursion is exactly the same as the *Virasoro constraint conditions* for the intersection numbers (1.10), and the semi-classical limit recovers the **Hitchin spectral curve** of the corresponding Higgs bundle. The information stored in the main part of the asymptotic expansion (1.7), as asked in Question 1.12, actually determines the spectral curve and its geometry. We can turn the story in the other way around: we will see that the functions $S_0(x)$ and $S_1(x)$ corresponding to (1.7) in the general context are indeed determined by the geometry of the Hitchin spectral curve of an appropriate Higgs bundle.

The stage setting is the following. As a base curve, we have \mathbb{P}^1 . On this curve we have a vector bundle

$$(1.20) \quad E = \mathcal{O}_{\mathbb{P}^1}(1) \oplus \mathcal{O}_{\mathbb{P}^1}(-1) = K_{\mathbb{P}^1}^{-\frac{1}{2}} \oplus K_{\mathbb{P}^1}^{\frac{1}{2}}$$

of rank 2. The main character of the Second Act is a *meromorphic* Higgs field

$$(1.21) \quad \phi = \begin{bmatrix} & 1 \\ x(dx)^2 & \end{bmatrix} : E \longrightarrow K_{\mathbb{P}^1}(4) \otimes E.$$

Here, x is an affine coordinate of $\mathbb{A}^1 \subset \mathbb{P}^1$, 1 on the (1,2)-component of ϕ is the natural morphism

$$1 : K_{\mathbb{P}^1}^{\frac{1}{2}} \xrightarrow{\sim} K_{\mathbb{P}^1}^{\frac{1}{2}} \longrightarrow K_{\mathbb{P}^1}^{\frac{1}{2}} \otimes \mathcal{O}_{\mathbb{P}^1}(4),$$

and

$$x(dx)^2 \in H^0(\mathbb{P}^1, K_{\mathbb{P}^1}^{\otimes 2}(4)) \cong \mathbb{C}$$

is the unique (up to a constant factor) meromorphic *quadratic differential* on \mathbb{P}^1 that has one zero at $x = 0$ and a pole of order 5 at $x = \infty$. We use K_C to denote the canonical sheaf on a projective algebraic curve C . The data (E, ϕ) is called a *Higgs pair*. Although ϕ contains a quadratic differential in its component, because of the shape of the vector bundle E , we see that

$$E = K_{\mathbb{P}^1}^{-\frac{1}{2}} \oplus K_{\mathbb{P}^1}^{\frac{1}{2}} \xrightarrow{\phi} \left(K_{\mathbb{P}^1}^{\frac{1}{2}} \oplus K_{\mathbb{P}^1}^{\frac{3}{2}} \right) \otimes \mathcal{O}_{\mathbb{P}^1}(4) = K_{\mathbb{P}^1}(4) \otimes \left(K_{\mathbb{P}^1}^{-\frac{1}{2}} \oplus K_{\mathbb{P}^1}^{\frac{1}{2}} \right),$$

hence

$$\phi \in H^0(\mathbb{P}^1, K_{\mathbb{P}^1}(4) \otimes \text{End}(E))$$

is indeed an $\text{End}(E)$ -valued meromorphic 1-form on \mathbb{P}^1 .

The cotangent bundle

$$\pi : T^*\mathbb{P}^1 \longrightarrow \mathbb{P}^1$$

is the total space of $K_{\mathbb{P}^1}$. Therefore, the pull-back bundle $\pi^*K_{\mathbb{P}^1}$ has a tautological section $\eta \in H^0(T^*\mathbb{P}^1, \pi^*K_{\mathbb{P}^1})$, which is a globally holomorphic 1-form on $T^*\mathbb{P}^1$. The global holomorphic 2-form $-d\eta$ defines the holomorphic symplectic structure, hence a hyper-Kähler structure, on $T^*\mathbb{P}^1$. If we trivialize the cotangent bundle on the affine neighborhood of \mathbb{P}^1 with a coordinate x , and use a fiber coordinate y , then $\eta = ydx$. As usual, we wish to define the spectral curve of the Higgs pair. Due to the fact that ϕ is singular at $x = \infty$, we cannot capture the whole story within the cotangent bundle. We note that the cotangent bundle $T^*\mathbb{P}^1$ has a natural compactification

$$\overline{T^*\mathbb{P}^1} := \mathbb{P}(K_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}) = \mathbb{F}_2 \xrightarrow{\pi} \mathbb{P}^1,$$

which is known as a Hirzebruch surface. The holomorphic 1-form η extends to $\overline{T^*\mathbb{P}^1}$ as a meromorphic 1-form with simple poles along the divisor at infinity.

Now we can consider the characteristic polynomial

$$\det(\eta - \pi^*\phi) \in H^0\left(\overline{T^*\mathbb{P}^1}, \pi^*(K_{\mathbb{P}^1}^{\otimes 2}(4))\right)$$

as a meromorphic section of the line bundle $\pi^*K_{\mathbb{P}^1}^{\otimes 2}$ on the compact space $\overline{T^*\mathbb{P}^1}$. It defines the Hitchin spectral curve

$$(1.22) \quad \Sigma = (\det(\eta - \pi^*\phi))_0 \subset \overline{T^*\mathbb{P}^1}$$

as a divisor. Again in terms of the local coordinate (x, y) of $\overline{T^*\mathbb{P}^1}$, the spectral curve Σ is simply given by

$$(1.23) \quad x = y^2.$$

It is a perfect parabola in the (x, y) -plane. But our Σ is in the Hirzebruch surface, not in the projective plane. Choose the coordinate $(u, w) \in \mathbb{F}_2$ around $(x, y) = (\infty, \infty)$ defined by

$$(1.24) \quad \begin{cases} u = \frac{1}{x} \\ \frac{1}{w} du = y dx \end{cases} .$$

Then the local expression of Σ around $(u, w) = (0, 0)$ becomes a quintic cusp

$$(1.25) \quad w^2 = u^5.$$

So the spectral curve Σ is indeed highly singular at the infinity!

1.2. Quantum curves, semi-classical limit, and the WKB analysis. At this stage we come to the notion of *quantization*. We wish to quantize the spectral curve Σ of (1.22). In terms of the affine coordinate (x, y) , the **quantum curve** of $y^2 - x = 0$ should be

$$(1.26) \quad \left(\left(\hbar \frac{d}{dx} \right)^2 - x \right) \Psi(x, \hbar) = 0,$$

which changes the commutative algebra $\mathbb{C}[x, y]$ to a Weyl algebra $\mathbb{C}[\hbar]\langle x, y \rangle$ defined by the commutation relation

$$(1.27) \quad [x, y] = -\hbar.$$

We consider $x \in \mathbb{C}[\hbar]\langle x, y \rangle$ as the multiplication operator by the coordinate x , and $y = \hbar \frac{d}{dx}$ as a differential operator.

How do we know that (1.26) is the right quantization of the spectral curve (1.23)? Apparently, the limit $\hbar \rightarrow 0$ of the differential operator does not reproduce the spectral curve. Let us now recall the WKB method for analyzing differential equations like (1.26). This is a method that relates classical mechanics and quantum mechanics. As we see below, the WKB method is not for finding a convergent analytic solution to the differential equation. Since the equation we wish to solve is considered to be a quantum equation, the corresponding classical problem, if it exists, should be recovered by taking $\hbar \rightarrow 0$. We denote by an unknown function $S_0(x)$ the ‘‘solution’’ to the corresponding classical problem. To emphasize the classical behavior at the $\hbar \rightarrow 0$ limit, we expand the solution to the quantum equation as

$$(1.28) \quad \Psi(x, \hbar) = \exp \left(\sum_{m=0}^{\infty} \hbar^{m-1} S_m(x) \right) := \exp \left(\frac{1}{\hbar} S_0(x) \right) \cdot \exp \left(\sum_{m=1}^{\infty} \hbar^{m-1} S_m(x) \right).$$

The idea is that as $\hbar \rightarrow 0$, the effect of $S_0(x)$ is magnified. But as a series in \hbar , (1.28) is ill defined because the coefficient of each power of \hbar is an infinite sum. It is also clear that $\hbar \rightarrow 0$ does not make sense for $\Psi(x, \hbar)$. Instead of expanding (1.28) immediately in \hbar and take its 0 limit, we use the following standard procedure. First we note that (1.26) is equivalent to

$$\left[\exp \left(-\frac{1}{\hbar} S_0(x) \right) \cdot \left(\left(\hbar \frac{d}{dx} \right)^2 - x \right) \cdot \exp \left(\frac{1}{\hbar} S_0(x) \right) \right] \exp \left(\sum_{m=1}^{\infty} \hbar^{m-1} S_m(x) \right) = 0.$$

Since the conjugate differential operator

$$\exp \left(-\frac{1}{\hbar} S_0(x) \right) \cdot \left(\left(\hbar \frac{d}{dx} \right)^2 - x \right) \cdot \exp \left(\frac{1}{\hbar} S_0(x) \right)$$

$$= \left(\hbar \frac{d}{dx} \right)^2 + 2\hbar S'_0(x) \frac{d}{dx} + (S'_0(x))^2 - x + \hbar S''_0(x)$$

is a well-defined differential operator, *and* its limit $\hbar \rightarrow 0$ makes sense, we interpret (1.26) as the following differential equation:

$$(1.29) \quad \left[\left(\hbar \frac{d}{dx} \right)^2 + 2\hbar S'_0(x) \frac{d}{dx} + (S'_0(x))^2 - x + \hbar S''_0(x) \right] \exp \left(\sum_{m=1}^{\infty} \hbar^{m-1} S_m(x) \right) = 0.$$

Here, $'$ indicates the x -derivative. This equation is equivalent to

$$(1.30) \quad \left(\sum_{m=0}^{\infty} \hbar^m S'_m(x) \right)^2 + \sum_{m=0}^{\infty} \hbar^{m+1} S''_m(x) - x = 0$$

for every $m \geq 0$. The coefficient of the \hbar^0 , or the $\hbar \rightarrow 0$ limit of (1.30), then gives

$$(1.31) \quad (S'_0(x))^2 - x = 0,$$

and that of \hbar^1 gives

$$(1.32) \quad S''_0(x) + 2S'_0(x)S'_1(x) = 0.$$

The \hbar^0 term is what we call the **semi-classical limit** of the differential equation (1.26). From (1.31) we obtain

$$(1.33) \quad S_0(x) = \pm \frac{2}{3} x^{\frac{3}{2}} + c_0,$$

with a constant of integration c_0 . Then plugging $S_0(x)$ into (1.32) we obtain

$$S_1(x) = -\frac{1}{4} \log x - \log(2\sqrt{\pi}) + c_1,$$

again with a constant of integration c_1 . Note that these solutions are consistent with (1.7). For $m \geq 1$, the coefficient of \hbar^{m+1} gives

$$(1.34) \quad S'_{m+1}(x) = -\frac{1}{2S'_0(x)} \left(S''_m(x) + \sum_{a=1}^m S'_a(x) S'_{m+1-a}(x) \right),$$

which can be solved recursively one by one from $S_0(x)$. This mechanism is the method of Wentzel-Kramers-Brillouin (WKB) approximation.

We can ignore the constants of integration when solving (1.34) because it is easy to restore them, if necessary, just by adding c_m to each $S_m(x)$ in (1.28). The solution then simply changes to another one

$$\left(\exp \left(\frac{1}{\hbar} c_0 \right) \exp \left(\frac{1}{\hbar} S_0(x) \right) \right) \cdot \exp \left(\sum_{m=1}^{\infty} \hbar^{m-1} c_m \right) \exp \left(\sum_{m=1}^{\infty} \hbar^{m-1} S_m(x) \right).$$

In terms of the main variable x , the above solution is a constant multiple of the original one. The two choices of the sign in (1.33) lead to two linearly independent solutions of (1.26). If we impose

$$(1.35) \quad \lim_{x \rightarrow \infty} S_m(x) = 0, \quad m \geq 2,$$

then the differential equation (1.34) uniquely determines all terms $S_m(x)$. Thus the choice of the negative sign in (1.33) and imposing $c_0 = c_1 = 0$ and (1.35), we obtain the unique exponentially decaying solution for $x \rightarrow \infty$ along the real line. This solution agrees with (1.9) and (1.16). Thus we obtain the second line of the Rainbow formula (1.17).

We also see from the semi-classical limit (1.31) that if we put $y = S'_0(x)$, then we recover the Hitchin spectral curve $x = y^2$. The functions $S_m(x)$ actually live on the spectral curve rather than the base \mathbb{P}^1 , because of the appearance of \sqrt{x} in (1.9).

Remark 1.13. One can ask a question: **Does (1.28) give a convergent solution?** The answer is a flat **No!** Suppose we solve the Airy differential equation with the WKB method explained above, and define a “solution” by (1.28). Expand the second factor of exponential as a power series in \hbar , and write the solution as

$$\Psi(x, \hbar) = \exp\left(\frac{1}{\hbar}S_0(x)\right) \sum_{n=0}^{\infty} \psi_n(x)\hbar^n.$$

Then for any compact subset $K \subset \mathbb{C} \setminus \{0\}$, there is a constant C_K such that

$$\sup_{x \in K} |\psi_n(x)| \leq C_K^n n!.$$

Therefore, unless we are in an extremely special case, the second exponential factor in the expression (1.28) does not converge as a power series in \hbar at all! Since

$$\Psi(x, \hbar) := Ai\left(x/\hbar^{\frac{2}{3}}\right)$$

is a solution that is entire in x and any \hbar for which $1/\hbar^{\frac{2}{3}}$ makes sense, the WKB method around $x \in \mathbb{C} \setminus \{0\}$ is the same as the asymptotic expansion of the Airy function $Ai(x)$ given in (1.14). There we see the factorial growth of the coefficients. Thus the WKB method is **not** for finding a convergent analytic solution.

1.3. The topological recursion as quantization. Then what is good about the WKB method and the purely formal solution (1.28)? Let us examine (1.9). We note that $S_m(x)$'s are one variable functions, and different values of g and n are summed in its definition. Therefore, knowing the solution $\Psi(x, \hbar)$ of (1.26) that decays exponentially as $x \rightarrow \infty$ along the real axis, assuming $\hbar > 0$, does not seem to possibly recover all values of the intersection numbers $\langle \tau_{d_1} \cdots \tau_{d_n} \rangle_{g,n}$ for all values of (d_1, \dots, d_n) and (g, n) . Then how much information does the quantum curve (1.26) really have?

Here comes the idea of **topological recursion** of Eynard and Orantin [34]. This mechanism gives a refined expression of each $S_m(x)$, and computes all intersection numbers. The solution Ψ of (1.28) is never holomorphic, and it makes sense only as the asymptotic expansion of a holomorphic solution at its essential singularity. The expansion of the Airy function $Ai(x)$ at its holomorphic point does not carry any interesting information. The function's key information is concentrated in the expansion at the essential singularity. The topological recursion is for obtaining this hidden information when applied at the essential singularity of the solution, by giving an explicit formula for the WBK expansion. And the WKB analysis is indeed a method that determines the relation between the quantum behavior and the classical behavior of a given system, i.e., the process of quantization.

As we have seen above, the quantum curve (1.26) recovers the spectral curve (1.23) by the procedure of semi-classical limit. We recall that the spectral curve lives in the Hirzebruch surface \mathbb{F}_2 , and it has a quintic cusp singularity (1.25) at $(x, y) = (\infty, \infty)$. It requires twice blow-ups of \mathbb{F}_2 to resolve the singularity of Σ . Let us denote this minimal resolution by $\tilde{\Sigma}$. The proper transform is a smooth curve of genus 0, hence it is a \mathbb{P}^1 . Let $B \cong \mathbb{P}^1$ be the 0-section, and $F \cong \mathbb{P}^1$ a fiber, of \mathbb{F}_2 . Then after two blow-ups, $\tilde{\Sigma} \subset Bl(\mathbb{F}_2)$ is identified as a divisor by the equation

$$\tilde{\Sigma} = 2B + 5F - 4E_2 - 2E_1 \in \text{Pic}(Bl(\mathbb{F}_2)),$$

where E_i is the exceptional divisor introduced at the i -th blow-up [25, Section 5].




FIGURE 1.3. Twice blow-up of \mathbb{F}_2 . The parabola in \mathbb{F}_2 has a quintic cusp singularity at the infinity (top). After the second blow-up, the proper transform becomes non-singular (bottom). Since all fibers of $\mathbb{F}_2 \rightarrow \mathbb{P}^1$ are equivalent, the toric picture has only one representative of the fiber $F = \mathbb{P}^1$. It is a stretch to place Σ and $\tilde{\Sigma}$ in this diagram, because the spectral curve is a double cover of the base B , rather than a degree 4 covering that the picture may suggest.

Since the desingularization $\tilde{\Sigma}$ is just a copy of a \mathbb{P}^1 , we can choose a **normalization coordinate** t on it so that the map $\tilde{\pi} : \tilde{\Sigma} \rightarrow \mathbb{P}^1$ to the base curve \mathbb{P}^1 is given by

$$(1.36) \quad \begin{cases} x = \frac{4}{t^2} \\ y = -\frac{2}{t} \end{cases}, \quad \begin{cases} u = \frac{t^2}{4} \\ w = \frac{t^5}{32} \end{cases}.$$

With respect to the normalization coordinate, define a homogeneous polynomial of degree $6g - 6 + 3n$ for $2g - 2 + n > 0$ by

$$(1.37) \quad F_{g,n}^A(t_1, \dots, t_n) := \frac{(-1)^n}{2^{2g-2+n}} \sum_{\substack{d_1 + \dots + d_n \\ = 3g-3+n}} \langle \tau_{d_1} \dots \tau_{d_n} \rangle_{g,n} \prod_{i=1}^n (2d_i - 1)!! \left(\frac{t_i}{2}\right)^{2d_i+1}$$

as a function on $(\tilde{\Sigma})^n$, and an n -linear differential form

$$(1.38) \quad W_{g,n}^A(t_1, \dots, t_n) := dt_1 \dots dt_n F_{g,n}^A(t_1, \dots, t_n).$$

For *unstable geometries* $(g, n) = (0, 1)$ and $(0, 2)$, we only define differential forms separately:

$$(1.39) \quad W_{0,1}^A(t) := \eta = ydx = \frac{16}{t^4} dt,$$

$$(1.40) \quad W_{0,2}^A(t_1, t_2) := \frac{dt_1 \cdot dt_2}{(t_1 - t_2)^2}.$$

The definition of $W_{0,1}^A$ encodes the geometry of the singular spectral curve $\Sigma \subset \mathbb{F}_2$ embedded in the Hirzebruch surface, and $W_{0,2}^A$ depends only on the intrinsic geometry of the normalization $\tilde{\Sigma}$. Then we have

Theorem 1.14 (Topological recursion for the intersection numbers, [27]).

$$(1.41) \quad W_{g,n}^A(t_1, \dots, t_n) = -\frac{1}{2\pi i} \int_{\gamma} \left(\frac{1}{t+t_1} + \frac{1}{t-t_1} \right) \frac{t^4}{64} \frac{1}{dt} dt_1 \left[W_{g-1, n+1}^A(t, -t, t_2, \dots, t_n) + \sum_{\substack{\text{no } (0,1) \text{ terms} \\ g_1+g_2=g \\ I \sqcup J = \{2, \dots, n\}}} W_{g_1, |I|+1}^A(t, t_I) W_{g_2, |J|+1}^A(-t, t_J) \right],$$

where the integral is taken with respect to the contour of Figure 1.4, and the sum is over all partitions of g and set partitions of $\{2, 3, \dots, n\}$ without including $g_1 = 0$ and $I = \emptyset$, or $g_2 = 0$ and $J = \emptyset$. The notation $\frac{1}{dt}$ represents the contraction of the quadratic differential proportional to $(dt)^2$ by the vector field $\frac{\partial}{\partial t}$.




FIGURE 1.4. The integration contour γ . This contour encloses an annulus bounded by two concentric circles centered at the origin. The outer one is a small circle around $\infty \in \tilde{\Sigma} = \mathbb{P}^1$, and the inner one a small circle around the origin. Both circles are positively oriented on $\tilde{\Sigma}$. Using the t -plane as the affine chart, the small loop around ∞ becomes a circle of a large radius $r \gg \infty$ with the opposite orientation, as in the figure. Geometrically, $t = 0$ and $t = \infty$ correspond to the two simple ramification points of the Galois covering $\tilde{\pi} : \tilde{\Sigma} \rightarrow \mathbb{P}^1$, and the contour γ consists of two small loops around these ramification points.

This is an example of the axiomatic mechanism of [34] called the *topological recursion*. We give a more geometric interpretation of the formula in the later sections. The derivation of the topological recursion (1.41) from (1.37) is the subject of Section 2 where the origin of the recursion formula is identified via graph enumeration. The power of this formula is

that all $W_{g,n}^A$ for $2g - 2 + n > 0$ are calculated from the initial values (1.39) and (1.40). For example,

$$\begin{aligned} W_{1,1}^A(t_1) &= - \left[\frac{1}{2\pi i} \int_{\gamma} \left(\frac{1}{t+t_1} + \frac{1}{t-t_1} \right) \frac{t^4}{64} \frac{1}{dt} W_{0,2}^A(t, -t) \right] dt_1 \\ &= - \left[\frac{1}{2\pi i} \int_{\gamma} \left(\frac{1}{t+t_1} + \frac{1}{t-t_1} \right) \frac{t^4}{64} \frac{(-dt)}{4t^2} \right] dt_1 \\ &= - \frac{1}{128} t_1^2 dt_1 \\ &= - \frac{3}{16} \langle \tau_1 \rangle_{1,1} t_1^2 dt_1. \end{aligned}$$

Thus we find $\langle \tau_1 \rangle_{1,1} = \frac{1}{24}$.

The functions $F_{g,n}^A$ for $2g - 2 + n > 0$ can be calculated by integration:

$$(1.42) \quad F_{g,n}^A(t_1, \dots, t_n) = \int_0^{t_1} \cdots \int_0^{t_n} W_{g,n}^A(t_1, \dots, t_n).$$

Note that

$$t \rightarrow 0 \iff x \rightarrow \infty.$$

Therefore, we are considering the expansion of quantities at the essential singularity of $Ai(x)$. It is surprising to see that the topological recursion indeed determines all intersection numbers (1.10)! Now define

$$(1.43) \quad S_m(x) = \sum_{2g-2+n=m-1} \frac{1}{n!} F_{g,n}^A(t(x), \dots, t(x)),$$

where we choose a branch of $\pi : \Sigma \rightarrow \mathbb{P}^1$, and consider $t = t(x)$ as a function in x . It coincides with (1.9). Since the *moduli* spaces $\overline{\mathcal{M}}_{0,1}$ and $\overline{\mathcal{M}}_{0,2}$ do not exist, we do not have an expression (1.37) for these unstable geometries. So let us formally apply (1.42) to (1.39):

$$(1.44) \quad F_{0,1}^A(t) := \int_0^t \frac{16}{t^4} dt = -\frac{16}{3} t^{-3} = -\frac{2}{3} x^{\frac{3}{2}} = S_0(x).$$

We do not have this type of integration procedure to produce $S_1(x)$ from $W_{0,2}^A$. So we simply define $S_1(x)$ by solving (1.32). Then the residues in (1.41) can be concretely computed, and produce a system of recursive partial differential equations among $F_{g,n}^A$'s. Their **principal specialization** (1.43) produces (1.34)! Therefore, we obtain (1.17).

In this context, the topological recursion is the process of quantization, because it actually constructs the function $\Psi(x, \hbar)$ by giving a closed formula for the WKB analysis, and hence the differential operator (1.26) that annihilates it, all from the classical curve $x = y^2$.

1.4. Non-Abelian Hodge correspondence and quantum curves. Then what is the quantum curve? Since it is a second order differential equation with a deformation parameter \hbar , and its semi-classical limit is the spectral curve of a Higgs bundle, which is a rank 2 bundle in our example, we can easily *guess* that it should be the result of the **non-Abelian Hodge correspondence**. Let us consider an \hbar -connection

$$(1.45) \quad \nabla^{\hbar} = \hbar d + \phi : E \rightarrow K_{\mathbb{P}^1}(4) \otimes E,$$

which gives the Higgs field at $\hbar = 0$. A flat section with respect to ∇^{\hbar} for $\hbar \neq 0$ can be obtained by solving

$$(1.46) \quad \left(\hbar \frac{d}{dx} + \begin{bmatrix} & 1 \\ x & \end{bmatrix} \right) \begin{bmatrix} \Psi(x, \hbar) \\ -\hbar \Psi(x, \hbar)' \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix},$$

where $\Psi(x, \hbar)'$ is the x -derivative. Clearly, (1.46) is equivalent to (1.26). For $\hbar = 1$, (1.45) is a holomorphic flat connection on $E|_{\mathbb{A}^1}$, the restriction of the vector bundle E on the affine coordinate neighborhood. Therefore, $(E, \nabla^{\hbar=1})$ defines a \mathcal{D} -module on $E|_{\mathbb{A}^1}$. As a holomorphic \mathcal{D} -module over \mathbb{A}^1 , we have an isomorphism

$$(E, \nabla^{\hbar=1})|_{\mathbb{A}^1} \cong \mathcal{D}/(\mathcal{D} \cdot P(x, 1)),$$

where

$$(1.47) \quad P(x, \hbar) := \left(\hbar \frac{d}{dx} \right)^2 - x,$$

and \mathcal{D} denotes the sheaf of linear differential operators with holomorphic coefficients on \mathbb{A}^1 .

Hitchin's original idea [46] of constructing Higgs bundles is to solve a system of differential equations, now known as *Hitchin's equations*,

$$(1.48) \quad \begin{aligned} F(D) - [\phi, \phi] &= 0 \\ D\phi &= 0 \\ D * \phi &= 0, \end{aligned}$$

defined on a **Hermitian** vector bundle $E \rightarrow C$ over a compact Riemann surface C . Here, D is a unitary connection in E , $F(D)$ is its curvature 2-form, $*$ is the Hodge $*$ -operator with respect to the metric data on both C and E , and ϕ is a differentiable Higgs field on E . Solving Hitchin's equation is equivalent to constructing a 1-parameter family of flat connections of the form

$$(1.49) \quad \nabla(\zeta) = \frac{\phi}{\zeta} + D + \phi^\dagger \zeta$$

that satisfies the adjoint condition

$$(1.50) \quad \nabla(\zeta) = \nabla(-1/\bar{\zeta})^\dagger$$

(see [38, 70, 79]). Here, ϕ^\dagger is the Hermitian conjugation of ϕ , and ∇^\dagger is the adjoint connection with respect to the Hermitian metric in E . The non-Abelian Hodge correspondence is the association of this 1-parameter family of flat connections (called a *real twister line*) to a given spectral curve.

A new idea that relates the non-Abelian Hodge correspondence,opers, and the topological recursion, is emerging [23]. For the case of $SU(2)$ -Higgs bundle E over $\mathbb{A}^1 \subset \mathbb{P}^1$, let us use the matrix ϕ of (1.21) as a Higgs field, and

$$(1.51) \quad H_R = \begin{bmatrix} |x|^{\frac{1}{2}} e^{h_R(|x|)} & \\ & |x|^{-\frac{1}{2}} e^{-h_R(|x|)} \end{bmatrix}$$

as the Hermitian metric in E . Here, $h_R(t)$ is a solution to the Painlevé III equation

$$(1.52) \quad \frac{1}{4} \left(h_R''(t) + \frac{1}{t} h_R'(t) \right) = 2R^2 t \sinh(2h_R(t)),$$

and $R \in \mathbb{R}_+$ is a positive real scaling parameter of the Higgs field. Then the real twister line $\nabla(\zeta)$ for a scaled Higgs field $R\phi$ is given by

$$(1.53) \quad \nabla(\zeta) = \frac{R}{\zeta}\phi + d - H_R^{-1}\bar{\partial}H_R + \zeta R\phi^\dagger.$$

Now we consider the **scaling limit**

$$(1.54) \quad R \longrightarrow 0, \quad \zeta \longrightarrow 0, \quad \frac{R}{\zeta} = \hbar.$$

At this scaling limit, the right-hand side of (1.52) vanishes, so we can choose

$$h_R(|x|) = -\frac{1}{2}\log|x|$$

for defining the Hermitian metric H_R at $R = 0$. Then from (1.51), we have a scaled connection

$$\hbar\nabla(0) = \hbar\frac{d}{dx} + \phi.$$

This is the \hbar -connection of (1.45)! It is also known as an **oper**.

The above consideration suggests the following:

Conjecture 1.15. *The topological recursion gives a particular scaling limit of the non-Abelian Hodge correspondence, which is the oper corresponding to the starting spectral curve.*

For a particular section of the Hitchin fibration known as the Hitchin section, the non-Abelian Hodge correspondence and the oper are indeed connected by a natural scaling limit [23]. The relation with the topological recursion beyond rank 2 case is an uncharted territory.

So far, we have considered $\Psi(x, \hbar)$ as a formal “function.” What is it geometrically? Since our vector bundle E is of the form (1.20), the shape of the equation (1.46) suggests that

$$(1.55) \quad \Psi(x, \hbar) \in \widehat{H}^0(\mathbb{P}^1, K_{\mathbb{P}^1}^{-\frac{1}{2}}),$$

where the hat sign indicates that we really do not have any good space to store the formal wave function $\Psi(x, \hbar)$. The reason for the appearance of $K_{\mathbb{P}^1}^{-\frac{1}{2}}$ as its home is understood as follows. Since $W_{g,n}^A$ for $2g - 2 + n$ is an n -linear differential form, (1.42) tells us that $F_{g,n}^A$ is just a number. Therefore, $S_m(x)$ that is determined by (1.43) for $m \geq 2$ is also just a number. Here, by a “number” we mean a genuine function in x . The differential equation (1.31) should be written

$$(1.56) \quad (dS_0(x))^{\otimes 2} = x(dx)^2 = q(x)$$

as an equation of quadratic differentials. Here,

$$q(x) \in K_{\mathbb{P}^1}^{\otimes 2}$$

can be actually any meromorphic quadratic differential on \mathbb{P}^1 , including $x(dx)^2$, so that

$$\phi = \begin{bmatrix} & 1 \\ q & \end{bmatrix} \in K_{\mathbb{P}^1} \otimes \text{End}(E)$$

is a meromorphic Higgs field on the vector bundle E in the same way. Similarly, (1.32) should be interpreted as

$$(1.57) \quad S_1(x) = -\frac{1}{2} \int d \log(dS_0) = -\frac{1}{2} \log \sqrt{q(x)} = -\frac{1}{4} \log q(x).$$

Now recall the conjugate differential equation (1.29). Its solution takes the form

$$(1.58) \quad \frac{1}{\sqrt[4]{q(x)}} \exp \left(\sum_{m=2}^{\infty} S_m(x) \hbar^{m-1} \right),$$

and as we have noted, the exponential factor is just a number. Therefore, the geometric behavior of this solution is determined by the factor

$$\frac{1}{\sqrt[4]{q(x)}} \in K_{\mathbb{P}^1}^{-\frac{1}{2}},$$

which is a meromorphic section of the negative half-canonical sheaf.

We recall that $\Psi(x, \hbar)$ has another factor $\exp(S_0(x)/\hbar)$. It should be understood as a “number” defined on the spectral curve Σ , because $dS_0(x) = \eta$ is the tautological 1-form on $T^*\mathbb{P}^1$ restricted on the spectral curve, and $S_0(x)$ is its integral on the spectral curve. Therefore, this factor tells us that the equation (1.26) should be considered on Σ . Yet its local x -dependence is indeed determined by $K_{\mathbb{P}^1}^{-\frac{1}{2}}$.

We have thus a good answer for Question 1.12 now. The main part of the asymptotic expansion (1.7) tells us what geometry we should consider. It tells us what the Hitchin spectral curve should be, and it also includes the information of Higgs bundle (E, ϕ) itself.

Remark 1.16. The Airy example, and another example we consider in the next section, are in many ways special, in the sense that the $S_2(x)$ -term of the WKB expansion is given by integrating the solutions $W_{1,1}$ and $W_{0,3}$ of the topological recursion (1.41). In general, the topological recursion mechanism of computing $W_{1,1}$ and $W_{0,3}$ from $W_{0,2}$ does not correspond to the WKB equation for S_2 . As discovered in [24], the topological recursion in its PDE form is equivalent to the WKB equations for all $S_m(x)$ in the range of $m \geq 3$. But the PDE recursion, which we discuss in detail in the later sections, does not determine S_2 . This fact reflects the algebraic geometry of Higgs bundles for the case of Airy function. Over \mathbb{P}^1 , the vector bundle E of (1.20) does not deform locally. For a higher genus situation, the S_2 term contains the information of E as an extension of $K_C^{\frac{1}{2}}$. Thus it cannot be obtained from the geometry of (E, ϕ) alone.

1.5. The Lax operator for Witten-Kontsevich KdV equation. Surprisingly, the operator $P(x, 1)$ of (1.47) at $\hbar = 1$ is the initial value of the *Lax operator* for the KdV equations that appears in the work of Witten [84] and Kontsevich [55]. Witten considered a different generating function of the intersection numbers (1.10) given by

$$(1.59) \quad \begin{aligned} F(s_0, s_1, s_2, \dots) &= \left\langle \exp \left(\sum_{d=0}^{\infty} s_d \tau_d \right) \right\rangle \\ &= \sum_{k_0, k_1, k_2, \dots=0}^{\infty} \left\langle \tau_0^{k_0} \tau_1^{k_1} \tau_2^{k_2} \dots \right\rangle \prod_{j=0}^{\infty} \frac{s_j^{k_j}}{k_j!} \\ &= \sum_{g=0}^{\infty} \sum_{n=1}^{\infty} \frac{1}{n!} \sum_{\substack{d_1 + \dots + d_n \\ = 3g - 3 + n}} \langle \tau_{d_1} \dots \tau_{d_n} \rangle_{g,n} s_{d_1} \dots s_{d_n} \\ &= \langle \tau_0^3 \rangle_{0,3} \frac{s_0^3}{3!} + \langle \tau_1 \rangle_{1,1} s_1 + \langle \tau_0^4 \rangle_{0,4} \frac{s_0^4}{4!} + \dots \end{aligned}$$

Define

$$t_{2j+1} := \frac{s_j}{(2j+1)!!},$$

and

$$(1.60) \quad u(t_1, t_3, t_5, \dots) := \left(\frac{\partial}{\partial s_0} \right)^2 F(s_0, s_1, s_2, \dots).$$

Then $u(t_1, t_3, t_5, \dots)$ satisfies the system of KdV equations, whose first equation is

$$(1.61) \quad ut_3 = \frac{1}{4}u_{t_1 t_1 t_1} + 3uu_{t_1}.$$

The system of KdV equations are the deformation equation for the universal iso-spectral family of second order ordinary differential operators of the form

$$(1.62) \quad L(X, t) := \left(\frac{d}{dX} \right)^2 + 2u(X + t_1, t_3, t_5, \dots)$$

in X and deformation parameters $t = (t_1, t_3, t_5, \dots)$. The operator is often referred to as the *Lax operator* for the KdV equations. The expression

$$\sqrt{L} = \frac{d}{dX} + u \cdot \left(\frac{d}{dX} \right)^{-1} - \frac{1}{2}u' \cdot \left(\frac{d}{dX} \right)^{-2} + \dots$$

makes sense in the ring of pseudo-differential operators, where $'$ denotes the X -derivative. The KdV equations are the system of *Lax equations*

$$\frac{\partial L}{\partial t_{2m+1}} = \left[\left(\sqrt{L}^{2m+1} \right)_+, L \right], \quad m \geq 0,$$

where $_+$ denotes the differential operator part of a pseudo-differential operator. The commutator on the right-hand side explains the invariance of the eigenvalues of the Lax operator L with respect to the deformation parameter t_{2m+1} . The t_1 -deformation is the translation $u(X) \mapsto u(X + t_1)$, and the t_3 -deformation is given by the KdV equation (1.61).

For the particular function (1.60) defined by the intersection numbers (1.59), the *initial* value of the Lax operator is

$$\left(\frac{d}{dX} \right)^2 + 2u(X + t_1, t_3, t_5, \dots) \Big|_{t_1=t_3=t_5=\dots=0} = \left(\frac{d}{dX} \right)^2 + 2X.$$

In terms of yet another variable $x = -\frac{2}{\sqrt[3]{4}}X$, the initial value becomes

$$\frac{1}{\sqrt[3]{4}} \left(\left(\frac{d}{dX} \right)^2 + 2X \right) = \left(\frac{d}{dx} \right)^2 - x = P(x, \hbar) \Big|_{\hbar=1},$$

which is the Airy differential operator.

Kontsevich [55] used the matrix Airy function to obtain all intersection numbers. The topological recursion replaces the asymptotic analysis of matrix integrations with a series of residue calculations on the spectral curve $x = y^2$.

1.6. **All things considered.** What we have in front of us is an interesting example of a theory yet to be constructed.

- We have generating functions of quantum invariants, such as Gromov-Witten invariants. They are symmetric functions.
- We take the *principal specialization* of these functions, and form a generating function of the specialized generating functions.
- This function then solves a 1-dimensional stationary Schrödinger equation. The equation is what we call a *quantum curve*.
- From this Schrödinger equation (or a quantum curve), we construct an algebraic curve, a *spectral curve* in the sense of Hitchin, through the process of *semi-classical limit*.
- The *topological recursion* applied to the spectral curve then recovers the starting quantum invariants.
- The spectral curve can be also expressed as the Hitchin spectral curve of a meromorphic Higgs bundle.
- Then the quantum curve is equivalent to the \hbar -connection on the base curve on which the Higgs bundle is defined.
- The topological recursion therefore constructs a flat section, although formal, of the \hbar -connection from the Hitchin spectral curve. At least locally, the \hbar -connection itself is thus constructed by the topological recursion.

We do not have a general theory yet. In particular, we do not have a global definition of quantum curves. In these lecture notes, we present what we know as of now.

The idea of topological recursion was devised for a totally different context. In the authors' work [24], for the first time the formalism of Eynard and Orantin was placed in the Higgs bundle context. The formalism depends purely on the geometry of the Hitchin spectral curve. Therefore, quantities that the topological recursion computes should represent the geometric information. Then in [25], we have shown through examples that quantization of singular spectral curves are related to certain enumerative geometry problems, when the quantum curve is analyzed at its singular point and the function $\Psi(x, \hbar)$, which should be actually considered as a section of $K_C^{-\frac{1}{2}}$, is expanded at its essential singularity. In the example described above, the corresponding counting problem is computing the intersection numbers of certain cohomology classes on $\overline{\mathcal{M}}_{g,n}$. The original topological recursion of [34] is generalized to singular spectral curves in [25] for this purpose.

The question we still do not know its answer is how to directly connect the Higgs bundle information with the geometric structure whose quantum invariants are captured by the topological recursion.

Since the time of inception of the topological recursion [16, 34], numerous papers have been produced, in both mathematics and physics. It is far beyond the authors' ability to make any meaningful comments on this vast body of literature in the present article. Luckily, interested readers can find useful information in Eynard's talk at the ICM 2014 [32]. Instead of attempting the impossible, we review here a glimpse of **geometric developments** inspired by the topological recursion that have taken place in the last few years.

The geometry community's keen attention was triggered when a concrete **remodeling conjecture** was formulated by string theorists, first by Mariño [59], and then in a more precise and generalized framework by Bouchard, Klemm, Mariño and Pasquetti [11, 12]. The conjecture states that open Gromov-Witten invariants of an arbitrary toric Calabi-Yau orbifold of dimension 3 can be calculated by the topological recursion formulated on

the *mirror curve*. A physical argument for the correctness of the conjecture is pointed out in [75]. Bouchard and Mariño [13] then derived a new conjectural formula for simple Hurwitz numbers from the remodeling conjecture. The correctness of the Hurwitz number conjecture can be easily checked by a computer for many concrete examples. At the same time, it was clear that the conjectural formula was totally different from the combinatorial formula known as the **cut-and-join equation** of [42, 43, 82].

After many computer experiments, one of the authors noticed that the conjectural formula of Bouchard and Mariño was exactly the *Laplace transform* of a particular variant of the cut-and-join equation. Once the precise relation between the knowns and unknowns is identified, often the rest is straightforward, even though still a technical difficulty remains. The conjecture for simple Hurwitz numbers of [13] is solved in [33, 68]. Its generalization to the orbifold Hurwitz numbers is then established in [10]. In each case, the Laplace transform plays the role of the mirror symmetry, changing the combinatorial problem on the A-side to a complex analysis problem on the B-side. The first case of the remodeling conjecture for \mathbb{C}^3 was solved, using the same idea, shortly afterwards in [88]. The remodeling conjecture in its full generality is recently solved in its final form by Fang, Liu, and Zong [37], based on an earlier work of [35].

Independent of these developments, the relation between the topological recursion and combinatorics of enumeration of various graphs drawn on oriented topological surfaces has been studied by the Melbourne group of mathematicians, including Do, Leigh, Manesco, Norbury, and Scott (see, for example, [22, 71, 72, 73]). The authors' earlier papers [15, 27, 64] are inspired by their work. A surprising observation of the Melbourne group, formulated in a conjectural formula, is that the Gromov-Witten invariants of \mathbb{P}^1 themselves should satisfy the topological recursion. Since \mathbb{P}^1 is not a Calabi-Yau manifold, this conjecture does not follow from the remodeling conjecture.

The $GW(\mathbb{P}^1)$ conjecture of [73] is solved by the Amsterdam group of mathematicians, consisting of Dunin-Barkowski, Shadrin, and Spitz, in collaboration with Orantin [30]. Their discovery, that the topological recursion on a disjoint union of open discs as its spectral curve is equivalent to the Givental formalism, has become a key technique of many later work [3, 28, 37].

The technical difficulty of the topological recursion lies in the evaluation of residue calculations involved in the formula. When the spectral curve is an open disc, this difficulty does not occur. But if the global structure of the spectral curve has to be considered, then one needs a totally different idea. The work of [10, 33, 68] has overcome the complex analysis difficulty in dealing with simple and orbifold Hurwitz numbers. There, the key idea is the use of the (piecewise) *polynomiality* of these numbers through the ELSV formula [31] and its orbifold generalization [51]. The paper [28] proves the converse: they first prove the polynomiality of the Hurwitz numbers without assuming the relation to the intersection numbers over $\overline{\mathcal{M}}_{g,n}$, and then establish the ELSV formula from the topological recursion, utilizing a technique of [68].

The topological recursion is a byproduct of the study of random matrix theory/matrix models [16, 34]. The recursion of the same nature appeared earlier in the work of Mirzakhani [60, 61] on the Weil-Petersson volume of the moduli space of bordered hyperbolic surfaces. The Laplace transform of the Mirzakhani recursion is an example of the topological recursion. Its spectral curve, the sine curve, was first identified in [65] as the intertwiner of two representations of the Virasoro algebra.

The notion of quantum curves goes back to [1]. It is further developed in physics community by Dijkgraaf, Hollands, Sułkowski, and Vafa [18, 19, 48]. The geometry community was

piqued by [17, 44], which speculated the relation between the topological recursion, quantization of the $SL(2, \mathbb{C})$ -character variety of the fundamental group of a knot complement in S^3 , the AJ-conjecture due to [39, 40], and the K_2 -group of algebraic K -theory. Although it is tantalizingly interesting, so far no mathematical results have been established in this direction for hyperbolic knots, or even it may be impossible (see for example, [9]), mainly due to the reducible nature of the spectral curve in this particular context. For torus knots, see also [14]. A rigorous construction of quantization of spectral curves was established for a few examples in [67] for the first time, but without any relation to knot invariants. One of the examples there will be treated in these lectures below. By now, we have many more mathematical examples of quantum curves [10, 22, 29, 66, 89, 90]. We note that in many of these examples, the spectral curves have a global parameter, even the curves are not necessarily the rational projective curve. Therefore, the situation is in some sense still the “genus 0 case.” The difficulty of quantization lies in the dealing of complicated entire functions, and the fact that the quantum curves are **difference equations**, rather than differential equations of finite order.

We are thus led to another question.

Question 1.17. *What can we do when we have a different situation, where the spectral curve of the theory is global, compact, and of a high genus?*

It comes as a surprise that the global residue calculation of the formula can be performed when the spectral curve of the topological recursion is precisely a Hitchin spectral curve associated with a rank 2 Higgs bundle. The result of the calculation then leads us to a local construction of a quantum curve. In this way a connection between quantization of Hitchin spectral curves and the topological recursion is discovered in [24].

If we regard the topological recursion as a method of calculation of quantum invariants, then we need to allow singular spectral curves, as we have seen earlier. The simplest quantization of the singular Hitchin spectral curve is then obtained by the topological recursion again, but this time, it has to be applied to the normalization of the singular spectral curve constructed in a particular way of blow-ups of the ambient compactified cotangent bundle. This is the content of [25], and we obtain a **Rees \mathcal{D} -module** as the result of quantization. We note here that the result of quantization is not unique. Depending on the purpose, one needs to use a different guideline for quantization. The result would be a different differential equation, but still having the same semi-classical limit.

For example, it has been rigorously proved in [50] that surprisingly the quantization procedure of [24, 25], including the desingularization of the spectral curve, *automatically* leads to an iso-monodromic deformation family, for the case of the Painlevé I equation. Their global parameter is essentially the normalization coordinate of the singular elliptic curve. In their work [50], they ask what one obtains if the straightforward topological recursion is applied for the quantization of a singular elliptic curve with a prescribed parameter in a particular way. They then find that the quantum curve is a Schrödinger equation whose coefficients have nontrivial dependence on \hbar , yet it is an iso-spectral family with respect to the parameter. This work, and also the numerous mathematical examples of quantum curves that have been already constructed, suggest that the idea of using \mathcal{D} -modules for the definition of quantum curves ([18, 25]) is not the final word. Differential operators of an infinite order, or difference operators mixed with differential operators, have to be also considered.

For the mirror curves of toric Calabi-Yau orbifolds of dimension 3 appearing in the context of the remodeling conjecture [11, 12, 37, 59], the conjectural quantum curves acquire a

very different nature. It has deep connections to number theory and quantum dilogarithm functions [53].

A suggestion of using deformation quantization modules for quantum curves is made by Kontsevich in late 2013 in a private communication to the authors. An interesting work toward this direction is proposed by Petit [77].

The relation of the quantization discussed in these lectures to the **non-Abelian Hodge correspondence** and **opers** is being investigated as of now. It seems to be that a coordinate-free global definition of quantum curves is emerging, and a direct relationship among quantum curves, non-Abelian Hodge correspondence, and opers is being identified. See [23] for more detail.

The story is expanding its horizon. We have come to the other side of the rainbow. And there we find ourselves on Newton's seashore. So far we have found only a few smoother pebbles or prettier shells, whilst...

2. FROM CATALAN NUMBERS TO THE TOPOLOGICAL RECURSION

Let us consider a function $f(X)$ in one variable, where X is an $N \times N$ Hermitian matrix. One of the main problems of matrix integration theory is to calculate the *expectation value*

$$(2.1) \quad \langle \text{tr} f(X) \rangle := \frac{\int_{H_{N \times N}} \text{tr} f(X) e^{V(X)} dX}{\int_{H_{N \times N}} e^{V(X)} dX},$$

where the *potential function* $V(X)$ is given, such as the Gaussian potential

$$(2.2) \quad V(X) = -\frac{1}{2} \text{tr}(X^2),$$

so that

$$C_N = \int_{H_{N \times N}} e^{V(X)} dX$$

is finite. The integration measure dX is the standard $U(N)$ -invariant Lebesgue measure of the space $H_{N \times N} = \mathbb{R}^{N^2}$ of Hermitian matrices of size N . When a Gaussian potential (2.2) is chosen, $e^{V(X)} dX$ is a probability measure after an appropriate normalization, and $\langle \text{tr}(X^m) \rangle$ is the m -th *moment*. If we know all the moments, then we can calculate the expectation value of any polynomial function $f(X)$. Therefore, the problem changes into calculating a generating function of the moments

$$(2.3) \quad \frac{1}{C_N} \sum_{m=0}^{\infty} \frac{1}{z^{m+1}} \int_{H_{N \times N}} \text{tr}(X^m) e^{V(X)} dX.$$

For a norm bounded matrix X , we have

$$\text{tr} \left(\frac{1}{z - X} \right) = \sum_{m=0}^{\infty} \frac{1}{z^{m+1}} \text{tr}(X^m).$$

However, we need to be careful here because the quantity called *resolvent* of a random matrix X ,

$$(2.4) \quad \left\langle \text{tr} \left(\frac{1}{z - X} \right) \right\rangle = \frac{1}{C_N} \int_{H_{N \times N}} \text{tr} \left(\frac{1}{z - X} \right) e^{V(X)} dX,$$

is not equal to (2.3). For example, let us consider the $N = 1$ case and write $X = x \in \mathbb{R}$. Then the formula

$$\frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{1}{z-x} e^{-\frac{1}{2}x^2} dx = \sum_{m=0}^{\infty} \frac{(2m-1)!!}{z^{2m+1}} = \frac{1}{\sqrt{2\pi}} \sum_{m=0}^{\infty} \frac{1}{z^{m+1}} \int_{-\infty}^{\infty} x^m e^{-\frac{1}{2}x^2} dx$$

is valid only as the asymptotic expansion of the analytic function

$$\frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \frac{1}{z-x} e^{-\frac{1}{2}x^2} dx$$

in z for $Im(z) \neq 0$ and $Re(z) \rightarrow +\infty$. Still we can see that the information of the generating function of the moment (2.3) can be extracted from the resolvent (2.4) if we apply the technique of asymptotic expansion of a holomorphic function at its essential singularity.

More generally, we can consider *multi-resolvent correlation functions*

$$(2.5) \quad \left\langle \prod_{i=1}^n \text{tr} \left(\frac{1}{z_i - X} \right) \right\rangle = \frac{1}{C_N} \int_{H_{N \times N}} \prod_{i=1}^n \text{tr} \left(\frac{1}{z_i - X} \right) e^{V(X)} dX.$$

When we say ‘‘calculating’’ the expectation value (2.5), we wish to identify it as a *holomorphic* function in all the parameters, i.e., (z_1, \dots, z_n) , the coefficients of the potential V , and the matrix size N . In particular, the analytic dependence on the parameter N is an important feature we wish to determine.

It is quite an involved problem in analysis, and we do not attempt to go into this route in these lecture notes. The collaborative effort of random matrix community has devised a recursive method of solving this analysis problem (see, for example, [16, 32, 34]), which is now known as the **topological recursion**. One thing we can easily expect here is that since (2.5) is an analytic function in (z_1, \dots, z_n) , there must be an obvious relation between the topological recursion and algebraic geometry.

What is amazing is that the exact same recursion formula happens to appear in the context of many different enumerative geometry problems, again and again. Even the counting problems are different, the topological recursion always takes the same general formalism. Therefore, to understand the nature of this formalism, it suffices to give the simplest non-trivial example. This is what we wish to accomplish in this section.

2.1. Counting graphs on a surface. Let us start with the following problem:

Problem 2.1. *Find the number of distinct cell-decompositions of a given closed oriented topological surface of genus g , with the specified number of 0-cells and the number of 1-cells that are incident to each 0-cell.*

Denote by C_g a compact, 2-dimensional, oriented topological manifold of genus g without boundary. Two cell-decompositions of C_g are identified if there is an orientation-preserving homeomorphism of C_g onto itself that brings one to the other. If there is such a map for the same cell-decomposition, then it is an automorphism of the data. The 1-skeleton of a cell-decomposition, which we denote by γ , is a *graph* drawn of C_g . We call its 0-cell a *vertex*, 1-cell an *edge*, and 2-cell a *face*. The midpoint of an edge separates the edge into two *half-edges* joined together at the midpoint. The *degree* of a vertex is the number of half-edges incident to it. For the purpose of counting, we label all vertices. To be more precise, we give a total order to the set of vertices. Most of the time we simply use $[n] = \{1, \dots, n\}$ to label the set of n vertices.

The 1-skeleton γ is usually called a *ribbon graph*, which is a graph with a cyclic order assigned to incident half-edges at each vertex. The face-labeled ribbon graphs describe an

orbifold cell-decomposition of $\mathcal{M}_{g,n} \times \mathbb{R}_+^n$. Since we label vertices of γ , there is a slight difference as to what the graph represents. It is the *dual* graph of a ribbon graph, and its vertices are labeled. To emphasize the dual nature, we call γ a **cell graph**.

Most cell graphs do not have any non-trivial automorphisms. If there is one, then it induces a cyclic permutation of half-edges at a vertex, since we label all vertices. Therefore, if we pick one of the incident half-edges at each vertex, assign an outgoing arrow to it, and require that an automorphism also fix the arrowed half-edges, then the graph has no non-trivial automorphisms. For a counting problem, no automorphism is a desirable situation because the bijective counting method works better there. Let us call such a graph an **arrowed cell graph**. Now the refined problem:

Problem 2.2. Let $\vec{\Gamma}_{g,n}(\mu_1, \dots, \mu_n)$ denote the set of arrowed cell graphs drawn on a closed, connected, oriented surface of genus g with n labeled vertices of degrees μ_1, \dots, μ_n . Find its cardinality

$$(2.6) \quad C_{g,n}(\mu_1, \dots, \mu_n) := |\vec{\Gamma}_{g,n}(\mu_1, \dots, \mu_n)|.$$

When $\gamma \in \vec{\Gamma}_{g,n}(\mu_1, \dots, \mu_n)$, we say γ has *type* (g, n) . Denote by $c_\alpha(\gamma)$ the number of α -cells of the cell-decomposition associated with γ . Then we have

$$2 - 2g = c_0(\gamma) - c_1(\gamma) + c_2(\gamma), \quad c_0(\gamma) = n, \quad 2c_1(\gamma) = \mu_1 + \dots + \mu_n.$$

Therefore, $\vec{\Gamma}_{g,n}(\mu_1, \dots, \mu_n)$ is a finite set.




FIGURE 2.1. A cell graph of type $(2, 6)$.

Example 2.1. An arrowed cell graph of type $(0, 1)$ is a collection of loops drawn on a plane as in Figure 2.2. If we assign a pair of parenthesis to each loop, starting from the arrowed one as $($, and go around the unique vertex counter-clock wise, then we obtain the parentheses pattern $(((())))$. Therefore,

$$(2.7) \quad C_{0,1}(2m) = C_m = \frac{1}{m+1} \binom{2m}{m}$$

is the m -th Catalan number. Thus it makes sense to call $C_{g,n}(\mu_1, \dots, \mu_n)$ **generalized Catalan numbers**.




FIGURE 2.2. A cell graph of type $(0, 1)$ with a vertex of degree 6.

Theorem 2.3 (Catalan Recursion, [83]). *The generalized Catalan numbers satisfy the following equation.*

$$(2.8) \quad C_{g,n}(\mu_1, \dots, \mu_n) = \sum_{j=2}^n \mu_j C_{g,n-1}(\mu_1 + \mu_j - 2, \mu_2, \dots, \widehat{\mu_j}, \dots, \mu_n) \\ + \sum_{\alpha+\beta=\mu_1-2} \left[C_{g-1,n+1}(\alpha, \beta, \mu_2, \dots, \mu_n) + \sum_{\substack{g_1+g_2=g \\ I \sqcup J = \{2, \dots, n\}}} C_{g_1,|I|+1}(\alpha, \mu_I) C_{g_2,|J|+1}(\beta, \mu_J) \right],$$

where $\mu_I = (\mu_i)_{i \in I}$ for an index set $I \subset [n]$, $|I|$ denotes the cardinality of I , and the third sum in the formula is for all partitions of g and set partitions of $\{2, \dots, n\}$.

Proof. Let γ be an arrowed cell graph counted by the left-hand side of (2.8). Since all vertices of γ are labeled, we write the vertex set by $\{p_1, \dots, p_n\}$. We take a look at the half-edge incident to p_1 that carries an arrow.

Case 1. The arrowed half-edge extends to an edge E that connects p_1 and p_j for some $j > 1$.

In this case, we contract the edge and join the two vertices p_1 and p_j together. By this process we create a new vertex of degree $\mu_1 + \mu_j - 2$. To make the counting bijective, we need to be able to go back from the contracted graph to the original, provided that we know μ_1 and μ_j . Thus we place an arrow to the half-edge next to E around p_1 with respect to the counter-clockwise cyclic order that comes from the orientation of the surface. In this process we have μ_j different arrowed graphs that produce the same result, because we must remove the arrow placed around the vertex p_j in the original graph. This gives the first line of the right-hand side of (2.8). See Figure 2.3.




FIGURE 2.3. The process of contracting the arrowed edge E that connects vertices p_1 and p_j , $j > 1$.

Case 2. The arrowed half-edge at p_1 is a loop E that goes out from, and comes back to, p_1 .

The process we apply is again contracting the loop E . The loop E separates all other incident half-edges at p_1 into two groups, one consisting of α of them placed on one side of the loop, and the other consisting of β half-edges placed on the other side. It can happen that $\alpha = 0$ or $\beta = 0$. Contracting a loop on a surface causes pinching. Instead of creating a pinched (i.e., singular) surface, we separate the double point into two new vertices of degrees α and β . Here again we need to remember the place of the loop E . Thus we put an arrow to the half-edge next to the loop in each group. See Figure 2.4.

After the pinching and separating the double point, the original surface of genus g with n vertices $\{p_1, \dots, p_n\}$ may change its topology. It may have genus $g - 1$, or it splits into two pieces of genus g_1 and g_2 . The second line of (2.8) records all such cases. Normally we would have a factor $\frac{1}{2}$ in front of the second line of the formula. We do not have it here




FIGURE 2.4. The process of contracting the arrowed loop E that is attached to p_1 .

because the arrow on the loop could be in two different directions. Placing the arrow on the other half-edge of the loop is equivalent to interchanging α and β .

This completes the proof. \square

Remark 2.4. For $(g, n) = (0, 1)$, the above formula reduces to

$$(2.9) \quad C_{0,1}(\mu_1) = \sum_{\alpha+\beta=\mu_1-2} C_{0,1}(\alpha)C_{0,1}(\beta).$$

Since the degree of the unique vertex is always even for type $(0, 1)$ graphs, by defining $C_{0,1}(0) = 1$, (2.9) gives the Catalan recursion. Only for $(g, n) = (0, 1)$, this irregular case of $\mu_1 = 0$ happens, because a degree 0 single vertex is *connected*, and gives a cell-decomposition of S^2 . All other cases, if one of the vertices has degree 0, then the Catalan number $C_{g,n}(\mu_1, \dots, \mu_n)$ is simply 0 because there is no corresponding *connected* cell decomposition.

Remark 2.5. Eqn. (2.8) is a recursion with respect to

$$2g - 2 + n + \sum_{i=1}^n \mu_i.$$

The values are therefore completely determined by the initial value $C_{0,1}(0) = 1$. The formula does not give a recursion of a *function* $C_{g,n}(\mu_1, \dots, \mu_n)$, because the same type (g, n) appears on the right-hand side.

The classical Catalan recursion (2.9) determines all values of $C_{0,1}(2m)$, but the closed formula (2.7) requires a different strategy. Let us introduce a generating function

$$(2.10) \quad z = z(x) = \sum_{m=0}^{\infty} C_{0,1}(2m)x^{-2m-1}.$$

From (2.9) we have

$$z^2 = \sum_{m=0}^{\infty} \left(\sum_{a+b=m} C_{0,1}(2a)C_{0,1}(2b) \right) x^{-2m-2} = \sum_{m=0}^{\infty} C_{0,1}(2m+2)x^{-2m-2}.$$

Since

$$xz = \sum_{m=-1}^{\infty} C_{0,1}(2m+2)x^{-2m-2},$$

we obtain an equation $xz = z^2 + 1$, or

$$(2.11) \quad x = z + \frac{1}{z}.$$

This is the inverse function of the complicated-looking generating function $z(x)$ at the branch $z \rightarrow 0$ as $x \rightarrow +\infty$! Thus the curve (2.11) knows everything about the Catalan

numbers. For example, we can prove the closed formula (2.7). The solution of (2.11) as a quadratic equation for z that gives the above branch is given by

$$z(x) = \frac{x - \sqrt{x^2 - 4}}{2} = \frac{x}{2} \left(1 - \sqrt{1 - \frac{4}{x^2}} \right).$$

The binomial expansion of the square root

$$\sqrt{1 + X} = \sum_{m=0}^{\infty} \binom{\frac{1}{2}}{m} X^m,$$

$$\binom{\frac{1}{2}}{m} := \frac{\frac{1}{2}(\frac{1}{2} - 1)(\frac{1}{2} - 2) \cdots (\frac{1}{2} - m + 1)}{m!} = (-1)^{m-1} \frac{(2m-3)!!}{2^m m!} = \frac{(-1)^{m-1}}{4^m} \frac{1}{2m-1} \binom{2m}{m},$$

then gives the closed formula (2.7) for the Catalan numbers:

$$\begin{aligned} z(x) &= \sum_{m=0}^{\infty} C_{0,1}(2m) x^{-2m-1} \\ &= \frac{x}{2} \left(1 - \sqrt{1 - \frac{4}{x^2}} \right) \\ &= \frac{x}{2} \left(1 - \left(\sum_{m=0}^{\infty} \frac{(-1)^{m-1}}{4^m} \frac{1}{2m-1} \binom{2m}{m} (-1)^m \left(\frac{4}{x^2} \right)^m \right) \right) \\ &= \frac{1}{2} \sum_{m=1}^{\infty} \frac{1}{2m-1} \binom{2m}{m} x^{-2m+1} \\ &= \sum_{m=0}^{\infty} \frac{1}{m+1} \binom{2m}{m} x^{-2m-1}. \end{aligned}$$

Another information we obtain from (2.11) is the radius of convergence of the infinite series $z(x)$ of (2.10). Since

$$dx = \left(1 - \frac{1}{z^2} \right) dz,$$

the map (2.11) is critical (or ramified) at $z = \pm 1$. The critical values are $x = \pm 2$. On the branch we are considering, (2.10) is the inverse function of (2.11) for all values of $|x| > 2$. This means the series $z(x)$ is absolutely convergent on the same domain.

Remark 2.6. In combinatorics, we often consider a generating function of interesting quantities only as a formal power series. The idea of topological recursion tells us that we should consider the **Riemann surface** of the Catalan number generating function $z = z(x)$. We then recognize that there is a global algebraic curve hidden in the scene, which is the curve of Figure 2.5. The topological recursion mechanism then tells us how to calculate all $C_{g,n}(\vec{\mu})$ from this curve alone, known as the **spectral curve**.

2.2. The spectral curve of a Higgs bundle and its desingularization. To consider the quantization of the curve (2.11), we need to place it into a cotangent bundle. Here again, we use the base curve \mathbb{P}^1 and the same vector bundle E of (1.20) on it. As a Higgs field, we use

$$(2.12) \quad \phi := \begin{bmatrix} & 1 \\ -(dx)^2 & -xdx \end{bmatrix} : E \longrightarrow K_{\mathbb{P}^1}(4) \otimes E.$$




FIGURE 2.5. The **Riemann surface** of the Catalan generating function $z = z(x)$.

Here, $(dx)^2 \in H^0(\mathbb{P}^1, K_{\mathbb{P}^1}^{\otimes 2}(4))$ is the unique (up to a constant factor) quadratic differential with an order 4 pole at the infinity, and $xdx \in H^0(\mathbb{P}^1, K_{\mathbb{P}^1}(2))$ is the unique meromorphic differential with a zero at $x = 0$ and a pole of order 3 at $x = \infty$. In the affine coordinate (x, y) of the Hirzebruch surface \mathbb{F}^2 as before, the spectral curve Σ is given by

$$(2.13) \quad \det(\eta - \pi^*(\phi)) = (y^2 + xy + 1)(dx)^2 = 0,$$

where $\pi : \mathbb{F}_2 \rightarrow \mathbb{P}^1$ is the projection. Therefore, the generating function $z(x)$ of (2.10) gives a parametrization of the spectral curve

$$(2.14) \quad \begin{cases} x = z(x) + \frac{1}{z(x)} \\ y = -z(x). \end{cases}$$

In the other affine coordinate (u, w) of (1.24), the spectral curve is singular at $(u, w) = (0, 0)$:

$$(2.15) \quad u^4 - uw + w^2 = 0.$$




FIGURE 2.6. The spectral curve Σ of (2.13). The horizontal line is the divisor $w = 0$ at infinity, and the vertical line is the fiber class $u = 0$. The spectral curve intersects with $w = 0$ four times. One of the two curve germ components is given by $w = u$, and the other by $w = u^3$.

Blow up $\mathbb{F}_2 = \overline{T^*\mathbb{P}^1}$ once at the nodal singularity $(u, w) = (0, 0)$ of the spectral curve Σ , and let $\tilde{\Sigma} \rightarrow \Sigma$ be the proper transform of Σ .

$$(2.16) \quad \begin{array}{ccc} \tilde{\Sigma} & \xrightarrow{\tilde{i}} & Bl(\overline{T^*\mathbb{P}^1}) \\ \downarrow \nu & & \downarrow \nu \\ \Sigma & \xrightarrow{i} & \overline{T^*\mathbb{P}^1} \\ \downarrow \tilde{\pi} & \swarrow \pi & \downarrow \pi \\ \mathbb{P}^1 & & \mathbb{P}^1 \end{array}$$

In terms of the coordinate w_1 defined by $w = w_1 u$, (2.15) becomes

$$(2.17) \quad u^2 + \left(w_1 - \frac{1}{2}\right)^2 = \frac{1}{4},$$

which is the defining equation for $\tilde{\Sigma}$. From this equation we see that its geometric genus is 0, hence it is just another \mathbb{P}^1 . The covering $\tilde{\pi} : \tilde{\Sigma} \rightarrow \mathbb{P}^1$ is ramified at two points, corresponding to the original ramification points $(x, y) = (\pm 2, \mp 1)$ of $\pi : \Sigma \rightarrow \mathbb{P}^1$. The rational parametrization of (2.17) is given by

$$(2.18) \quad \begin{cases} u = \frac{1}{2} \cdot \frac{t^2 - 1}{t^2 + 1} \\ w_1 = \frac{1}{2} - \frac{t}{t^2 + 1}, \end{cases}$$

where t is the affine coordinate of $\tilde{\Sigma}$ so that $t = \pm 1$ gives $(u, w) = (0, 0)$. Indeed, the parameter t is a **normalization coordinate** of the spectral curve Σ :

$$(2.19) \quad \begin{cases} x = 2 + \frac{4}{t^2 - 1} \\ y = -\frac{t+1}{t-1}, \end{cases} \quad \begin{cases} u = \frac{1}{2} \cdot \frac{t^2 - 1}{t^2 + 1} \\ w = \frac{1}{4} \cdot \frac{(t-1)^3(t+1)}{(t^2 + 1)^2}. \end{cases}$$

Although the expression of x and y in terms of the normalization coordinate is more complicated than (2.14), it is important to note that the spectral curve $\tilde{\Sigma}$ is now non-singular.

2.3. The generating function, or the Laplace transform. For all (g, n) except for $(0, 1)$ and $(0, 2)$, let us introduce the generating function of (2.6) as follows:

$$(2.20) \quad F_{g,n}^C(x_1, \dots, x_n) := \sum_{\mu_1 \geq 1, \dots, \mu_n \geq 1} \frac{C_{g,n}(\mu_1, \dots, \mu_n)}{\mu_1 \cdots \mu_n} x_1^{-\mu_1} \cdots x_n^{-\mu_n}.$$

If we consider $x_i = e^{w_i}$, then the above sum is just the discrete Laplace transform of the function in n integer variables:

$$\frac{C_{g,n}(\mu_1, \dots, \mu_n)}{\mu_1 \cdots \mu_n}.$$

Our immediate goal is to compute the Laplace transform as a holomorphic function. Since the only information we have now is the generalized Catalan recursion (2.8), how much can we say about this function? Actually, the following theorem is proved, all from using the recursion alone!

Theorem 2.7. *Let us consider the generating function (2.20) as a function in the normalization coordinates (t_1, \dots, t_n) satisfying*

$$x_i = 2 + \frac{4}{t_i^2 - 1}, \quad i = 1, 2, \dots, n,$$

as identified in (2.19), and by abuse of notation, we simply write it as $F_{g,n}^C(t_1, \dots, t_n)$. For the range of (g, n) with the stability condition $2g - 2 + n > 0$, we have the following.

- *The generating function $F_{g,n}^C(t_1, \dots, t_n)$ is a **Laurent polynomial** in the t_i -variables of the total degree $3(2g - 2 + n)$.*
- *The reciprocity relation holds:*

$$(2.21) \quad F_{g,n}^C(1/t_1, \dots, 1/t_n) = F_{g,n}^C(t_1, \dots, t_n).$$

- *The special values at $t_i = -1$ are given by*

$$(2.22) \quad F_{g,n}^C(t_1, \dots, t_n)|_{t_i = -1} = 0$$

for each i .

- The diagonal value at $t_i = 1$ gives the orbifold Euler characteristic of the moduli space $\mathcal{M}_{g,n}$:

$$(2.23) \quad F_{g,n}^C(1, \dots, 1) = (-1)^n \chi(\mathcal{M}_{g,n}).$$

- The restriction of the Laurent polynomial $F_{g,n}^C(t_1, \dots, t_n)$ to its highest degree terms gives a homogeneous polynomial defined by

$$(2.24) \quad F_{g,n}^{C, \text{highest}}(t_1, \dots, t_n) = \frac{(-1)^n}{2^{2g-2+n}} \sum_{\substack{d_1 + \dots + d_n \\ = 3g-3+n}} \langle \tau_{d_1} \cdots \tau_{d_n} \rangle_{g,n} \prod_{i=1}^n (2d_i - 1)!! \left(\frac{t_i}{2}\right)^{2d_i+1}.$$

Thus the function $F_{g,n}^C(t_1, \dots, t_n)$ knows the orbifold Euler characteristic of $\mathcal{M}_{g,n}$, and all the cotangent class intersection numbers (1.10) for all values of (g, n) in the stable range! It is also striking that it is actually a Laurent polynomial, while the definition (2.20) is given only as a formal Laurent series. The reciprocity (2.21) is the reflection of the invariance of the spectral curve Σ under the rotation

$$(x, y) \mapsto (-x, -y).$$

This surprising theorem is a consequence of the Laplace transform of the Catalan recursion itself.

Theorem 2.8 (Differential recursion, [91]). *The Laplace transform $F_{g,n}^C(t_1, \dots, t_n)$ satisfies the following differential recursion equation for every (g, n) subject to $2g - 2 + n \geq 2$.*

$$(2.25) \quad \begin{aligned} & \frac{\partial}{\partial t_1} F_{g,n}^C(t_1, \dots, t_n) \\ &= -\frac{1}{16} \sum_{j=2}^n \left[\frac{t_j}{t_1^2 - t_j^2} \left(\frac{(t_1^2 - 1)^3}{t_1^2} \frac{\partial}{\partial t_1} F_{g,n-1}^C(t_1, \dots, \widehat{t}_j, \dots, t_n) \right. \right. \\ & \quad \left. \left. - \frac{(t_j^2 - 1)^3}{t_j^2} \frac{\partial}{\partial t_j} F_{g,n-1}^C(t_2, \dots, t_n) \right) \right] \\ & \quad - \frac{1}{16} \sum_{j=2}^n \frac{(t_1^2 - 1)^2}{t_1^2} \frac{\partial}{\partial t_1} F_{g,n-1}^C(t_1, \dots, \widehat{t}_j, \dots, t_n) \\ & \quad - \frac{1}{32} \frac{(t_1^2 - 1)^3}{t_1^2} \left[\frac{\partial^2}{\partial u_1 \partial u_2} F_{g-1, n+1}^C(u_1, u_2, t_2, t_3, \dots, t_n) \right] \Big|_{u_1 = u_2 = t_1} \\ & \quad - \frac{1}{32} \frac{(t_1^2 - 1)^3}{t_1^2} \sum_{\substack{\text{stable} \\ g_1 + g_2 = g \\ I \sqcup J = \{2, 3, \dots, n\}}} \frac{\partial}{\partial t_1} F_{g_1, |I|+1}^C(t_1, t_I) \frac{\partial}{\partial t_1} F_{g_2, |J|+1}^C(t_1, t_J). \end{aligned}$$

For a subset $I \subset \{1, 2, \dots, n\}$, we denote $t_I = (t_i)_{i \in I}$. The “stable” summation means $g_1 + |I| - 1 > 0$ and $g_2 + |J| - 1 > 0$.

The differential recursion uniquely determines all $F_{g,n}^C(t_1, \dots, t_n)$ by integrating the right-hand side of (2.25) from -1 to t_1 with respect to the variable t_1 . The initial conditions are

$$(2.26) \quad F_{1,1}^C(t) = -\frac{1}{384} \frac{(t+1)^4}{t^2} \left(t - 4 + \frac{1}{t} \right)$$

and

$$(2.27) \quad F_{0,3}^C(t_1, t_2, t_3) = -\frac{1}{16}(t_1 + 1)(t_2 + 1)(t_3 + 1) \left(1 + \frac{1}{t_1 t_2 t_3} \right).$$

Remark 2.9. Theorem 2.8 is proved by Mincheng Zhou in his senior thesis [91]. It is the result of a rather difficult calculation of the Laplace transform of the Catalan recursion (2.8).

Remark 2.10. Theorem 2.7 is never stated in this format before. Its proof follows from the results of [25, 27, 64, 67], based on the induction using (2.25). The essential point of the discovery of Theorem 2.7 is the use of the normalization coordinate t of (2.19). The authors almost accidentally found the coordinate transformation

$$z(x) = \frac{t + 1}{t - 1}$$

in [27]. Then in [25], we noticed that this coordinate was exactly the normalization coordinate that was naturally obtained in the blow-up process (2.16).

The uniqueness of the solution of (2.25) allows us to identify the solution $F_{g,n}^C$ with the Laplace transform of the number of lattice points in $\mathcal{M}_{g,n}$, as we see later in this section. Through this identification, (2.22) and (2.23) become obvious. The asymptotic behavior (2.24) follows from the lattice point approximation of the Euclidean volume of polytopes, and Kontsevich's theorem that identifies the Euclidean volume of $\mathcal{M}_{g,n}$ and the intersection numbers (1.10) on $\overline{\mathcal{M}}_{g,n}$.

2.4. The unstable geometries and the initial value of the topological recursion.

The actual computation of the Laplace transform equation (2.25) from (2.8) requires the evaluation of the Laplace transforms of $C_{0,1}(\mu_1)$ and $C_{0,2}(\mu_1, \mu_2)$. It is done as follows.

Since a degree 0 vertex is allowed for the $(g, n) = (0, 1)$ unstable geometry, we do not have the corresponding $F_{0,1}^C(x)$ in (2.20). Since

$$d_{x_1} \cdots d_{x_n} F_{g,n}^C = \sum_{\mu_1 \geq 1, \dots, \mu_n \geq 1} (-1)^n C_{g,n}(\mu_1, \dots, \mu_n) x_1^{-\mu_1-1} \cdots x_n^{-\mu_n-1} dx_1 \cdots dx_n,$$

we could choose

$$d_x F_{0,1}^C(x) = -(z(x) - x^{-1}) dx$$

as a defining equation for $F_{0,1}^C$.

In the light of (2.14), $ydx = -zdx$ is a natural global holomorphic 1-form on $T^*\mathbb{P}^1$, called the *tautological* 1-form. Its exterior differentiation $d(ydx) = dy \wedge dx$ defines the canonical holomorphic symplectic structure on $T^*\mathbb{P}^1$. Since we are interested in the *quantization* of the Hitchin spectral curve, we need a symplectic structure here, which is readily available for our use from ydx .

Therefore, it is reasonable for us to define the 'function' $F_{0,1}$ by

$$(2.28) \quad dF_{0,1} = ydx, \quad \text{or} \quad F_{0,1} = \int ydx.$$

Although this equation does not determine the content term, it does not play any role for our purposes. Here, we choose the constant of integration to be 0. Since the symplectic structure on $T^*\mathbb{P}^1$ is non-trivial, we need to interpret the equation (2.28) being defined on the spectral curve, and be prepared that we may not obtain any meromorphic function on

the spectral curve. For the Catalan case, we have to use the spectral curve coordinate z or t as a variable of $F_{g,n}^C$, instead of x . Since

$$-zdx = -zdz + \frac{dz}{z},$$

we conclude

$$(2.29) \quad \begin{aligned} F_{0,1}^C(z) &:= -\frac{1}{2}z^2 + \log z \\ &= -\frac{1}{2}\left(\frac{t+1}{t-1}\right)^2 + \log\left(\frac{t+1}{t-1}\right). \end{aligned}$$

We encounter a non-algebraic function here, indeed.

For the computation of the Laplace transform $F_{0,2}^C$, we need an explicit formula for $C_{0,2}(\mu_1, \mu_2)$. Luckily, such computation has been carried out in [54], fully utilizing the technique of the *dispersionless Toda lattice hierarchy*. It is surprising to see how much integrable system consideration is involved in computing such a simple quantity as $C_{0,2}(\mu_1, \mu_2)$! The result is the following.

Theorem 2.11 (Calculation of the 2-point Catalan numbers, [54]). *For every $\mu_1, \mu_2 > 0$, the genus 0, 2-point Catalan numbers are given by*

$$(2.30) \quad C_{0,2}(\mu_1, \mu_2) = 2 \frac{\lfloor \frac{\mu_1+1}{2} \rfloor \lfloor \frac{\mu_2+1}{2} \rfloor}{\mu_1 + \mu_2} \binom{\mu_1}{\lfloor \frac{\mu_1}{2} \rfloor} \binom{\mu_2}{\lfloor \frac{\mu_2}{2} \rfloor}.$$

We refer to [27] for the derivation of $F_{0,2}^C$. The result is

$$(2.31) \quad \begin{aligned} F_{0,2}^C(z_1, z_2) &= -\log(1 - z_1 z_2) \\ &= -\log(-2(t_1 + t_2)) + \log(t_1 - 1) + \log(t_2 - 1). \end{aligned}$$

For the purpose of later use, we note that

$$(2.32) \quad \begin{aligned} d_{t_1} d_{t_2} F_{0,2}^C(t_1, t_2) &= \frac{dt_1 \cdot dt_2}{(t_1 + t_2)^2} \\ &= \frac{dt_1 \cdot dt_2}{(t_1 - t_2)^2} - (\tilde{\pi} \times \tilde{\pi})^* \frac{dx_1 \cdot dx_2}{(x_1 - x_2)^2}, \end{aligned}$$

where $\tilde{\pi} : \tilde{\Sigma} \rightarrow \mathbb{P}^1$ is the projection of (2.16), i.e., the variable transformation (2.19).

2.5. Geometry of the topological recursion. Computation of a multi-resolvent (2.5) is one thing. What we have in front of us is a different story. We wish to compute an asymptotic expansion of a solution to the differential equation that is defined on the base curve C and gives the quantization of the Hitchin spectral curve of a meromorphic Higgs bundle (E, ϕ) . The expansion has to be done at the essential singularity of the solution.

Question 2.12. *Is there a mathematical framework suitable for such problems?*

The discovery of [24, 25] gives an answer: **Generalize the formalism of Eynard and Orantin of [34] to the geometric situation of meromorphic Higgs bundles. Then this generalized topological recursion computes the asymptotic expansion in question.**

We are now ready to present the topological recursion, continuing our investigation of the particular example of Catalan numbers. The point here is that **the topological recursion is a universal formula depending only on geometry**. Therefore, we can arrive at the general formula from any example. *One example rules them all.*

Theorem 2.13 (The topological recursion for the generalized Catalan numbers, [27]).
 Define symmetric n -linear differential forms on $(\tilde{\Sigma})^n$ for $2g - 2 + n > 0$ by

$$(2.33) \quad W_{g,n}^C(t_1, \dots, t_n) := d_{t_1} \cdots d_{t_n} F_{g,n}^C(t_1, \dots, t_n),$$

and for $(g, n) = (0, 2)$ by

$$(2.34) \quad W_{0,2}^C(t_1, t_2) := \frac{dt_1 \cdot dt_2}{(t_1 - t_2)^2}.$$

Then these differential forms satisfy the following integral recursion equation, called the **topological recursion**.

$$(2.35) \quad W_{g,n}^C(t_1, \dots, t_n) = -\frac{1}{64} \frac{1}{2\pi i} \int_{\gamma} \left(\frac{1}{t+t_1} + \frac{1}{t-t_1} \right) \frac{(t^2-1)^3}{t^2} \cdot \frac{1}{dt} \cdot dt_1 \\ \times \left[\sum_{j=2}^n \left(W_{0,2}^C(t, t_j) W_{g,n-1}^C(-t, t_2, \dots, \hat{t}_j, \dots, t_n) + W_{0,2}^C(-t, t_j) W_{g,n-1}^C(t, t_2, \dots, \hat{t}_j, \dots, t_n) \right) \right. \\ \left. + W_{g-1, n+1}^C(t, -t, t_2, \dots, t_n) + \sum_{\substack{\text{stable} \\ g_1+g_2=g \\ I \sqcup J = \{2,3,\dots,n\}}} W_{g_1, |I|+1}^C(t, t_I) W_{g_2, |J|+1}^C(-t, t_J) \right].$$

The last sum is restricted to the stable geometries, i.e., the partitions should satisfies $2g_1 - 1 + |I| > 0$ and $2g_2 - 1 + |J|$, as in (2.25). The contour integral is taken with respect to t on the exactly the same cycle defined by Figure 1.4 as before, where t is the normalization coordinate of (2.19). Note that the second and the third lines of (2.35) is a quadratic differential in the variable t . The notation $\frac{1}{dt}$ means the contraction with respect to the vector field $\frac{\partial}{\partial t}$ of the quadratic differential on $\tilde{\Sigma}$.

Remark 2.14. The recursion (2.35) is a genuine induction formula with respect to $2g - 2 + n$. Thus from $W_{0,2}^C$, we can calculate all $W_{g,n}^C$'s one by one. This is a big difference between (2.35) and (2.8). The latter relation contains the terms with $C_{g,n}$ in the right-hand side as well, therefore, $C_{g,n}$ is not determined as a function by an induction procedure.

Remark 2.15. If we apply (2.33) to $F_{0,2}^C$ of (2.31), then we obtain (2.32), not (2.34). The difference is the pull-back of the 2-form $\frac{dx_1 \cdot dx_2}{(x_1 - x_2)^2}$. This difference does not affect the recursion formula (2.35) for $2g - 2 + n > 1$. The only case affected is $(g, n) = (1, 1)$. The above recursion allows us to calculate $W_{1,1}^C$ from $W_{0,2}^C$ as we see below, but we cannot use (2.32) in place of $W_{0,2}^C$ for this case because the specialization $t_1 = t, t_2 = -t$ does not make sense in $d_{t_1} d_{t_2} F_{0,2}^C(t_1, t_2)$.

Remark 2.16. In Subsection 3.2, we will formulate the topological recursion as a universal formalism for the context of Hitchin spectral curves in a coordinate-free manner depending only on the geometric setting. There we will explain the meaning of $W_{0,2}$, and the formula for the topological recursion. At this moment, we note that the origin of the topological recursion is the edge-contraction mechanism of the Catalan recursion (2.8). There is a surprising relation between the edge-contraction operations and two dimensional topological quantum field theories. We refer to [26] for more detail.

Remark 2.17. The integral recursion (2.35) and the PDE recursion (2.25) are equivalent in the range of $2g - 2 + n \geq 2$, since if we know $W_{g,n}^C$ from the integral recursion, then we

can calculate $F_{g,n}^C$ by the integration

$$F_{g,n}^C(t_1, \dots, t_n) = \int_{-1}^{t_1} \cdots \int_{-1}^{t_n} W_{g,n}^C.$$

But the differential recursion does not provide any mechanism to calculate $F_{1,1}^C$ and $F_{0,3}^C$.

To see how the topological recursion works, let us compute $W_{1,1}^C(t_1)$ from (2.35).

$$\begin{aligned} W_{1,1}^C(t_1) &= -\frac{1}{64} \frac{1}{2\pi i} \left[\int_{\gamma} \left(\frac{1}{t+t_1} + \frac{1}{t-t_1} \right) \frac{(t^2-1)^3}{t^2} \cdot \frac{1}{dt} \cdot W_{0,2}^C(t, -t) \right] dt_1 \\ &= -\frac{1}{64} \frac{1}{2\pi i} \left[\int_{\gamma} \left(\frac{1}{t+t_1} + \frac{1}{t-t_1} \right) \frac{(t^2-1)^3}{t^2} \cdot \frac{1}{dt} \cdot \left(\frac{-(dt)^2}{(2t)^2} \right) \right] dt_1 \\ &= \frac{1}{256} \frac{1}{2\pi i} \left[\int_{\gamma} \left(\frac{1}{t+t_1} + \frac{1}{t-t_1} \right) \frac{(t^2-1)^3}{t^2} \cdot \frac{1}{t^2} \cdot dt \right] dt_1 \\ &= -\frac{1}{128} \frac{(t_1^2-1)^3}{t_1^4}. \end{aligned}$$

Here, we changed the contour integral to the negative of the residue calculations at $t = \pm t_1$, as indicated in Figure 1.4. From (2.26), we find that indeed

$$F_{1,1}^C(t_1) = \int_{-1}^{t_1} W_{1,1}^C.$$

As explained in [27], we can calculate $W_{0,3}^C$ from $W_{0,2}^C$ as well, which recovers the initial condition (2.27).

To understand the geometry behind the topological recursion, we need to identify each term of the formula. First, recall the normalization morphism

$$\tilde{\pi} : \tilde{\Sigma} \longrightarrow \mathbb{P}^1$$

of (2.16). We note that the transformation $t \mapsto -t$ appearing in the recursion formula is the **Galois conjugation** of the global Galois covering $\tilde{\pi} : \tilde{\Sigma} \longrightarrow \mathbb{P}^1$. From (2.19), we see that this transformation is induced by the the involution $y \mapsto -y$ of $T^*\mathbb{P}^1$. The fixed point of the Galois conjugation is the ramification points of the covering $\tilde{\pi}$, and the residue integration of (2.35) is taken around the two ramification points.

We claim that $W_{0,2}^C(t_1, t_2)$ is the *Cauchy differentiation kernel* on $\tilde{\Sigma}$. This comes from the intrinsic geometry of the curve $\tilde{\Sigma}$. The Cauchy differentiation kernel on \mathbb{P}^1 is the unique meromorphic symmetric bilinear differential form

$$(2.36) \quad c(t_1, t_2) := \frac{dt_1 \cdot dt_2}{(t_1 - t_2)^2}$$

on $\mathbb{P}^1 \times \mathbb{P}^1$ such that

$$(2.37) \quad df(t_2) = q_*(c(t_1, t_2)p^*f(t_1))$$

for every rational function f on \mathbb{P}^1 . Here, p and q are the projection maps

$$(2.38) \quad \begin{array}{ccc} & \mathbb{P}^1 \times \mathbb{P}^1 & \\ p \swarrow & & \searrow q \\ \mathbb{P}^1 & & \mathbb{P}^1 \end{array}$$

to the first and second factors. The push-forward q_* is defined by the integral

$$q_*(c(t_1, t_2)p^*f(t_1)) = \frac{1}{2\pi i} \oint c(t_1, t_2)f(t_1)$$

along a small loop in the fiber $q^*(t_2)$ that is centered at its intersection with the diagonal of $\Delta \subset \mathbb{P}^1 \times \mathbb{P}^1$. In terms of the other affine coordinate $u_i = 1/t_i$ of \mathbb{P}^1 , we have

$$\frac{dt_1 \cdot dt_2}{(t_1 - t_2)^2} = \frac{du_1 \cdot du_2}{(u_1 - u_2)^2} = -\frac{dt_1 \cdot du_2}{(t_1 u_2 - 1)^2} = -\frac{du_1 \cdot dt_2}{(u_1 t_2 - 1)^2}.$$

Therefore, $c(t_1, t_2)$ is a globally defined bilinear meromorphic form on $\mathbb{P}^1 \times \mathbb{P}^1$ with poles only along the diagonal Δ .

Remark 2.18. The Cauchy integration formula

$$f(w) = \frac{1}{2\pi i} \oint_{|z-w|<\epsilon} f(z) \frac{dz}{z-w}$$

is the most useful formula in complex analysis. The Cauchy integration kernel $dz/(z-w)$ is a globally defined meromorphic 1-form on \mathbb{C} that has only one pole, and its order is 1. It has to be noted that on a compact Riemann surface Σ , we do not have such a form. The best we can do is the meromorphic 1-form $\omega_\Sigma^{a-b}(t)$ with the following properties:

- It is a globally defined meromorphic 1-form with a pole of order 1 and residue 1 at a and a pole of order 1 and residue -1 at b for any pair (a, b) of distinct points of Σ .
- It is holomorphic in t except for $t = a$ and $t = b$.

Since we can add any holomorphic 1-form to $\omega_\Sigma^{a-b}(t)$ without violating the characteristic properties, the ambiguity of this form is exactly $H^0(\Sigma, K_\Sigma)$. Therefore, it is unique on $\Sigma = \mathbb{P}^1$, which is given by

$$(2.39) \quad \omega_{\mathbb{P}^1}^{a-b}(t) = \left(\frac{1}{t-a} - \frac{1}{t-b} \right) dt.$$

Note that

$$d_{t_2} \omega_{\mathbb{P}^1}^{t_2-b}(t_1) = c(t_1, t_2).$$

for any b .

Let us go back to the topological recursion for the Catalan numbers (2.35). Since all $W_{g,n}^C$ are determined by $W_{0,2}^C$ using the recursion, the remaining quantity we need to identify is the *integration kernel*. Recall the tautological 1-form

$$\eta_{\mathbb{P}^1} := ydx$$

on the tangent bundle $T^*\mathbb{P}^1$. We can see from (2.16) that its pull-back to the normalized spectral curve $\tilde{\Sigma}$ is given by

$$(2.40) \quad \begin{aligned} \eta(t) &= - \left(\frac{t+1}{t-1} \right) d \left(\frac{4}{t^2-1} \right) \\ &= \frac{8t}{(t+1)(t-1)^3} dt. \end{aligned}$$

The integration kernel is then given by

$$(2.41) \quad \frac{1}{2} \frac{\omega_{\mathbb{P}^1}^{(-t)-t}(t_1)}{\eta(-t) - \eta(t)} = -\frac{1}{64} \left(\frac{1}{t+t_1} + \frac{1}{t-t_1} \right) \frac{(t^2-1)^3}{t^2} \cdot \frac{1}{dt} \cdot dt_1.$$

2.6. The quantum curve for Catalan numbers. There is yet another important role the differential recursion (2.25) plays. This is the derivation of the quantum curve equation for the Catalan case. In terms of the coordinates (x, y) of (2.13), the spectral curve is $y^2 + xy + 1 = 0$. The generating function $z(x)$ of (2.10), the normalization coordinate t of (2.19), and the base curve coordinate x on \mathbb{P}^1 are related by

$$(2.42) \quad t = t(x) := \frac{z(x) + 1}{z(x) - 1},$$

which we consider as a function of x that gives the branch of $\tilde{\Sigma}$ determined by

$$\lim_{x \rightarrow +\infty} t(x) = -1.$$

Theorem 2.19 (Quantum curve for generalized Catalan numbers, [67]). *Define*

$$(2.43) \quad \Psi(t, \hbar) := \exp \left(\sum_{g=0}^{\infty} \sum_{n=1}^{\infty} \frac{1}{n!} \hbar^{2g-2+n} F_{g,n}^C(t, t, \dots, t) \right),$$

incorporating (2.25), (2.29), and (2.31). Then as a function in x through (2.42), $\Psi(t(x), \hbar)$ satisfies the following differential equation

$$(2.44) \quad \left[\left(\hbar \frac{d}{dx} \right)^2 + x \left(\hbar \frac{d}{dx} \right) + 1 \right] \Psi(t(x), \hbar) = 0.$$

The semi-classical limit of (2.44) using $S_0(x) = F_{0,1}^C(t(x))$ coincides with the spectral curve

$$y^2 + xy + 1 = 0$$

of (2.13).

Remark 2.20. Since $t = t(x)$ is a complicated function, it is surprising to see that $\Psi(t, \hbar)$ satisfies such a simple equation as (2.44).

Remark 2.21. The definition (2.43) and the meaning of (2.44) is the same as in the situation we have explained in Introduction. First, we define

$$(2.45) \quad S_m(x) := \sum_{2g-2+n=m-1} \frac{1}{n!} F_{g,n}^C(t(x), t(x), \dots, t(x))$$

for every $m \geq 0$. We then interpret (2.44) as

$$(2.46) \quad \left[e^{-\frac{1}{\hbar} S_0(x)} \left(\left(\hbar \frac{d}{dx} \right)^2 + x \left(\hbar \frac{d}{dx} \right) + 1 \right) e^{\frac{1}{\hbar} S_0(x)} \right] \exp \left(\sum_{m=1}^{\infty} \hbar^{m-1} S_m(x) \right) = 0.$$

To prove Theorem 2.19, let us recall a lemma from [67]:

Lemma 2.22. *Consider an open coordinate neighborhood $U \subset \Sigma$ of a projective algebraic curve Σ with a coordinate t . Let $f(t_1, \dots, t_n)$ be a meromorphic symmetric function in n variables defined on Σ^n . Then on the coordinate neighborhood U^n , we have*

$$(2.47) \quad \begin{aligned} \frac{d}{dt} f(t, t, \dots, t) &= n \left[\frac{\partial}{\partial u} f(u, t, \dots, t) \right] \Big|_{u=t}; \\ \frac{d^2}{dt^2} f(t, t, \dots, t) &= n \left[\frac{\partial^2}{\partial u^2} f(u, t, \dots, t) \right] \Big|_{u=t} \\ &\quad + n(n-1) \left[\frac{\partial^2}{\partial u_1 \partial u_2} f(u_1, u_2, t, \dots, t) \right] \Big|_{u_1=u_2=t}. \end{aligned}$$

For a meromorphic function in one variable $f(t)$ on Σ , we have

$$(2.48) \quad \lim_{t_2 \rightarrow t_1} \left[\omega_{\Sigma}^{t_2-b}(t_1)(f(t_1) - f(t_2)) \right] = d_1 f(t_1),$$

where $\omega_{\Sigma}^{t_2-b}(t_1)$ is the 1-form of Remark 2.18.

Proof of Theorem 2.19. Differential forms can be restricted to a subvariety, but partial differential equations cannot be restricted to a subvariety in general. Therefore, it is non-trivial that (2.25) has any meaningful restriction to the diagonal of $(\mathbb{P}^1)^n$. Our strategy is the following. First, we expand (2.46) with respect to powers of \hbar , and derive an infinite system of ordinary differential equations for a finite collection of $S_m(x)$'s. (This method is known as the WKB analysis.) We then prove that these ODEs are exactly the **principal specialization** of (2.25), using (2.47).

Let

$$(2.49) \quad F(x, \hbar) := \sum_{m=0}^{\infty} \hbar^{m-1} S_m(x).$$

Unlike the ill-defined expression (2.43), $F(x, \hbar)$ is just a generating function of $S_m(x)$'s. Then (2.44), interpreted as (2.46), is equivalent to

$$(2.50) \quad \hbar^2 \frac{d^2}{dx^2} F + \hbar^2 \frac{dF}{dx} \frac{dF}{dx} + x \hbar \frac{dF}{dx} + 1 = 0.$$

The \hbar -expansion of (2.50) gives

$$(2.51) \quad \hbar^0\text{-terms : } (S'_0(x))^2 + x S'_0(x) + 1 = 0,$$

$$(2.52) \quad \hbar^1\text{-terms : } 2S'_0(x)S'_1(x) + S''_0(x) + x S'_1(x) = 0,$$

$$(2.53) \quad \hbar^{m+1}\text{-terms : } S''_m(x) + \sum_{a+b=m+1} S'_a(x)S'_b(x) + x S'_{m+1}(x) = 0, \quad m \geq 1,$$

where $'$ denotes the x -derivative. The WKB method is to solve these equations iteratively and find $S_m(x)$ for all $m \geq 0$. Here, (2.51) is the **semi-classical limit** of (2.44), and (2.52) is the *consistency condition* we need for solving the WKB expansion. Since $dS_0(x) = y dx$, we have $y = S'_0(x)$. Thus (2.51) follows from the spectral curve equation (2.13) and the definition of $F_{0,1}^C(t)$ of (2.29).

Recalling $x = z + 1/z$, we obtain

$$\frac{d}{dx} = \frac{z^2}{z^2 - 1} \frac{d}{dz}.$$

In z -coordinate, $F_{0,2}^C(z, z) = -\log(1 - z^2)$, which follows from (2.31). Therefore,

$$S'_1(x) = -\frac{1}{2} \frac{z^2}{z^2 - 1} \frac{d}{dz} \log(1 - z^2) = -\frac{z^3}{(z^2 - 1)^2}.$$

We can then calculate

$$\begin{aligned} 2S'_0(x)S'_1(x) + S''_0(x) + x S'_1(x) &= 2z \frac{z^3}{(z^2 - 1)^2} - \frac{z^2}{z^2 - 1} - \left(z + \frac{1}{z} \right) \frac{z^3}{(z^2 - 1)^2} \\ &= \frac{1}{(z^2 - 1)^2} (2z^4 - z^2(z^2 - 1) - z^4 - z^2) \\ &= 0. \end{aligned}$$

Therefore, (2.52) holds. We refer to [25] for the proof of (2.53) from (2.25) and (2.22). \square

Remark 2.23. The solution $\Psi(t(x), \hbar)$ is a formal section of $K_{\mathbb{P}^1}^{-\frac{1}{2}}$, as before. Note that the quantum curve (2.44) has an irregular singularity at $x = \infty$, and hence its solution has an essential singularity at the infinity. The expression (2.43) gives the asymptotic expansion of a solution around

$$t \rightarrow -1 \iff x \rightarrow \infty.$$

The expansion in $1/x$ for $\hbar > 0$ is given in (3.50) using the Tricomi confluent hypergeometric function.

2.7. Counting lattice points on the moduli space $\mathcal{M}_{g,n}$. The topological recursion (2.35) is a consequence of (2.25), and the PDE recursion (2.25) is essentially the Laplace transform of the combinatorial formula (2.8). Then from where does the relation to the intersection numbers (2.24) arise?

To see this relation, we need to consider the dual of the cell graphs. They are commonly known as Grothendieck's dessins d'enfants (see for example, [78]), or ribbon graphs, as mentioned earlier. Recall that a ribbon graph has unlabeled vertices and edges, but faces are labeled. A *metric* ribbon graph is a ribbon graph with a positive real number (the length) assigned to each edge. For a given ribbon graph Γ with $e = e(\Gamma)$ edges, the space of metric ribbon graphs is $\mathbb{R}_+^{e(\Gamma)}/\text{Aut}(\Gamma)$, where the automorphism group acts by permutations of edges (see [63, Section 1]). We restrict ourselves to the case that $\text{Aut}(\Gamma)$ fixes each 2-cell of the cell-decomposition. We also require that every vertex of a ribbon graph has degree (i.e., valence) 3 or more. Using the canonical holomorphic coordinate systems on a topological surface of [63, Section 4] and the Strebel differentials [81], we have an isomorphism of topological orbifolds

$$(2.54) \quad \mathcal{M}_{g,n} \times \mathbb{R}_+^n \cong RG_{g,n}.$$

Here

$$RG_{g,n} = \coprod_{\substack{\Gamma \text{ ribbon graph} \\ \text{of type } (g,n)}} \frac{\mathbb{R}_+^{e(\Gamma)}}{\text{Aut}(\Gamma)}$$

is the orbifold consisting of metric ribbon graphs of a given topological type (g,n) . The gluing of orbi-cells is done by making the length of a non-loop edge tend to 0. The space $RG_{g,n}$ is a smooth orbifold (see [63, Section 3] and [80]). We denote by $\pi : RG_{g,n} \rightarrow \mathbb{R}_+^n$ the natural projection via (2.54), which is the assignment of the collection of perimeter length of each boundary to a given metric ribbon graph.

Consider a ribbon graph Γ whose faces are labeled by $[n] = \{1, 2, \dots, n\}$. For the moment let us give a label to each edge of Γ by an index set $[e] = \{1, 2, \dots, e\}$. The edge-face incidence matrix is then defined by

$$A_\Gamma = [a_{i\eta}]_{i \in [n], \eta \in [e]};$$

$a_{i\eta} =$ the number of times edge η appears in face i .

Thus $a_{i\eta} = 0, 1$, or 2 , and the sum of the entries in each column is always 2. The Γ contribution of the space $\pi^{-1}(p_1, \dots, p_n) = RG_{g,n}(\mathbf{p})$ of metric ribbon graphs with a prescribed perimeter $\mathbf{p} = (p_1, \dots, p_n)$ is the orbifold polytope

$$P_\Gamma(\mathbf{p})/\text{Aut}(\Gamma), \quad P_\Gamma(\mathbf{p}) = \{\mathbf{x} \in \mathbb{R}_+^e \mid A_\Gamma \mathbf{x} = \mathbf{p}\},$$

where $\mathbf{x} = (\ell_1, \dots, \ell_e)$ is the collection of edge lengths of the metric ribbon graph Γ . We have

$$\sum_{i \in [n]} p_i = \sum_{i \in [n]} \sum_{\eta \in [e]} a_{i\eta} \ell_\eta = 2 \sum_{\eta \in [e]} \ell_\eta.$$

We recall the topological recursion for the number of metric ribbon graphs $RG_{g,n}^{\mathbb{Z}_+}$ whose edges have integer lengths, following [15]. We call such a ribbon graph an *integral ribbon graph*. We can interpret an integral ribbon graph as Grothendieck's *dessin d'enfant* by considering an edge of integer length as a chain of edges of length one connected by bivalent vertices, and reinterpreting the notion of $\text{Aut}(\Gamma)$ suitably. Since we do not go into the number theoretic aspects of dessins, we stick to the more geometric notion of integral ribbon graphs.

Definition 2.24. The weighted number $|RG_{g,n}^{\mathbb{Z}_+}(\mathbf{p})|$ of integral ribbon graphs with prescribed perimeter lengths $\mathbf{p} \in \mathbb{Z}_+^n$ is defined by

$$(2.55) \quad N_{g,n}(\mathbf{p}) = |RG_{g,n}^{\mathbb{Z}_+}(\mathbf{p})| = \sum_{\substack{\Gamma \text{ ribbon graph} \\ \text{of type } (g,n)}} \frac{|\{\mathbf{x} \in \mathbb{Z}_+^{e(\Gamma)} \mid A_\Gamma \mathbf{x} = \mathbf{p}\}|}{|\text{Aut}(\Gamma)|}.$$

Since the finite set $\{\mathbf{x} \in \mathbb{Z}_+^{e(\Gamma)} \mid A_\Gamma \mathbf{x} = \mathbf{p}\}$ is a collection of lattice points in the polytope $P_\Gamma(\mathbf{p})$ with respect to the canonical integral structure $\mathbb{Z} \subset \mathbb{R}$ of the real numbers, $N_{g,n}(\mathbf{p})$ can be thought of counting the number of *lattice points* in $RG_{g,n}(\mathbf{p})$ with a weight factor $1/|\text{Aut}(\Gamma)|$ for each ribbon graph. The function $N_{g,n}(\mathbf{p})$ is a symmetric function in $\mathbf{p} = (p_1, \dots, p_n)$ because the summation runs over all ribbon graphs of topological type (g, n) .

Remark 2.25. Since the integral vector \mathbf{x} is restricted to take strictly positive values, we would have $N_{g,n}(\mathbf{p}) = 0$ if we were to substitute $\mathbf{p} = 0$. This normalization is natural from the point of view of lattice point counting and Grothendieck's *dessins d'enphants*. However, we do not make such a substitution in these lectures because we consider \mathbf{p} as a strictly positive integer vector. This situation is similar to Hurwitz theory [33, 68], where a partition μ is a strictly positive integer vector that plays the role of our \mathbf{p} . We note that a different assignment of values was suggested in [71, 72].

For brevity of notation, we denote by $p_I = (p_i)_{i \in I}$ for a subset $I \in [n] = \{1, 2, \dots, n\}$. The cardinality of I is denoted by $|I|$. The following topological recursion formula was proved in [15] using the idea of ciliation of a ribbon graph.

Theorem 2.26 ([15]). *The number of integral ribbon graphs with prescribed boundary lengths satisfies the topological recursion formula*

$$(2.56) \quad p_1 N_{g,n}(p_{[n]}) = \frac{1}{2} \sum_{j=2}^n \left[\sum_{q=0}^{p_1+p_j} q(p_1+p_j-q) N_{g,n-1}(q, p_{[n] \setminus \{1,j\}}) \right. \\ \left. + H(p_1-p_j) \sum_{q=0}^{p_1-p_j} q(p_1-p_j-q) N_{g,n-1}(q, p_{[n] \setminus \{1,j\}}) \right. \\ \left. - H(p_j-p_1) \sum_{q=0}^{p_j-p_1} q(p_j-p_1-q) N_{g,n-1}(q, p_{[n] \setminus \{1,j\}}) \right]$$

$$\begin{aligned}
& + \frac{1}{2} \sum_{0 \leq q_1 + q_2 \leq p_1} q_1 q_2 (p_1 - q_1 - q_2) \left[N_{g-1, n+1}(q_1, q_2, p_{[n] \setminus \{1\}}) \right. \\
& \qquad \qquad \qquad \left. + \sum_{\substack{\text{stable} \\ g_1 + g_2 = g \\ I \sqcup J = [n] \setminus \{1\}}} N_{g_1, |I|+1}(q_1, p_I) N_{g_2, |J|+1}(q_2, p_J) \right].
\end{aligned}$$

Here

$$H(x) = \begin{cases} 1 & x > 0 \\ 0 & x \leq 0 \end{cases}$$

is the Heaviside function, and the last sum is taken for all partitions $g = g_1 + g_2$ and $I \sqcup J = \{2, 3, \dots, n\}$ subject to the stability conditions $2g_1 - 1 + I > 0$ and $2g_2 - 1 + |J| > 0$.

For a fixed (g, n) in the stable range, i.e., $2g - 2 + n > 0$, we choose n variables t_1, t_2, \dots, t_n , and define the function

$$(2.57) \quad z(t_i, t_j) = \frac{(t_i + 1)(t_j + 1)}{2(t_i + t_j)}.$$

An edge η of a ribbon graph Γ bounds two faces, say i_η and j_η . These two faces may be actually the same. Now we define the *Poincaré polynomial* [64] of $RG_{g,n}$ in the z -variables by

$$(2.58) \quad F_{g,n}^P(t_1, \dots, t_n) = \sum_{\substack{\Gamma \text{ ribbon graph} \\ \text{of type } (g,n)}} \frac{(-1)^{e(\Gamma)}}{|\text{Aut}(\Gamma)|} \prod_{\substack{\eta \text{ edge} \\ \text{of } \Gamma}} z(t_{i_\eta}, t_{j_\eta}),$$

which is a polynomial in $z(t_i, t_j)$ but actually a symmetric rational function in t_1, \dots, t_n .

Let us consider the *Laplace transform*

$$(2.59) \quad L_{g,n}(w_1, \dots, w_n) \stackrel{\text{def}}{=} \sum_{\mathbf{p} \in \mathbb{Z}_+^n} N_{g,n}(\mathbf{p}) e^{-\langle \mathbf{p}, w \rangle}$$

of the number of integral ribbon graphs $N_{g,n}(\mathbf{p})$, where $\langle \mathbf{p}, w \rangle = p_1 w_1 + \dots + p_n w_n$, and the summation is taken over all integer vectors $\mathbf{p} \in \mathbb{Z}_+^n$ of strictly positive entries. We shall prove that after the coordinate change of [15] from the w -coordinates to the t -coordinates defined by

$$(2.60) \quad e^{-w_j} = \frac{t_j + 1}{t_j - 1}, \quad j = 1, 2, \dots, n,$$

the Laplace transform $L_{g,n}(w_{[n]})$ becomes the Poincaré polynomial

$$F_{g,n}^P(t_1, \dots, t_n) = L_{g,n}(w_1(t), \dots, w_n(t)).$$

The Laplace transform $L_{g,n}(w_N)$ can be evaluated using the definition of the number of integral ribbon graphs (2.55). Let a_η be the η -th column of the incidence matrix A_Γ so that

$$A_\Gamma = [a_1 | a_2 | \dots | a_{e(\Gamma)}].$$

Then

$$(2.61) \quad L_{g,n}(w_{[n]}) = \sum_{\mathbf{p} \in \mathbb{Z}_+^n} N_{g,n}(\mathbf{p}) e^{-\langle \mathbf{p}, w \rangle}$$

$$\begin{aligned}
 &= \sum_{\substack{\Gamma \text{ ribbon graph} \\ \text{of type } (g,n)}} \sum_{\mathbf{p} \in \mathbb{Z}_+^n} \frac{1}{|\text{Aut}(\Gamma)|} |\{\mathbf{x} \in \mathbb{Z}_+^{e(\Gamma)} \mid A_\Gamma \mathbf{x} = \mathbf{p}\}| e^{-\langle \mathbf{p}, w \rangle} \\
 &= \sum_{\substack{\Gamma \text{ ribbon graph} \\ \text{of type } (g,n)}} \frac{1}{|\text{Aut}(\Gamma)|} \sum_{\mathbf{x} \in \mathbb{Z}_+^{e(\Gamma)}} e^{-\langle A_\Gamma \mathbf{x}, w \rangle} \\
 &= \sum_{\substack{\Gamma \text{ ribbon graph} \\ \text{of type } (g,n)}} \frac{1}{|\text{Aut}(\Gamma)|} \prod_{\substack{\eta \text{ edge} \\ \text{of } \Gamma}} \sum_{\ell_\eta=1}^{\infty} e^{-\langle a_\eta, w \rangle \ell_\eta} \\
 &= \sum_{\substack{\Gamma \text{ ribbon graph} \\ \text{of type } (g,n)}} \frac{1}{|\text{Aut}(\Gamma)|} \prod_{\substack{\eta \text{ edge} \\ \text{of } \Gamma}} \frac{e^{-\langle a_\eta, w \rangle}}{1 - e^{-\langle a_\eta, w \rangle}}.
 \end{aligned}$$

Every edge η bounds two faces, which we call face i_η^+ and face i_η^- . When $a_{i\eta} = 2$, these faces are the same. We then calculate

$$(2.62) \quad \frac{e^{-\langle a_\eta, w \rangle}}{1 - e^{-\langle a_\eta, w \rangle}} = -z\left(t_{i_\eta^+}, t_{i_\eta^-}\right).$$

This follows from (2.60) and

$$\begin{aligned}
 \frac{e^{-(w_i+w_j)}}{1 - e^{-(w_i+w_j)}} &= \frac{\frac{(t_i+1)(t_j+1)}{(t_i-1)(t_j-1)}}{1 - \frac{(t_i+1)(t_j+1)}{(t_i-1)(t_j-1)}} = -\frac{(t_i+1)(t_j+1)}{2(t_i+t_j)} = -z(t_i, t_j), \\
 \frac{e^{-2w_i}}{1 - e^{-2w_i}} &= -\frac{(t_i+1)^2}{4t_i} = -z(t_i, t_i).
 \end{aligned}$$

Note that since $z(t_i, t_j)$ is a symmetric function, which face is named i_η^+ or i_η^- does not matter. From (2.61) and (2.62), we have established

Theorem 2.27 (The Poincaré polynomials and the Laplace transform, [64]). *The Laplace transform $L_{g,n}(w_{[n]})$ in terms of the t -coordinates (2.60) is the Poincaré polynomial*

$$(2.63) \quad F_{g,n}^P(t_{[n]}) = \sum_{\substack{\Gamma \text{ ribbon graph} \\ \text{of type } (g,n)}} \frac{(-1)^{e(\Gamma)}}{|\text{Aut}(\Gamma)|} \prod_{\substack{\eta \text{ edge} \\ \text{of } \Gamma}} z\left(t_{i_\eta^+}, t_{i_\eta^-}\right).$$

Corollary 2.28. *The evaluation of $F_{g,n}^P(t_{[n]})$ at $t_1 = \dots = t_n = 1$ gives the Euler characteristic of $RG_{g,n}$*

$$(2.64) \quad F_{g,n}^P(1, 1, \dots, 1) = \chi(RG_{g,n}) = (-1)^n \chi(\mathcal{M}_{g,n}).$$

Furthermore, if we evaluate at $t_j = -1$ for any j , then we have

$$(2.65) \quad F_{g,n}^P(t_1, t_2, \dots, t_n) \Big|_{t_j=-1} = 0$$

as a function in the rest of the variables $t_{[n] \setminus \{j\}}$.

Proof. The Euler characteristic calculation immediately follows from $z(1, 1) = 1$.

Consider a ribbon graph Γ of type (g, n) . Its j -th face has at least one edge on its boundary. Therefore,

$$\prod_{\eta \text{ edge of } \Gamma} z\left(t_{i_\eta^+}, t_{i_\eta^-}\right)$$

has a factor $(t_j + 1)$ by (2.57). It holds for every ribbon graph Γ in the summation of (2.63). Therefore, (2.65) follows. \square

The following theorem is established in [64]. Its proof is the computation of the Laplace transform of the lattice point recursion (2.56).

Theorem 2.29 (Differential recursion for the Poincaré polynomials, [64]). *The Poincaré polynomials $F_{g,n}^P(t_1, \dots, t_n)$ satisfy exactly the same differential recursion (2.25) with the same initial values of $F_{1,1}^P$ and $F_{0,3}^P$.*

Since the recursion uniquely determines all $F_{g,n}$'s for $2g - 2 + n > 0$, we have the following:

Corollary 2.30. *For every (g, n) in the stable range $2g - 2 + n > 0$, the two functions are identical:*

$$(2.66) \quad F_{g,n}^C(t_1, \dots, t_n) = F_{g,n}^P(t_1, \dots, t_n).$$

Because of this identification, we see that (2.64) implies (2.23), and (2.22) follows from (2.65). We can also see how (2.24) holds.

The limit $t_i \rightarrow \infty$ corresponds to $x_i \rightarrow 2$ through the normalization coordinate of (2.19), and $x_i = 2$ corresponds to a branch point of $\tilde{\pi} : \tilde{\Sigma} \rightarrow \mathbb{P}^1$ of (2.16). The defining equation (2.20) of $F_{g,n}^C$ does not tell us what we obtain by taking the limit $x_i \rightarrow 2$. The geometric meaning of the $t_i \rightarrow \infty$ limit becomes crystal clear by the equality (2.66). Let us take a look at the definition of the Poincaré polynomial (2.58). The fact that it is a Laurent polynomial follows from the recursion (2.25) by induction. If $|t_i| > |t_j| \gg 1$, then

$$z(t_i, t_j) = \frac{(t_i + 1)(t_j + 1)}{2(t_i + t_j)} \sim \frac{1}{2}t_j.$$

Therefore, the highest degree part of $F_{g,n}^P$ comes from the graphs of type (g, n) with the largest number of edges. Denoting the number of vertices of a ribbon graph Γ by $v(\Gamma)$, we have

$$2 - 2g - n = v(\Gamma) - e(\Gamma).$$

To maximize $e(\Gamma)$, we need to maximize $v(\Gamma)$, which is achieved by taking a trivalent graph (since we do not allow degree 1 and 2 vertices in our ribbon graph). By counting the number of half-edges of a trivalent graph, we obtain $2e(\Gamma) = 3v(\Gamma)$. Hence we have

$$(2.67) \quad e(\Gamma) = 6g - 6 + 3n.$$

This is the degree of $F_{g,n}^P$, which agrees with the degree of (2.24), and also consistent with the dimension of (2.54).

Fix a point $\mathbf{p} \in \mathbb{Z}_+^n$, and scale it by a large integer $\lambda \gg 1$. Then from (2.55) we see that the number of lattice points in the polytope $P_\Gamma(\lambda\mathbf{p})/\text{Aut}(\Gamma)$ that is counted as a part of $N_{g,n}(\lambda\mathbf{p})$ is the same as the number of scaled lattice points $\mathbf{x} \in \frac{1}{\lambda}\mathbb{Z}_+^n$ in $P_\Gamma(\mathbf{p})/\text{Aut}(\Gamma)$. As $\lambda \rightarrow \infty$, the number of lattice points can be approximated by the Euclidean volume of the polytope (cf. theory of Ehrhart polynomials).

For a fixed (w_1, \dots, w_n) with $\text{Re}(w_j) > 0$, the contribution from large \mathbf{p} 's in the Laplace transform $L_{g,n}(w_1, \dots, w_n)$ of (2.59) is small. The asymptotic behavior of $L_{g,n}$ as $w \rightarrow 0$ picks up the large perimeter contribution of $N_{g,n}(\mathbf{p})$, or the counting of the lattice points of smaller and smaller mesh size. Since

$$t_j \rightarrow \infty \iff w_j \rightarrow 0,$$

the large t_j behavior of $F_{g,n}^P$, which is a homogeneous polynomial of degree $6g - 6 + 3n$, reflects the information of the volume of $\mathcal{M}_{g,n}$ in its coefficients. From Kontsevich [55], we learn that the volume is exactly the intersection number appearing in (2.24).

The topological recursion for the Airy case (1.41) is the $t \rightarrow \infty$ limit of the Catalan topological recursion (2.35), as we see from the limit

$$\frac{(t^2 - 1)^3}{t^2} \rightarrow t^4.$$

Since the integrand of (1.41) has no poles at $t = 0$, the small circle of the contour γ does not contribute any residue. Thus we have derived the Airy topological recursion from the Catalan topological recursion.

3. QUANTIZATION OF SPECTRAL CURVES

Quantum curves assemble information of quantum invariants in a compact manner. The global nature of quantum curves is not well understood at this moment of writing. In this section, we focus on explaining the relation between the topological recursion and the *local* expression of quantum curves, suggested for example, in [44]. We give the precise definition of the topological recursion in a geometric and coordinate-free manner. Our formulation depends only on the geometric data we start with for the context of Higgs bundles. Global definition of the quantum curves are not established as of now. We refer to [23] for a more recent development.

3.1. Geometry of non-singular Hitchin spectral curves of rank 2. We need to make some choices to determine the differential form $W_{0,2}$ that gives the initial value of the topological recursion. It can reflect many aspects of the Hitchin spectral curve. In these lectures, we choose the Cauchy differentiation kernel as $W_{0,2}$, that depends only on the intrinsic geometry of the Hitchin spectral curve, for our purpose. Of course one can make other choices for different purposes, such as the kernel associated with a connection of a spectral line bundle on the Hitchin spectral curve.

Even for the Cauchy differentiation kernel, there is no canonical choice. It depends on a symplectic basis for the homology of the spectral curve. We give one particular choice in this subsection.

Throughout this section, we denote by C a smooth projective curve of genus $g(C)$ defined over \mathbb{C} . As before, K_C is the canonical bundle on C . The cotangent bundle $\pi : T^*C \rightarrow C$ is the total space of K_C , and there is the tautological section

$$(3.1) \quad \eta \in H^0(T^*C, \pi^*K_C),$$

which is a globally defined holomorphic 1-form on T^*C .

$$(3.2) \quad \begin{array}{ccc} & \pi^*(K_C) & \\ & \swarrow & \\ T^*C & & K_C \\ & \searrow \pi & \swarrow \\ & C & \end{array}$$

Choose a *generic* quadratic differential $s \in H^0(C, K_C^{\otimes 2})$, so that the spectral curve

$$(3.3) \quad \Sigma_s \subset T^*C$$

that is defined by

$$(3.4) \quad \eta^{\otimes 2} + \pi^*s = 0$$

is non-singular. Our spectral curve $\Sigma = \Sigma_s$ is a double sheeted ramified covering of C defined by (3.4). The genus of the spectral curve is $g(\Sigma) = 4g(C) - 3$. This is because a generic s has $\deg(K_C^{\otimes 2}) = 4g(C) - 4$ simple zeros, which correspond to branch points of the covering $\pi : \Sigma \rightarrow C$. Thus the genus is calculated by the Riemann Hurwitz formula

$$2(2 - 2g(C) - (4g(C) - 4)) = 2 - 2g(\Sigma) - (4g(C) - 4).$$

The cotangent bundle T^*C has a natural involution

$$(3.5) \quad \sigma : T^*C \supset T_x^*C \ni (x, y) \mapsto (x, -y) \in T_x^*C \subset T^*C,$$

which preserves the spectral curve Σ . The action of this involution is the deck-transformation. Indeed, the covering $\pi : \Sigma \rightarrow C$ is a **Galois covering** with the Galois group $\mathbb{Z}/2\mathbb{Z} = \langle \sigma \rangle$.

Let $R \subset \Sigma$ denote the ramification divisor of this covering. Because Σ is non-singular, R is supported at $4g(C) - 4$ distinct points that are determined by $s = 0$ on C . We consider C as the 0-section of T^*C . Thus both C and Σ are divisors of T^*C . Hence R is also defined as $C \cdot \Sigma$ in T^*C supported on $C \cap \Sigma$. Note that η vanishes only along $C \subset T^*C$. As a holomorphic 1-form on Σ , $\iota^*\eta$ has $2g(\Sigma) - 2 = 8g(C) - 8$ zeros on Σ . Thus it has a degree 2 zero at each point of $\text{supp}(R)$.

As explained in the earlier sections using examples, we wish to choose a differential form $W_{0,2}$ for our geometric situation. For this purpose, let us recall the differential form ω_Σ^{a-b} of Remark 2.18. Since we can add any holomorphic 1-form to ω_Σ^{a-b} , we need to impose $g(\Sigma)$ independent conditions to make it unique. If we have a principal polarization of the period matrix for Σ , then one obvious choice would be to impose

$$(3.6) \quad \oint_{a_j} \omega_\Sigma^{a-b} = 0$$

for all “ A -cycles” a_j , $j = 1, 2, \dots, g(\Sigma)$, following Riemann himself. The reason for this canonical choice is that we can make it for a family of spectral curves

$$\{\Sigma_s\}_{s \in U},$$

where $U \subset H^0(C, K_C^{\otimes 2})$ is a contractible open neighborhood of s . Since the base curve C is fixed, we can choose and fix a symplectic basis for $H_1(C, \mathbb{Z})$:

$$(3.7) \quad \langle A_1, \dots, A_g; B_1, \dots, B_g \rangle = H_1(C, \mathbb{Z}).$$

From this choice, we construct a canonical symplectic basis for $H^1(\Sigma, \mathbb{Z})$ as follows. Let us label points of $R = \{p_1, p_2, \dots, p_{4g-4}\}$, where $g = g(C)$. We can connect p_{2i} and p_{2i+1} , $i = 1, \dots, 2g - 3$, with a simple path on Σ that is mutually non-intersecting so that $\pi^*(\overline{p_{2i}p_{2i+1}})$, $i = 1, \dots, 2g - 3$, form a part of the basis for $H_1(\Sigma, \mathbb{Z})$. We denote these cycles by $\alpha_1, \dots, \alpha_{2g-3}$. Since π is locally homeomorphic away from R , we have g cycles a_1, \dots, a_g on Σ_s so that $\pi_*(a_j) = A_j$ for $j = 1, \dots, g$, where A_j 's are the A -cycles of C chosen as (3.7). We define the A -cycles of Σ to be the set

$$(3.8) \quad \{a_1, \dots, a_g, \sigma_*(a_1), \dots, \sigma_*(a_g), \alpha_1, \dots, \alpha_{2g-3}\} \subset H_1(\Sigma, \mathbb{Z}),$$

where σ is the Galois conjugation (see Figure 3.1). Clearly, this set can be extended into a symplectic basis for $H_1(\Sigma, \mathbb{Z})$. This choice of the symplectic basis trivializes the homology bundle

$$\{H_1(\Sigma_s, \mathbb{Z})\}_{s \in U} \rightarrow U \subset H^0(C, K_C^{\otimes 2})$$

locally on the contractible neighborhood U .




FIGURE 3.1. The choice of a symplectic basis for $H_1(\Sigma, \mathbb{Z})$ from that of $H_1(C, \mathbb{Z})$.

3.2. The generalized topological recursion for the Hitchin spectral curves. The topological recursion of [11, 34], that has initiated the explosive developments on this subject in recent years, is restricted to non-singular spectral curves in \mathbb{C}^2 or $(\mathbb{C}^*)^2$. A systematic generalization of the formalism to the case of Hitchin spectral curves are given for the first time in [24, 25]. First, we gave the definition for non-singular Hitchin spectral curves associated with holomorphic Higgs bundles in [24]. We then extended the consideration to meromorphic Higgs bundles and singular spectral curves in [25]. In each case, however, the actual evaluation of the generalized topological recursion for the purpose of quantization of the Hitchin spectral curves is limited to Higgs bundles of rank 2. This is due to many technical difficulties, and at this point we still do not have a better understanding of the theory in its full generality.

The purpose of this subsection is thus to present the theory in the way we know as of now, with the scope limited to what seems to work. Many aspects of the story can be immediately generalized. Mainly for the sake of simplicity of presentation, we concentrate on the case of rank 2 Higgs bundles.

The geometric setup is the following. We have a smooth projective curve C defined over \mathbb{C} of an arbitrary genus, and a meromorphic Higgs bundle (E, ϕ) . Here, the vector bundle E of rank 2 is a special one of the form

$$(3.9) \quad E = K_C^{-\frac{1}{2}} \oplus K_C^{\frac{1}{2}}.$$

As a meromorphic Higgs field, we use

$$(3.10) \quad \phi = \begin{bmatrix} & 1 \\ -a_2 & -a_1 \end{bmatrix} : E \longrightarrow K_C(D) \otimes E$$

with poles at an effective divisor D of C , where

$$\begin{aligned} a_1 &\in H^0(C, K_C(D)) \\ a_2 &\in H^0(C, K_C^{\otimes 2}(D)). \end{aligned}$$

Although ϕ involves a quadratic differential in its coefficient, since

$$\begin{aligned} \begin{bmatrix} & 1 \\ -a_2 & a_1 \end{bmatrix} : K_C^{-\frac{1}{2}} \oplus K_C^{\frac{1}{2}} &\longrightarrow K_C^{\frac{1}{2}} \oplus \left(a_2 K_C^{-\frac{1}{2}} + a_1 K_C^{\frac{1}{2}} \right) \\ &\longrightarrow K_C^{\frac{1}{2}} \oplus \left(K_C^{\frac{3}{2}} \otimes \mathcal{O}_C(D) \right) = K_C(D) \otimes \left(K_C^{-\frac{1}{2}} \oplus K_C^{\frac{1}{2}} \right), \end{aligned}$$

we have

$$\phi \in H^0(C, K_C(D) \otimes \text{End}(E)).$$

Remark 3.1. We use a particular section (E, ϕ) of the *Hitchin fibration* given by the form of (3.10). This section is often called a *Hitchin section*. This choice is suitable for the WKB analysis explained below. The result of our quantization through WKB constructs a **formal** section of $K_C^{-\frac{1}{2}}$, and the relation to \hbar -connection on C makes our choice necessary. The theory being developed as of now (relation to opers) [23] also requires the choice of a Hitchin section.

Denote by

$$(3.11) \quad \overline{T^*C} := \mathbb{P}(K_C \oplus \mathcal{O}_C) \xrightarrow{\pi} C$$

the compactified cotangent bundle of C (see [5, 57]), which is a ruled surface on the base C . The Hitchin spectral curve

$$(3.12) \quad \begin{array}{ccc} \Sigma & \xrightarrow{i} & \overline{T^*C} \\ & \searrow \pi & \downarrow \pi \\ & & C \end{array}$$

for a meromorphic Higgs bundle is defined as the divisor of zeros on $\overline{T^*C}$ of the characteristic polynomial of ϕ :

$$(3.13) \quad \Sigma = (\det(\eta - \pi^*\phi))_0,$$

where $\eta \in H^0(T^*C, \pi^*K_C)$ is the tautological 1-form on T^*C extended as a meromorphic 1-form on the compactification $\overline{T^*C}$. Our main theorem of [25] is the following.

Theorem 3.2 (Main results of [25]). • *The integral topological recursion of [24, 34] is extended to the curve Σ , as (3.24). For this purpose, we blow up $\overline{T^*C}$ several times as in (3.16) to construct the normalization $\tilde{\Sigma}$. The blow-up $Bl(\overline{T^*C})$ is the minimal resolution of the support $\Sigma \cup C_\infty$ of the total divisor*

$$(3.14) \quad \Sigma - 2C_\infty = (\det(\eta - \pi^*\phi))_0 - (\det(\eta - \pi^*\phi))_\infty$$

of the characteristic polynomial, where

$$(3.15) \quad C_\infty := \mathbb{P}(K_C \oplus \{0\}) = \overline{T^*C} \setminus T^*C$$

is the divisor at infinity. Therefore, in $Bl(\overline{T^*C})$, the proper transform $\tilde{\Sigma}$ of Σ is smooth and does not intersect with the proper transform of C_∞ .

$$(3.16) \quad \begin{array}{ccccc} \tilde{\Sigma} & \xrightarrow{\tilde{i}} & Bl(\overline{T^*C}) & & \\ \downarrow \tilde{\nu} & & \searrow \nu & & \\ \Sigma & \xrightarrow{i} & \overline{T^*C} & & \\ \downarrow \pi & \nearrow \pi & \downarrow \pi & \nearrow \pi & \\ C & & C & & \end{array}$$

- The genus of the normalization $\tilde{\Sigma}$ is given by

$$g(\tilde{\Sigma}) = 2g(C) - 1 + \frac{1}{2}\delta,$$

where δ is the sum of the number of cusp singularities of Σ and the ramification points of $\pi : \Sigma \rightarrow C$.

- The generalized topological recursion Definition 3.4 requires a globally defined meromorphic 1-form $W_{0,1}$ on $\tilde{\Sigma}$ and a symmetric meromorphic 2-form $W_{0,2}$ on the product $\tilde{\Sigma} \times \tilde{\Sigma}$ as the initial data. We choose

$$(3.17) \quad \begin{cases} W_{0,1} = \tilde{i}^* \nu^* \eta \\ W_{0,2} = d_1 d_2 \log E_{\tilde{\Sigma}}, \end{cases}$$

where $E_{\tilde{\Sigma}}$ is a normalized Riemann prime form on $\tilde{\Sigma}$. The form $W_{0,2}$ depends only on the intrinsic geometry of the smooth curve $\tilde{\Sigma}$. The geometry of (3.16) is encoded in $W_{0,1}$. The integral topological recursion produces a symmetric meromorphic n -linear differential form $W_{g,n}(z_1, \dots, z_n)$ on $\tilde{\Sigma}$ for every (g, n) subject to $2g - 2 + n > 0$ from the initial data (3.17).

- The residue evaluation of the integral topological recursion (3.24) is explicitly performed as in [24, (4.7)], and we obtain a differential recursion (3.25). It determines the free energy $F_{g,n}(z_1, \dots, z_n)$, a symmetric meromorphic function on \mathcal{U}^n for $2g - 2 + n \geq 2$, up to a constant. Here, $\varpi : \mathcal{U} \rightarrow \tilde{\Sigma}$ is the universal covering of $\tilde{\Sigma}$.
- The quantum curve associated with the Hitchin spectral curve Σ is defined as a differential equation on C . On each coordinate neighborhood $U \subset C$ with coordinate x , a generator of the quantum curve is given by

$$P(x, \hbar) = \left(\hbar \frac{d}{dx} \right)^2 - \text{tr} \phi(x) \left(\hbar \frac{d}{dx} \right) + \det \phi(x).$$

In particular, the **semi-classical limit** of the quantum curve recovers the singular spectral curve Σ , not its normalization $\tilde{\Sigma}$.

- We show that the **all-order WKB expansion**

$$(3.18) \quad \Psi(x, \hbar) = \exp \left(\sum_{m=0}^{\infty} \hbar^{m-1} S_m(x) \right)$$

of a solution to the Schrödinger equation

$$(3.19) \quad \left(\left(\hbar \frac{d}{dx} \right)^2 - \text{tr} \phi(x) \left(\hbar \frac{d}{dx} \right) + \det \phi(x) \right) \Psi(x, \hbar) = 0,$$

near each critical value of $\pi : \Sigma \rightarrow C$, can be obtained by the **principal specialization** of the differential recursion (3.25), after determining the first three terms. The procedure is the following. We determine S_0, S_1 , and S_2 by solving

$$(3.20) \quad \begin{aligned} (S_0'(x))^2 - \text{tr} \phi(x) S_0'(x) + \det \phi(x) &= 0, \\ 2S_0'(x) S_1'(x) + S_0''(x) - \text{tr} \phi(x) S_1'(x) &= 0, \\ S_1''(x) + \sum_{a+b=2} S_a'(x) S_b'(x) - \text{tr} \phi(x) S_2'(x) &= 0. \end{aligned}$$

Then find $F_{1,1}(z)$ and $F_{0,3}(z_1, z_2, z_3)$ so that

$$S_2(x) = F_{1,1}(z(x)) + \frac{1}{6} F_{0,3}(z(x), z(x), z(x)).$$

With these $F_{1,1}$ and $F_{0,3}$ as the initial values, solve (3.25), and put

$$(3.21) \quad S_m(x) = \sum_{2g-2+n=m-1} \frac{1}{n!} F_{g,n}(z(x)), \quad m \geq 3,$$

where $F_{g,n}(z(x))$ is the principal specialization of $F_{g,n}(z_1, \dots, z_n)$ evaluated at a local section $z = z(x)$ of $\tilde{\pi} : \tilde{\Sigma} \rightarrow C$. The **wave function** $\Psi(x, \hbar)$ is a formal section of the line bundle $K_C^{-\frac{1}{2}}$ on C .

- The canonical ordering of the quantization of the local functions on T^*C is automatically chosen in the process of the integration from (3.24) to (3.25) and the principal specialization (3.21). This selects the canonical ordering in (3.19).

Remark 3.3. We do not have a closed formula for $F_{1,1}$ and $F_{0,3}$ from the given geometric data. In general, they are **not** given by integrating $W_{1,1}$ and $W_{0,3}$ of the integral topological recursion defined below.

Definition 3.4 (Integral topological recursion for a degree 2 covering). Let $\tilde{\pi} : \tilde{\Sigma} \rightarrow C$ be a degree 2 covering of C by a non-singular curve $\tilde{\Sigma}$. We denote by R the ramification divisor of $\tilde{\pi}$. In this case the covering $\tilde{\pi}$ is a Galois covering with the Galois group $\mathbb{Z}/2\mathbb{Z} = \langle \tilde{\sigma} \rangle$, and R is the fixed-point divisor of the involution $\tilde{\sigma}$. The **integral topological recursion** is an inductive mechanism of constructing meromorphic differential forms $W_{g,n}$ on the Hilbert scheme $\tilde{\Sigma}^{[n]}$ of n -points on $\tilde{\Sigma}$ for all $g \geq 0$ and $n \geq 1$ in the *stable range* $2g - 2 + n > 0$, from given initial data $W_{0,1}$ and $W_{0,2}$.

- $W_{0,1}$ is a meromorphic 1-form on $\tilde{\Sigma}$.
- $W_{0,2}$ is defined to be

$$(3.22) \quad W_{0,2}(z_1, z_2) = d_1 d_2 \log E_{\tilde{\Sigma}}(z_1, z_2),$$

where $E_{\tilde{\Sigma}}(z_1, z_2)$ is the normalized Riemann prime form on $\tilde{\Sigma} \times \tilde{\Sigma}$ (see [24, Section 2]).

Let $\omega^{a-b}(z)$ be a normalized Cauchy kernel on $\tilde{\Sigma}$ of Remark 2.18, which has simple poles at $z = a$ of residue 1 and at $z = b$ of residue -1 . Then

$$d_1 \omega^{z_1-b}(z_2) = W_{0,2}(z_1, z_2).$$

Define

$$(3.23) \quad \Omega := \tilde{\sigma}^* W_{0,1} - W_{0,1}.$$

Then $\tilde{\sigma}^* \Omega = -\Omega$, hence $\text{supp}(R) \subset \text{supp}(\Omega)$, where $\text{supp}(\Omega)$ denotes the support of both zero and pole divisors of Ω . The inductive formula of the topological recursion is then given by the following:

$$(3.24) \quad W_{g,n}(z_1, \dots, z_n) = \frac{1}{2} \frac{1}{2\pi\sqrt{-1}} \sum_{p \in \text{supp}(\Omega)} \oint_{\gamma_p} \omega^{\tilde{z}-z}(z_1) \\ \times \frac{1}{\Omega(z)} \left[W_{g-1, n+1}(z, \tilde{z}, z_2, \dots, z_n) + \sum_{\substack{\text{No } (0,1) \\ g_1+g_2=g \\ I \sqcup J = \{2, \dots, n\}}} W_{g_1, |I|+1}(z, z_I) W_{g_2, |J|+1}(\tilde{z}, z_J) \right].$$

Here,

- γ_p is a positively oriented small loop around a point $p \in \text{supp}(\Omega)$;
- the integration is taken with respect to $z \in \gamma_p$ for each $p \in \text{supp}(\Omega)$;
- $\tilde{z} = \tilde{\sigma}(z)$ is the Galois conjugate of $z \in \tilde{\Sigma}$;
- the operation $1/\Omega$ denotes the contraction of the meromorphic vector field dual to the 1-form Ω , considered as a meromorphic section of $K_{\tilde{\Sigma}}^{-1}$;

- “No $(0, 1)$ ” means that $g_1 = 0$ and $I = \emptyset$, or $g_2 = 0$ and $J = \emptyset$, are excluded in the summation;
- the sum runs over all partitions of g and set partitions of $\{2, \dots, n\}$, other than those containing the $(0, 1)$ geometry;
- $|I|$ is the cardinality of the subset $I \subset \{2, \dots, n\}$; and
- $z_I = (z_i)_{i \in I}$.

3.3. Quantization of Hitchin spectral curves. The passage from the topological recursion (3.24) to the quantum curve (3.19) is the evaluation of the residues in the formula.

Definition 3.5 (Free energies). The **free energy** of type (g, n) is a function $F_{g,n}(z_1, \dots, z_n)$ defined on the universal covering \mathcal{U}^n of $\tilde{\Sigma}^n$ such that

$$d_1 \cdots d_n F_{g,n} = W_{g,n}.$$

Remark 3.6. The free energies may contain logarithmic singularities, since it is an integral of a meromorphic function. For example, $F_{0,2}$ is the Riemann prime form itself considered as a function on \mathcal{U}^2 , which has logarithmic singularities along the diagonal [24, Section 2].

Definition 3.7 (Differential recursion for a degree 2 covering). The **differential recursion** is the following partial differential equation for all (g, n) subject to $2g - 2 + n \geq 2$:

$$(3.25) \quad d_1 F_{g,n}(z_1, \dots, z_n) = \sum_{j=2}^n \left[\frac{\omega^{z_j - \sigma(z_j)}(z_1)}{\Omega(z_1)} \cdot d_1 F_{g,n-1}(z_{[\hat{j}]}) - \frac{\omega^{z_j - \sigma(z_j)}(z_1)}{\Omega(z_j)} \cdot d_j F_{g,n-1}(z_{[\hat{1}]}) \right] + \frac{1}{\Omega(z_1)} d_{u_1} d_{u_2} \left[F_{g-1, n+1}(u_1, u_2, z_{[\hat{1}]}) + \sum_{\substack{\text{stable} \\ g_1 + g_2 = g \\ I \sqcup J = [\hat{1}]}} F_{g_1, |I|+1}(u_1, z_I) F_{g_2, |J|+1}(u_2, z_J) \right] \Bigg|_{\substack{u_1 = z_1 \\ u_2 = z_1}}.$$

Here, $1/\Omega$ is again the contraction operation, and the index subset $[\hat{j}]$ denotes the exclusion of $j \in \{1, 2, \dots, n\}$.

Remark 3.8. As pointed out in [24, Remark 4.8], (3.25) is a coordinate-free equation, written in terms of exterior differentiations and the contraction operation on the universal covering of $\tilde{\Sigma}$.

Theorem 3.9 (Differential recursion implies integral recursion, [25]). *Let $\varpi : \mathcal{U} \rightarrow \tilde{\Sigma}$ be the universal covering of $\tilde{\Sigma}$. Suppose that $F_{g,n}$ for $2g - 2 + n > 0$ are globally meromorphic on $\mathcal{U}^{[n]}$ with poles located only along the divisor of $\mathcal{U}^{[n]}$ when one of the factors lies in the pull-back divisor $\varpi^*(\Omega)_0$ of zeros of Ω . Define $W_{g,n} := d_1 \cdots d_n F_{g,n}$ for $2g - 2 + n > 0$, and use (3.22) and (3.23) for (g, n) in the unstable range. If $F_{g,n}$'s satisfy the differential recursion (3.25), and if $W_{1,1}$ and $W_{0,3}$ satisfy the initial equations of the integral topological recursion (3.24), then $W_{g,n}$'s for all valued of (g, n) satisfy the integral topological recursion.*

Although the context of the statement is slightly different, the proof is essentially the same as that of [24, Theorem 4.7]. The crucial assumption we have made is that $\tilde{\pi} : \tilde{\Sigma} \rightarrow \mathcal{C}$ is a Galois covering. Therefore, the Galois conjugation $\tilde{\sigma} : \tilde{\Sigma} \rightarrow \tilde{\Sigma}$ is a globally defined holomorphic mapping. To calculate the residues in the integral recursion (3.24), we need the global analysis of

$$\omega^{\tilde{z}-z}(z_1) \in H^0(\tilde{\Sigma} \times \tilde{\Sigma}, q_2^* K_{\tilde{\Sigma}} \boxtimes q_2^* \mathcal{O}_{\tilde{\Sigma}}(\tilde{z} + z)),$$

where q_1 and q_2 are projections

$$(3.26) \quad \begin{array}{ccc} & \tilde{\Sigma} \times \tilde{\Sigma} & \\ q_1 \swarrow & & \searrow q_2 \\ z \in \tilde{\Sigma} & & \tilde{\Sigma} \ni z_1. \end{array}$$

The residue integration is done at each point $p \in \text{supp}(\Omega)$. The poles of the integrand of (3.24) that are enclosed in the union

$$\gamma = \bigcup_{p \in \text{supp}(\Omega)} \gamma_p$$

of the contours on the complement of $\text{supp}(\Omega)$ are located at

- (1) $z = z_1, z = \tilde{z}_1$ from $\omega^{\tilde{z}-z}(z_1)$; and
- (2) $z = z_1, z = \tilde{z}_j, j = 2, \dots, n$, from $W_{0,2}(z, z_j)$ and $W_{0,2}(\tilde{z}, z_j)$ that appear in the second line of (3.24).

The integrand has other poles at $\text{supp}(\Omega)$ that includes the ramification divisor R , but they are not enclosed in γ . The local behavior of $\omega^{\tilde{z}-z}(z_1)$ at $z = z_1, z = \tilde{z}_1$ is well understood, and residues of the integrand of (3.24) are simply the evaluation of the differential form at $z = z_1, z = \tilde{z}_1$. The double poles coming from $W_{0,2}(z, z_j)$ and $W_{0,2}(\tilde{z}, z_j)$ contribute as differentiation of the factor it is multiplied to. Adding all contributions from the poles, we obtain (3.25).

Now let us consider a spectral curve $\Sigma \subset \overline{T^*C}$ of (3.12) defined by a pair of meromorphic sections $a_1 = -\text{tr}\phi$ of K_C and $a_2 = \det\phi$ of $K_C^{\otimes 2}$. Let $\tilde{\Sigma}$ be the desingularization of Σ in (2.16). We apply the topological recursion (3.24) to the covering $\tilde{\pi} : \tilde{\Sigma} \rightarrow C$. The geometry of the spectral curve Σ provides us with a canonical choice of the initial differential forms (3.17). At this point we pay a special attention that the topological recursions (3.24) and (3.25) are both defined on the spectral curve $\tilde{\Sigma}$, while we wish to construct a differential equation on C . Since the free energies are defined on the universal covering of $\tilde{\Sigma}$, we need to have a mechanism to relate a coordinate on the desingularized spectral curve and that of the base curve C .

To analyze the singularity structure of Σ , let us consider the discriminant of the defining equation (3.4) of the spectral curve.

Definition 3.10 (Discriminant divisor). The **discriminant divisor** of the spectral curve

$$(3.27) \quad \eta^{\otimes 2} + \pi^* a_1 \eta + \pi^* a_2 = 0$$

is a divisor on C defined by

$$(3.28) \quad \Delta := \left(\frac{1}{4} a_1^2 - a_2 \right) = \Delta_0 - \Delta_\infty.$$

Here,

$$(3.29) \quad \Delta_0 = \sum_{i=1}^m m_i q_i, \quad m_i > 0, \quad q_i \in C,$$

is the divisor of zeros, and

$$(3.30) \quad \Delta_\infty = \sum_{j=1}^n n_j p_j, \quad n_j > 0, \quad p_j \in C,$$

is the divisor of ∞ .

Since $\frac{1}{4}a_1^2 - a_2$ is a meromorphic section of $K_C^{\otimes 2}$, we have

$$(3.31) \quad \deg \Delta = \sum_{i=1}^m m_i - \sum_{j=1}^n n_j = 4g - 4.$$

Theorem 3.11 (Geometric genus formula, [25]). *Let us define an invariant of the discriminant divisor by*

$$(3.32) \quad \delta = |\{i \mid m_i \equiv 1 \pmod{2}\}| + |\{j \mid n_j \equiv 1 \pmod{2}\}|.$$

Then the geometric genus of the spectral curve Σ of (3.27) is given by

$$(3.33) \quad g(\tilde{\Sigma}) := p_g(\Sigma) = 2g - 1 + \frac{1}{2}\delta.$$

We note that (3.31) implies $\delta \equiv 0 \pmod{2}$.

Take an arbitrary point $p \in C \setminus \text{supp}(\Delta)$, and a local coordinate x around p . By choosing a small disc V around p , we can make the inverse image of $\tilde{\pi} : \tilde{\Sigma} \rightarrow C$ consisting of two isomorphic discs. Since V is away from the critical values of $\tilde{\pi}$, the inverse image consists of two discs in the original spectral curve Σ . Note that we choose an eigenvalue α of ϕ on V in Theorem 3.2. We are thus specifying one of the inverse image discs here. Let us name the disc V_α that corresponds to α .

At this point apply the WKB analysis to the differential equation (3.19) with the WKB expansion of the solution

$$(3.34) \quad \Psi^\alpha(x, \hbar) = \exp\left(\sum_{m=0}^{\infty} \hbar^{m-1} S_m(x(z))\right) = \exp F^\alpha(x, \hbar),$$

where we choose a coordinate z of V_α so that the function $x = x(z)$ represents the projection $\pi : V_\alpha \rightarrow V$. The equation $P\Psi^\alpha = Pe^{F^\alpha} = 0$ reads

$$(3.35) \quad \hbar^2 \frac{d^2}{dx^2} F^\alpha + \hbar^2 \frac{dF^\alpha}{dx} \frac{dF^\alpha}{dx} + a_1 \hbar \frac{dF^\alpha}{dx} + a_2 = 0.$$

The \hbar -expansion of (3.35) gives

$$(3.36) \quad \hbar^0\text{-terms : } (S'_0(x))^2 + a_1 S'_0(x) + a_2 = 0,$$

$$(3.37) \quad \hbar^1\text{-terms : } 2S'_0(x)S'_1(x) + S''_0(x) + a_1 S'_1(x) = 0,$$

$$(3.38) \quad \hbar^{m+1}\text{-terms : } S''_m(x) + \sum_{a+b=m+1} S'_a(x)S'_b(x) + a_1 S'_{m+1}(x) = 0, \quad m \geq 1,$$

where $'$ denotes the x -derivative. The WKB method is to solve these equations iteratively and find $S_m(x)$ for all $m \geq 0$. Here, (3.36) is the **semi-classical limit** of (3.19), and (3.37) is the *consistency condition* we need to solve the WKB expansion, the same as before. Since the 1-form $dS_0(x)$ is a local section of T^*C , we identify $y = S'_0(x)$. Then (3.36) is the local expression of the spectral curve equation (3.4). This expression is the same everywhere for $p \in C \setminus \text{supp}(\Delta)$. We note a_1 and a_2 are globally defined. Therefore, we recover the spectral curve Σ from the differential operator of (3.19).

The topological recursion provides a closed formula for each $S_m(x)$, $m \geq 2$.

Theorem 3.12 (Topological recursion and WKB, [25]). *Let us determine $S_0(x)$ and $S_1(x)$ from the semi-classical limit and the consistency condition. Then the **principal specialization** of (3.25) is equivalent to (3.38) for all $m \geq 2$.*

Remark 3.13. The differential recursion (3.25) assumes $F_{1,1}$ and $F_{0,3}$ as the initial values. The equation itself does not give any condition for them. The discovery of [24, 25] is that the WKB equations for $S_m(x)$'s are equivalent to (3.25) for all $m \geq 2$. We note that no mechanism is established for deducing S_2 from S_0 and S_1 , other than solving the WKB equations directly. This situation reflects the following. Although $W_{0,1}$ and $W_{0,2}$ are well defined for the integral topological recursion (3.24), we do not have any systematic way to define $F_{0,1}$ and $F_{0,2}$ from them. Once $F_{1,1}$ and $F_{0,3}$ are given, then the differential recursion gives all $F_{g,n}$'s in the stable range of (g, n) . However, there is so far no unique mechanism that gives these two initial values from the unstable geometries.

3.4. Classical differential equations. Riemann and Poincaré found the interplay between algebraic geometry of curves in a ruled surface and the asymptotic expansion of an analytic solution to a differential equation defined on the base curve of the ruled surface. We look at these classical subjects from a new point of view. Let us now recall the definition of regular and irregular singular points of a second order differential equation.

Definition 3.14. Let

$$(3.39) \quad \left(\frac{d^2}{dx^2} + a_1(x) \frac{d}{dx} + a_2(x) \right) \Psi(x) = 0$$

be a second order differential equation defined around a neighborhood of $x = 0$ on a small disc $|x| < \epsilon$ with meromorphic coefficients $a_1(x)$ and $a_2(x)$ with poles at $x = 0$. Denote by k (reps. ℓ) the order of the pole of $a_1(x)$ (resp. $a_2(x)$) at $x = 0$. If $k \leq 1$ and $\ell \leq 2$, then (3.39) has a **regular singular point** at $x = 0$. Otherwise, consider the *Newton polygon* of the order of poles of the coefficients of (3.39). It is the upper part of the convex hull of three points $(0, 0)$, $(1, k)$, $(2, \ell)$. As a convention, if $a_j(x)$ is identically 0, then we assign $-\infty$ as its pole order. Let $(1, r)$ be the intersection point of the Newton polygon and the line $x = 1$. Thus

$$(3.40) \quad r = \begin{cases} k & 2k \geq \ell, \\ \frac{\ell}{2} & 2k \leq \ell. \end{cases}$$

The differential equation (3.39) has an **irregular singular point of class $r - 1$** at $x = 0$ if $r > 1$.

To illustrate the scope of interrelations among the geometry of meromorphic Higgs bundles, their spectral curves, the singularities of quantum curves, \hbar -connections, and the quantum invariants, let us tabulate five examples here (see Table 3.4). The differential operators of these equations are listed on the third column. In the first three rows, the quantum curves are examples of classical differential equations known as Airy, Hermite, the Gauß hypergeometric equations. The fourth and the fifth rows are added to show that it is *not* the singularity of the spectral curve that determines the singularity of the quantum curve. In each example, the Higgs bundle (E, ϕ) we are considering consists of the base curve $C = \mathbb{P}^1$ and the rank 2 vector bundle E on \mathbb{P}^1 of (1.20).

The first column of the table shows the Higgs field $\phi : E \rightarrow K_{\mathbb{P}^1}(4) \otimes E$. Here, x is the affine coordinate of $\mathbb{P}^1 \setminus \{\infty\}$. Since our vector bundle is a specific direct sum of line bundles, the non-Abelian Hodge correspondence is simple in each case. Except for the Gauß hypergeometric case, it is given by

$$(3.41) \quad \nabla^{\hbar} = \hbar d - \phi,$$

Higgs Field	Spectral Curve	Quantum Curve
$\begin{bmatrix} & 1 \\ x(dx)^2 & \end{bmatrix}$	$y^2 - x = 0$ $w^2 - u^5 = 0$ $\Sigma = 2C_0 + 5F$ $p_a = 2, p_g = 0$	Airy $(\hbar \frac{d}{dx})^2 - x$ Class $\frac{3}{2}$ irregular singularity at ∞
$\begin{bmatrix} & 1 \\ -(dx)^2 & -dx \end{bmatrix}$	$y^2 + xy + 1 = 0$ $w^2 - uw + u^4 = 0$ $\Sigma = 2C_0 + 4F$ $p_a = 1, p_g = 0$	Hermite $(\hbar \frac{d}{dx})^2 + x\hbar \frac{d}{dx} + 1$ Class 2 irregular singularity at ∞
$\begin{bmatrix} & \frac{1}{x} \\ \frac{(dx)^2}{4(1-x)} & \frac{2x-1}{x(1-x)} dx \end{bmatrix}$	$y^2 + \frac{2x-1}{x(x-1)}y + \frac{1}{4x(x-1)} = 0$ $w^2 + 4(u-2)uw$ $-4u^2(u-1) = 0$ $\Sigma = 2C_0 + 4F$ $p_a = 1, p_g = 0$	Gauß Hypergeometric $(\hbar \frac{d}{dx})^2 + \frac{2x-1}{x(x-1)}\hbar \frac{d}{dx} + \frac{1}{4x(x-1)}$ Regular singular points at $x = 0, 1, \infty$
$\begin{bmatrix} & 1 \\ -\frac{(dx)^2}{x+1} & -dx \end{bmatrix}$	$y^2 + y + \frac{1}{x+1} = 0$ $w^2 - u(u+1)w$ $+u^3(u+1) = 0$ $\Sigma = 2C_0 + 4F$ $p_a = 1, p_g = 0$	$(\hbar \frac{d}{dx})^2 + \hbar \frac{d}{dx} + \frac{1}{x+1}$ Regular singular point at $x = -1$ and a class 1 irregular singularity at $x = \infty$
$\begin{bmatrix} & 1 \\ \frac{(dx)^2}{x^2-1} & -\frac{2x^2}{x^2-1} dx \end{bmatrix}$	$(x^2 - 1)y^2 + 2x^2y - 1 = 0$ non-singular $\Sigma = 2C_0 + 4F$ $p_a = p_g = 1$	$(\hbar \frac{d}{dx})^2 + 2\frac{x^2}{x^2-1}\hbar \frac{d}{dx} - \frac{1}{x^2-1}$ Regular singular points at $x = \pm 1$ and a class 1 irregular singularity at $x = \infty$

TABLE 1. Examples of quantum curves.

where d is the exterior differentiation operator acting on sections of $E|_{\mathbb{A}^1}$ restricted to the affine part $\mathbb{A}^1 \subset \mathbb{P}^1$. The form of (3.41) is valid because of our choice, $(0, 1)$, as the first row of the Higgs field.

For the third example of a Gauß hypergeometric equation, we use a particular choice of parameters so that the \hbar -connection becomes an \hbar -deformed Gauß-Manin connection of (3.43). More precisely, for every $x \in \mathcal{M}_{0,4}$, we consider the elliptic curve $E(x)$ ramified over \mathbb{P}^1 at four points $\{0, 1, x, \infty\}$, and its two periods given by the elliptic integrals [58]

$$(3.42) \quad \omega_1(x) = \int_1^\infty \frac{ds}{\sqrt{s(s-1)(s-x)}}, \quad \omega_2(x) = \int_x^1 \frac{ds}{\sqrt{s(s-1)(s-x)}}.$$

The quantum curve in this case is an \hbar -deformed **meromorphic Gauß-Manin connection**

$$(3.43) \quad \nabla_{GM}^\hbar = \hbar d - \begin{bmatrix} & \frac{1}{x} \\ -\frac{(dx)^2}{4(x-1)} & \left(-\frac{2x-1}{x(x-1)} + \frac{\hbar}{x}\right) dx \end{bmatrix}$$

in the vector bundle $K_{\mathcal{M}_{0,4}}^{-\frac{1}{2}} \oplus K_{\mathcal{M}_{0,4}}^{\frac{1}{2}}$ of rank 2 over $\overline{\mathcal{M}}_{0,4}$. Here, d denotes the exterior differentiation acting on the local sections of this vector bundle on the affine part. The restriction ∇_{GM}^1 of the connection at $\hbar = 1$ is equivalent to the Gauß-Manin connection

that characterizes the two periods of (3.42), and the Higgs field is the classical limit of the connection matrix at $\hbar \rightarrow 0$:

$$(3.44) \quad \phi = \begin{bmatrix} & \frac{1}{x} \\ -\frac{(dx)^2}{4(x-1)} & -\frac{2x-1}{x(x-1)} dx \end{bmatrix}.$$

The spectral curve $\Sigma \subset \overline{T^*\mathcal{M}_{0,4}}$ as a moduli space consists of the data $(E(x), \alpha_1(x), \alpha_2(x))$, where $\alpha_1(x)$ and $\alpha_2(x)$ are the two eigenvalues of the Higgs field ϕ . The spectral curve $\Sigma \subset T^*\overline{\mathcal{M}_{0,4}} = \mathbb{F}^2$ as a divisor in the Hirzebruch surface is determined by the characteristic equation

$$(3.45) \quad y^2 + \frac{2x-1}{x(x-1)}y + \frac{1}{4x(x-1)} = 0$$

of the Higgs field. Geometrically, Σ is a singular rational curve with one ordinary double point at $x = \infty$. The quantum curve is a **quantization** of the characteristic equation (3.45) for the eigenvalues $\alpha_1(x)$ and $\alpha_2(x)$ of $\phi(x)$. It is an \hbar -deformed **Picard-Fuchs equation**

$$\left(\left(\hbar \frac{d}{dx} \right)^2 + \frac{2x-1}{x(x-1)} \left(\hbar \frac{d}{dx} \right) + \frac{1}{4x(x-1)} \right) \omega_i(x, \hbar) = 0,$$

and its semi-classical limit agrees with the singular spectral curve Σ . As a second order differential equation, the quantum curve has two independent solutions corresponding to the two eigenvalues. At $\hbar = 1$, these solutions are exactly the two periods $\omega_1(x)$ and $\omega_2(x)$ of the Legendre family of elliptic curves $E(x)$. The topological recursion produces asymptotic expansions of these periods as functions in $x \in \overline{\mathcal{M}_{0,4}}$, at which the elliptic curve $E(x)$ degenerates to a nodal rational curve.

This is a singular connection with simple poles at $0, 1, \infty$, and has an explicit \hbar -dependence in the connection matrix. The Gauß-Manin connection ∇_{GM}^1 at $\hbar = 1$ is equivalent to the Picard-Fuchs equation that characterizes the periods (3.42) of the Legendre family of elliptic curves $E(x)$ defined by the cubic equation

$$(3.46) \quad t^2 = s(s-1)(s-x), \quad x \in \mathcal{M}_{0,4} = \mathbb{P}^1 \setminus \{0, 1, \infty\}.$$

The second column gives the spectral curve of the Higgs bundle (E, ϕ) . Since the Higgs fields have poles, the spectral curves are no longer contained in the cotangent bundle $T^*\mathbb{P}^1$. We need the compactified cotangent bundle

$$\overline{T^*\mathbb{P}^1} = \mathbb{P}(K_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}) = \mathbb{F}_2,$$

which is a Hirzebruch surface. The parameter y is the fiber coordinate of the cotangent line $T_x^*\mathbb{P}^1$. The first line of the second column is the equation of the spectral curve in the (x, y) affine coordinate of \mathbb{F}_2 . All but the last example produce a singular spectral curve. Let (u, w) be a coordinate system on another affine chart of \mathbb{F}_2 defined by

$$\begin{cases} x = 1/u \\ ydx = vdu, & w = 1/v. \end{cases}$$

The singularity of Σ in the (u, w) -plane is given by the second line of the second column. The third line of the second column gives $\Sigma \in \text{NS}(\mathbb{F}_2)$ as an element of the Néron-Severi group of \mathbb{F}_2 . Here, C_0 is the class of the zero-section of $T^*\mathbb{P}^1$, and F represents the fiber class of $\pi : \mathbb{F}_2 \rightarrow \mathbb{P}^1$. We also give the arithmetic and geometric genera of the spectral curve.

A solution $\Psi(x, \hbar)$ to the first example is given by the **Airy function**

$$(3.47) \quad Ai(x, \hbar) = \frac{1}{2\pi} \hbar^{-\frac{1}{6}} \int_{-\infty}^{\infty} \exp\left(\frac{ipx}{\hbar^{2/3}} + i\frac{p^3}{3}\right) dp,$$

which is an entire function in x for $\hbar \neq 0$, as discussed earlier in these lectures. The expansion coordinate $x^{\frac{3}{2}}$ of (1.37) indicates the class of the irregular singularity of the Airy differential equation.

The solutions to the second example are given by confluent hypergeometric functions, such as ${}_1F_1\left(\frac{1}{2\hbar}; \frac{1}{2}; -\frac{x^2}{2\hbar}\right)$, where

$$(3.48) \quad {}_1F_1(a; c; z) := \sum_{n=0}^{\infty} \frac{(a)_n}{(c)_n} \frac{z^n}{n!}$$

is the **Kummer confluent hypergeometric function**, and the **Pochhammer symbol** $(a)_n$ is defined by

$$(3.49) \quad (a)_n := a(a+1)(a+2)\cdots(a+n-1).$$

For $\hbar > 0$, the topological recursion determines the asymptotic expansion of a particular entire solution known as a **Tricomi confluent hypergeometric function**

$$\begin{aligned} & \Psi^{\text{Catalan}}(x, \hbar) \\ &= \left(-\frac{1}{2\hbar}\right)^{\frac{1}{2\hbar}} \left(\frac{\Gamma[\frac{1}{2}]}{\Gamma[\frac{1}{2\hbar} + \frac{1}{2}]} {}_1F_1\left(\frac{1}{2\hbar}; \frac{1}{2}; -\frac{x^2}{2\hbar}\right) + \frac{\Gamma[-\frac{1}{2}]}{\Gamma[\frac{1}{2\hbar}]} \sqrt{-\frac{x^2}{2\hbar}} {}_1F_1\left(\frac{1}{2\hbar} + \frac{1}{2}; \frac{3}{2}; -\frac{x^2}{2\hbar}\right) \right). \end{aligned}$$

The expansion is given in the form

$$(3.50) \quad \begin{aligned} \Psi^{\text{Catalan}}(x, \hbar) &= \left(\frac{1}{x}\right)^{\frac{1}{\hbar}} \sum_{n=0}^{\infty} \frac{\hbar^n \left(\frac{1}{\hbar}\right)_{2n}}{(2n)!!} \cdot \frac{1}{x^{2n}} \\ &= \exp\left(\sum_{g=0}^{\infty} \sum_{n=1}^{\infty} \frac{1}{n!} \hbar^{2g-2+n} F_{g,n}^C(x, \dots, x)\right), \end{aligned}$$

where $F_{g,n}^C$ is defined by (2.20), in terms of generalized Catalan numbers. The expansion variable x^2 in (3.50) indicates the class of irregularity of the Hermite differential equation at $x = \infty$. The cases for $(g, n) = (0, 1)$ and $(0, 2)$ require again a special treatment, as we discussed earlier.

The Hermite differential equation becomes simple for $\hbar = 1$, and we have the asymptotic expansion

$$(3.51) \quad \begin{aligned} i\sqrt{\frac{\pi}{2}} e^{-\frac{1}{2}x^2} \left[1 - \operatorname{erf}\left(\frac{ix}{\sqrt{2}}\right)\right] &= \sum_{n=0}^{\infty} \frac{(2n-1)!!}{x^{2n+1}} \\ &= \exp\left(\sum_{2g-2+n \geq -1} \frac{1}{n!} \sum_{\mu_1, \dots, \mu_n > 0} \frac{C_{g,n}(\mu_1, \dots, \mu_n)}{\mu_1 \cdots \mu_n} \prod_{i=1}^n x^{-(\mu_1 + \dots + \mu_n)}\right). \end{aligned}$$

Here, $\operatorname{erf}(x) := \frac{2}{\sqrt{\pi}} \int_0^x e^{-z^2} dz$ is the Gauß error function.




FIGURE 3.2. The imaginary part and the real part of $\Psi^{\text{Catalan}}(x, 1)$. For $x \gg 0$, the imaginary part dies down, and only the real part has a non-trivial asymptotic expansion. Thus (3.51) is a series with real coefficients.

One of the two independent solutions to the third example, the Gauß hypergeometric equation, that is holomorphic around $x = 0$, is given by

$$(3.52) \quad \Psi^{\text{Gauß}}(x, \hbar) = {}_2F_1 \left(-\frac{\sqrt{(h-1)(h-3)}}{2h} + \frac{1}{h} - \frac{1}{2}, \frac{\sqrt{(h-1)(h-3)}}{2h} + \frac{1}{h} - \frac{1}{2}; \frac{1}{h}; x \right),$$

where

$$(3.53) \quad {}_2F_1(a, b; c; x) := \sum_{n=0}^{\infty} \frac{(a)_n (b)_n}{(c)_n} \frac{x^n}{n!}$$

is the **Gauß hypergeometric function**. The topological recursion calculates the B-model genus expansion of the periods of the Legendre family of elliptic curves (3.46) at the point where the elliptic curve degenerates to a nodal rational curve. For example, the procedure applied to the spectral curve

$$y^2 + \frac{2x-1}{x(x-1)}y + \frac{1}{4x(x-1)} = 0$$

with a choice of

$$\eta = \frac{-(2x-1) - \sqrt{3x^2 - 3x + 1}}{2x(x-1)} dx,$$

which is an eigenvalue $\alpha_1(x)$ of the Higgs field ϕ , gives a genus expansion at $x = 0$:

$$(3.54) \quad \Psi^{\text{Gauß}}(x, \hbar) = \exp \left(\sum_{g=0}^{\infty} \sum_{n=1}^{\infty} \frac{1}{n!} \hbar^{2g-2+n} F_{g,n}^{\text{Gauß}}(x) \right).$$

At $\hbar = 1$, we have a topological recursion expansion of the period $\omega_1(x)$ defined in (3.42):

$$(3.55) \quad \frac{\omega_1(x)}{\pi} = \Psi^{\text{Gauß}}(x, 1) = \exp \left(\sum_{g=0}^{\infty} \sum_{n=1}^{\infty} \frac{1}{n!} F_{g,n}^{\text{Gauß}}(x) \right).$$

A subtle point we notice here is that while the Gauß hypergeometric equation has regular singular points at $x = 0, 1, \infty$, the Hermite equation has an irregular singular point of class 2 at ∞ . The spectral curve of each case has an ordinary double point at $x = \infty$. But the crucial difference lies in the intersection of the spectral curve Σ with the divisor C_∞ . For

the Hermite case we have $\Sigma \cdot C_\infty = 4$ and the intersection occurs all at once at $x = \infty$. For the Gauß hypergeometric case, the intersection $\Sigma \cdot C_\infty = 4$ occurs once each at $x = 0, 1$, and twice at $x = \infty$. This *confluence* of regular singular points is the source of the irregular singularity in the Hermite differential equation.

The fourth row indicates an example of a quantum curve that has one regular singular point at $x = -1$ and one irregular singular point of class 1 at $x = \infty$. The spectral curve has an ordinary double point at $x = \infty$, the same as the Hermite case. As Figure 3.3 shows, the class of the irregular singularity at $x = \infty$ is determined by how the spectral curve intersects with C_∞ .




FIGURE 3.3. The spectral curves of the second and the fourth examples. The horizontal line is the divisor C_∞ , and the vertical line is the fiber class F at $x = \infty$. The spectral curve intersects with C_∞ a total of four times. The curve on the right has a triple intersection at $x = \infty$, while the one on the left intersects all at once.

The existence of the irregular singularity in the quantum curve associated with a spectral curve has nothing to do with the singularity of the spectral curve. The fifth example shows a non-singular spectral curve of genus 1 (Figure 3.4), for which the quantum curve has a class 1 irregular singularity at $x = \infty$.




FIGURE 3.4. The spectral curve of the fifth example, which is non-singular. The corresponding quantum curve has two regular singular points at $x = \pm 1$, and a class 1 irregular singular point at $x = \infty$.

4. DIFFERENCE OPERATORS AS QUANTUM CURVES

Quantum curves often appear as infinite-order differential operators, or *difference operators*. In this section we present three typical examples: simple Hurwitz numbers, a special double Hurwitz numbers, and the Gromov-Witten invariants of \mathbb{P}^1 . These examples do not come from the usual Higgs bundle framework, because the rank of the Higgs bundle corresponds to the order of the quantum curves as a differential operator. Therefore, we ask:

Question 4.1. *What is the geometric structure generalizing the Hitchin spectral curves that correspond to difference operators as their quantization?*

In these lectures, we do not address this question, leaving it for a future investigation. We are content with giving examples here. We refer to [53] for a new and different perspective for the notion of quantum curves for difference operators.

4.1. Simple and orbifold Hurwitz numbers. The *simple Hurwitz number* $H_{g,n}(\vec{\mu})$ counts the automorphism weighted number of the topological types of simple Hurwitz covers of \mathbb{P}^1 of type $(g, \vec{\mu})$. A holomorphic map $\varphi : C \rightarrow \mathbb{P}^1$ is a *simple Hurwitz cover* of type $(g, \vec{\mu})$ if C is a complete nonsingular algebraic curve defined over \mathbb{C} of genus g , φ has n labeled poles of orders $\vec{\mu} = (\mu_1, \dots, \mu_n)$, and all other critical points of φ are unlabeled simple ramification points.

In a similar way, we consider the *orbifold Hurwitz number* $H_{g,n}^{(r)}(\vec{\mu})$ for every positive integer $r > 0$ to be the automorphism weighted count of the topological types of smooth orbifold morphisms $\varphi : C \rightarrow \mathbb{P}^1[r]$ with the same pole structure as the simple Hurwitz number case. Here, C is a connected 1-dimensional orbifold (a *twisted curve*) modeled on a nonsingular curve of genus g with $(\mu_1 + \dots + \mu_n)/r$ stacky points of the type $[p/(\mathbb{Z}/r\mathbb{Z})]$. We impose that the inverse image of the morphism φ of the unique stacky point $[0/(\mathbb{Z}/r\mathbb{Z})] \in \mathbb{P}^1[r]$ coincides with the set of stacky points of C . When $r = 1$ we recover the simple Hurwitz number: $H_{g,n}^{(1)}(\vec{\mu}) = H_{g,n}(\vec{\mu})$.

Theorem 4.2 (Cut-and-join equation, [10]). *The orbifold Hurwitz numbers $H_{g,n}^{(r)}(\mu_1, \dots, \mu_n)$ satisfy the following equation.*

$$(4.1) \quad sH_{g,n}^{(r)}(\mu_1, \dots, \mu_n) = \frac{1}{2} \sum_{i \neq j} (\mu_i + \mu_j) H_{g,n-1}^{(r)}(\mu_i + \mu_j, \mu_{[\hat{i}, \hat{j}]}) \\ + \frac{1}{2} \sum_{i=1}^n \sum_{\alpha+\beta=\mu_i} \alpha\beta \left[H_{g-1,n+1}^{(r)}(\alpha, \beta, \mu_{[\hat{i}]}) + \sum_{\substack{g_1+g_2=g \\ I \sqcup J = [\hat{i}]}} H_{g_1,|I|+1}^{(r)}(\alpha, \mu_I) H_{g_2,|J|+1}^{(r)}(\beta, \mu_J) \right].$$

Here

$$(4.2) \quad s = s(g, \vec{\mu}) = 2g - 2 + n + \frac{\mu_1 + \dots + \mu_n}{r}$$

is the number of simple ramification point given by the Riemann-Hurwitz formula. As before, we use the convention that for any subset $I \subset [n] = \{1, 2, \dots, n\}$, $\mu_I = (\mu_i)_{i \in I}$. The hat notation \hat{i} indicates that the index i is removed. The last summation is over all partitions of g and set partitions of $[\hat{i}] = \{1, \dots, i-1, i+1, \dots, n\}$.

Remark 4.3. There is a combinatorial description, in the same manner we have done for the Catalan numbers in Section 2, for simple and orbifold Hurwitz numbers. The cut-and-join equation is derived as the edge-contraction formula, exactly in the same way for the Catalan recursion (2.8). See [26] for more detail.

We regard $H_{g,n}^{(r)}(\vec{\mu})$ as a function in n integer variables $\vec{\mu} \in \mathbb{Z}_+^n$. Following the recipe of [27, 33, 67] that is explained in the earlier sections, we define the *free energies* as the Laplace transform

$$(4.3) \quad F_{g,n}^{(r)}(z_1, \dots, z_n) = \sum_{\vec{\mu} \in \mathbb{Z}_+^n} H_{g,n}^{(r)}(\vec{\mu}) e^{-\langle \vec{w}, \vec{\mu} \rangle}.$$

Here, $\vec{w} = (w_1, \dots, w_n)$ is the vector of the Laplace dual coordinates of $\vec{\mu}$, $\langle \vec{w}, \vec{\mu} \rangle = w_1 \mu_1 + \dots + w_n \mu_n$, and variables z_i and w_i for each i are related by the r -Lambert function

$$(4.4) \quad e^{-w} = ze^{-z^r}.$$

It is often convenient to use a different variable $x = ze^{-z^r}$, with which the plane analytic curve called r -**Lambert curve** is given by

$$(4.5) \quad \begin{cases} x = ze^{-z^r} \\ y = z^r. \end{cases}$$

Then the free energies $F_{g,n}^{(r)}$ of (4.3) are generating functions of the orbifold Hurwitz numbers. By abuse of notation, we also write

$$(4.6) \quad F_{g,n}^{(r)}(x_1, \dots, x_n) = \sum_{\vec{\mu} \in \mathbb{Z}_+^n} H_{g,n}^{(r)}(\vec{\mu}) \prod_{i=1}^n x_i^{\mu_i} = \sum_{\vec{\mu} \in \mathbb{Z}_+^n} H_{g,n}^{(r)}(\vec{\mu}) \prod_{i=1}^n (z_i e^{-z_i^r})^{\mu_i}.$$

For every (g, n) , the power series (4.6) in (x_1, \dots, x_n) is convergent and defines an analytic function.

Theorem 4.4 (Differential recursion for Hurwitz numngers, [10]). *In terms of the z -variables, the free energies are calculated as follows.*

$$(4.7) \quad F_{0,1}^{(r)}(z) = \frac{1}{r} z^r - \frac{1}{2} z^{2r},$$

$$(4.8) \quad F_{0,2}^{(r)}(z_1, z_2) = \log \frac{z_1 - z_2}{x_1 - x_2} - (z_1^r + z_2^r),$$

where $x_i = z_i e^{-z_i^r}$. For (g, n) in the stable range, i.e., when $2g - 2 + n > 0$, the free energies satisfy the differential recursion equation

$$(4.9) \quad \begin{aligned} & \left(2g - 2 + n + \frac{1}{r} \sum_{i=1}^n z_i \frac{\partial}{\partial z_i} \right) F_{g,n}^{(r)}(z_1, \dots, z_n) \\ &= \frac{1}{2} \sum_{i \neq j} \frac{z_i z_j}{z_i - z_j} \left[\frac{1}{(1 - rz_i^r)^2} \frac{\partial}{\partial z_i} F_{g,n-1}^{(r)}(z_{[\hat{j}]}) - \frac{1}{(1 - rz_j^r)^2} \frac{\partial}{\partial z_j} F_{g,n-1}^{(r)}(z_{[\hat{i}]}) \right] \\ & \quad + \frac{1}{2} \sum_{i=1}^n \frac{z_i^2}{(1 - rz_i^r)^2} \frac{\partial^2}{\partial u_1 \partial u_2} F_{g-1,n+1}^{(r)}(u_1, u_2, z_{[\hat{i}]}) \Big|_{u_1=u_2=z_i} \\ & \quad + \frac{1}{2} \sum_{i=1}^n \frac{z_i^2}{(1 - rz_i^r)^2} \sum_{\substack{\text{stable} \\ g_1+g_2=g \\ I \sqcup J = [\hat{i}]}} \left(\frac{\partial}{\partial z_i} F_{g_1,|I|+1}^{(r)}(z_i, z_I) \right) \left(\frac{\partial}{\partial z_i} F_{g_2,|J|+1}^{(r)}(z_i, z_J) \right). \end{aligned}$$

Remark 4.5. Since $F_{g,n}^{(r)}(z_1, \dots, z_n) \Big|_{z_i=0} = 0$ for every i , the differential recursion (4.9), which is a linear first order partial differential equation, uniquely determines $F_{g,n}^{(r)}$ one by one inductively for all (g, n) subject to $2g - 2 + n > 0$. This is a generalization of the result of [68] to the orbifold case.

Remark 4.6. The differential recursion of Theorem 4.4 is obtained by taking the Laplace transform of the cut-and-join equation for $H_{g,n}^{(r)}(\vec{\mu})$. The r -Lambert curve itself, (4.5), is obtained by computing the Laplace transform of $H_{0,1}^{(r)}(\mu)$, and solving the differential equation that arises from the cut-and-join equation. See also [26] for a different formulation of the

r -Hurwitz numbers using the graph enumeration formulation and a universal mechanism to obtain the spectral curve.

The differential recursion produces two results, as we have seen for the case of the Catalan numbers. One is the quantum curve by taking the principal specialization, and the other the topological recursion of [34].

Theorem 4.7 (Quantum curves for r -Hurwitz numbers, [10]). *We introduce the partition function, or the wave function, of the orbifold Hurwitz numbers as*

$$(4.10) \quad \Psi^{(r)}(z, \hbar) = \exp \left(\sum_{g=0}^{\infty} \sum_{n=1}^{\infty} \frac{1}{n!} \hbar^{2g-2+n} F_{g,n}^{(r)}(z, z, \dots, z) \right).$$

It satisfies the following system of (an infinite-order) linear differential equations.

$$(4.11) \quad \left(\hbar D - e^{r(-w + \frac{r-1}{2}\hbar)} e^{r\hbar D} \right) \Psi^{(r)}(z, \hbar) = 0,$$

$$(4.12) \quad \left(\frac{\hbar}{2} D^2 - \left(\frac{1}{r} + \frac{\hbar}{2} \right) D - \hbar \frac{\partial}{\partial \hbar} \right) \Psi^{(r)}(z, \hbar) = 0,$$

where

$$D = \frac{z}{1 - rz^r} \frac{\partial}{\partial z} = x \frac{\partial}{\partial x} = - \frac{\partial}{\partial w}.$$

Let the differential operator of (4.11) (resp. (4.12)) be denoted by P (resp. Q). Then we have the commutator relation

$$(4.13) \quad [P, Q] = P.$$

The semi-classical limit of each of the equations (4.11) or (4.12) recovers the r -Lambert curve (4.5).

Remark 4.8. The Schrödinger equation (4.11) is first established in [66].

Remark 4.9. The above theorem is a generalization of [67, Theorem 1.3] for an arbitrary $r > 0$. The restriction $r = 1$ reduces to the simple Hurwitz case.

Remark 4.10. Unlike the situation of Hitchin spectral curves, the results of the quantization of the analytic spectral curves are a **difference-differential equation**, and a PDE containing the differentiation with respect to the deformation parameter \hbar .

Now let us define

$$(4.14) \quad W_{g,n}^{(r)}(z_1, \dots, z_n) := d_1 d_2 \cdots d_n F_{g,n}^{(r)}(z_1, \dots, z_n).$$

Then we have

Theorem 4.11 (Topological recursion for orbifold Hurwitz numbers, [10]). *For the stable range $2g - 2 + n > 0$, the symmetric differential forms (4.14) satisfy the following integral recursion formula.*

$$(4.15) \quad W_{g,n}^{(r)}(z_1, \dots, z_n) = \frac{1}{2\pi i} \sum_{j=1}^r \oint_{\gamma_j} K_j(z, z_1) \left[W_{g-1, n+1}^{(r)}(z, s_j(z), z_2, \dots, z_n) \right. \\ \left. + \sum_{i=2}^n \left(W_{0,2}^{(r)}(z, z_i) \otimes W_{g, n-1}^{(r)}(s_j(z), z_{[\hat{1}, \hat{i}]} \right) + W_{0,2}^{(r)}(s_j(z), z_i) \otimes W_{g, n-1}^{(r)}(z, z_{[\hat{1}, \hat{i}]}) \right)$$

$$+ \left. \sum_{\substack{\text{stable} \\ g_1+g_2=g \\ I \sqcup J = \{2, \dots, n\}}} W_{g_1, |I|+1}^{(r)}(z, z_I) \otimes W_{g_2, |J|+1}^{(r)}(s_J(z), z_J) \right].$$

Here, the integration is taken with respect to z along a small simple closed loop γ_j around p_j , and $\{p_1, \dots, p_r\}$ are the critical points of the r -Lambert function $x(z) = ze^{-z^r}$ at $1 - rz^r = 0$. Since $dx = 0$ has a simple zero at each p_j , the map $x(z)$ is locally a double-sheeted covering around $z = p_j$. We denote by s_j the deck transformation on a small neighborhood of p_j . Finally, the integration kernel is defined by

$$(4.16) \quad K_j(z, z_1) = \frac{1}{2} \frac{1}{W_{0,1}^{(r)}(s_j(z_1)) - W_{0,1}^{(r)}(z_1)} \otimes \int_z^{s_j(z)} W_{0,2}^{(r)}(\cdot, z_1).$$

Remark 4.12. As mentioned earlier, the significance of the integral formalism is its universality. The differential equation (4.9) takes a different form depending on the counting problem, whereas the integral formula (4.15) depends only on the choice of the spectral curve.

Remark 4.13. The proof is based on the idea of [33]. The notion of the *principal part* of meromorphic differentials plays a key role in converting the Laplace transform of the cut-and-join equation into a residue formula.

4.2. Gromov-Witten invariants of the projective line. Hurwitz numbers and Gromov-Witten invariants of \mathbb{P}^1 are closely related [74]. However, their relations to the topological recursion is rather different. For example, the topological recursion for stationary Gromov-Witten invariants of \mathbb{P}^1 was conjectured by Norbury and Scott [73] as a concrete formula, but its proof [30] is done in a very different way than that of [33, 68]. This is based on the fact that we do not have a counterpart of the cut-and-join equation for the case of the Gromov-Witten invariants of \mathbb{P}^1 . Nonetheless, the quantum curve exists.

Let $\overline{\mathcal{M}}_{g,n}(\mathbb{P}^1, d)$ denote the moduli space of stable maps of degree d from an n -pointed genus g curve to \mathbb{P}^1 . This is an algebraic stack of dimension $2g - 2 + n + 2d$. The dimension reflects the fact that a generic map from an algebraic curve to \mathbb{P}^1 has only simple ramifications, and the number of such ramification points, which we can derive from the Riemann-Hurwitz formula, gives the dimension of this stack. The descendant Gromov-Witten invariants of \mathbb{P}^1 are defined by

$$(4.17) \quad \left\langle \prod_{i=1}^n \tau_{b_i}(\alpha_i) \right\rangle_{g,n}^d := \int_{[\overline{\mathcal{M}}_{g,n}(\mathbb{P}^1, d)]^{vir}} \prod_{i=1}^n \psi_i^{b_i} ev_i^*(\alpha_i),$$

where $[\overline{\mathcal{M}}_{g,n}(\mathbb{P}^1, d)]^{vir}$ is the virtual fundamental class of the moduli space,

$$ev_i : \overline{\mathcal{M}}_{g,n}(\mathbb{P}^1, d) \longrightarrow \mathbb{P}^1$$

is a natural morphism defined by evaluating a stable map at the i -th marked point of the source curve, $\alpha_i \in H^*(\mathbb{P}^1, \mathbb{Q})$ is a cohomology class of the target \mathbb{P}^1 , and ψ_i is the tautological cotangent class in $H^2(\overline{\mathcal{M}}_{g,n}(\mathbb{P}^1, d), \mathbb{Q})$. We denote by 1 the generator of $H^0(\mathbb{P}^1, \mathbb{Q})$, and by $\omega \in H^2(\mathbb{P}^1, \mathbb{Q})$ the Poincaré dual to the point class. We assemble the Gromov-Witten invariants into particular generating functions as follows. For every (g, n) in the stable sector $2g - 2 + n > 0$, we define the *free energy* of type (g, n) by

$$(4.18) \quad F_{g,n}(x_1, \dots, x_n) := \left\langle \prod_{i=1}^n \left(-\frac{\tau_0(1)}{2} - \sum_{b=0}^{\infty} \frac{b! \tau_b(\omega)}{x_i^{b+1}} \right) \right\rangle_{g,n}.$$

Here the degree d is determined by the dimension condition of the cohomology classes to be integrated over the virtual fundamental class. We note that (4.18) contains the class $\tau_0(1)$. For unstable geometries, we introduce two functions

$$(4.19) \quad S_0(x) := x - x \log x + \sum_{d=1}^{\infty} \left\langle -\frac{(2d-2)! \tau_{2d-2}(\omega)}{x^{2d-1}} \right\rangle_{0,1}^d,$$

$$(4.20) \quad S_1(x) := -\frac{1}{2} \log x + \frac{1}{2} \sum_{d=0}^{\infty} \left\langle \left(-\frac{\tau_0(1)}{2} - \sum_{b=0}^{\infty} \frac{b! \tau_b(\omega)}{x^{b+1}} \right)^2 \right\rangle_{0,2}^d,$$

utilizing an earlier work of [27]. Then we have

Theorem 4.14 (The quantum curve for the Gromov-Witten invariants of \mathbb{P}^1 , [29]). *The wave function*

$$(4.21) \quad \Psi(x, \hbar) := \exp \left(\frac{1}{\hbar} S_0(x) + S_1(x) + \sum_{2g-2+n>0} \frac{\hbar^{2g-2+n}}{n!} F_{g,n}(x, \dots, x) \right)$$

satisfies the quantum curve equation of an infinite order

$$(4.22) \quad \left[\exp \left(\hbar \frac{d}{dx} \right) + \exp \left(-\hbar \frac{d}{dx} \right) - x \right] \Psi(x, \hbar) = 0.$$

Moreover, the free energies $F_{g,n}(x_1, \dots, x_n)$ as functions in n -variables, and hence all the Gromov-Witten invariants (4.17), can be recovered from the equation (4.22) alone, using the mechanism of the **topological recursion** of [34].

Remark 4.15. The appearance of the extra terms in S_0 and S_1 , in particular, the $\log x$ terms, is trickier than the cases studied in these lectures. We refer to [29, Section 3].

Remark 4.16. Put

$$(4.23) \quad S_m(x) := \sum_{2g-2+n=m-1} \frac{1}{n!} F_{g,n}(x, \dots, x).$$

Then our wave function is of the form

$$(4.24) \quad \Psi(x, \hbar) = \exp \left(\sum_{m=0}^{\infty} \hbar^{m-1} S_m(x) \right),$$

which provides the WKB approximation of the quantum curve equation (4.22). Thus the significance of (4.18) is that the exponential generating function (4.21) of the descendant Gromov-Witten invariants of \mathbb{P}^1 gives the solution to the WKB analysis in a closed formula for the difference equation (4.22).

Remark 4.17. For the case of Hitchin spectral curves [24, 25], the Schrödinger-like equation (4.22) is a direct consequence of the generalized topological recursion. In the $GW(\mathbb{P}^1)$ context, the topological recursion does not play any role in establishing (4.22).

We can recover the classical mechanics corresponding to (4.22) by taking its semi-classical limit, which is the singular perturbation limit

$$(4.25) \quad \lim_{\hbar \rightarrow 0} \left(e^{-\frac{1}{\hbar} S_0(x)} \left[\exp \left(\hbar \frac{d}{dx} \right) + \exp \left(-\hbar \frac{d}{dx} \right) - x \right] e^{\frac{1}{\hbar} S_0(x)} e^{\sum_{m=1}^{\infty} \hbar^{m-1} S_m(x)} \right) \\ = \left(e^{S'_0(x)} + e^{-S'_0(x)} - x \right) e^{S_1(x)} = 0.$$

In terms of new variables $y(x) = S'_0(x)$ and $z(x) = e^{y(x)}$, the semi-classical limit gives us an equation for the spectral curve

$$z \in \Sigma = \mathbb{C}^* \subset \mathbb{C} \times \mathbb{C}^* \xleftarrow{\text{exp}} T^*\mathbb{C} = \mathbb{C}^2 \ni (x, y)$$

by

$$(4.26) \quad \begin{cases} x = z + \frac{1}{z} \\ y = \log z \end{cases}.$$

This is the reason we consider (4.22) as the quantization of the Laudau-Ginzburg model

$$x = z + \frac{1}{z}.$$

It was conjectured in [73] that the stationary Gromov-Witten theory of \mathbb{P}^1 should satisfy the topological recursion with respect to the spectral curve (4.26). The conjecture is solved in [30] as a corollary to its main theorem, which establishes the correspondence between the topological recursion and the Givental formalism.

The key discovery of [29] is that the quantum curve equation (4.22) is equivalent to a recursion equation

$$(4.27) \quad \frac{x}{\hbar} \left(e^{-\hbar \frac{d}{dx}} - 1 \right) X_d(x, \hbar) + \frac{1}{1 + \frac{x}{\hbar}} e^{\hbar \frac{d}{dx}} X_{d-1}(x, \hbar) = 0$$

for a rational function

$$(4.28) \quad X_d(x, \hbar) = \sum_{\lambda \vdash d} \left(\frac{\dim \lambda}{d!} \right)^2 \prod_{i=1}^{\ell(\lambda)} \frac{x + (i - \lambda_i)\hbar}{x + i\hbar}.$$

Here λ is a partition of $d \geq 0$ with parts λ_i and $\dim \lambda$ denotes the dimension of the irreducible representation of the symmetric group S_d characterized by λ .

Acknowledgement. The present article is based on the series of lectures that the authors have given in Singapore, Kobe, Hannover, Hong Kong, and Leiden in 2014–2015. They are indebted to Richard Wentworth and Graeme Wilkin for their kind invitation to the Institute for Mathematical Sciences at the National University of Singapore, where these lectures were first delivered at the IMS Summer Research Institute, *The Geometry, Topology and Physics of Moduli Spaces of Higgs Bundles*, in July 2014. The authors are also grateful to Masa-Hiko Saito for his kind invitation to run the Kobe Summer School consisting of an undergraduate and a graduate courses on the related topics at Kobe University, Japan, in July–August, 2014. A part of these lectures was also delivered at the *Advanced Summer School: Modern Trends in Gromov-Witten Theory*, Leibniz Universität Hannover, in September 2014.

The authors' special thanks are due to Laura P. Schaposnik, whose constant interest in the subject made these lecture notes possible.

The authors are grateful to the American Institute of Mathematics in California, the Banff International Research Station, Max-Planck-Institut für Mathematik in Bonn, and the Lorentz Center for Mathematical Sciences, Leiden, for their hospitality and financial support for the collaboration of the authors. They also thank Jørgen Andersen, Philip Boalch, Gaëtan Borot, Vincent Bouchard, Andrea Brini, Leonid Chekhov, Bertrand Eynard, Laura Fredrickson, Tamás Hausel, Kohei Iwaki, Maxim Kontsevich, Andrew Neitzke, Paul Norbury, Alexei Oblomkov, Brad Safnuk, Albert Schwarz, Sergey Shadrin, Yan Soibelman, and Piotr Sułkowski for useful discussions. O.D. thanks the Perimeter Institute for Theoretical Physics, Canada, and M.M. thanks the University of Amsterdam, l'Institut

des Hautes Études Scientifiques, Hong Kong University of Science and Technology, and the Simons Center for Geometry and Physics, for financial support and hospitality. The research of O.D. has been supported by Leibniz Universität Hannover and MPIM in Bonn. The research of M.M. has been supported by MPIM in Bonn, NSF grants DMS-1104734 and DMS-1309298, and NSF-RNMS: Geometric Structures And Representation Varieties (GEAR Network, DMS-1107452, 1107263, 1107367).

REFERENCES

- [1] M. Aganagic, R. Dijkgraaf, A. Klemm, M. Mariño, and C. Vafa, *Topological Strings and Integrable Hierarchies*, [arXiv:hep-th/0312085], Commun. Math. Phys. **261**, 451–516 (2006).
- [2] G.B. Airy, *On the intensity of light in the neighbourhood of a caustic*, Transactions of the Cambridge Philosophical Society **6**, 379–402 (1838).
- [3] J. Andersen, G. Borot, and N. Orantin, *Modular functors, cohomological field theories and topological recursion*, arXiv:1509.01387 (2015).
- [4] D. Arinkin, *On λ -connections on a curve where λ is a formal parameter*, Mathematical Research Letters **12**, 551–565 (2005).
- [5] A. Beauville, M. S. Narasimhan, and S. Ramanan, *Spectral curves and the generalized theta divisor*, J. reine angew. Math. **398**, 169–179 (1989).
- [6] O. Biquard and P. Boalch, *Wild non-Abelian Hodge theory on curves*, Compos. Math. **140**, 179–204 (2004).
- [7] P. Boalch, *Symplectic manifolds and isomonodromic deformations*, Adv. in Math. **163**, 137–205 (2001).
- [8] P. Boalch, *Hyperkähler manifolds and nonabelian Hodge theory of (irregular) curves*, arXiv:1203.6607v1 (2012).
- [9] G. Borot and B. Eynard, *All-order asymptotics of hyperbolic knot invariants from non-perturbative topological recursion of A -polynomials*, arXiv:1205.2261v1 [math-ph] (2012).
- [10] V. Bouchard, D. Hernández Serrano, X. Liu, and M. Mulase, *Mirror symmetry for orbifold Hurwitz numbers*, arXiv:1301.4871 [math.AG] (2013).
- [11] V. Bouchard, A. Klemm, M. Mariño, and S. Pasquetti, *Remodeling the B-model*, Commun. Math. Phys. **287**, 117–178 (2009).
- [12] V. Bouchard, A. Klemm, M. Mariño, and S. Pasquetti, *Topological open strings on orbifolds*, Commun. Math. Phys. **296**, 589–623 (2010).
- [13] V. Bouchard and M. Mariño, *Hurwitz numbers, matrix models and enumerative geometry*, Proc. Symposia Pure Math. **78**, 263–283 (2008).
- [14] A. Brini, B. Eynard, and M. Mariño, *Torus knots and mirror symmetry* arXiv:1105.2012 (2011).
- [15] K. Chapman, M. Mulase, and B. Safnuk, *The Kontsevich constants for the volume of the moduli of curves and topological recursion*, Communications in Number Theory and Physics **5**, 643–698 (2011).
- [16] L. Chekhov, B. Eynard, and N. Orantin, *Free energy topological expansion for the 2-matrix model*, Journal of High Energy Physics **0612:053** (2006).
- [17] R. Dijkgraaf, H. Fuji, and M. Manabe, *The Volume Conjecture, Perturbative Knot Invariants, and Recursion Relations for Topological Strings*, Nucl. Phys. **B849**, 166–211 (2011).
- [18] R. Dijkgraaf, L. Hollands, and P. Sułkowski, *Quantum curves and \mathcal{D} -modules*, Journal of High Energy Physics **0810.4157**, 1–58 (2009).
- [19] R. Dijkgraaf, L. Hollands P. Sułkowski, and C. Vafa, *Supersymmetric gauge theories, intersecting branes and free Fermions*, Journal of High Energy Physics **0802.106**, (2008).
- [20] R. Dijkgraaf, E. Verlinde, and H. Verlinde, *Loop equations and Virasoro constraints in non-perturbative two-dimensional quantum gravity*, Nucl. Phys. **B348**, 435–456 (1991).
- [21] N. Do, O. Leigh, and P. Norbury, *Orbifold Hurwitz numbers and Eynard-Orantin invariants*, arXiv:1212.6850 (2012).
- [22] N. Do and D. Manescu, *Quantum curves for the enumeration of ribbon graphs and hypermaps*, arXiv:1312.6869 [math.GT] (2013).
- [23] O. Dumitrescu, L. Fredrickson, G. Kydonakis, R. Mazzeo, M. Mulase, and A. Neitzke, *Opers versus nonabelian Hodge*, in preparation.
- [24] O. Dumitrescu and M. Mulase, *Quantum curves for Hitchin fibrations and the Eynard-Orantin theory*, Lett. Math. Phys. **104**, 635–671 (2014).
- [25] O. Dumitrescu and M. Mulase, *Quantization of spectral curves for meromorphic Higgs bundles through topological recursion*, arXiv:1411.1023 (2014).
- [26] O. Dumitrescu and M. Mulase, *Edge-contraction on dual ribbon graphs, 2D TQFT, and the mirror of orbifold Hurwitz numbers*, arXiv:1508.05922 (2015).
- [27] O. Dumitrescu, M. Mulase, A. Sorkin and B. Safnuk, *The spectral curve of the Eynard-Orantin recursion via the Laplace transform*, in “Algebraic and Geometric Aspects of Integrable Systems and Random Matrices,” Dzhamay, Maruno and Pierce, Eds. Contemporary Mathematics **593**, 263–315 (2013).

- [28] P. Dunin-Barkowski, M. Kazarian, N. Orantin, S. Shadrin, and L. Spitz, *Polynomiality of Hurwitz numbers, Bouchard-Mario conjecture, and a new proof of the ELSV formula*, arXiv:1307.4729 [math.AG (physics.hep-th physics.math-ph)] (2013).
- [29] P. Dunin-Barkowski, M. Mulase, P. Norbury, A. Popolitov, and S. Shadrin, *Quantum spectral curve for the Gromov-Witten theory of the complex projective line*, arXiv:1312.5336 [math-ph] (2013).
- [30] P. Dunin-Barkowski, N. Orantin, S. Shadrin, and L. Spitz, *Identification of the Givental formula with the spectral curve topological recursion procedure*, arXiv:1211.4021 [physics.math-ph (math.AG, physics.hep-th)] (2012).
- [31] T. Ekedahl, S. Lando, M. Shapiro, A. Vainshtein, *Hurwitz numbers and intersections on moduli spaces of curves*, Invent. Math. **146**, 297–327 (2001) [arXiv:math/0004096].
- [32] B. Eynard, *A short overview of the “Topological recursion”*, arXiv:1412.3286 (2014).
- [33] B. Eynard, M. Mulase and B. Safnuk, *The Laplace transform of the cut-and-join equation and the Bouchard-Mariño conjecture on Hurwitz numbers*, Publications of the Research Institute for Mathematical Sciences **47**, 629–670 (2011).
- [34] B. Eynard and N. Orantin, *Invariants of algebraic curves and topological expansion*, Communications in Number Theory and Physics **1**, 347–452 (2007).
- [35] B. Eynard and N. Orantin, *Computation of open Gromov-Witten invariants for toric Calabi-Yau 3-folds by topological recursion, a proof of the BKMP conjecture*, arXiv:1205.1103v2 [math-ph] (2013).
- [36] C. Faber, *A conjectural description of the tautological ring of the moduli space of curves*, in “Moduli of curves and abelian varieties,” 109–129, Aspects Math., **E33**, Vieweg, Braunschweig, 1999.
- [37] B. Fang, C.-C. M. Liu, and Z. Zong, *All genus open-closed mirror symmetry for affine toric Calabi-Yau 3-orbifolds*, arXiv:1310.4818 [math.AG] (2013).
- [38] D. Gaiotto, G.W. Moore, and A. Neitzke, *Wall-crossing, Hitchin systems, and the WKB approximation*, arXiv:0907.3987 [hep-th] (2009).
- [39] S. Garoufalidis, *On the characteristic and deformation varieties of a knot*, Geometry & Topology Monographs **7**, 291–309 (2004).
- [40] S. Garoufalidis and T.T.Q. Lê, *The colored Jones function is qholonomic*, Geometry and Topology **9**, 1253–1293 (2005).
- [41] E. Getzler and M.M. Kapranov, *Modular operad*, Compositio Mathematica **110**, 65–126 (1998).
- [42] I.P. Goulden, *A differential operator for symmetric functions and the combinatorics of multiplying transpositions*, Trans. A.M.S., **344**, 421–440 (1994).
- [43] I.P. Goulden and D.M. Jackson, *Transitive factorisations into transpositions and holomorphic mappings on the sphere*, Proc. A.M.S., **125**, 51–60 (1997).
- [44] S. Gukov and P. Sulkowski, *A-polynomial, B-model, and quantization*, arXiv:1108.0002v1 [hep-th] (2011).
- [45] J. Harer and D. Zagier, *The Euler characteristic of the moduli space of curves*, Inventiones Mathematicae **85**, 457–485 (1986).
- [46] N.J. Hitchin, *The self-duality equations on a Riemann surface*, Proc. London Math. Soc. (Ser. 3) **55**, 59–126 (1987).
- [47] N.J. Hitchin, *Stable bundles and integrable systems*, Duke Math. J. **54**, 91–114 (1987).
- [48] L. Hollands, *Topological strings and quantum curves*, Ph.D. Thesis, University of Amsterdam arXiv:0911.3413 [hep-th] (2009).
- [49] E. Ionel, *Relations in the tautological ring of \mathcal{M}_g* , Duke Math. J. **129**, 157–186 (2005).
- [50] K. Iwaki and A. Saens-Rodriguez, *Quantum Curve and the First Painlevé Equation*, arXiv:1507.06557 (2015).
- [51] P. Johnson, R. Pandharipande, and H.H. Tseng, *Abelian Hurwitz-Hodge integrals*, Michigan Math. J. **60**, 171–198 (2011) [arXiv:0803.0499].
- [52] A. Kapustin and E. Witten, *Electric-magnetic duality and the geometric Langlands program*, Communications in Number Theory and Physics **1**, 1–236 (2007).
- [53] R. Kashaev and M. Mariño, *Operators from mirror curves and the quantum dilogarithm*, arXiv:1501.01014 [hep-th], (2015).
- [54] Y. Kodama and V.U. Pierce, *Combinatorics of dispersionless integrable systems and universality in random matrix theory*, Communications in Mathematical Physics **292**, 529–568 (2009).
- [55] M. Kontsevich, *Intersection theory on the moduli space of curves and the matrix Airy function*, Communications in Mathematical Physics **147**, 1–23 (1992).
- [56] M. Kontsevich and Yuri Manin, *Gromov-Witten classes, quantum cohomology, and enumerative geometry*, Communications in Mathematical Physics **164**, 525–562 (1994).
- [57] M. Kontsevich and Y. Soibelman, *Wall-crossing structures in Donaldson-Thomas invariants, integrable systems and Mirror Symmetry*, arXiv:1303.3253 [math.AG] (2013).
- [58] M. Kontsevich and D. Zagier, *Periods*, in “Mathematics unlimited–2001 and beyond”, 771–808, Springer-verlag, Berlin, (2001).
- [59] M. Mariño, *Open string amplitudes and large order behavior in topological string theory*, J. High Energy Physics **0803-060**, 1–33 (2008).
- [60] M. Mirzakhani, *Simple geodesics and Weil-Petersson volumes of moduli spaces of bordered Riemann surfaces*, Invent. Math. **167**, 179–222 (2007).

- [61] M. Mirzakhani, *Weil-Petersson volumes and intersection theory on the moduli space of curves*, J. Amer. Math. Soc. **20**, 1–23 (2007).
- [62] M. Mulase, *Algebraic theory of the KP equations*, in “Perspectives in Mathematical Physics,” R. Penner and S.-T. Yau, Editors, International Press Company, 157–223 (1994).
- [63] M. Mulase and M. Penkava, *Ribbon graphs, quadratic differentials on Riemann surfaces, and algebraic curves defined over \mathbb{Q}* , The Asian Journal of Mathematics **2** (4), 875–920 (1998).
- [64] M. Mulase and M. Penkava, *Topological recursion for the Poincaré polynomial of the combinatorial moduli space of curves*, Advances in Mathematics **230**, 1322–1339 (2012).
- [65] M. Mulase and B. Safnuk, *Mirzakhani’s Recursion Relations, Virasoro Constraints and the KdV Hierarchy*, Indian Journal of Mathematics **50**, 189–228 (2008).
- [66] M. Mulase, S. Shadrin, and L. Spitz, *The spectral curve and the Schrödinger equation of double Hurwitz numbers and higher spin structures*, Communications in Number Theory and Physics **7**, 1–19 (2013).
- [67] M. Mulase and P. Sulkowski, *Spectral curves and the Schrödinger equations for the Eynard-Orantin recursion*, arXiv:1210.3006 (2012).
- [68] M. Mulase and N. Zhang, *Polynomial recursion formula for linear Hodge integrals*, Communications in Number Theory and Physics **4**, 267–294 (2010).
- [69] D. Mumford, *Towards an enumerative geometry of the moduli space of curves*, in “Arithmetic and Geometry,” M. Artin and J. Tate, eds., Part II, 271–328, Birkhäuser, 1983.
- [70] A. Neitzke, *Hitchin systems in $\mathcal{N} = 2$ field theory*, Preprint, November 2013.
- [71] P. Norbury, *Counting lattice points in the moduli space of curves*, arXiv:0801.4590 (2008).
- [72] P. Norbury, *String and dilaton equations for counting lattice points in the moduli space of curves*, arXiv:0905.4141 (2009).
- [73] P. Norbury and N. Scott, *Gromov-Witten invariants of \mathbb{P}^1 and Eynard-Orantin invariants*, arXiv:1106.1337 (2011).
- [74] A. Okounkov and R. Pandharipande, *Gromov-Witten theory, Hurwitz numbers, and matrix models, I*, Proc. Symposia Pure Math. **80**, 325–414 (2009).
- [75] H. Ooguri, P. Sulkowski, M. Yamazaki, *Wall Crossing As Seen By Matrix Models*, arXiv:1005.1293 (2010).
- [76] R. Pandharipande and A. Pixton, *Relations in the tautological ring of the moduli space of curves*, arXiv:1301.4561 [math.AG] (2013).
- [77] F. Petit, *Quantization of spectral curves and DQ-modules*, arXiv:1507.04315 (2015).
- [78] L. Schneps and P. Lochak, *Geometric Galois actions*, London Mathematical Society Lecture Notes Series **242**, 1997.
- [79] C.T. Simpson, *Higgs bundles and local systems*, Publications Mathématiques de l’I.H.E.S. **75**, 5–95 (1992).
- [80] D. D. Sleator, R. E. Tarjan, and W. P. Thurston, *Rotation distance, triangulations, and hyperbolic geometry*, Journal of the American Mathematical Society **1**, 647–681 (1988).
- [81] K. Strebel, *Quadratic differentials*, Springer-Verlag, 1984.
- [82] R. Vakili, Harvard Thesis 1997.
- [83] T.R.S. Walsh and A.B. Lehman, *Counting rooted maps by genus. I*, Journal of Combinatorial Theory **B-13**, 192–218 (1972).
- [84] E. Witten, *Two dimensional gravity and intersection theory on moduli space*, Surveys in Differential Geometry **1**, 243–310 (1991).
- [85] E. Witten, *Gauge theory and wild ramification*, Analysis and Applications **6**, 429–501 (2008).
- [86] E. Witten, *Mirror symmetry, Hitchin’s equations, and Langlands duality*, arXiv:0802.0999 (2008).
- [87] E. Witten, *Analytic continuation of Chern-Simons theory*, arXiv:1001.2933v4 [hep-th] (2010).
- [88] J. Zhou, *Local Mirror Symmetry for the Topological Vertex* arXiv:0911.2343 (2009).
- [89] J. Zhou, *Intersection numbers on Deligne-Mumford moduli spaces and quantum Airy curve*, arXiv:1206.5896 [math.AG] (2012).
- [90] J. Zhou, *Quantum Mirror Curves for \mathbb{C}^3 and the Resolved Conifold*, arXiv:1207.0598v1 [math.AG] (2012).
- [91] M. Zhou, *The differential recursion equation for the generalized Catalan numbers*, Senior Thesis, University of California, Davis (2013).

OLIVIA DUMITRESCU: MAX-PLANCK-INSTITUT FÜR MATHEMATIK, VIVATSGASSE 7, 53111 BONN, GERMANY,

INSTITUT FÜR ALGEBRAISCHE GEOMETRIE, FAKULTÄT FÜR MATHEMATIK UND PHYSIK, LEIBNIZ UNIVERSITÄT HANNOVER, WELFENGARTEN 1, 30167 HANNOVER, GERMANY

AND SIMION STOILOW INSTITUTE OF MATHEMATICS, ROMANIAN ACADEMY, 21 CALEA GRIVITEI STREET, 010702 BUCHAREST, ROMANIA

E-mail address: dumitrescu@mpim-bonn.mpg.de

MOTOHICO MULASE: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, DAVIS, CA 95616–8633, U.S.A.,

AND KAVLI INSTITUTE FOR PHYSICS AND MATHEMATICS OF THE UNIVERSE, THE UNIVERSITY OF TOKYO, KASHIWA,
JAPAN

E-mail address: `mulase@math.ucdavis.edu`