

UCSF

UC San Francisco Previously Published Works

Title

Giardavirus Internal Ribosome Entry Site Has an Apparently Unique Mechanism of Initiating Translation

Permalink

<https://escholarship.org/uc/item/8649s7zq>

Journal

PLOS ONE, 4(10)

ISSN

1932-6203

Authors

Garlapati, Srinivas
Wang, Ching C

Publication Date

2009

DOI

10.1371/journal.pone.0007435

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Giardavirus Internal Ribosome Entry Site Has an Apparently Unique Mechanism of Initiating Translation

Srinivas Garlapati, Ching C. Wang*

Department of Pharmaceutical Chemistry, University of California San Francisco, San Francisco, California, United States of America

Abstract

Giardavirus (GLV) utilizes an internal ribosome entry site (IRES) for translation initiation in the early branching eukaryote *Giardia lamblia*. Unlike most of the viral IRESs among higher eukaryotes, which localize primarily within the 5'-untranslated region (UTR), the GLV IRES comprises 253 nts of 5'UTR and the initial 264 nts in the open-reading-frame (ORF). To test if GLV IRES also functions in higher eukaryotic systems, we examined it in rabbit reticulocyte lysate (RRL) and found that it functions much less efficiently than the IRES from the Encephalomyocarditis virus (EMCV) or Cricket paralysis virus (CrPV). In contrast, both EMCV-IRES and CrPV-IRESs were inactive in transfected *Giardia* cells. Structure-function analysis indicated that only the stem-loop U5 from the 5'UTR and the stem-loop I plus the downstream box (Dbox) from the ORF of GLV IRES are required for limited IRES function in RRL. Edeine, a translation initiation inhibitor, did not significantly affect the function of GLV IRES in either RRL or *Giardia*, indicating that a pre-initiation complex is not required for GLV IRES-mediated translation initiation. However, the small ribosomal subunit purified from *Giardia* did not bind to GLV IRES, indicating that additional protein factors may be necessary. A member of the helicase family IBP1 and two known viral IRES binding proteins La autoantigen and SRp20 have been identified in *Giardia* that bind to GLV IRES *in vitro*. These three proteins could be involved in facilitating small ribosome recruitment for initiating translation.

Citation: Garlapati S, Wang CC (2009) Giardavirus Internal Ribosome Entry Site Has an Apparently Unique Mechanism of Initiating Translation. PLoS ONE 4(10): e7435. doi:10.1371/journal.pone.0007435

Editor: Alexander S. Spirin, Institute of Protein Research, Russian Academy of Sciences, Russian Federation

Received: May 20, 2009; **Accepted:** September 12, 2009; **Published:** October 14, 2009

Copyright: © 2009 Garlapati, Wang. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by Grant AI-30475 from the National Institutes of Health to Ching C. Wang. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: ccwang@cgl.ucsf.edu

Introduction

Internal ribosome entry site (IRES) mediated translation is an alternative mechanism of translation initiation adopted by many viruses and some cellular mRNAs among higher eukaryotes [1–3]. Among the members of *Picornaviridae* and *Flaviviridae* families, the IRESs are primarily located in the 5'-untranslated regions (UTRs) of the transcripts [3]. However, some rare exceptions such as the Dicistroviruses with IRES elements in the intergenic regions (IGR) [4,5] and HIV 2 having an IRES element entirely in the downstream coding region have been identified [6]. For the cellular mRNAs, IRESs are primarily located in the 5' UTRs and often function in a cell-cycle-dependent manner [7,8].

In the cap-dependent translation initiation, the 40S ribosomal subunit complexed with initiation factors eIF3 and eIF2-GTP-Met.tRNA_i (43S pre-initiation complex), binds to the 5' cap structure of the mRNA via the eIF4F complex and subsequently scans for the start codon [9]. In contrast, the IRES mediated translation initiation involves direct recruitment of the translation machinery that positions the 40S small ribosomal subunit onto the start codon [1]. The process of 40S ribosome binding to IRES varies with different types of IRES and also with the protein factors involved [10,11]. For example, the poliovirus (PV) IRES and Encephalomyocarditis virus (EMCV) IRES require all the canonical initiation factors except for the cap binding protein eIF4E to recruit the 43S pre-initiation complex [12–14], whereas the Hepatitis C virus (HCV) IRES requires only the binding of initiation factor eIF3 for efficient recruitment of naked 40S

ribosome [15–17]. In contrast, the IRESes present in the intergenic regions (IGR) of Dicistroviruses do not require any initiation factors for binding to the 40S ribosome [18–20].

In addition to the initiation factors, some of the viral IRESs also bind non-canonical protein factors known as IRES *trans*-acting factors (ITAFs) that have been shown to stimulate IRES activity. For instance, La autoantigen was shown to enhance the IRES activity of PV IRES [21,22], HCV IRES [23–25], EMCV IRES [26] and Coxsackievirus B3 IRES [27], whereas the polypyrimidine tract binding protein (PTB) was found to bind and induce conformational changes in EMCV IRES [28–30], PV IRES [31], and Foot-and-mouth disease virus (FMDV) IRES [30]. Recently, a splicing factor SRp20 was also implicated in PV IRES activity via an interaction with Poly (rC) binding protein 2 (PCBP2) [32]. A distinct class of RNA binding proteins containing three K-homologous (KH) domains such as heterogeneous nuclear ribonuclear proteins hnRNPE1 and hnRNPE2 have also been shown to bind to PV IRES [33,34] and Hepatitis A virus IRES [35]. For many cellular IRESes, several proteins that are involved in mRNA splicing and transport were identified as *trans*-acting factors [2,3,36,37].

Viruses of the *Totiviridae* family represent a small group of double stranded RNA viruses that infect protozoan parasites and lower fungi [38]. Giardavirus (GLV), a member of the *Totiviridae* family, inhabits the cytoplasm of an early branching protozoan parasite *Giardia lamblia* [39]. Its transcript encodes two proteins: a major capsid protein of 100 kDa and a minor 190 kDa gag-pol fusion protein produced via a –1 ribosomal frame-shift [40,41]. It

lacks a 5' cap structure but contains a highly structured 5' untranslated region (UTR) [42]. The 5'UTR alone is not sufficient to initiate the translation of the viral transcript. It needs to combine with a 264 base stretch of the downstream coding sequence to function as an IRES in *Giardia* [43,44]. This unusual IRES has been subjected to a thorough structure-function analysis that identified several complex secondary structures essential for IRES function (Fig. 1) [45–47]. However, how GLV IRES recruits the host translation machinery to initiate protein synthesis in *Giardia* is poorly understood.

The viral host *Giardia* is known to possess the translation machinery with many unusual features, such as the involvement of a 70S instead of a 80S ribosome [48] and the translation initiation factors that are either missing or structurally divergent as compared to that of the higher eukaryotes [49]. To learn whether the GLV IRES may function in a novel mechanism of translation initiation due to the unusual translation machinery in *Giardia*, we tested its function in rabbit reticulocyte lysates (RRL) and found significantly reduced IRES activity with very simple structural requirement. Further analysis indicated that GLV IRES may not require a pre-initiation complex for initiating translation both in *Giardia* and in RRL, nor does it bind to *Giardia* small ribosomal subunit *in vitro*. A member of the helicase family GILBP1 is identified from the affinity purified IRES-protein complexes formed in *Giardia* lysate. In addition, two ITAF homologues La autoantigen and SRp20 are identified in *Giardia* that exhibited binding to the GLV IRES *in vitro*. These results suggest a mechanism of GLV IRES binding to the ITAF protein factors prior to ribosomal recruitment.

Results

The function of GLV IRES in RRL

We used *in vitro* synthesized dicistronic viral transcripts that contained two consecutive reporters Rluc and Fluc, and monitored their expressions in RRL and compared the results obtained with those from the *Giardia* trophozoites transfected with the same transcripts [44] (Fig. 2). For the control transcript pC631Rluc-Fluc (with Rluc and Fluc separated by 10 nts), a significant Rluc activity of $5,018,431.5 \pm 284,394.9$ RLU, and an approximately 100-fold lower Fluc activity of $52,848.2 \pm 7,173.2$ RLU was observed, resulting in a Fluc/Rluc ratio of $10.3 \pm 0.9 \times 10^{-3}$ (Fig. 2A). When the GLV 5'UTR sequence was inserted between the two cistrons

of the control transcript, the F/R ratio became $5.3 \pm 1.2 \times 10^{-3}$ (pC631Rluc-UTR-Fluc, Fig. 2D), whereas an insertion of the 264 nt downstream coding region from the GLV transcript (pC631Rluc-Cod-Fluc) resulted in a ratio of $9.9 \pm 0.2 \times 10^{-3}$ (Fig. 2E), suggesting that neither the 5'UTR nor the 264 nts of coding region alone has the IRES activity. However, when the entire GLV IRES was placed in the inter-cistronic region (pC631Rluc-UTRCod-Fluc), the F/R ratio was $28.0 \pm 0.5 \times 10^{-3}$, representing a 2 to 3-fold increase from the control value (Fig. 2F). These results indicate that, as in *Giardia*, the combination of the 5'UTR with 264 nts of capsid coding region are required for the relatively low IRES activity in RRL. When EMCV IRES or CrPV IRES was placed in between the two cistrons, the F/R ratios in RRL were raised to $591.1 \pm 25.1 \times 10^{-3}$ and $89.9 \pm 5.3 \times 10^{-3}$, respectively (Fig. 2, B and C), indicating that they function much more efficiently than GLV IRES. In contrast, both EMCV IRES and CrPV IRES were unable to drive Fluc expression from the dicistronic transcripts in transfected *Giardia* cells (Fig. 2, B and C), suggesting that these IRESs are inefficient in recruiting the translation machinery of *Giardia*.

A structure-function analysis of GLV IRES in RRL

In *Giardia*, the functional GLV-IRES spans from nucleotide #114 to #631 in the viral transcript [44]. It includes essential structures such as a pseudoknot U3 (nt #134–176), stem-loops U4a (nt #204–219), U4b (nt #221–261), U4c (nt #263–292) and U5 (nt #314–344) in the 5'UTR, and stem-loop I (nt #378–402), a downstream box (Dbox) sequence (nt #433–445), and another pseudoknot (nts 511–587) in the coding region (Fig. 1). To determine if these secondary structures are also required for the limited GLV-IRES function in RRL, various deletion mutants of GLV-IRES were tested [45,47]. The results indicated that the U3 pseudoknot, U4a, b and c stem-loops from the 5'UTR (Fig. 3B–D) and the pseudoknot from the downstream capsid coding region (Fig. 3E) are not required for IRES function. The two stem-loop structures, U5 from the 5'UTR and I from the coding region, flanking a 31 nt un-structured sequence with the initiation codon AUG localized at the center was sufficient to function as an IRES in RRL. These structures were originally postulated to accommodate the recruited 40S ribosomal subunit in *Giardia* [44] and may still serve the same function in RRL. Interestingly, the results also indicated that the Dbox is apparently involved in the IRES




Figure 1. The essential secondary structures of GLV IRES. Secondary structures identified in the GLV IRES by chemical/enzymatic structure probing and site-directed mutagenesis. doi:10.1371/journal.pone.0007435.g001




Figure 2. GLV IRES activity in RRL and in transfected *Giardia* WB trophozoites. Transcripts were synthesized from various dicistronic cDNA constructs each consisting of the Renilla luciferase Rluc (shaded box) and Photinus luciferase Fluc (hatched box) genes flanked by the 5' and 3' portions of GLV cDNA (the coding region, dash-hatched box; 5' and 3' UTRs, black bars) and located downstream from the T7 promoter (black box). The Rluc and Fluc cistrons are separated by 10 nucleotides in the control pC631Rluc-Fluc transcript (indicated by small black box). Schematic diagrams of dicistronic cDNA constructs (1) pC631Rluc-Fluc, (2) pC631Rluc-UTR-Fluc, (3) pC631Rluc-Cod-Fluc, (4) pC631Rluc-UTRCod-Fluc, (5) pC631Rluc-EMCV-Fluc, and (6) pC631Rluc-CrPV-Fluc are presented. Reaction products from RRL and transfected *Giardia* trophozoites were assayed for Ruc and Fluc activities and the IRES activity observed with each transcript was expressed as a ratio between the two luciferase activities (Fluc/Rluc × 10³). doi:10.1371/journal.pone.0007435.g002

activity in RRL (Fig. 3, compare E and F). Since Dbox was postulated to bind the 3'-end of the 16S-like ribosomal RNA in *Giardia* [50], it is not immediately clear what function it may perform in RRL. This significantly shortened GLV IRES in RRL is only 210 nts and is approximately twice as active as the full length GLV IRES in RRL (Fig. 3, G). To rule out the possibility that this enhanced activity is due to read-through of the first cistron, it was inserted into the dicistronic transcript in an inverted orientation (Fig. 3, H). The Fluc activity was completely lost, suggesting a real, albeit limited, function of this truncated GLV IRES in RRL.

GLV IRES does not require binding of the pre-initiation complex to initiate translation

A pre-initiation complex consisting of 40S ribosome complexed with eIF3, eIF2-GTP-Met.tRNA_i, is required for cap mediated as well as EMCV IRES mediated translation initiation [9,12,13]. In contrast, CrPV IRES [51] and HCV IRES [52] do not require

this pre-initiation complex as they can directly bind the 40S ribosome to initiate translation [16,18]. These two modes of ribosome recruitment are distinguishable by using the translation initiation inhibitor edeine that prevents the recognition of the initiation codon by a pre-initiation complex in RRL [53,54], thereby inhibits the cap-mediated and EMCV IRES driven but not the CrPV IRES and HCV IRES mediated translation initiation [51,52,55]. To determine if GLV IRES requires a pre-initiation complex for translation initiation, we tested 5' capped dicistronic constructs in RRL in the presence of edeine and we found that the GLV IRES mediated translation was unaffected up to 0.5 μM of the drug (Fig. 4C), whereas the cap-mediated as well as the EMCV IRES mediated translation was inhibited by edeine in a dose dependent manner (Fig. 4A). As expected, the CrPV IRES mediated translation was also unaffected by edeine up to 0.5 μM (Fig. 4B). Interestingly, the truncated GLV IRES (nt #276–487) was also found to be insensitive to the drug up to 1 μM (Fig. 4D).




Figure 3. Structural requirements for GLV IRES activity in RRL and in *Giardia*. Effect of varying deletions in the 5'UTR and in the 264 nt coding region on GLV-IRES activity observed in RRL and in transfected *G. lamblia* WB trophozoites. The extent of deletion in each region is mentioned in the text and referred by the number of the transcript in the figure. The Rluc and Fluc activities were assayed from RRL reaction mixture and *Giardia* lysates. ND, not determined.

doi:10.1371/journal.pone.0007435.g003




Figure 4. Effect of edeine on IRES functions in RRL. The translation of dicistronic transcripts (A) Cap-Rluc-EMCV IRES-Fluc, (B) Cap-Rluc-CrPV IRES-Fluc, (C) Cap-Rluc-GLV IRES (1–631 nts) Fluc and (D) Cap-Rluc-GLV IRES (276–487) Fluc in RRL, in the presence of varying concentrations of edeine was assayed. The luciferase activity of Rluc (dotted bars) and Fluc (gray bars) are presented as percentage (%) relative luciferase activities as compared to the untreated control samples (without edeine). Error bars represent standard errors from assaying triplicate samples in the same experiment. doi:10.1371/journal.pone.0007435.g004

To determine if GLV IRES is also resistant to edeine while functioning in the transfected *Giardia*, the 5' capped dicistronic transcript in combination with varying concentrations of edeine were co-electroporated into *Giardia* cells. Cells lysed 5 hr post-transfection were assayed for Rluc and Fluc activities. As in RRL, cap-mediated translation of the Rluc was significantly inhibited by edeine at concentrations above 0.25 μ M, with a 30% inhibition achieved at 1 μ M (Fig. 5). In contrast, GLV IRES activity was not inhibited even up to 1 μ M of edeine (Fig. 5). These results indicate that GLV IRES does not require formation of a pre-initiation complex for initiating translation both in *Giardia* and in RRL.

GLV-IRES does not bind to the small ribosomal subunit purified from *Giardia*

Since the function of GLV IRES was resistant to edeine, we tested its possible binding to the purified small ribosomal subunit in the absence of any initiation factors as CrPV IRES or HCV IRES are known to do with the mammalian small ribosomal subunit [16,18]. Purified 40S ribosomal subunit from RRL was incubated with radiolabeled GLV IRES RNA or CrPV IRES RNA and the 40S-IRES RNA complexes were separated in a 10–30% sucrose density gradient. Only 10% of the labeled GLV IRES RNA was detected in the fractions containing the small ribosome subunit, indicating limited binding of GLV IRES RNA to the rabbit small ribosomal subunit. In contrast, more than 50%




Figure 5. Effect of edeine on the GLV IRES activity in transfected *G. lamblia* WB trophozoites. The dicistronic transcript Cap-Rluc-GLV IRES (1–631 nt)-Fluc was electroporated into *Giardia* cells in combination with varying concentrations of edeine. The Rluc (dotted bars) and Fluc (gray bars) activities were assayed after 5 hours post transfection and expressed as relative luciferase activities compared to the no-drug control. The error bars represent the standard errors from assaying triplicate samples in the same experiment. doi:10.1371/journal.pone.0007435.g005

of the labeled CrPV IRES RNA was bound to the small subunit (Fig. 6A). To investigate if the small ribosomal subunit from *Giardia* binds to CrPV or GLV IRES RNA, we tested the latter on the small ribosomal subunit purified from *Giardia* by the same procedure as that from RRL, but no apparent binding of either IRES was detectable in the sucrose gradient (Fig. 6B) or in the filter binding assay (data not shown). The failure of CrPV-IRES in binding to *Giardia* small ribosomal subunit agrees with its failure in initiating translation in *Giardia* (see Figure 2). The lack of binding between GLV IRES and purified small ribosomal subunit from *Giardia* suggests that protein factors other than those required for pre-initiation complex formation could be required prior to the ribosomal recruitment by GLV IRES.

Formation and analysis of IRES-protein complexes

In order to biochemically identify the potential factors in *Giardia* that may bind to GLV IRES to enable the latter to recruit the ribosomal small subunit, we incubated the radiolabeled GLV IRES with *Giardia* cell lysates and analyzed the potential RNA-protein complexes thus formed by sucrose density gradient centrifugation. The GLV IRES-lysate mixture was separated into two distinct peaks (peaks 1 and 2) both heavier than the single GLV IRES peak in the no lysate control (Fig. 7A). The two heavier peaks were significantly reduced by a 5-fold excess of unlabeled GLV IRES but remained unchanged when an equivalent amount of random yeast RNA fragments of 300–500 nts (Ambion) was added to the mixture (Fig. 7D), suggesting that specific complexes between GLV IRES and certain components in the *Giardia* lysates were formed that constituted peaks 1 and 2. To further analyze these complexes, individual fractions collected from the gradient were each concentrated and separated on composite agarose (0.5%): acrylamide (2.75%) gels [19]. A steadily increasing shift of the radiolabeled GLV IRES RNA band toward slower mobility was clearly demonstrated from fractions #6 to #13, suggesting complexes of different sizes between GLV IRES RNA and the lysates (Fig. 7B). Peak 1 consisted of fractions #6 and #7, which had the highest radioactivity and well-defined shift in mobility, but there was no detectable ribosomal RNA in it. It may be the dominant form of GLV IRES RNA-lysate complex formed under the current experimental conditions and could be the GLV IRES RNA-ITAFs complex formed in *Giardia* prior to ribosomal recruitment. When peak 2, consisting of fractions #10 and 11, was analyzed for ribosomal RNA in an agarose gel, the 16S small ribosomal RNA band was detected primarily in fractions #10 and 11 (Fig. 7C). The peak could thus represent the complex between peak 1 and the small ribosomal subunit, i.e., the GLV IRES RNA-ITAFs-small ribosomal subunit complex, the likely initiator formed on GLV IRES.

Affinity purification of the GLV IRES-binding proteins using StreptoTag

In order to purify the GLV IRES-protein complexes formed in the *Giardia* lysate, and observed as peak 1 in sucrose density gradient (Figure 7A), we adopted a method of affinity purifying StreptoTagged RNA-protein complex through a dihydrostreptomycin coupled Sepharose 6B column [56], successfully used previously in purifying the 48S translation initiation complexes from RRL [57,58]. Hybrid RNA containing the GLV IRES with the StreptoTag at its 3' end through a linker was synthesized. The linker contained a primer-binding site and three repeats of CU's [57] to prevent potential base pairings between GLV IRES and the StreptoTag sequence (checked by MFOLD program). The radiolabeled and StreptoTagged GLV IRES was then incubated

with *Giardia* lysates and loaded onto the streptomycin-Sepharose 6B column. After several washes with the column buffer, the bound complexes were eluted with 10 μ M of streptomycin. The peak fractions (#10–12, Fig. 8A) from the eluted samples were pooled, concentrated and subjected to SDS-PAGE analysis. Two major protein bands of molecular sizes \sim 45 and \sim 90 kDa were identified (Fig. 8B), which could be components of the peak 1 complex detected in the previous sucrose density gradient (Fig. 7B).

The 45 and 90 kDa protein bands were subjected to mass spectrometry using the MADLI-TOF procedure and the resulting peptides were used to identify the protein in the *Giardia* genome database. MS-FIT program matched 47 peptides in the tryptic digest of the 45 kDa band to a protein (accession # XP_770230) that belongs to the super-family I of DNA and RNA helicases, with a MOWSE score of $6.22 \times 10^{+6}$ ($p < 0.05$) (Fig. S1). This protein hereafter is referred to as the IRES binding protein 1 (IBP1). IBP1 contains two COG1112 domains that are characteristic of Super-family I of DNA and RNA helicases, however, it does not share sequence homology with any of the RNA binding proteins nor contains conserved RNA binding motifs. The encoding DNA sequence of IBP1 was amplified by PCR, cloned, expressed as a maltose binding protein (MBP) fusion protein in transformed *Escherichia coli* and affinity purified (Fig. 9A). The purified protein was then tested for binding to GLV IRES RNA in gel-shift analysis. The mobility of radiolabeled GLV IRES was significantly reduced by increasing concentrations of MBP-IBP1 (Fig. 9B, lanes 2–4) but unaffected by MBP (Fig. 9B, lane 7). Moreover, this binding was effectively competed off by a 10-fold excess of unlabeled GLV IRES (Fig. 9B, lane 6) but not by a 100-fold excess of yeast RNA of random 300–400 nt sequences (Ambion) (Fig. 9B, lane 5). These results indicate that IBP1 specifically binds to GLV IRES. It was the first *Giardia* protein identified to bind to GLV IRES.

The peptides from the tryptic digest of the other GLV IRES-binding 90 kDa protein could not be matched to any of the existing coding sequences in the *Giardia* genome database and it was not pursued further.

Identification in *Giardia* of homologues of trans-acting factors known to bind to other viral IRESs

There have been many non-canonical *trans*-acting protein factors (ITAFs) identified in higher eukaryotes that are found to bind and stimulate the activity of various viral IRESs [3]. These ITAFs were used to search for homologues in the *Giardia* genome database (<http://giardiadb.org/giardiadb/>). A homologue of La autoantigen, which was found to bind HCV, EMCV, PV and Coxsackie virus IRES elements [21,23,26,27], a homologue of SRp20, which is known to bind to Poliovirus IRES [32] and a homologue of hnRNP E2 involved in binding to PV IRES [34] were identified in *Giardia*. The La homologue in *Giardia* contains the characteristic La domain within the first 100 amino acids, followed by the RNA recognition motifs (RRM) 2 and 3, and a divergent C-terminal domain (Fig. S2), whereas SRp20 shares significant sequence identity with mouse and human SRp20 only at the N terminus (Fig. S3). In contrast to human hnRNPE2, *Giardia* hnRNPE2 contained only a single KH domain (data not shown). DNA fragments encoding the three homologues (GILa, accession # XP_001705495; GISRp20, accession # XP_001708843; GlnRNPE2, accession # XP_00170795) were cloned, expressed in the form of 6X-His tagged fusion proteins in transformed *E. coli*, and affinity purified (Fig. S4). The purified proteins were tested for their binding to GLV IRES in the gel shift assays. GILa showed a strong binding to GLV IRES RNA in a dose dependent manner (Fig. 10, lanes 2–4). This binding was significantly reduced in the presence of 5 to 10-fold excess of

A. RRL



B. *Giardia*

Figure 6. Binding of small ribosomal subunits to IRESs. Small ribosomal subunits were purified from (A) RRL and (B) *Giardia* trophozoites. They were incubated with radiolabeled GLV IRES and CrPV IRES respectively and fractionated in 10–30% sucrose density gradient centrifugations. Fractions were collected from each gradient and percent radioactivity in each fraction was recorded.
doi:10.1371/journal.pone.0007435.g006




Figure 7. Formation of IRES-associated complexes in *Giardia* cell lysates. A. IRES-associated complexes were resolved into two peaks (Peaks 1 and 2) by 10–50% linear sucrose density gradient centrifugation. B. Analysis of sucrose density gradient fractions from #6 to #13 by composite agarose:acrylamide gel electrophoresis followed by autoradiography. C. The total RNA from fraction # 7–13 were concentrated and analyzed by agarose gel electrophoresis to visualize the presence of ribosomal RNA. D. Analysis of IRES-associated complexes formed in the *Giardia* lysate in the presence of unlabeled IRES RNA (5-fold excess) or non-specific yeast RNA (5-fold excess) (Ambion). doi:10.1371/journal.pone.0007435.g007

unlabeled GLV IRES (Fig. 10, lanes 5–6) but unaffected by an excessive amount of yeast RNA (Fig. 10, lanes 7–8). Similarly, GISRp20 also exhibited binding to radiolabeled GLV IRES in a dose dependent manner (Fig. 11A, lanes 2–4), which was significantly reduced by 15-fold excess of unlabeled GLV-IRES (Fig. 11B, lanes 3–5) but unaffected by excessive yeast RNA (Ambion) (Fig. 11B, lanes 6–8) indicating specific binding of GISRp20 to GLV IRES. In contrast to GISRp20 and GILa proteins, recombinant GlhnRNPE2 did not show detectable binding to radiolabeled GLV IRES RNA (data not shown). Thus, two additional proteins in *Giardia*, GISRp20 and GILa, demonstrating specific binding to GLV IRES were identified. Based on the known functions of their homologues in higher eukaryotes, the potential functions of GILa and GISRp20 on GLV IRES activity in *Giardia* could be postulated.

To determine the potential roles of the three putative trans-acting protein factors GIBP1, GILa and GISRp20 in GLV IRES mediated translation initiation, we tested whether their bindings to GLV IRES would help recruit the *Giardia* small ribosomal subunit. The result shown in Figure 12 indicates that the RNA-protein complex is incapable of recruiting the small ribosomal subunit.

Apparently, the three proteins do not constitute the entire spectrum of ITAFs needed for ribosomal recruitment. But the shift of the RNA-protein complex toward heavier fractions in the sucrose density gradient suggests that additional proteins may be needed to join the complex to make the complex we observed in peak 1 in Figure 7A between GLV IRES and *Giardia* lysates.

Discussion

Our study shows that GLV IRES is functional in RRL system despite the evolutionary divergence between *Giardia* and rabbit. However, it functions at a much lower efficiency than EMCV IRES or CrPV IRES. This GLV IRES function in RRL requires a significantly truncated structure involving only the upstream stem-loop U5 and the downstream stem-loop I plus the Dbox, flanking the AUG codon in the center. This is the structure amid a highly complex assembly of secondary structures when the original IRES was found to function in *Giardia* [44]. Requirement of additional structures for optimal IRES activity in *Giardia* could reflect the functional differences between the translation machineries of *Giardia* and RRL.




Figure 8. Purification of GLV IRES-associated complexes using streptomycin-sepharose 6B column chromatography. A. The elution profile of a mixture of radiolabeled Strepto-tagged IRES RNA and *Giardia* lysates through a streptomycin-sepharose 6B column. B. Sybroruby stained 10% SDS-PAGE gel of the pooled fractions #10-#12 of *Giardia* lysates alone (lane 1), and the corresponding fractions of *Giardia* lysates plus the Strepto-tagged GLV-IRES RNA (lane 2). doi:10.1371/journal.pone.0007435.g008

Our data also indicate that GLV IRES may follow a distinct pathway for forming an initiation complex. Resistance of GLV IRES to the translation initiation inhibitor, edeine, indicates that a 40S-eIF2-GTP/Met-tRNA_i pre-initiation complex is not required for initiating translation. Edeine has been shown to bind between the P- and the E-sites of the ribosome and interfere with the recognition of the AUG codon by a scanning pre-initiation complex [53,54,59]. The resistance of CrPV IRES to edeine inhibition is attributed to its ability to bind directly to the small ribosomal subunit and position pseudoknot structure PKI in the P-site of the ribosome, and initiate translation from the non-canonical GCU codon positioned at the ribosomal A-site [55,60,61]. Similarly, HCV IRES has also been shown to resist edeine inhibition at relatively low concentrations [52,62] and yet it initiates translation from the AUG codon located at the P-site of the ribosome [17,52], by first binding directly to the small

ribosomal subunit in the absence of initiation factors and then recruiting the eIF3 and eIF2-GTP-tRNA ternary complex [16,17,63,64]. Since GLV IRES does not share sequence or structural similarity with CrPV IRES or HCV IRES, the mechanism of its resistance to edeine could be quite different. GLV IRES utilizes the single AUG located between stem-loop U5 and I to initiate translation [46]. It may be employing a mechanism resembling that of the HCV IRES, involving initial binding of the small ribosomal subunit to form initiation complexes that bypasses the inhibitory effect of edeine. But the failure of GLV IRES in directly binding to *Giardia* small ribosomal subunit suggested a more complex situation. It is well established that the ability of CrPV IRES and HCV IRES to directly bind to small ribosomal subunit is due to their inherent nature to fold into a compact three-dimensional structure containing specific ribosome binding domains [16,18,65,66]. The lack of GLV IRES




Figure 9. Purification of the MBP-IBP1 fusion protein expressed in *E. coli* and examination of its binding to GLV IRES RNA in gel-shift assays. A. SDS-PAGE analysis of the MBP-IBP1 fusion protein (indicated by an arrow) purified from the *E. coli* lysates using an amylose agarose column. B. Binding of varying amounts of MBP-IBP1 to ³²P-labeled GLV IRES RNA that was analyzed by composite agarose (0.5%):acrylamide (2.75%) gel electrophoresis and autoradiography. The RNA-protein complexes have slower gel mobility than the free RNA. Lane 1: free IRES RNA, Lanes 2–4: IRES RNA incubated with 1.5, 3.5, and 4.5 μg of MBP-IBP1 fusion protein, Lane 5, IRES RNA binding to MBP-IBP1 is unaffected by an excess of non-specific yeast RNA, Lane 6, IRES RNA binding to MBP-IBP1 is competed off by unlabeled IRES RNA, Lane 7, IRES RNA does not bind to purified MBP. doi:10.1371/journal.pone.0007435.g009

binding to the *Giardia* small ribosomal subunit could be attributed to an inability to fold into such a compact three-dimensional structure. The secondary and tertiary structures in the 5'UTR




Figure 10. GLV IRES RNA binds to recombinant GILa protein. Varying amounts of purified GILa protein was incubated with radiolabeled GLV IRES RNA (lanes 2–4) and the RNA-protein complexes were separated by composite agarose (0.5%): acrylamide (2.75%) gel electrophoresis and monitored by autoradiography. The binding was reduced by 5 to 10 fold excess of unlabeled IRES RNA (lanes 5 and 6), but unaffected by 5 to 10-fold excess of non-specific yeast RNA (lanes 7 and 8). doi:10.1371/journal.pone.0007435.g010

region of GLV IRES are each separated by single stranded regions [47]. They may not form a compact high-order structure on their own but may bind to protein factors known to promote structural organization of an IRES to form a structure capable of binding directly to the small ribosomal subunit [10,11]. The detection of peaks 1 and 2 from a GLV IRES and *Giardia* lysate mixture in a sucrose gradient is consistent with this possibility. IBP1, identified in this mixture, is most likely playing a role in the binding between the small ribosomal subunit and GLV IRES.

In addition to IBP1, we identified a homologue of La autoantigen in *Giardia* that binds to GLV IRES specifically *in vitro*. Human La autoantigen is a dimer and contains several RNA binding domains [24]. It binds to a specific region (nt position 559–624) in the PV IRES and stimulates the 48S complex formation both *in vitro* and *in vivo* [22,24]. It also binds near the initiation codon of the HCV IRES to stimulate the IRES activity [23], presumably through stabilizing the IRES structure and thus facilitating the interaction with the translation machinery [67]. Since, La-motif containing proteins are highly conserved among eukaryotes [68], GILa could have similar functions as human La. It could be stabilizing the structure of GLV IRES and thus facilitating the binding of small ribosomal subunit.

SRp20 has been reported to stimulate PV IRES activity by interacting with hnRNPE2 [32] and to co-sediment with the 80S ribosome and polysomes [69]. GISRp20, showing specific binding to GLV IRES *in vitro*, could have a similar function on GLV IRES through a direct interaction with the small ribosomal subunit.

In conclusion, our data indicate that GLV IRES utilizes a distinctive pathway of recruiting the small ribosomal subunit that does not require a pre-initiation complex. Inability of the three ITAFs to stimulate recruitment of the small ribosomal subunit to




Figure 11. GLV IRES RNA binds to recombinant GISRp20 protein. A. Varying amounts of purified GISRp20 protein were incubated with radiolabeled GLV IRES RNA and analyzed by composite agarose (0.5%); acrylamide (2.75%) gel electrophoresis and autoradiography (lanes 1–4). B. The binding was competed by 5, 10 and 15 fold-excess of unlabeled GLV IRES RNA (lanes 3–5) but not by 5, 10, and 15 fold excess of non-specific yeast RNA (Ambion) (lanes 6–8).
doi:10.1371/journal.pone.0007435.g011

GLV IRES indicates that additional protein factors are needed. Further studies to identify other components in the peaks 1 and 2 of GLV IRES and lysate mixture (Fig. 7) are required for elucidating the mechanism of GLV IRES mediated translation initiation.




Figure 12. The lack of effect of binding of the three ITAFs on recruiting Giardia small ribosomal subunit. Radiolabeled GLV IRES alone (diamond), GLV IRES incubated with the small ribosomal subunit (triangle), GLV IRES incubated with the ITAFs GILBP1, GILa, and GISRp20 (square) or plus the small ribosomal subunit (star) were each fractionated in a 10–30% sucrose density gradient centrifugation. Each gradient was collected in fractions, and the radioactivity was counted.
doi:10.1371/journal.pone.0007435.g012

Materials and Methods

Construction of dicistronic vectors

Construction of plasmids pC631Rluc-Fluc, pC631Rluc-UTR-Fluc, pC631Rluc-Cod-Fluc, pC631Rluc-UTRCod-Fluc, pC631Rluc-114-367Cod-Fluc, pC631Rluc-126-367Cod-Fluc, pC631Rluc-176-367Cod-Fluc and pC631Rluc-278-367Cod-Fluc has been described previously [44]. For construction of pC631Rluc-EMCV-Fluc ~600 bp *XhoI/HindIII* fragment containing EMCV IRES was excised from pIRES (Clontech) and inserted into the *XhoI/HindIII* site upstream of the Fluc gene in pC631Rluc-Fluc. For pC631Rluc-CrPV-Fluc, CrPV IRES was amplified by PCR as a *XhoI/HindIII* fragment using CrPV1-1 (Eric Jan, University of British Columbia) as template and cloned into the *XhoI/HindIII* site of pC631Rluc-Fluc. Similar strategies were employed for construction of pC631Rluc-UTRCodΔ1-Fluc, pC631Rluc-UTRCodΔ2-Fluc and pC631Rluc-278CodΔ2-Fluc. The UTRCodΔ1, UTRCodΔ2 and 278-CodΔ2 sequences were amplified as *XhoI/HindIII* fragments and cloned into pC631Rluc-Fluc. To generate pC631Rluc-278CodΔ2rev-Fluc, a 278-CodΔ2 sequence was amplified as *HindIII/XhoI* fragment and was ligated with *XhoI/HindIII* digested pC631Rluc-Fluc.

For generation of capped transcripts, Renilla luciferase gene (*Rluc*) was PCR amplified using a pNull-Rluc plasmid (Promega) as template and the product was inserted downstream of T7 promoter using the *NheI* and *XhoI* sites in pIRES vector (Clontech) to generate pIRES-Rluc. A UTRCod-Fluc region was PCR amplified from pC631Rluc-UTRCod-Fluc as a *XhoI/XbaI* fragment, cloned into pGEM-T vector (Promega), excised as a *XhoI/NotI* fragment and inserted into the *XhoI/NotI* site located downstream of the Rluc gene in pIRES-Rluc. To generate transcripts containing a 50 polyA tail, annealed oligonucleotides 5' CTAGA₅₁G 3' and 5' AATTCT₅₁ 3' were inserted into the *XbaI/EcoRI* site downstream of the Fluc gene in the pIRES-Rluc-UTRCod-Fluc plasmid.

In vitro transcription

The dicistronic constructs in pC631 plasmid were linearized using *Nru* I and used as templates for *in vitro* synthesis of transcripts using MegaScript T7 transcription kit (Ambion). For synthesis of capped and polyadenylated transcripts, the dicistronic constructs in pIRES plasmids were linearized with *Eco*RI and used as templates in the mMessage mMachine T7 transcription kit (Ambion).

In vitro translation assays

Dicistronic transcripts were expressed in rabbit reticulocyte lysate (RRL) by using Flexi-Rabbit reticulocyte lysate system according to the manufacturer's instructions (Promega). Briefly, 0.5–1.0 μ g of the transcript was added to a 25 μ L reaction mixture and was incubated at 37°C for 90 min. Equal amounts of transcripts were used for each experiment. Reaction products (2.5 μ L) were assayed for Fluc and Rluc activities using Dual-luciferase reporter assay system (Promega). To test the effect of edeine on translation, different concentrations of the drug were added to the reaction mixture and incubated on ice for 5 minutes prior to the addition of the transcript.

Transfection of *Giardia* trophozoites and the luciferase assay

Transient transfection of *Giardia* trophozoites was carried out as previously described [43,45]. The cells were harvested after 5 or 16 hours of post-transfection and the cell lysate was assayed for Fluc and Rluc activities by using a Dual luciferase assay system (Promega) [44]. To test the effect of edeine on IRES activity, the drug was mixed with the *in vitro* synthesized transcripts and electroporated into *Giardia* cells. The concentrations of drug indicated in Figure 6, represent its concentration in the electroporation mix of 400 μ L.

Purification of 40S ribosomal subunits and sucrose gradient centrifugation

The 40S ribosomal subunits from RRL were prepared as described [16]. Polysomes from *Giardia* trophozoites were prepared [70] by gently lysing the cell by dounce homogenizer and pelleted at 78,000xg for 4 hours in a TLA 100.3 rotor (Beckman), and resuspended in buffer A (20 mM Tris-Cl, pH 7.5, 4 mM MgCl₂, 50 mM KCl, 2 mM DTT, complete protease inhibitor) to a final concentration of 100 OD₂₆₀/mL. Puromycin A was added to 1 mM and incubated on ice for 10 min followed by another 10 min at 37°C. To the treated polysomes, 2.5 M KCl was slowly added to 0.5 M and layered on to 10–30% sucrose gradient in buffer B (20 mM Tris-HCl, pH 7.5, 0.5 M KCl, 3 mM MgCl₂, 2 mM DTT) and centrifuged at 64,000xg in a SW 55 rotor at 4°C for 17 hrs. Fractions (0.2 mL) were collected and checked for rRNA by agarose gel electrophoresis. The fractions containing separated 40S and 60S subunits were pooled and concentrated using a centricon-30 concentrator and exchanged with buffer C (0.24 M sucrose, 20 mM Tris, pH 7.5, 10 mM KCl, 1 mM MgCl₂, 1 mM DTT, 0.1 mM EDTA).

The purified mammalian 40S subunits or *Giardia* small ribosomal subunits (150 nM) were incubated with 5' end labeled GLV IRES RNA (10–60 ng) in buffer E (20 mM Tris-HCl, pH 7.5, 100 mM KOAc, 2.5 mM MgOAc, 0.25 mM spermidine, 2 mM DTT) and incubated at room temperature for 30 min. The IRES-ribosomal complexes were then layered onto 10–30% linear sucrose gradient in buffer E and centrifuged at 200,000 x g in a SW41 rotor at 4°C for 2 hours and 10 min. The gradient was fractionated (500 μ L each) and the radioactivity was

counted. As control, labeled IRES RNA was incubated under similar conditions in the absence of mammalian 40S ribosomal subunits or *Giardia* small ribosomal subunit and was centrifuged through the gradient along with the test samples. To test the role of trans-acting protein factors, 4 μ g each of GIBP1, GILa and GISRp20 were pre-incubated with radiolabeled GLV IRES for 5 min at room temperature and then further incubated with purified *Giardia* small ribosomal subunit for additional 15 minutes. The reaction mixtures were then separated on a 10–30% sucrose gradient.

Preparation of cell extracts and analysis of IRES-protein complexes

Giardia cell lysate was prepared as described by Bergamini et al., [71]. Briefly, 200 mL of logarithmic *Giardia* trophozoites were harvested by centrifugation at 5,000xg at 4°C for 10 min and washed three times with phosphate buffered saline (PBS) and suspended in 200 μ L of lysis buffer (10 mM HEPES, pH 7.6, 10 mM KOAc, 0.5 mM MgOAc, 5 mM DTT, complete protease inhibitors minus EDTA). The cells were lysed by gentle sonication on ice for 5 minutes with 10-second pulses at 20-second intervals. The cell lysate was centrifuged at 10,000xg for 10 min. at 4°C and the cleared supernatant was used for the complex formation. The 10 μ L (1.25 OD at 260 nm) cleared supernatant was pre-incubated with 0.5 U/ μ L of RNasin (Promega) at 30°C for 10 minutes to inhibit ribonucleases and then incubated with 3 μ g of uniformly radiolabeled GLV IRES RNA (631 nts) in incubation buffer (16 mM HEPES, pH 7.6, 50 mM KOAc, 2.5 mM MgOAc, 0.1 mM spermidine, 1 mM DTT) [71] at 30°C for an additional 20 min in a final volume of 50 μ L. The reaction mixture was layered on top of a 10 mL 10–40% linear sucrose density gradient (in reaction buffer) and centrifuged at 78,000xg at 4°C in a SW41 rotor (Beckman) for 16 hours. The gradients were fractionated and the radioactivity in each fraction was counted using liquid scintillation counting.

Purification of the IRES-associated complexes by the StreptoTag method

Two complementary primers containing the aptamer sequence [56,57] were annealed and inserted into Hind III/*Eco*RI sites located at the 3' end of the IRES sequence in pC631 plasmid. The recombinant plasmid linearized by *Eco*RI was used as a template for *in vitro* transcription reactions. The tagged IRES RNA molecules were made using T7 megascript transcription kit (Ambion) in the presence of trace amounts of ³²P UTP. The radiolabeled RNA was purified using RNeasy purification kit (Qiagen) and the purified RNA was suspended in folding buffer (Tris-HCl, pH 7.5, NaCl 150 mM, 3 mM MgCl₂). To allow the proper folding of the IRES and the streptoTag aptamer, the hybrid RNA molecules were heated at 65°C for 5 min, followed by 10 min at 37°C and then cooled to room temperature.

The IRES-protein complexes were loaded onto the dihydro-streptomycin coupled Sepharose 6B column (1 mL bead volume) and washed with 10 column volumes of wash buffer (16 mM HEPES, pH 7.4, 50 mM KOAc, 2.5 mM MgCl₂, 6.8% sucrose) at room temperature [56]. The complexes were then eluted with five column volumes of wash buffer containing 10 μ M streptomycin. Since hybrid RNA molecules were radiolabeled, the entire purification process was monitored using a radioactive counter. The streptomycin-eluted fractions (1 mL each) with high radioactive counts were pooled and the purified complexes were pelleted at 100,000xg in a TLA100.3 rotor at 4°C for 12–16 hours. The pellets were suspended in wash buffer and analyzed by SDS-

PAGE. The protein bands were stained with Sypro Ruby (Invitrogen) and visualized by UV-light.

Protein identification by Mass Spectrometry (MALDI-TOF)

The Sypro Ruby stained protein bands were excised from the SDS-PAGE gels and subjected to in-gel trypsin digestion [49]. The digested peptides were cleaned with ZipTipC18 and analyzed with a matrix-assisted laser desorption ionization-time of flight (MALDI-TOF) mass spectrometry instrument (Voyager DE-STR mass spectrometer, Applied Biosystems). The resulting peptide masses were used to query the *Giardia* genome database using MS-FIT of the ProteinProspector program (<http://prospector.ucsf.edu/prospector/4.0.7/hm/msfit/hm>).

Cloning and purification of IRES binding proteins

The coding sequence for the IBP1 protein was amplified and cloned into expression vector pMAL-C2X (New England Biolabs) and was expressed as Maltose binding protein (MBP) fusion protein in *E. coli* cells. The expressed MBP-IBP1 fusion protein was purified by amylose-agarose chromatography following the manufacturer's instructions (New England Biolabs). The GISRp20 and GLIa coding sequences were cloned into expression vector pET28b (Qiagen), and expressed as 6XHis tagged proteins in *E. coli*. The expressed proteins were purified from the cell lysates using Ni-NTA agarose affinity chromatography (Qiagen).

Gel Shift assays

Approximately 100 ngs of 5' radiolabeled IRES RNA was mixed with varying amounts of purified protein in the binding buffer (20 mM Tris, pH 7.6, 50 mM KCl, 2.5 mM MgOAc, 0.05% NP40, 1 mM DTT and 1 µg/µL of yeast tRNA) and incubated at 30°C for 20 min. The IRES-protein complexes were separated in a composite agarose (0.5%): acrylamide (2.75%) gel (Jan, Kinzy and Sarnow, 2003) and visualized by phosphorimager (Amersham).

References

- Hellen CUT, Sarnow P (2001) Internal ribosome entry sites in eukaryotic mRNA molecules. *Genes and Dev* 15: 1593–1612.
- Stoneley M, Willis AE (2004) Cellular internal ribosome entry segments: structures, trans-acting factors and regulation of gene expression. *Oncogene* 23: 3200–3207.
- Baird SD, Turcotte M, Korneluk RG, Holcik M (2006) Searching for IRES. *RNA* 12: 1755–1785.
- Sasaki J, Nakashima N (1999) Translation initiation at the CUU codon is mediated by the internal ribosome entry site of an insect picorna-like virus *in vitro*. *J Virol* 73: 1219–1226.
- Wilson JE, Powell MJ, Hoover SE, Sarnow P (2000) Naturally occurring dicistronic cricket paralysis virus RNA is regulated by two internal ribosome entry sites. *Mol Cell Biol* 20: 4990–4999.
- Herbreteau CH, Weill L, Décimo D, Prévôt D, Darlix JL, et al. (2005) HIV-2 genomic RNA contains a novel type of IRES located downstream of its initiation codon. *Nat Struct Mol Biol* 12: 1001–1007.
- Cornelis S, Bruynooghe Y, Denecker G, Van Huffel S, Tinton S, et al. (2000) Identification and characterization of a novel cell cycle-regulated internal ribosome entry site. *Mol Cell* 5: 597–605.
- Pyronnet S, Pradayrol L, Sonenberg N (2000) A cell cycle dependent internal ribosome entry site. *Mol. Cell* 5: 607–616.
- Merrick WC, Hershey JWB (1996) The pathway and mechanism of eukaryotic protein synthesis. In: Hershey JWB, Mathews MB, Sonenberg N, eds. *Translational control*. Cold Spring Harbor: Cold Spring Harbor Laboratory Press. pp 31–69.
- Martinez-Salas E, Ramos R, Lafuente E, Lopez de Quinto S (2001) Functional interactions in internal translation initiation directed by viral and cellular IRES elements. *J Gen Virol* 82: 973–984.
- Martinez-Salas E, Pacheco A, Serrano P, Fernandez N (2008) New insights into internal ribosome entry site elements relevant from viral gene expression. *J Gen Virol* 89: 611–626.
- Pestova TV, Hellen CUT, Shatsky IN (1996) Canonical eukaryotic initiation factors determine initiation of translation by internal ribosome entry. *Mol Cell Biol* 16: 6859–6869.
- Pestova TV, Shatsky IN, Hellen CUT (1996) Functional dissection of eukaryotic initiation factor 4F: the 4A subunit and the central domain of the 4G subunit are

Supporting Information

Figure S1 The amino acid sequence of IBP1 identified by Mass spectrometry. The underlined sequences represent the peptides that were identified in mass spectrometry and used to identify the protein in the *Giardia* genome database.

Found at: doi:10.1371/journal.pone.0007435.s001 (0.09 MB TIF)

Figure S2 Amino acid sequence alignment of *Giardia* La protein with homologues from *Trypanosoma brucei*, *Drosophila melanogaster*, human and yeast. The La motif, RRM2 and RRM3 domains are marked with colored lines.

Found at: doi:10.1371/journal.pone.0007435.s002 (0.88 MB PDF)

Figure S3 Amino acid sequence alignment of *Giardia* SRp20 protein with homologues from human and mouse. The RRM motif is indicated by colored line.

Found at: doi:10.1371/journal.pone.0007435.s003 (0.26 MB PDF)

Figure S4 SDS-PAGE analysis of the purified 6xHis tagged GiLa (lane 2) and GISRp20 (lane 3) from *E. coli*.

Found at: doi:10.1371/journal.pone.0007435.s004 (0.07 MB PDF)

Acknowledgments

We would like to thank Dr. Eric Jan, University of British Columbia, for providing CrPV1-1 plasmid and edeine for our study and also for testing of GLV IRES binding to 40S subunit. We also thank Ashesh Saraiya and Wei Li for suggestions and critical reading of the manuscript.

Author Contributions

Conceived and designed the experiments: SG CCW. Performed the experiments: SG. Analyzed the data: SG CCW. Contributed reagents/materials/analysis tools: SG. Wrote the paper: SG CCW.

sufficient to mediate internal entry of 43S preinitiation complexes. *Mol Cell Biol* 16: 6870–6878.

- Kolupaeva VG, Lomakin IB, Pestova TV, Hellen CUT (2003) Eukaryotic initiation factors 4G and 4A mediate conformational changes downstream of the initiation codon of encephalomyocarditis virus internal ribosome entry site. *Mol Cell Biol* 23: 687–698.
- Pestova TV, Shatsky IN, Fletcher SP, Jackson RJ, Hellen CUT (1998) A prokaryotic-like mode of cytoplasmic eukaryotic ribosome binding to the initiation codon during internal translation initiation of hepatitis C and classical swine fever virus RNAs. *Genes and Dev* 12: 67–83.
- Kieft JS, Zhou K, Jubin R, Doudna JA (2001) Mechanism of ribosome recruitment by hepatitis C IRES RNA. *RNA* 7: 194–206.
- Ji H, Fraser CS, Yu Y, Leary J, Doudna JA (2004) Coordinated assembly of human translation initiation complexes by the hepatitis C virus internal ribosome entry site RNA. *Proc Natl Acad Sci USA* 101: 16990–16995.
- Jan E, Sarnow P (2002) Factorless ribosome assembly on the internal ribosome entry site of cricket paralysis virus. *J Mol Biol* 324: 889–902.
- Jan E, Kinzy TG, Sarnow P (2003) Divergent tRNA-like element supports initiation, elongation, and termination of protein biosynthesis. *Proc Natl Acad Sci USA* 100: 15410–15415.
- Jan E (2006) Divergent IRES elements in invertebrates. *Virus Res* 119: 16–28.
- Meerowitch K, Pelletier J, Sonenberg N (1989) A cellular protein that binds to the 5' noncoding region of poliovirus RNA: implications for internal translation initiation. *Genes Dev* 3: 1026–1034.
- Craig AW, Svitkin YV, Lee HS, Belsham GJ, Sonenberg N (1997) The La autoantigen contains a dimerization domain that is essential for enhancing translation. *Mol Cell Biol* 17: 163–169.
- Ali N, Siddiqui A (1997) The La autoantigen binds 5' noncoding region of the hepatitis C virus in the context of the initiator AUG codon and stimulates internal ribosome entry site-mediated translation. *Proc Natl Acad Sci USA* 94: 2249–2254.
- Costa-Mattioli M, Svitkin Y, Sonenberg N (2004) La autoantigen is necessary for optimal function of the Poliovirus and Hepatitis C virus internal ribosome entry site *in vivo* and *in vitro*. *Mol Cell Biol* 24: 6861–6870.
- Izumi RE, Das S, Barat B, Ray Chaudhuri S, Dasgupta A (2004) A peptide from autoantigen La blocks Poliovirus and Hepatitis C virus cap-independent

- translation and reveals a single tyrosine critical for La RNA binding and translation stimulation. *J Virol* 78: 3763–3776.
26. Kim YK, Jang SK (1999) La protein is required for efficient translation driven by encephalomyocarditis virus internal ribosomal entry site. *J Gen Virol* 80: 3159–3166.
 27. Ray PS, Das S (2002) La autoantigen is required for the internal ribosome entry site-mediated translation of the Cocksackievirus B3 RNA. *Nucl Acids Res* 30: 4500–4508.
 28. Borovojagin A, Petova T, Shatsky T (1994) Pyrimidine tract binding protein strongly stimulates in vitro encephalomyocarditis virus RNA translation at the level of pre-initiation complex formation. *FEBS Lett* 351: 299–302.
 29. Kaminski A, Hunt SL, Patton JG, Jackson RJ (1995) Direct evidence that polypyrimidine tract binding protein (PTB) is essential for internal initiation of translation of encephalomyocarditis virus RNA. *RNA* 1: 924–938.
 30. Kolupaeva VG, Hellen CV, Shatsky IN (1996) Structural analysis of the interaction of the pyrimidine tract binding protein with the internal ribosome entry site of encephalomyocarditis virus and foot and mouth disease virus RNAs. *RNA* 2: 1199–1212.
 31. Hellen CUT, Witherell GW, Schmid M, Shin SH, Pestova TV, et al. (1993) A cytoplasmic 57-kDa protein that is required for translation of picornavirus RNA by internal ribosome entry is identical to the nuclear pyrimidine-tract binding protein. *Proc Natl Acad Sci USA* 15: 7642–7646.
 32. Bedard KM, Daijogo S, Semler B (2007) A nucleocytoplasmic SR protein functions in viral IRES-mediated translation initiation. *EMBO J* 26: 459–467.
 33. Gamarnik AV, Andino R (1997) Two functional complexes formed by KH domain containing proteins with the 5' noncoding region of Poliovirus RNA. *RNA* 3: 882–892.
 34. Blyn LB, Swiderek KM, Richards O, Stahl DC, Semler BL, et al. (1996) Poly (rC) binding protein 2 binds to stem-loop IV of the Poliovirus RNA 5' noncoding region: identification by automated liquid chromatography-tandem mass spectrometry. *Proc Natl Acad Sci USA* 93: 11115–11120.
 35. Graff J, Cha J, Blyn LB, Ehrenfeld E (1998) Interaction of poly (rC) binding protein 2 with the 5' noncoding region of hepatitis A virus RNA and its effects on translation. *J Virol* 72: 9668–9675.
 36. Komar AA, Hatzoglou M (2005) Internal ribosome entry site in cellular mRNAs: mystery of their existence. *J Biol Chem* 280: 23425–23428.
 37. Semler BL, Waterman ML (2008) IRES-mediated pathways to polysomes: nuclear versus cytoplasmic routes. *Trends Microbiol* 16: 1–5.
 38. Wang AL, Wang CC (1991) Viruses of parasitic protozoa. *Parasitol Today* 7: 76–80.
 39. Wang AL, Wang CC (1986) Discovery of a specific double-stranded RNA virus in *Giardia lamblia*. *Mol Biochem Parasitol* 21: 269–276.
 40. Wang AL, Yang HM, Shen KA, Wang CC (1993) Giardavirus double-stranded RNA genome encodes a capsid polypeptide and a gag-pol-like fusion protein by a translation frameshift. *Proc Natl Acad Sci USA* 90: 8595–8599.
 41. Li L, Wang AL, Wang CC (2001) Structural analysis of the –1 ribosomal frameshift elements in giardavirus mRNA. *J Virol* 75: 10612–10622.
 42. Yu DC, Wang AL, Wu CH, Wang CC (1995) Virus-mediated expression of firefly luciferase in the parasitic protozoan *Giardia lamblia*. *Mol Cell Biol* 15: 4867–4872.
 43. Yu DC, Wang CC (1996) Identification of cis-acting signals in the giardavirus. (GLV) genome required for expression of firefly luciferase in *Giardia lamblia*. *RNA* 2: 824–834.
 44. Garlapati S, Wang CC (2004) Identification of a novel internal ribosome entry site in giardavirus that extends to both sides of the initiation codon. *J Biol Chem* 279: 3389–3397.
 45. Garlapati S, Chou J, Wang CC (2001) Specific secondary structures in capsid-coding region of Giardavirus transcript are required for its translation in *Giardia lamblia*. *J Mol Biol* 308: 623–638.
 46. Garlapati S, Wang CC (2002) Identification of an essential pseudoknot in the putative downstream internal ribosome entry site in giardavirus transcript. *RNA* 8: 601–611.
 47. Garlapati S, Wang CC (2005) Structural elements in the 5' untranslated region of Giardavirus transcript essential for internal ribosome entry site-mediated translation initiation. *Eukary Cell* 4: 742–754.
 48. Champney WS, Chittum HS, Samuels R (1992) Ribosomes from trichomonad protozoa have prokaryotic characteristics. *Int J Biochem* 24: 1125–1133.
 49. Li L, Wang CC (2004) Capped mRNA with a single nucleotide leader is optimally translated in a primitive eukaryote, *Giardia lamblia*. *J Biol Chem* 279: 14656–14664.
 50. Yu DC, Wang AL, Botka CW, Wang CC (1998) Protein synthesis in *Giardia lamblia* may involve interaction between a downstream box (DB) in mRNA and an anti-DB in the 16S-like ribosomal RNA. *Mol Biochem Parasitol* 96: 151–165.
 51. Thompson SR, Gulyas KD, Sarnow P (2001) Internal initiation in *Saccharomyces cerevisiae* mediated by an initiator tRNA/eIF2-independent internal ribosome entry site element. *Proc Natl Acad Sci USA* 98: 12972–12977.
 52. Lancaster AM, Jan E, Sarnow P (2006) Initiation factor-independent translation mediated by the hepatitis C virus internal ribosome entry site. *RNA* 12: 894–902.
 53. Kozak M, Shatkin AJ (1978) Migration of 40S ribosomal subunits on messenger RNA in the presence of edeine. *J Biol Chem* 253: 6568–6577.
 54. Odon OW, Kramer G, Henderson AB, Pinphanichakarn P, Hardesty B (1978) GTP hydrolysis during methionyl-tRNA^f binding to 40S ribosomal subunits and the site of edeine inhibition. *J Biol Chem* 253: 1807–1816.
 55. Wilson J, Pestova TV, Hellen CUT, Sarnow P (2000) Initiation of protein synthesis from the A-site of the ribosome. *Cell* 102: 511–520.
 56. Bachler M, Schroeder R, von Ashen U (1999) StreptoTag: a novel method for the isolation of RNA-binding proteins. *RNA* 5: 1509–1516.
 57. Locker N, Easton LE, Lukavsky PJ (2006) Affinity purification of eukaryotic 48S initiation complexes. *RNA* 12: 683–690.
 58. Locker N, Lukavsky PJ (2007) A practical approach to isolate 48S complexes: affinity purification and analyses. *Methods Enzymol* 429: 83–104.
 59. Dinos G, Wilson DN, Teraoka Y, Szafarski W, Fucini P, et al. (2004) Dissecting the ribosomal inhibition mechanisms of edeine and pactamycin: the universally conserved residues G693 and C795 regulate P-site RNA binding. *Mol Cell* 13: 113–124.
 60. Spahn CM, Jan E, Mulder A, Grassucci RA, Sarnow P, Frank J (2004) Cryo-EM visualization of a viral internal ribosome entry site bound to human ribosomes: the IRES functions as an RNA-based translation factor. *Cell* 118: 465–475.
 61. Pflugstein JS, Constantino DA, Kieft JS (2006) Structural basis for ribosome recruitment and manipulation by a viral IRES RNA. *Science* 314: 1450–1454.
 62. Lytle JB, Wu L, Robertson HD (2001) The ribosome binding site of Hepatitis C virus mRNA. *J Virol* 75: 7629–7636.
 63. Spahn CM, Kieft JS, Grassucci RA, Penczek PA, Zhou K, et al. (2001) Hepatitis C virus IRES RNA-induced changes in the conformation of the 40S ribosomal subunit. *Science* 291: 1959–1962.
 64. Siridechadilok B, Fraser CS, Hall RJ, Doudna JA, Nogales E (2005) Structural roles for human translation factor eIF3 in initiation of protein synthesis. *Science* 310: 1513–1515.
 65. Kieft JS, Zhou K, Jubin R, Murray MG, Lau JY, et al. (1999) The hepatitis C virus internal ribosome entry site adopts an ion-dependent tertiary fold. *J Mol Biol* 292: 513–529.
 66. Constantino D, Kieft JS (2005) A preformed compact ribosome-binding domain in the cricket paralysis-like virus IRES RNAs. *RNA* 11: 332–343.
 67. Belsham GJ, Sonenberg N (2000) Picornavirus RNA translation: roles for cellular proteins. *Trends Microbiol* 8: 330–335.
 68. Bousquet-Antonelli C, Deragon JM (2009) A comprehensive analysis of the Lamotif protein superfamily. *RNA* 15: 750–764.
 69. Sanford JR, Gray NK, Beckmann K, Caceres JF (2004) A novel role for shuttling SR proteins in mRNA translation. *Genes Dev* 18: 755–768.
 70. Shirakura T, Maki Y, Hideji Y, Nobuko A, Wada A, et al. (2001) Characterization of the ribosomal proteins of the amitochondriate protist, *Giardia lamblia*. *Mol Biochem Parasitol* 112: 153–156.
 71. Bergamini G, Preiss T, Hentze MW (2000) Picornavirus IRESes and the poly(A) tail jointly promote cap-independent translation in a mammalian cell-free system. *RNA* 6: 1781–1790.