

UC Davis

UC Davis Previously Published Works

Title

Functional informed genome-wide interaction analysis of body mass index, diabetes and colorectal cancer risk

Permalink

<https://escholarship.org/uc/item/86q5m86b>

Journal

Cancer Medicine, 9(10)

ISSN

2045-7634

Authors

Xia, Zhiyu

Su, Yu-Ru

Petersen, Paneen

et al.

Publication Date

2020-05-01

DOI

10.1002/cam4.2971





Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

ORIGINAL RESEARCH

Functional informed genome-wide interaction analysis of body mass index, diabetes and colorectal cancer risk

Zhiyu Xia¹  | Yu-Ru Su² | Paneen Petersen² | Lihong Qi³ | Andre E. Kim⁴  | Jane C. Figueiredo^{4,5} | Yi Lin² | Hongmei Nan⁶ | Lori C. Sakoda^{2,7} | Demetrius Albanes⁸ | Sonja I. Berndt⁸ | Stéphane Bézieau⁹ | Stephanie Bien² | Daniel D. Buchanan^{10,11,12}  | Graham Casey¹³ | Andrew T. Chan^{14,15,16,17,18,19} | David V. Conti⁴ | David A. Drew^{14,16} | Steven J. Gallinger²⁰ | W. James Gauderman⁴ | Graham G. Giles^{21,22} | Stephen B. Gruber⁴ | Marc J. Gunter²³ | Michael Hoffmeister²⁴  | Mark A. Jenkins²² | Amit D. Joshi^{16,18} | Loic Le Marchand²⁵ | Juan P. Lewinger⁴ | Li Li²⁶ | Noralane M. Lindor²⁷ | Victor Moreno^{28,29,30,31} | Neil Murphy²³ | Rami Nassir³² | Polly A. Newcomb^{1,2} | Shuji Ogino^{18,33,34} | Gad Rennert^{35,36,37} | Mingyang Song^{14,16,38} | Xiaoliang Wang² | Alicja Wolk³⁹ | Michael O. Woods⁴⁰ | Hermann Brenner^{24,41} | Emily White^{1,2} | Martha L. Slattery⁴² | Edward L. Giovannucci^{15,18,38} | Jenny Chang-Claude^{43,44} | Paul D. P. Pharoah⁴⁵ | Li Hsu^{2,46} | Peter T. Campbell⁴⁷ | Ulrike Peters^{1,2}

¹Department of Epidemiology, University of Washington, Seattle, WA, USA

²Public Health Sciences, Fred Hutchinson Cancer Research Center, Seattle, WA, USA

³Department of Public Health Sciences, University of California Davis, Davis, CA, USA

⁴Department of Preventive Medicine &, USC Norris Comprehensive Cancer Center, Keck School of Medicine, University of Southern California, Los Angeles, CA, USA

⁵Department of Medicine, Samuel Oschin Comprehensive Cancer Institute, Cedars-Sinai Medical Center, Los Angeles, CA, USA

⁶Department of Epidemiology, Richard M. Fairbanks School of Public Health, Indiana University, Indianapolis, IN, USA

⁷Division of Research, Kaiser Permanente Northern California, Oakland, CA, USA

⁸Division of Cancer Epidemiology and Genetics, National Cancer Institute, National Institutes of Health, Bethesda, MD, USA

⁹Service de Génétique Médicale, Centre Hospitalier Universitaire (CHU) Nantes, Nantes, France

¹⁰Colorectal Oncogenomics Group, Department of Clinical Pathology, The University of Melbourne, Parkville, Victoria, Australia

¹¹University of Melbourne Centre for Cancer Research, Victorian Comprehensive Cancer Centre, Parkville, Victoria, Australia

¹²Genetic Medicine and Family Cancer Clinic, The Royal Melbourne Hospital, Parkville, Victoria, Australia

¹³Center for Public Health Genomics, University of Virginia, Charlottesville, VA, USA

¹⁴Division of Gastroenterology, Massachusetts General Hospital, Harvard Medical School, Boston, MA, USA

¹⁵Channing Division of Network Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA

¹⁶Clinical and Translational Epidemiology Unit, Massachusetts General Hospital, Harvard Medical School, Boston, MA, USA

¹⁷Broad Institute of Harvard and MIT, Cambridge, MA, USA

¹⁸Department of Epidemiology, Harvard T.H. Chan School of Public Health, Harvard University, Boston, MA, USA

¹⁹Department of Immunology and Infectious Diseases, Harvard T.H. Chan School of Public Health, Harvard University, Boston, MA, USA

²⁰Lunenfeld Tanenbaum Research Institute, Mount Sinai Hospital, University of Toronto, Toronto, Ontario, Canada

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2020 The Authors. *Cancer Medicine* published by John Wiley & Sons Ltd.

²¹Cancer Epidemiology Division, Melbourne, Victoria, Australia

²²Centre for Epidemiology and Biostatistics, Melbourne School of Population and Global Health, The University of Melbourne, Melbourne, Victoria, Australia

²³Section of Nutrition and Metabolism, International Agency for Research on Cancer, Lyon, France

²⁴Division of Clinical Epidemiology and Aging Research, German Cancer Research Center (DKFZ), Heidelberg, Germany

²⁵University of Hawaii Cancer Center, Honolulu, HI, USA

²⁶Department of Family Medicine, University of Virginia, Charlottesville, VA, USA

²⁷Department of Health Science Research, Mayo Clinic, Scottsdale, AZ, USA

²⁸Oncology Data Analytics Program, Catalan Institute of Oncology-IDIBELL, L'Hospitalet de Llobregat, Barcelona, Spain

²⁹CIBER Epidemiología y Salud Pública (CIBERESP), Madrid, Spain

³⁰Department of Clinical Sciences, Faculty of Medicine, University of Barcelona, Barcelona, Spain

³¹ONCOBEL Program, Bellvitge Biomedical Research Institute (IDIBELL), L'Hospitalet de Llobregat, Barcelona, Spain

³²Department of Pathology, School of Medicine, Umm Al-Qura University, Makkah, Saudi Arabia

³³Department of Pathology, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA

³⁴Department of Oncologic Pathology, Dana-Farber Cancer Institute, Boston, MA, USA

³⁵Department of Community Medicine and Epidemiology, Lady Davis Carmel Medical Center, Haifa, Israel

³⁶Ruth and Bruce Rappaport Faculty of Medicine, Technion-Israel Institute of Technology, Haifa, Israel

³⁷Clalit National Cancer Control Center, Haifa, Israel

³⁸Department of Nutrition, Harvard T.H. Chan School of Public Health, Harvard University, Boston, MA, USA

³⁹Institute of Environmental Medicine, Karolinska Institutet, Stockholm, Sweden

⁴⁰Memorial University of Newfoundland, Discipline of Genetics, St. John's, Canada

⁴¹Division of Preventive Oncology, German Cancer Research Center (DKFZ), National Center for Tumor Diseases (NCT), Heidelberg, Germany

⁴²Department of Internal Medicine, University of Utah, Salt Lake City, UT, USA

⁴³Division of Cancer Epidemiology, German Cancer Research Center (DKFZ), Heidelberg, Germany

⁴⁴University Medical Centre Hamburg-Eppendorf, University Cancer Centre Hamburg (UCCH), Hamburg, Germany

⁴⁵Department of Public Health and Primary Care, University of Cambridge, Cambridge, UK

⁴⁶Department of Biostatistics, University of Washington, Seattle, WA, USA

⁴⁷Behavioral and Epidemiology Research Group, American Cancer Society, Atlanta, GA, USA

Correspondence

Ulrike Peters, Department of Epidemiology, University of Washington, Seattle, WA, USA.
Email: upeters@fredhutch.org

Abstract

Background: Body mass index (BMI) and diabetes are established risk factors for colorectal cancer (CRC), likely through perturbations in metabolic traits (e.g. insulin resistance and glucose homeostasis). Identification of interactions between variation in genes and these metabolic risk factors may identify novel biologic insights into CRC etiology.

Methods: To improve statistical power and interpretation for gene-environment interaction ($G \times E$) testing, we tested genetic variants that regulate expression of a gene together for interaction with BMI (kg/m^2) and diabetes on CRC risk among 26 017 cases and 20 692 controls. Each variant was weighted based on PrediXcan analysis of gene expression data from colon tissue generated in the Genotype-Tissue Expression Project for all genes with heritability $\geq 1\%$. We used a mixed-effects model to jointly measure the $G \times E$ interaction in a gene by partitioning the interactions into the predicted gene expression levels (fixed effects), and residual $G \times E$ effects (random effects). $G \times \text{BMI}$ analyses were stratified by sex as BMI-CRC associations differ by sex. We used false discovery rates to account for multiple comparisons and reported all results with $\text{FDR} < 0.2$.

Results: Among 4839 genes tested, genetically predicted expressions of *FOXA1* ($P = 3.15 \times 10^{-5}$), *PSMC5* ($P = 4.51 \times 10^{-4}$) and *CD33* ($P = 2.71 \times 10^{-4}$) modified the association of BMI on CRC risk for men; *KIAA0753* ($P = 2.29 \times 10^{-5}$) and *SCN1B* ($P = 2.76 \times 10^{-4}$) modified the association of BMI on CRC risk for women; and *PTPN2* modified the association between diabetes and CRC risk in both sexes ($P = 2.31 \times 10^{-5}$).

Conclusions: Aggregating G \times E interactions and incorporating functional information, we discovered novel genes that may interact with BMI and diabetes on CRC risk.

KEYWORDS

BMI, colorectal cancer, diabetes, gene expression, gene-environmental interaction

1 | BACKGROUND

Colorectal cancer (CRC) is a major source of cancer morbidity and mortality worldwide. According to the World Health Organization (WHO), CRC is the third most common cancer

worldwide and accounts for approximately 10% of global cancer incidence and mortality (<http://globocan.iarc.fr/>). Genetic factors play an important role in the etiology of both familial and sporadic CRC.¹ CRC is a complex, multifactorial disease with many genetic and modifiable lifestyle factors including

Funding information

National Cancer Institute, National Institutes of Health, US Department of Health and Human Services (U01 CA164930, U01 CA137088, R01 CA059045, R01201407); Center for Inherited Disease Research (CIDR) (X01-HG008596 and X-01-HG007585); NIH/NCI Cancer Center Support Grant P30 CA015704; Hospital Clinical Research Program (PHRC-BRD09/C) from the University Hospital Center of Nantes (CHU de Nantes); Regional Council of Pays de la Loire, the Groupement des Entreprises Françaises dans la Lutte contre le Cancer (GEFLUC), the Association Anne de Bretagne Génétique and the Ligue Régionale Contre le Cancer (LRCC). Intramural Research Program of the US National Cancer Institute, National Institutes of Health, and by US Public Health Service contract HHSN261201500005C from the National Cancer Institute, Department of Health and Human Services. COLO2&3: National Institutes of Health (R01 CA60987); National Cancer Institute (NCI), National Institutes of Health (NIH) (grant numbers U01 CA122839, R01 CA143247); NCI/NIH (grant number U01 CA167551); Australasian Colorectal Cancer Family Registry (grant numbers U01 CA074778 and U01/U24 CA097735); USC Consortium Colorectal Cancer Family Registry (grant numbers U01/U24 CA074799); Mayo Clinic Cooperative Family Registry for Colon Cancer Studies (grant number U01/U24 CA074800), Ontario Familial Colorectal Cancer Registry (grant number U01/U24 CA074783), Seattle Colorectal Cancer Family Registry (grant number U01/U24 CA074794); University of Hawaii Colorectal Cancer Family Registry (grant number U01/U24 CA074806; National Cancer Institute, National Institutes of Health (NCI/NIH), US Department of Health and Human Services (grant numbers U19 CA148107, R01 CA81488, P30 CA014089, R01 CA197350; P01 CA196569; R01 CA201407); National Institutes of Environmental Health Sciences, National Institutes of Health (grant number T32 ES013678); The American Cancer Society; German Research Council (BR 1704/6-1, BR 1704/6-3, BR 1704/6-4, CH 117/1-1, HO 5117/2-1, HE 5998/2-1, KL 2354/3-1, RO 2270/8-1 and BR 1704/17-1); Interdisciplinary Research Program of the National Center for Tumor Diseases (NCT), Germany; German Federal Ministry of Education and Research (01KH0404, 01ER0814, 01ER0815, 01ER1505A and 01ER1505B); National Institutes of Health (R01 CA48998 to M. L. Slattery); National Institutes of Health (P01 CA055075, U01 CA167552, U01 CA167552, R01 CA137178, R01 CA151993, and R35 CA197735); National Institutes of Health (R01 CA137178, P01 CA087969, U01 CA186107, R01 CA151993, and R35 CA197735) and PHS by the National Institutes of Health (R01 CA042182); Clinical Investigator Award from Damon Runyon Cancer Research Foundation (CI-8); NCI R01CA136726); VicHealth and Cancer Council Victoria; Australian NHMRC grants 509348, 209057, 251553 and 504711; Cancer Council Victoria; National Institutes of Health (R37 CA54281, P01 CA033619, and R01 CA063464); National Institutes of Health, US Department of Health and Human Services (R01 CA81488; Interdisciplinary Health Research Team award from the Canadian Institutes of Health Research (CRT 43821); the National Institutes of Health, US Department of Health and Human Services (U01 CA74783); National Cancer Institute of Canada grants (18223 and 18226); Canadian Cancer Society Research Institute; National Institutes of Health, through funding allocated to the Ontario Registry for Studies of Familial Colorectal Cancer (U01 CA074783); Ontario Research Fund, the Canadian Institutes of Health Research, and the Ontario Institute for Cancer Research, through generous support from the Ontario Ministry of Research and Innovation; Intramural Research Program of the Division of Cancer Epidemiology and Genetics and supported by contracts from the Division of Cancer Prevention, National Cancer Institute, NIH, DHHS; National Institutes of Health (NIH), Genes, Environment and Health Initiative (GEI) Z01 CP 010200, NIH U01 HG004446, and NIH GEI U01 HG 004438; NHS in the East of England through the Clinical Academic Reserve. Cancer Research UK (C490/A16561); the UK National Institute for Health Research Biomedical Research Centres at the University of Cambridge; Swedish Research Council/Infrastructure grant, the Swedish Cancer Foundation, and the Karolinska Institute's Distinguished Professor Award; National Institutes of Health (K05 CA154337); National Heart, Lung, and Blood Institute, National Institutes of Health, US Department of Health and Human Services through contracts HHSN268201100046C, HHSN268201100001C, HHSN268201100002C, HHSN268201100003C, HHSN268201100004C, and HHSN271201100004C; Colorectal Cancer Genetics & Genomics; Instituto de Salud Carlos III, co-funded by FEDER funds –a way to build Europe– (grants PI14-613 and PI09-1286); Agency for Management of University and Research Grants (AGAUR) of the Catalan Government (grant 2017SGR723); Junta de Castilla y León (grant LE22A10-2); Xarxa de Bancs de Tumors de Catalunya sponsored by Pla Director d'Oncologia de Catalunya (XBTC); Plataforma Biobancos PT13/0010/0013 and ICObIOBANC, sponsored by the Catalan Institute of Oncology; National Institutes of Health (R01 CA076366 and U01 CA074794).

diet,² obesity,³ physical activity,⁴ and diabetes⁵ among others contributing to its etiology.

Obesity, compared to normal weight, is associated with greater risk of CRC⁶ in a dose-response manner.³ According to a recent World Health Organization report, 39% of adults (1.9 billion) aged 18 years and older were overweight, and 13% (650 million) were obese (<https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight>). With a growing obesity epidemic, the number of people with diabetes has also dramatically increased from 108 million in 1980 to around 500 million in 2018 worldwide (<https://www.who.int/news-room/fact-sheets/detail/diabetes>). These trends are expected to continue, and will likely continue to contribute to the burden imposed by CRC in the coming decades.⁷ Sex has been found to modify the association between obesity and CRC risk: the risk of developing CRC is often higher in obese men compared to obese women.⁸ These sex differences may be contributed to differences in sex hormones whereby estrogens, in particular, derived from adipose tissue offset the risk otherwise mitigated by obesity more so for women than men.⁹ Obesity and diabetes are interrelated risk factors for CRC that may impact CRC risk through metabolic abnormalities including pathways related to inflammation, insulin, and glucose homeostasis.¹⁰ Findings from meta-analyses have indicated that diabetes is associated with an approximately 30% increased relative risk of developing colorectal cancer compared to nondiabetes, after adjusting for BMI.⁵

It has been estimated that genetic variants explain up to 35% of the heritability in CRC risk.¹¹ To date, genome-wide association studies (GWAS) have identified more than 100 independent common genetic variants that are robustly associated with CRC.¹² However, these variants only explain a fraction of total heritability. Given the complexity of CRC etiology, it can be expected that a closer investigation of gene-environment (G × E) interactions may help identify additional novel loci and biological interactions that give insight to the pathogenesis of CRC. A few studies have investigated G × E interactions with BMI and diabetes for CRC risk, mainly focusing on the single nucleotide polymorphisms (SNPs) that had been previously identified by GWAS.^{13,14} The candidate G × E analysis by Sainz et al indicated that SNPs in *IGF2BP2* (rs4402960) and *PPARγ* (rs1801282) may interact with diabetes on CRC risk.¹⁵ As statistical power remains a major concern for G × E analysis, conducting set-based G × E testing and incorporating functional genomic information may improve power and help to interpret the underlying biology.

In this study, we conducted a novel set-based genome-wide approach to test interactions between genetic predicted gene expression and BMI and diabetes with CRC risk. We applied MiSTi, a set-based G × E testing framework which allows for incorporation of functional information.¹⁶

2 | METHODS

2.1 | Study participants

We used epidemiological and genetic data of 26 017 incident CRC cases and 20 692 controls from 33 participating studies in three international CRC consortia: the Genetics and Epidemiology of Colorectal Cancer Consortium (GECCO), the Colorectal Transdisciplinary Study (CORECT) and the Colon Cancer Family Registry (CCFR). Details have been published previously.^{13,17} Participants with non-European ancestry were excluded because of small sample sizes. All studies were approved by their respective Institutional Review Boards.

2.2 | Genotype data

Details on genotyping and imputation have been reported previously.^{17,18} In brief, DNA was mostly obtained from blood/buccal samples. Several platforms (the Illumina HumanHap 300 k, 240 k, 550 k and OncoArray 610 k BeadChip Array system, or Affymetrix platform) were used for genotyping.¹⁹ Samples were excluded based on sample call rates $\leq 97\%$, heterozygosity, unexpected duplicates or relative pairs, gender discrepancy and principal component analysis (PCA) outlier of HapMap2 CEU cluster. SNPs were excluded on the basis of inconsistency across platforms, call rate $< 98\%$, and out of Hardy-Weinberg equilibrium (HWE) in controls ($P < .0001$).¹⁹ SNPs were imputed to the Haplotype Reference Consortium (HRC version r1.0),²⁰ and restricted by imputation accuracy ($R^2 > 0.3$ for SNPs with MAF $> 1\%$, $R^2 > 0.5$ for SNPs with MAF $> 0.5\%$ and $< 1\%$, and $R^2 > 0.99$ for SNPs with MAF $< 0.05\%$).

2.3 | Estimation of gene expression levels

Functional information was generated from PrediXcan,²¹ which used an elastic net penalized regression to select eQTLs that jointly predict gene expression levels based on genome-wide genotypes and transcriptome data from 169 colon tissue samples from the GTEx project (GTEx v6). While GTEx measured gene expression at two different locations in the colon (sigmoid and transverse), we restricted analysis to the transverse. The transverse colon samples were done on the entire colonic wall while the analysis of the sigmoid colon was restricted to the muscularis tissue, which is less relevant for CRC development. The heritability of gene expression levels explained by the SNPs (predictive R^2) were calculated using a mixed-effects model.^{21,22} The estimated weights and predictive R^2 for eQTL sets associated with individual genes were downloaded from the publicly available PredictDB

Repository (<http://hakyimlab.org/predictdb/>). Genes with $R^2 \geq 0.01$ were selected for interaction analyses. A total of 4839 genes were included.

2.4 | Exposure assessment

Demographics and environmental exposures were self-reported at either in-person interview or via structured self-administered questionnaires, based on each participating study. A multistep, iterative data harmonization procedure was applied, reconciling each study's unique protocols and data collection instruments.^{13,23} Numerous quality-control checks were performed, and outlying values of variables were truncated to the minimum or maximum value of an established range for each variable. Variables were combined into a single dataset with common definition, standardized coding, and standardized permissible values. For the main exposure variables (BMI and diabetes), continuous measurement of BMI (per 5 kg/m², excluded participants below 18.5 kg/m²) as well as a binary self-reported diagnosis of diabetes were used.

2.5 | Statistical analysis

Individual level genotyping and environmental data were used for statistical analysis. We used MiSTi,¹⁶ a set-based statistical framework providing mixed effects score tests for $G \times E$ interaction, to identify genes with eQTLs that interact with BMI or diabetes on CRC risk. MiSTi models the $G \times E$ interaction effects with two components, the fixed- and random-effect components. The fixed-effect component incorporates the weights from PrediXcan to calculate the genetically predicted gene expression levels for samples in our data, and then assesses the interaction between the predicted expression of each gene and BMI as well as diabetes. The random-effects component quantifies residual interaction effects that are not accounted for by the fixed-effect component. To combine the fixed and random effects components, we used the data-adaptive weighted combination approach under MiSTi (aMiSTi).¹⁶ All BMI analyses were stratified by sex as BMI-CRC associations differ by sex. As associations for diabetes and CRC have been reported to be similar for men and women, we analyzed diabetes-by-SNP associations in men and women combined and included sex as a covariable. Genes with false discovery rate (FDR) <0.2 were considered statistically significant.

For $G \times E$ interactions discovered in our analysis, we conducted secondary analyses with multivariable-adjusted generalized linear regression models to assess main effects and interactions between individual eQTL variants and BMI/diabetes on CRC risk. A sequential analysis was conducted where we started with the most significant SNP with the

$G \times E$ interaction, then took the next significant one while adjusting for the first one, and so forth until the SNP's P -value was greater than .05. Age, study and PCs were adjusted for BMI models, and sex was additionally included in the sequential analysis for diabetes-by-SNP associations. eQTL variants which drive the significant interaction effects were identified and reported. All statistical analyses were performed using R (version 3.4.4).

3 | RESULTS

The demographic characteristics and exposures of interest are summarized in Table S1. Compared to controls, cases had a higher BMI (27.4 vs 26.7 kg/m²) and a higher prevalence of diabetes (13.5% vs 10.7%). The main effects of BMI and diabetes on CRC risk for each study individually and combined are summarized in Figures S1A, S1B, and S2, respectively. Among men, each 5 kg/m² increase in BMI was associated with 24% higher risk of CRC (fixed effect OR = 1.24, 95% CI 1.19-1.29, P -value < .01); among women, each 5 kg/m² increase in BMI was associated with 12% higher risk of CRC (fixed effect OR = 1.12, 95% CI 1.09-1.15, P -value < .01). In addition, diabetes was associated with a 22% higher risk of CRC (fixed effect OR = 1.22, 95% CI 1.14-1.30, P -value < .01) compared to those without diabetes.

Among the 4839 genes tested, we observed interactions between genetically predicted expression and BMI for CRC risk at FDR <0.2 for three genes among men, and two genes among women (Table 1; Figure S3A and S3B). All genetic signals in the five genes were detected by the random-effect components (Table 1). Among men, the most significant gene was *FOXA1* ($p_{\text{interaction}} = 3.15 \times 10^{-5}$) located on 14q21.1. A total of 43 eQTLs were included in this gene. The second most significant gene was *CD33* ($p_{\text{interaction}} = 2.71 \times 10^{-4}$, located on 19q13.3) with 30 eQTL included in the interaction test for this gene. Another gene, *PSMC5*, located on 17q23.3 also surpassed the FDR threshold and interacted with BMI among men ($p_{\text{interaction}} = 4.51 \times 10^{-4}$), with 32 eQTL tested. Among women, *KIAA0753* ($p_{\text{interaction}} = 2.29 \times 10^{-5}$, located on 17p13.1) and *SCN1B* ($p_{\text{interaction}} = 2.76 \times 10^{-4}$, located on 19q13.11) showed interactions with BMI on CRC risk. There were 30 and 11 eQTLs included in the interaction testing of these two genes, respectively.

When studying interactions with diabetes, we observed that the eQTLs of *PTPN2* located at 18p11.21 modified the association between diabetes and CRC risk ($p_{\text{interaction}} = 2.31 \times 10^{-5}$) (Table 1; Figure S4). The interaction signal was mainly from the interaction of genetically predicted gene expression with diabetes ($p_{\text{interaction}} = 1.03 \times 10^{-5}$), detected by the fixed-effect components. A total of 95 eQTLs predicted the expression level of this gene.

TABLE 1 Results for interactions of eQTLs with BMI and diabetes and risk of colorectal cancer with a FDR <0.2

Gene name	CHR	# of SNPs	^a R ²	P-values		
				Fixed-effect component	Random-effect component	Adaptive weight
BMI						
Male						
<i>FOXA1</i>	14q21.1	43	1.90%	0.125	1.03 × 10 ⁻⁵	3.15 × 10 ⁻⁵
<i>CD33</i>	19q13.3	30	1.87%	0.963	1.28 × 10 ⁻⁴	2.71 × 10 ⁻⁴
<i>PSMC5</i>	17q23.3	32	10.50%	0.719	9.96 × 10 ⁻⁵	4.51 × 10 ⁻⁴
Female						
<i>KIAA0753</i>	17p13.1	30	14.49%	0.029	5.91 × 10 ⁻⁵	2.29 × 10 ⁻⁵
<i>SCN1B</i>	19q13.11	11	2.27%	0.426	6.33 × 10 ⁻⁵	2.76 × 10 ⁻⁴
Diabetes						
<i>PTPN2</i>	18p11.21	95	7.87%	1.03 × 10 ⁻⁵	0.605	2.31 × 10 ⁻⁵

^aR² is the heritability in gene expression.

In secondary analysis we assessed main effects and interactions of individual variants with BMI or diabetes on CRC risk for each gene set that showed FDR <0.2. The results suggested that associations observed for *FOXA1*, *CD33*, *PSMC5*, and *PTPN2* might be largely driven by some specific individual variants (Table S2). In Tables S3A and S3B, we demonstrated the associations of the predicted gene expression on CRC risk stratified by BMI at quartiles and diabetes.

4 | DISCUSSION

In this large genome-wide investigation using transcription data to inform G × E interaction testing, we observed suggested interactions between genetic predicted gene expression and BMI and diabetes for risk of CRC. The strongest association was for *FOXA1* and BMI with CRC risk for men, and the predicted gene expression level of *PTPN2* interacted with diabetes on CRC risk among men and women combined. Our findings also suggested that the gene expression levels of *CD33* and *PSMC5* may interact with BMI on CRC risk among men, and the gene expression levels of *KIAA0753* and *SCN1B* may interact with BMI on CRC risk among women. The identified genes are novel in modifying BMI-CRC and diabetes-CRC associations.

Our most significant result in the BMI interaction analysis was for the *FOXA1* (Forkhead Box A1) gene-BMI interaction for CRC risk, among men. *FOXA1* is a transcription factor that belongs to the *FOX* gene superfamily,²⁴ which is responsible for various biological processes, including cell proliferation, apoptosis and differentiation.²⁵ Several studies have found that *FOXA1* expression in cancer tissues was associated with multiple types of human cancers,^{26,27} reflecting its crucial roles in cellular processes. According to Sahu et al,²⁷ besides pioneering the androgen receptor (AR)

pathway, *FOXA1* depletion elicited extensive redistribution of AR-binding sites on LNCaP-1F5 cell chromatin that was commensurate with changes in androgen-dependent gene expression signature. They also found that the role of *FOXA1* in androgen signaling is distinctly different from that in estrogen signaling, providing evidence to the results that *FOXA1* was associated with CRC and interacted with BMI only among men, but not women. A recent study detected the expression of *FOXA1* in samples of CRC tissues and matched noncancerous tissues using immunohistochemistry to determine the clinical significance of *FOXA1* and its role in CRC.²⁸ Their research demonstrated that the *FOXA1* expression level in cancer tissues was significantly higher among CRC cases compared to noncancer specimens, and positive expression of *FOXA1* in cancer tissues was associated with poor clinicopathological characteristics as well as poor prognosis of CRC.²⁸ Though many existing studies indicated the strong associations between *FOXA1* expression in cancer tissues and human cancers, so far none of them focused on interactions between *FOXA1* and BMI on cancers. In our genome-wide G × BMI interaction scans, we demonstrated that genetic variants in *FOXA1* interacted with BMI among men. Further exploration suggested that multiple genetic variants in the tested gene-set contributed to the *FOXA1*-BMI interaction effect. This interaction may be explained by the observation that *FOXA1* binds to four distinct intronic regions of the *FTO* (fat mass and obesity associated) gene,²⁹ which has a known predisposing role to obesity.³⁰ However, functional follow up analysis are needed to shed further light on this interaction effect of this gene. Overall, previous literature provides strong support for a potential role of *FOXA1* in CRC which may be mediated through the *FTO* gene that could explain the observed interaction with obesity.

We identified an interaction between *PTPN2* (protein tyrosine phosphatase nonreceptor-type 2) and diabetes with

CRC risk. *PTPN2* gene encodes the T-cell-specific protein tyrosine phosphatase, and functions as a negative regulator of inflammation by inhibiting the transcription factor *STAT1* in the *IFN- γ* signaling pathway.³¹ Previous studies demonstrated that several variants located in *PTPN2* were significantly associated with inflammatory bowel disease,³² celiac disease,³³ rheumatoid arthritis³⁴ and diabetes.³⁵ A recent study identified the dual role for *PTPN2* in directly regulating inflammasome activation and IL-1 β production to suppress pro-inflammatory responses during colitis but promote intestinal tumor development.³⁶ Since diabetes is also an inflammation-related disorder,³⁷ it might suggest a mechanism on how *PTPN2* interacts with diabetes on CRC risk. *PTPN2* was also found to be associated with activation of PI3K/AKT pathway and tamoxifen resistance in breast cancer.³⁸ PI3K/AKT is a well-documented pathway associated with human cancer risk that heavily regulates glucose and IGF signaling³⁹; therefore, it is possible that the expression of *PTPN2* interacts with diabetes through regulation of a variety of tyrosine kinases, given that tyrosine kinase activity of the insulin receptor is associated with human diabetes.⁴⁰ Again, functional follow up will be needed to better understand the interaction of this strong candidate gene.

We also observed a suggestive interaction between the *CD33* gene and BMI on CRC risk among men. We identified two genetic variants that were likely at least in part driving the significance. The *CD33* gene encodes a differentiation antigen of acute myeloid leukemia (AML) progenitor cells and is a very well-known pathological marker of AML.⁴¹ *CD33* is a transmembrane receptor, and was found to express on myeloid and lymphoid cells in about 85%-90% of patients with acute myeloid leukemia.⁴² *CD33* gene expression was also discovered to inhibit Ca²⁺ flux, cell growth and apoptosis.⁴³ Even though the association between *CD33* and AML were well-studied, how *CD33* is associated with CRC risk and how it interacts with BMI on CRC risk is yet to be identified. Previous research indicated that the frequency of *CD33* + cells in blood, as a subset of myeloid-derived suppressor cells, was significantly higher in obese subjects compared to nonobese individuals,⁴⁴ providing some evidence for a potential modifying effect of obesity. Our study provides preliminary results for a potential interaction between *CD33* and BMI on CRC risk that will require additional follow up analysis.

We identified that the *SCN1B* (Sodium Voltage-Gated Channel Beta Subunit 1) gene interacted with BMI on CRC risk among women at FDR <0.2. Voltage-gated sodium channels (Na_v)⁴⁵ are composed of one large pore-forming principal subunit and one or two smaller transmembrane subunits considered as auxiliary, and the *SCN1B* gene generates one of such subunits.⁴⁵ Multiple studies have demonstrated that its expression regulated cellular functions such as migration, differentiation, endosome acidification, phagocytosis, and podosome formation.⁴⁶ In addition, Na_v

are found abnormally expressed in carcinoma cells and tumor biopsies, and their activity is associated with aggressive features and cancer progression.⁴⁷ Expression of the Na_v1.5 isoform in breast tumors was found to be correlated with metastases development and patients' death,⁴⁸ and *SCN1B* mRNA was also discovered to be more abundant in highly invasive prostate cancer cell lines.⁴⁹ Our study revealed that the gene *SCN1B* significantly interacted with BMI on CRC risk, though the interaction became non-significant after Bonferroni correction. It is possible that obesity interacts with Na_v and further affects the regulations of cellular functions, resulting in various types of human cancers including CRC; further follow up studies are warranted.

Several strengths and limitations need to be considered when interpreting the findings. In our study, which is the largest to date to investigate gene-BMI and gene-diabetes interactions on CRC risk, we integrated colon-specific gene expression data to inform interaction testing. However, the tissue collection of the gene expression is suboptimal given that the transverse colon tissue samples from the GTEx Project covers the entire colonic wall, which not only included the epithelial cells of the mucosa from which CRC derived, but also all other tissue layers, which dilutes the epithelial gene expression profile. For this reason, we used the gene expression data from the transverse colon tissues instead of the sigmoid colon tissues in the GTEx project, as the sigmoid colon tissue samples were collected from the muscle tissues only. We did not include other tissues, as we have observed that colorectal cancer risk loci are enriched of colorectal tissue specific enhancer marks, which are key regulators for gene expression.⁵⁰ We also did not narrow down our analysis further using marginal eQTLs that have achieved transcriptome-wide association study (TWAS) significance level, because we were concerned that we might miss novel interactions. We combined colon and rectal cancer cases together, because the comprehensive analysis of The Cancer Genome Atlas (TCGA) demonstrated that colon and rectal cancer are very similar.⁵¹ To improve statistical power, we used our novel statistical set-based G \times E mixed effects score tests, MiSTi, which allows testing of both fixed- and random-effects of the interaction. As expected, the predicted heritability in gene expression differs between the genes. Accordingly, our statistical power to detect interaction between genetic-defined gene expression (fixed effects) varies and is higher for more heritable genes, given the effect size are the same. In other words, if there is little evidence for E and predicted gene expression, it could be due to low R^2 for the gene. Since expression levels of different genes may differ across populations and our analysis was limited to those of European descent, our results may not be necessarily generalizable to other race/ethnicity groups. Furthermore, the self-reported BMI and diabetes measurements might be

subjected to recall bias, however, we found similar effects for prospective cohort studies and case-control studies. Lastly, future independent replications are warranted, since FDR <0.2 is not stringent.

In summary, by incorporating functional information and conducting aggregated gene-based testing, the most significant interactions we observed were between *FOXA1* and BMI among men, and between *PTPN2* and diabetes for CRC risk. Other suggested genes interacting with BMI at FDR <0.2 included *CD33* and *PSMC5* among males, and *SCN1B* and *KIAA0753* among females. These findings provide support for potential new biological insights that could help in understanding the underlying mechanisms of BMI and diabetes on CRC. Independent replication and functional follow-up studies are warranted to confirm the functions of these genes in relation to BMI/diabetes and CRC development.

ACKNOWLEDGMENTS

ASTERISK: We are very grateful to Dr Bruno Buecher without whom this project would not have existed. We also thank all those who agreed to participate in this study, including the patients and the healthy control persons, as well as all the physicians, technicians and students. **CPS-II:** The authors thank the CPS-II participants and Study Management Group for their invaluable contributions to this research. The authors acknowledge the contribution to this study from the central cancer registries supported through the Centers for Disease Control and Prevention National Program of Cancer Registries, and cancer registries supported by the National Cancer Institute Surveillance Epidemiology and End Results program. **DACHS:** We thank all participants and cooperating clinicians, and Ute Handte-Daub, Utz Benschaid, Muhabbet Celik and Ursula Eilber for excellent technical assistance. **Harvard cohorts (HPFS, NHS, PHS):** The study protocol was approved by the institutional review boards of the Brigham and Women's Hospital and Harvard TH Chan School of Public Health, and those of participating registries as required. We thank the participants and staff of the HPFS, NHS, and PHS for their valuable contributions as well as the following state cancer registries for their help: AL, AZ, AR, CA, CO, CT, DE, FL, GA, ID, IL, IN, IA, KY, LA, ME, MD, MA, MI, NE, NH, NJ, NY, NC, ND, OH, OK, OR, PA, RI, SC, TN, TX, VA, WA, and WY. The authors assume full responsibility for analyses and interpretation of these data. **Kentucky:** We acknowledge the staff at the Kentucky Cancer Registry. **PLCO:** The authors thank the PLCO Cancer Screening Trial screening center investigators and the staff from Information Management Services Inc and Westat Inc. Most importantly, we thank the study participants for their contributions that made this study possible. **PMH-SCCFR:** The authors would like to thank the study participants and staff of the Hormones and

Colon Cancer and Seattle Cancer Family Registry studies (CORE Studies). **SEARCH:** We thank the SEARCH team. **WHI:** The authors thank the WHI investigators and staff for their dedication, and the study participants for making the program possible. A full listing of WHI investigators can be found at: <http://www.whi.org/researchers/Documents%20%20Write%20a%20Paper/WHI%20Investigator%20Short%20List.pdf>. **Disclaimer:** Where authors are identified as personnel of the International Agency for Research on Cancer / World Health Organization, the authors alone are responsible for the views expressed in this article and they do not necessarily represent the decisions, policy or views of the International Agency for Research on Cancer / World Health Organization.

CONFLICT OF INTEREST

None.

FUNDING INFORMATION

Genetics and Epidemiology of Colorectal Cancer Consortium (GECCO): National Cancer Institute, National Institutes of Health, U.S. Department of Health and Human Services (U01 CA164930, U01 CA137088, R01 CA059045, R01201407). Genotyping/Sequencing services were provided by the Center for Inherited Disease Research (CIDR) (X01-HG008596 and X-01-HG007585). CIDR is fully funded through a federal contract from the National Institutes of Health to The Johns Hopkins University, contract number HHSN268201200008I. This research was funded in part through the NIH/NCI Cancer Center Support Grant P30 CA015704. **ASTERISK:** a Hospital Clinical Research Program (PHRC-BRD09/C) from the University Hospital Center of Nantes (CHU de Nantes) and supported by the Regional Council of Pays de la Loire, the Groupement des Entreprises Françaises dans la Lutte contre le Cancer (GEFLUC), the Association Anne de Bretagne Génétique and the Ligue Régionale Contre le Cancer (LRCC). The ATBC Study is supported by the Intramural Research Program of the U.S. National Cancer Institute, National Institutes of Health, and by U.S. Public Health Service contract HHSN261201500005C from the National Cancer Institute, Department of Health and Human Services. **COLO2&3:** National Institutes of Health (R01 CA60987). The Colon Cancer Family Registry (CCFR) Illumina GWAS was supported by funding from the National Cancer Institute (NCI), National Institutes of Health (NIH) (grant numbers U01 CA122839, R01 CA143247). The CCFR participant recruitment and collection of data and biospecimens used in this study were supported by the NCI/NIH (grant number U01 CA167551) and through NCI/NIH cooperative agreements with the following Colon CFR centers: Australasian Colorectal Cancer Family Registry (grant numbers U01 CA074778 and U01/U24 CA097735), USC Consortium Colorectal Cancer Family Registry (grant

numbers U01/U24 CA074799), Mayo Clinic Cooperative Family Registry for Colon Cancer Studies (grant number U01/U24 CA074800), Ontario Familial Colorectal Cancer Registry (grant number U01/U24 CA074783), Seattle Colorectal Cancer Family Registry (grant number U01/U24 CA074794), and University of Hawaii Colorectal Cancer Family Registry (grant number U01/U24 CA074806) The content of this manuscript does not necessarily reflect the views or policies of the NCI, NIH or any of the collaborating centers in the Colon Cancer Family Registry (CCFR), nor does mention of trade names, commercial products, or organizations imply endorsement by the US Government, any cancer registry, or the CCFR. Colorectal Cancer Transdisciplinary (CORECT) Study: The CORECT Study was supported by the National Cancer Institute, National Institutes of Health (NCI/NIH), U.S. Department of Health and Human Services (grant numbers U19 CA148107, R01 CA81488, P30 CA014089, R01 CA197350; P01 CA196569; R01 CA201407) and National Institutes of Environmental Health Sciences, National Institutes of Health (grant number T32 ES013678). CPS-II: The American Cancer Society funds the creation, maintenance, and updating of the Cancer Prevention Study-II (CPS-II) cohort. This study was conducted with Institutional Review Board approval. DACHS: This work was supported by the German Research Council (BR 1704/6-1, BR 1704/6-3, BR 1704/6-4, CH 117/1-1, HO 5117/2-1, HE 5998/2-1, KL 2354/3-1, RO 2270/8-1 and BR 1704/17-1), the Interdisciplinary Research Program of the National Center for Tumor Diseases (NCT), Germany, and the German Federal Ministry of Education and Research (01KH0404, 01ER0814, 01ER0815, 01ER1505A and 01ER1505B). DALs: National Institutes of Health (R01 CA48998 to M. L. Slattery). Harvard cohorts (HPFS, NHS, PHS): HPFS is supported by the National Institutes of Health (P01 CA055075, UM1 CA167552, U01 CA167552, R01 CA137178, R01 CA151993, and R35 CA197735), NHS by the National Institutes of Health (R01 CA137178, P01 CA087969, UM1 CA186107, R01 CA151993, and R35 CA197735) and PHS by the National Institutes of Health (R01 CA042182). Kentucky: This work was supported by the following grant support: Clinical Investigator Award from Damon Runyon Cancer Research Foundation (CI-8); NCI R01CA136726. MCCS cohort recruitment was funded by VicHealth and Cancer Council Victoria. The MCCS was further supported by Australian NHMRC grants 509348, 209057, 251553 and 504711 and by infrastructure provided by Cancer Council Victoria. Cases and their vital status were ascertained through the Victorian Cancer Registry (VCR) and the Australian Institute of Health and Welfare (AIHW), including the National Death Index and the Australian Cancer Database. MEC: National Institutes of Health (R37 CA54281, P01 CA033619, and R01 CA063464). MECC: This work was supported by the National Institutes of

Health, U.S. Department of Health and Human Services (R01 CA81488 to SBG and GR). NFCCR: This work was supported by an Interdisciplinary Health Research Team award from the Canadian Institutes of Health Research (CRT 43821); the National Institutes of Health, U.S. Department of Health and Human Services (U01 CA74783); and National Cancer Institute of Canada grants (18223 and 18226). The authors wish to acknowledge the contribution of Alexandre Belisle and the genotyping team of the McGill University and Génome Québec Innovation Centre, Montréal, Canada, for genotyping the Sequenom panel in the NFCCR samples. Funding was provided to Michael O. Woods by the Canadian Cancer Society Research Institute. OFCCR: National Institutes of Health, through funding allocated to the Ontario Registry for Studies of Familial Colorectal Cancer (U01 CA074783); see CCFR section above. Additional funding toward genetic analyses of OFCCR includes the Ontario Research Fund, the Canadian Institutes of Health Research, and the Ontario Institute for Cancer Research, through generous support from the Ontario Ministry of Research and Innovation. PLCO: Intramural Research Program of the Division of Cancer Epidemiology and Genetics and supported by contracts from the Division of Cancer Prevention, National Cancer Institute, NIH, DHHS. Funding was provided by National Institutes of Health (NIH), Genes, Environment and Health Initiative (GEI) Z01 CP 010200, NIH U01 HG004446, and NIH GEI U01 HG 004438. SEARCH: The University of Cambridge has received salary support in respect of PDPP from the NHS in the East of England through the Clinical Academic Reserve. Cancer Research UK (C490/A16561); the UK National Institute for Health Research Biomedical Research Centres at the University of Cambridge. Swedish Mammography Cohort and Cohort of Swedish Men: This work is supported by the Swedish Research Council /Infrastructure grant, the Swedish Cancer Foundation, and the Karolinska Institute's Distinguished Professor Award to Alicja Wolk. VITAL: National Institutes of Health (K05 CA154337). WHI: The WHI program is funded by the National Heart, Lung, and Blood Institute, National Institutes of Health, U.S. Department of Health and Human Services through contracts HHSN268201100046C, HHSN268201100001C, HHSN268201100002C, HHSN268201100003C, HHSN268201100004C, and HHSN271201100004C. SPAIN: Colorectal Cancer Genetics & Genomics, Spanish study was supported by Instituto de Salud Carlos III, co-funded by FEDER funds –a way to build Europe– (grants PI14-613 and PI09-1286), Agency for Management of University and Research Grants (AGAUR) of the Catalan Government (grant 2017SGR723), and Junta de Castilla y León (grant LE22A10-2). Sample collection of this work was supported by the Xarxa de Bancs de Tumors de Catalunya sponsored by Pla Director d'Oncologia de Catalunya (XBTC), Plataforma Biobancos PT13/0010/0013

and ICOBIOBANC, sponsored by the Catalan Institute of Oncology. PMH-SCCFR: National Institutes of Health (R01 CA076366 to P. Newcomb and U01 CA074794 to J. Potter).

AUTHOR CONTRIBUTION

ZX, Y-RS, PP, YL, SB, XW, LH, PTC, and UP were involved in data analysis and interpretation, JCF, YL, DA, SIB, SB, DDB, GC, ATC, SJG, GGG, SBG, MJG, MH, MAJ, ADJ, LLM, LL, NML, VM, NM, PAN, GR, AW, MOW, HB, EW, MLS, ELG, JC-C, PDPP, PTC, and UP were involved in recruitment of study participants, data collection (questionnaire, biospecimens and genotyping), and data harmonization, Y-RS, HS and UP were involved in study design; all authors have been involved in manuscript writing and review.

DATA AVAILABILITY STATEMENT

Research data are not shared.

ORCID

Zhiyu Xia  <https://orcid.org/0000-0002-8340-8508>

Andre E. Kim  <https://orcid.org/0000-0003-1217-5249>

Daniel D. Buchanan  <https://orcid.org/0000-0003-2225-6675>

Michael Hoffmeister  <https://orcid.org/0000-0002-8307-3197>

REFERENCES

- Tanskanen T, van den Berg L, Välimäki N, et al. Genome-wide association study and meta-analysis in Northern European populations replicate multiple colorectal cancer risk loci. *Int J Cancer*. 2018;142(3):540-546.
- Terry P, Giovannucci E, Michels KB, et al. Fruit, vegetables, dietary fiber, and risk of colorectal cancer. *J Natl Cancer Inst*. 2001;93(7):525-533.
- Moghaddam AA, Woodward M, Huxley R. Obesity and risk of colorectal cancer: A meta-analysis of 31 studies with 70,000 events. *Cancer Epidemiol Biomarkers Prev*. 2007;16(12):2533-2547.
- Samad AKA, Taylor RS, Marshall T, Chapman MAS. A meta-analysis of the association of physical activity with reduced risk of colorectal cancer. *Color Dis*. 2005;7(3):204-213.
- Larsson SC, Orsini N, Wolk A. Diabetes mellitus and risk of colorectal cancer: a meta-analysis. *J Natl Cancer Inst*. 2005;97(22):1679-1687.
- Bergström A, Pisani P, Tenet V, Wolk A, Adami HO. Overweight as an avoidable cause of cancer in Europe. *Int J Cancer*. 2001;91(3):421-430.
- Meester RGS, Mannalithara A, Lansdorp-Vogelaar I, Ladabaum U. Trends in incidence and stage at diagnosis of colorectal cancer in adults aged 40 through 49 years, 1975–2015. *JAMA*. 2019;321(19):1933-1934.
- Slattery ML, Ballard-Barbash R, Edwards S, Caan BJ, Potter JD. Body mass index and colon cancer: an evaluation of the modifying effects of estrogen (United States). *Cancer Causes Control*. 2003;14(1):75-84.
- Kim H, Giovannucci EL. Sex differences in the association of obesity and colorectal cancer risk. *Cancer Causes Control*. 2017;28(1):1-4.
- Cohen DH, LeRoith D. Obesity, type 2 diabetes, and cancer: the insulin and IGF connection. *Endocr Relat Cancer*. 2012;19(5):F27-F45.
- Jiao S, Peters U, Berndt S, et al. Estimating the heritability of colorectal cancer. *Hum Mol Genet*. 2014;23(14):3898-3905.
- Yang T, Li X, Montazeri Z, et al. Gene–environment interactions and colorectal cancer risk: An umbrella review of systematic reviews and meta-analyses of observational studies. *Int J Cancer*. 2019;145(9):2315-2329.
- Hutter CM, Chang-Claude J, Slattery ML, et al. Characterization of gene-environment interactions for colorectal cancer susceptibility loci. *Cancer Res*. 2012;72(8):2036-2044.
- Kantor ED, Hutter CM, Minnier J, et al. Gene-environment interaction involving recently identified colorectal cancer susceptibility loci. *Cancer Epidemiol Biomarkers Prev*. 2014;23(9):1824-1833.
- Sainz J, Rudolph A, Hoffmeister M, et al. Effect of type 2 diabetes predisposing genetic variants on colorectal cancer risk. *J Clin Endocrinol Metab*. 2012;97(5):E845-E851.
- Su YR, Di CZ, Hsu L. A unified powerful set-based test for sequencing data analysis of G × E interactions. *Biostatistics*. 2017;18(1):119-131.
- Huyghe JR, Bien SA, Harrison TA, et al. Discovery of common and rare genetic risk variants for colorectal cancer. *Nat Genet*. 2019;51(1):76-87.
- Peters U, Jiao S, Schumacher FR, et al. Identification of genetic susceptibility loci for colorectal tumors in a genome-wide meta-analysis. *Gastroenterology*. 2013;144(4):799-807.
- Peters U, Hutter CM, Hsu L, et al. Meta-analysis of new genome-wide association studies of colorectal cancer risk. *Hum Genet*. 2012;131(2), 217-234.
- McCarthy S, Das S, Kretzschmar W, et al. A reference panel of 64,976 haplotypes for genotype imputation. *Nat Genet*. 2016;48(10):1279-1283.
- Gamazon ER, Wheeler HE, Shah KP, et al. A gene-based association method for mapping traits using reference transcriptome data. *Nat Genet*. 2015;47(9):1091-1098.
- Torres JM, Gamazon ER, Parra EJ, et al. Cross-tissue and tissue-specific eQTLs: Partitioning the heritability of a complex trait. *Am J Hum Genet*. 2014;95(5):521-534.
- Jeon J, Du M, Schoen RE, et al. Determining risk of colorectal cancer and starting age of screening based on lifestyle, environmental, and genetic factors. *Gastroenterology*. 2018;154(8):2152-2164.e19.
- Bernardo GM, Keri RA. FOXA1: a transcription factor with parallel functions in development and cancer. *Biosci Rep*. 2012;32(2):113-130.
- Augello MA, Hickey TE, Knudsen KE. FOXA1: Master of steroid receptor function in cancer. *EMBO J*. 2011;30(19):3885-3894.
- Dou C, Wang Y, Li C, et al. MicroRNA-212 suppresses tumor growth of human hepatocellular carcinoma by targeting FOXA1. *Oncotarget*. 2015;6(15):13216-13228.
- Sahu B, Laakso M, Ovaska K, et al. Dual role of FoxA1 in androgen receptor binding to chromatin, androgen signalling and prostate cancer. *EMBO J*. 2011;30(19):3962-3976.
- Ma W, Jiang J, Li M, et al. The clinical significance of forkhead box protein A1 and its role in colorectal cancer. *Mol Med Rep*. 2016;14(3):2625-2631.
- Bochkis IM, Schug J, Ye DZ, et al. Genome-wide location analysis reveals distinct transcriptional circuitry by paralogous regulators Foxa1 and Foxa2. *PLoS Genet*. 2012;8(6):e1002770.

30. Ehrlich AC, FriedenberG FK. Genetic Associations of Obesity: The Fat-Mass and Obesity-Associated (FTO) Gene. *Clin Transl Gastroenterol*. 2016;7(1):e140.
31. Peng H, Li J, Chen X, Zhou X, Zhu W, Li F. Genetic variants of PTPN2 gene in Chinese children with type 1 diabetes mellitus. *Monit Med Sci*. 2015;21:2653-2658.
32. Scharl M, Mwinyi J, Fischbeck A, et al. Crohn's disease-associated polymorphism within the PTPN2 gene affects muramyl-dipeptide-induced cytokine secretion and autophagy. *Inflamm Bowel Dis*. 2012;18(5):900-912.
33. Festen EAM, Goyette P, Green T, et al. A meta-analysis of genome-wide association scans identifies IL18RAP, PTPN2, TAGAP, and PUS10 as shared risk loci for crohn's disease and celiac disease. *PLoS Genet*. 2011;7(1):e1001283.
34. Burton PR, Clayton DG, Cardon LR, et al. Genome-wide association study of 14,000 cases of seven common diseases and 3,000 shared controls. *Nature*. 2007;447(7145):661-678.
35. Todd JA, Walker NM, Cooper JD, et al. Robust associations of four new chromosome regions from genome-wide analyses of type 1 diabetes. *Nat Genet*. 2007;39(7):857-864.
36. Spalinger MR, Manzini R, Hering L, et al. PTPN2 regulates inflammasome activation and controls onset of intestinal inflammation and colon cancer. *Cell Rep*. 2018;22(7):1835-1848.
37. Hotamisligil GS. Inflammation and metabolic disorders. *Nature*. 2006;444:860-867.
38. Karlsson E, Veenstra C, Emin S, et al. Loss of protein tyrosine phosphatase, non-receptor type 2 is associated with activation of AKT and tamoxifen resistance in breast cancer. *Breast Cancer Res Treat*. 2015;153(1):31-40.
39. Luo J, Manning BD, Cantley LC. Targeting the PI3K-Akt pathway in human cancer: rationale and promise. *Cancer Cell*. 2003;4(4):257-262.
40. Hotamisligil GS, Budavari A, Murray D, Spiegelman BM. Reduced tyrosine kinase activity of the insulin receptor in obesity-diabetes. *J Clin Invest*. 1994;94(4):1543-1549.
41. Minagawa K, Jamil MO, Al-Obaidi M, et al. In vitro pre-clinical validation of suicide gene modified anti-CD33 redirected chimeric antigen receptor T-cells for acute myeloid leukemia. *PLoS One*. 2016;11(12):e0166891.
42. Sperr WR, Florian S, Hauswirth AW, Valent P. CD33 as a target of therapy in acute myeloid leukemia: Current status and future perspectives. *Leuk Lymphoma*. 2005;46(8):1115-1120.
43. Peiper SC, Ashmun RA, Look AT. Molecular cloning, expression, and chromosomal localization of a human gene encoding the CD33 myeloid differentiation antigen. *Blood*. 1988;72(1):314-321.
44. Klimcakova E, Roussel B, Kovacova Z, et al. Macrophage gene expression is related to obesity and the metabolic syndrome in human subcutaneous fat as well as in visceral fat. *Diabetologia*. 2011;54(4):876-887.
45. Catterall WA. Voltage-gated sodium channels: structure, function, and pathophysiology, in: encyclopedia of biological chemistry: second edition. 2013. <https://doi.org/10.1016/B978-0-12-378630-2.00208-5>
46. Black JA, Waxman SG. Noncanonical roles of voltage-gated sodium channels. *Neuron*. 2013;80(2):280-291.
47. Besson P, Driffort V, Bon É, Gradek F, Chevalier S, Roger S. How do voltage-gated sodium channels enhance migration and invasiveness in cancer cells? *Biochim Biophys Acta – Biomembr*. 2015;1848:2493-2501.
48. Fraser SP, Diss JKJ, Chioni AM, et al. Voltage-gated sodium channel expression and potentiation of human breast cancer metastasis. *Clin Cancer Res*. 2005;11(15):5381-5389.
49. Diss JKJ, Fraser SP, Walker MM, Patel A, Latchman DS, Djamgoz MBA. β -Subunits of voltage-gated sodium channels in human prostate cancer: Quantitative in vitro and in vivo analyses of mRNA expression. *Prostate Cancer Prostatic Dis*. 2008;11(4):325-333.
50. Bien SA, Auer PL, Harrison TA, et al. Enrichment of colorectal cancer associations in functional regions: Insight for using epigenomics data in the analysis of whole genome sequence-imputed GWAS data. *PLoS One*. 2017;12(11):e0186518.
51. Muzny DM, Bainbridge MN, Chang K, et al. Comprehensive molecular characterization of human colon and rectal cancer. *Nature*. 2012; 487(7407):330-337.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

How to cite this article: Xia Z, Su Y-R, Petersen P, et al. Functional informed genome-wide interaction analysis of body mass index, diabetes and colorectal cancer risk. *Cancer Med*. 2020;9:3563–3573. <https://doi.org/10.1002/cam4.2971>