## **UC Irvine**

# **UC Irvine Previously Published Works**

#### **Title**

Erratum: "1.5D quasilinear model and its application on beams interacting with Alfvén eigenmodes in DIII-D" [Phys. Plasmas 19, 092511 (2012)]

#### **Permalink**

https://escholarship.org/uc/item/89d0t4b1

### Journal

Physics of Plasmas, 21(4)

#### **ISSN**

1070-664X

#### **Authors**

Ghantous, K Gorelenkov, NN Berk, HL et al.

#### **Publication Date**

2014-04-01

#### DOI

10.1063/1.4870636

## **Copyright Information**

This work is made available under the terms of a Creative Commons Attribution License, available at <a href="https://creativecommons.org/licenses/by/4.0/">https://creativecommons.org/licenses/by/4.0/</a>

Peer reviewed



# Erratum: "1.5D quasilinear model and its application on beams interacting with Alfvén eigenmodes in DIII-D" [Phys. Plasmas 19, 092511 (2012)]

K. Ghantous, N. N. Gorelenkov, H. L. Berk, W. W. Heidbrink, and M. A. Van Zeeland

<sup>1</sup>Princeton Plasma Physics Laboratory, P.O. Box 451, Princeton, New Jersey 08543-0451, USA

<sup>2</sup>Institute for Fusion Studies, University of Texas, 2100 San Jacinto Blvd, Austin, Texas 78712-1047, USA

<sup>3</sup>Department of Physics and Astronomy, University of California, Irvine, Irvine, California 92697, USA

<sup>4</sup>General Atomics, P.O. Box 85608, San Diego, California 92186-560, USA

(Received 24 March 2014; accepted 26 March 2014; published online 1 April 2014)

[http://dx.doi.org/10.1063/1.4870636]

We make the following corrections to typos we found in the original paper.<sup>1</sup>

In the abstract, the reference is to shot #112117 not #127112 [W. W. Heidbrink *et al.*, Nucl. Fusion **48**, 084001 (2008)].

Equation (13)

For consistency, we replace the variable q by e for the charge, since q is used as safety factor elsewhere. Equation (13) therefore reads as

$$\gamma_L = \frac{\pi}{2} \int \frac{e}{c} dP_{\phi} dE \left| \left\langle \frac{v_d \cdot \delta E}{\omega} \right\rangle \right|^2 \delta(\Omega) \left( \omega \frac{\partial}{\partial E} + n \frac{\partial}{\partial P_{\phi}} \right) f,$$

where  $D_k = W_k |\langle \frac{e}{c\omega} v_d \cdot \delta E \rangle|^2 \delta(\Omega)$  is the diffusion coefficient at the resonances in phase space.

Equation (17)

Equation (17) should be written as

$$\left. \frac{\partial \beta_{EP}}{\partial r} \right|_{\text{crt}} = \frac{\gamma_d}{\gamma_L'}.$$

Equation (18)

 $\rho$  in Eq. (18) is  $\rho_s$ 

$$\frac{\gamma_{rad}}{\omega} = -3 \left| \frac{\sqrt{\rho_s^2}}{r\sqrt{2}} \frac{\mathrm{sm}(m+1)}{2m+1} \right|^{0.67},$$

where s is the local shear at the location of the mode, m is the poloidal mode number,  $\rho_s = c_s/\Omega_{ci}$  with  $c_s = \sqrt{\gamma ZkT_e/m_i}$  is the ion sound speed, and  $\Omega_{ci}$  is the ion gyro-frequency.

Equation (20)

The exponential in Eq. (20) for the electron landau damping is -1/s not 1/s. We also correct the expression for  $G(\hat{\epsilon})$  and further explain some variables

$$\frac{\gamma_{eL}}{\omega} = -\frac{\pi^{3/2}}{6} q^2 \beta_e \frac{v_A}{v_e} \left(\frac{5}{2} \epsilon\right) G(\hat{\epsilon}) e^{-1/s}$$

and  $G(\hat{\epsilon}) \approx 4.47 + 0.42\hat{\epsilon} + 0.02\hat{\epsilon}^2$  not  $G(\hat{\epsilon}) \approx 4.47 - 0.42\hat{\epsilon} + 0.02\hat{\epsilon}^2$  as typed in the manuscript. s is the local shear  $s = (r/q) \, dq/dr$ , where r is the radius and q is the safety factor.  $\hat{\epsilon} = 2\epsilon/(1-\epsilon)$ , where  $\epsilon = r/a$  with a minor radius of the plasma last close flux surface.

Equation (21)

There is a missing factor of  $(\pi/2)^2$ . Equation (21) should be

$$\begin{split} \frac{\gamma_{eColl}}{\omega} &= -\frac{1}{4} \left(\frac{\pi}{2}\right)^2 \left[ I_1 \left(\frac{8snq\rho_s}{5r\epsilon}\right)^2 + I_2 q^2 \frac{8\beta_{pc}}{1+\sigma} \right] \sqrt{\frac{\nu}{\omega}} \\ &\times \left[ \ln \left(16\sqrt{\frac{\omega\epsilon}{\nu}}\right) \right]^{-3/2}. \end{split}$$

Equation (25)

The diamagnetic frequency in Eq. (25) is

$$\omega^* = \frac{nq_m v_0^2}{r_m \Omega_0} \frac{\partial \ln \beta}{\partial r},$$

where  $q_m$  and  $r_m$  are the safety factor and radius at the location of the toroidal Alfvenic eigenmodes mode. n is the toroidal mode number and  $\Omega_c$  is the fast ion cyclotron frequency.

<sup>&</sup>lt;sup>1</sup>K. Ghantous, N. N. Gorelenkov, H. L. Berk, W. W. Heidbrink, and M. A. Van Zeeland, Phys. Plasmas 19, 092511 (2012).