

UCLA

Proceedings of UCLA Health

Title

Bicuspid Aortic Valve Infective Endocarditis: Is There a Need to Widen the Indications for Antibiotic Prophylaxis

Permalink

<https://escholarship.org/uc/item/89h0k1mv>

Journal

Proceedings of UCLA Health, 23(1)

Authors

Anousheh, Ramtin

Mehrinfar-Zadeh, Ramona

Publication Date

2020-03-13

CLINICAL VIGNETTE

Bicuspid Aortic Valve Infective Endocarditis: Is There a Need to Widen the Indications for Antibiotic Prophylaxis

Ramtin Anousheh, MD, MPH and Ramona Mehrinfar-Zadeh, MD

Introduction

Bicuspid aortic valve (BAV), is the most common congenital cardiac valvular disease with prevalence ranging from 0.5% to 2%.¹ It increases the risk of infective endocarditis, thoracic aortic aneurysm and dissection, and sudden cardiac death.² Patients with bicuspid aortic valves do not routinely need antibiotics before dental and surgical procedures and currently the only indication for prophylaxis is prior history of endocarditis.³ We present a case of sub-acute infective endocarditis in an otherwise healthy female.

Case Presentation

A 51-year-old female initially presented to primary care clinic with dizziness and feeling unwell. She had prior diagnosis of acute sinusitis, documented on MR and had received several courses of antibiotics. Her symptoms did not improve over three months and she developed anemia, 30 lbs unintentional weight loss, intermittent palpitations and visual disturbances prompting multispecialty referrals including cardiology. At her outpatient cardiology evaluation a loud diastolic murmur was detected and a transthoracic echocardiogram (TTE) showed a large vegetation on a bicuspid aortic valve with moderate to severe aortic regurgitation with preserved left ventricular dimensions and function.

She was admitted to the hospital and diagnosed with culture-negative infective endocarditis and treated with six-weeks of intravenous vancomycin. Follow up TEE showed substantial decrease in the size of vegetation, ventricular dilatation and decreased systolic function. She subsequently underwent aortic valve replacement and repair of thoracic aortic aneurysm.

Discussion

Bicuspid aortic valve is primarily inherited in an autosomal-dominant pattern. Studies have found a 15% rate of familial clustering.⁴ In a study of 142 patients with bicuspid aortic valve, 20% of first-degree relatives had some cardiac abnormality found on screening, of whom 68% had bicuspid aortic valve. Of these, 71% were newly detected abnormalities.⁵

A systolic ejection click is followed by an early peaking systolic murmur, most commonly at the apex. Bicuspid valves are typically initially asymptomatic but are commonly associated with progressive valvulopathy and aortopathy.

Sabet et al studied 542 patients with congenital bicuspid aortic valves undergoing aortic valve replacement. Seventy-five percent of the patients had isolated aortic stenosis, 10% had aortic stenosis with some degree of aortic insufficiency, and 13% had isolated aortic insufficiency. Increased risk of aneurysmal dilatation, coarctation, and dissection of thoracic aorta was also found.²

Several observational studies have shown that the patients with a native BAV are at markedly increased risk of aortic valve endocarditis compared with endocarditis of other native heart valves. The relative risk of IE of a BAV was markedly greater than for a tricuspid aortic valve. Recent estimates of the incidence of infective endocarditis (IE) occurring in BAV patients are 2% or 0.3%/year.⁶

The etiology of BAV association with increased frequency of IE is not established. Altered flow patterns across the BAV can result in valvular endothelial injury, including platelet and fibrinogen deposition. This has been postulated to allow seeding of hematogenous bacteria and fungi.⁷

Older age, intravenous drug abuse and multiple comorbidities including end-stage renal disease and indwelling cardiac devices are independent risk factors for endocarditis in patients with BAV.

In the current era, TTE can usually confirm the diagnosis of IE. If there is uncertainty in the diagnosis, TEE can improve the sensitivity and specificity. Due to the high rate of negative blood cultures caused by empiric antibacterial therapy administered prior to the hospitalization in United States, the role of TTE or TEE is particularly important.

For aortic valve IE, surgeons should excise the diseased aortic valve, and clear calcified plaque and infective tissues. Presence of peri-valvular abscesses increases the technical difficulties and risks of surgical treatment.

Current guidelines do not recommend prophylactic antibiotics in patients with BAV undergoing procedures prone to bacteremia. As the majority of patients with BAV are unaware of their BAV status until the occurrence of significant morbidity, estimating the risk of IE in patients with BAV, compared with individuals with tricuspid valves is challenging.

Randomized trials of prophylactic antibiotics in patients with BAV undergoing procedures prone to bacteremia have potential value.

Considering the overwhelming evidence based on multiple published case reports and series, we recommend antibiotic prophylaxis prior to high risk procedures in patients with known structurally or functionally abnormal bicuspid aortic valves.

REFERENCES

1. **Michelena HI, Prakash SK, Della Corte A, Bissell MM, Anavekar N, Mathieu P, Bossé Y, Limongelli G, Bossone E, Benson DW, Lancellotti P, Isselbacher EM, Enriquez-Sarano M, Sundt TM 3rd, Pibarot P, Evangelista A, Milewicz DM, Body SC; BAVCon Investigators.** Bicuspid aortic valve: identifying knowledge gaps and rising to the challenge from the International Bicuspid Aortic Valve Consortium (BAVCon). *Circulation*. 2014 Jun 24;129(25):2691-704. doi: 10.1161/CIRCULATIONAHA.113.007851. Review. PubMed PMID: 24958752; PubMed Central PMCID: PMC4145814.
2. **Sabet HY, Edwards WD, Tazelaar HD, Daly RC.** Congenitally bicuspid aortic valves: a surgical pathology study of 542 cases (1991 through 1996) and a literature review of 2,715 additional cases. *Mayo Clin Proc*. 1999 Jan;74(1):14-26. PubMed PMID: 9987528.
3. **Wilson W, Taubert KA, Gewitz M, Lockhart PB, Baddour LM, Levison M, Bolger A, Cabell CH, Takahashi M, Baltimore RS, Newburger JW, Strom BL, Tani LY, Gerber M, Bonow RO, Pallasch T, Shulman ST, Rowley AH, Burns JC, Ferrieri P, Gardner T, Goff D, Durack DT; American Heart Association Rheumatic Fever, Endocarditis, and Kawasaki Disease Committee; American Heart Association Council on Cardiovascular Disease in the Young; American Heart Association Council on Clinical Cardiology; American Heart Association Council on Cardiovascular Surgery and Anesthesia; Quality of Care and Outcomes Research Interdisciplinary Working Group.** Prevention of infective endocarditis: guidelines from the American Heart Association: a guideline from the American Heart Association Rheumatic Fever, Endocarditis, and Kawasaki Disease Committee, Council on Cardiovascular Disease in the Young, and the Council on Clinical Cardiology, Council on Cardiovascular Surgery and Anesthesia, and the Quality of Care and Outcomes Research Interdisciplinary Working Group. *Circulation*. 2007 Oct 9;116(15):1736-54. Epub 2007 Apr 19. Erratum in: *Circulation*. 2007 Oct 9;116(15):e376-7. PubMed PMID: 17446442.
4. **Shah SY, Higgins A, Desai MY.** Bicuspid aortic valve: Basics and beyond. *Cleve Clin J Med*. 2018 Oct;85(10):779-784. doi: 10.3949/ccjm.85a.17069. Review. PubMed PMID: 30289756.
5. **Giusti B, Sticchi E, De Cario R, Magi A, Nistri S, Pepe G.** Genetic Bases of Bicuspid Aortic Valve: The Contribution of Traditional and High-Throughput Sequencing Approaches on Research and Diagnosis. *Front Physiol*. 2017 Aug 24;8:612. doi: 10.3389/fphys.2017.00612. eCollection 2017. Review. PubMed PMID: 28883797; PubMed Central PMCID: PMC5573733.
6. **Kiyota Y, Della Corte A, Montiero Vieira V, Habchi K, Huang CC, Della Ratta EE, Sundt TM, Shekar P, Muehlschlegel JD, Body SC.** Risk and outcomes of aortic valve endocarditis among patients with bicuspid and tricuspid aortic valves. *Open Heart*. 2017 May 16;4(1):e000545. doi: 10.1136/openhrt-2016-000545. eCollection 2017. PubMed PMID: 28674620; PubMed Central PMCID: PMC5471870.
7. **Veloso TR, Chaouch A, Roger T, Giddey M, Vouillamoz J, Majcherczyk P, Que YA, Rousson V, Moreillon P, Entenza JM.** Use of a human-like low-grade bacteremia model of experimental endocarditis to study the role of Staphylococcus aureus adhesins and platelet aggregation in early endocarditis. *Infect Immun*. 2013 Mar;81(3):697-703. doi: 10.1128/IAI.01030-12. Epub 2012 Dec 17. PubMed PMID: 23250949; PubMed Central PMCID: PMC3584864.