

UC Berkeley

UC Berkeley Previously Published Works

Title

Cav1.3 channels control D2-autoreceptor responses via NCS-1 in substantia nigra dopamine neurons

Permalink

<https://escholarship.org/uc/item/8b9974t4>

Journal

Brain, 137(8)

ISSN

0006-8950

Authors

Dragicevic, Elena
Poetschke, Christina
Duda, Johanna
et al.

Publication Date

2014-08-01

DOI

10.1093/brain/awu131

Peer reviewed

Ca_v1.3 channels control D2-autoreceptor responses via NCS-1 in substantia nigra dopamine neurons

Elena Dragicevic,¹ Christina Poetschke,^{1,*} Johanna Duda,^{1,*} Falk Schlaudraff,¹ Stephan Lammel,² Julia Schiemann,³ Michael Fauler,¹ Andrea Hetzel,⁴ Masahiko Watanabe,⁵ Rafael Lujan,⁶ Robert C. Malenka,² Joerg Striessnig⁷ and Birgit Liss¹

1 Institute of Applied Physiology, University of Ulm, Ulm, Germany

2 Nancy Pritzker Laboratory, Department of Psychiatry and Behavioural Sciences, Stanford University School of Medicine, Palo Alto, USA

3 Institute of Neurophysiology, Goethe University Frankfurt, Germany

4 Rosalind Franklin University of Medicine and Science, North Chicago, USA

5 Department of Anatomy, Hokkaido University School of Medicine, Sapporo, Japan

6 Instituto de Investigación en Discapacidades Neurológicas (IDINE), Departamento de Ciencias Médicas, Universidad Castilla-La Mancha, Albacete, Spain

7 Institute of Pharmacy, Department of Pharmacology and Toxicology, Centre of Molecular Biosciences, University of Innsbruck, Austria

*These authors contributed equally to this work.

Correspondence to: Birgit Liss,
Institute of Applied Physiology, University of Ulm,
Albert Einsteinallee 11, 89081, Ulm,
Germany
E-mail: birgit.liss@uni-ulm.de

See doi:10.1093/brain/awu150 for the scientific commentary on this article.

Dopamine midbrain neurons within the substantia nigra are particularly prone to degeneration in Parkinson's disease. Their selective loss causes the major motor symptoms of Parkinson's disease, but the causes for the high vulnerability of SN DA neurons, compared to neighbouring, more resistant ventral tegmental area dopamine neurons, are still unclear. Consequently, there is still no cure available for Parkinson's disease. Current therapies compensate the progressive loss of dopamine by administering its precursor L-DOPA and/or dopamine D2-receptor agonists. D2-autoreceptors and Ca_v1.3-containing L-type Ca²⁺ channels both contribute to Parkinson's disease pathology. L-type Ca²⁺ channel blockers protect SN DA neurons from degeneration in Parkinson's disease and its mouse models, and they are in clinical trials for neuroprotective Parkinson's disease therapy. However, their physiological functions in SN DA neurons remain unclear. D2-autoreceptors tune firing rates and dopamine release of SN DA neurons in a negative feedback loop through activation of G-protein coupled potassium channels (GIRK2, or KCNJ6). Mature SN DA neurons display prominent, non-desensitizing somatodendritic D2-autoreceptor responses that show pronounced desensitization in PARK-gene Parkinson's disease mouse models. We analysed surviving human SN DA neurons from patients with Parkinson's disease and from controls, and detected elevated messenger RNA levels of D2-autoreceptors and *GIRK2* in Parkinson's disease. By electrophysiological analysis of postnatal juvenile and adult mouse SN DA neurons in *in vitro* brain-slices, we observed that D2-autoreceptor desensitization is reduced with postnatal maturation. Furthermore, a transient high-dopamine state *in vivo*, caused by one injection of either L-DOPA or cocaine, induced adult-like, non-desensitizing D2-autoreceptor responses, selectively in juvenile SN DA neurons, but not ventral tegmental area dopamine neurons. With pharmacological and genetic tools, we identified that the expression of this sensitized D2-autoreceptor phenotype required Ca_v1.3 L-type Ca²⁺ channel activity, internal Ca²⁺, and the interaction of the neuronal calcium sensor NCS-1 with D2-autoreceptors. Thus, we identified a first physiological function of Ca_v1.3 L-type Ca²⁺ channels in SN DA neurons for

Received October 4, 2013. Revised March 31, 2014. Accepted April 6, 2014. Advance Access publication June 16, 2014

© The Author (2014). Published by Oxford University Press on behalf of the Guarantors of Brain.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

homeostatic modulation of their D2-autoreceptor responses. L-type Ca^{2+} channel activity however, was not important for pacemaker activity of mouse SN DA neurons. Furthermore, we detected elevated substantia nigra dopamine messenger RNA levels of NCS-1 (but not $\text{Ca}_v1.2$ or $\text{Ca}_v1.3$) after cocaine in mice, as well as in remaining human SN DA neurons in Parkinson's disease. Thus, our findings provide a novel homeostatic functional link in SN DA neurons between $\text{Ca}_v1.3$ -L-type- Ca^{2+} channels and D2-autoreceptor activity, controlled by NCS-1, and indicate that this adaptive signalling network ($\text{Ca}_v1.3$ /NCS-1/D2/GIRK2) is also active in human SN DA neurons, and contributes to Parkinson's disease pathology. As it is accessible to pharmacological modulation, it provides a novel promising target for tuning substantia nigra dopamine neuron activity, and their vulnerability to degeneration.

Keywords: D2-autoreceptor; isradipine; Parkinson's disease; L-DOPA; cocaine

Introduction

Substantia nigra dopaminergic neurons project to the dorsal striatum and are important for voluntary movement. In contrast with neighbouring dopamine midbrain neurons in the ventral tegmental area, SN DA neurons are particularly prone to degeneration in Parkinson's disease (Bjorklund and Dunnett, 2007; Schultz, 2010). The cause for this selective vulnerability of SN DA neurons is still unclear, and the typical motor symptoms of Parkinson's disease occur when ~70% of SN DA neurons are already lost (Collier *et al.*, 2011; Shulman *et al.*, 2011; Sulzer and Surmeier, 2012). Consequently, there is still no cure for Parkinson's disease available, and current Parkinson's disease therapies try to compensate the progressive loss of striatal dopamine by administering its precursor L-DOPA (still the gold standard in Parkinson's disease therapy) and/or dopamine D2 receptor agonists (Collier *et al.*, 2011; Gazewood *et al.*, 2013; Olanow and Schapira, 2013). Understanding the molecular mechanisms of substantia nigra dopamine function and signalling in health and disease is the prerequisite for the development of novel neuroprotective Parkinson's disease therapies. Dopamine D2-autoreceptors and $\text{Ca}_v1.3$ L-type- Ca^{2+} channels, calcium load and metabolic stress in SN DA neurons have already been identified as trigger factors, contributing to the progressive loss of SN DA neurons, striatal dopamine, and thus to Parkinson's disease pathology (Liss and Roeper, 2010; Surmeier *et al.*, 2012; Ford, 2014), but their interplay is still unclear.

Dopamine release from striatal axon terminals (and from soma and dendrites) of SN DA neurons is controlled by their electrical activity (Beckstead *et al.*, 2007; Rice *et al.*, 2011; Ford, 2014). In synaptic isolation, SN DA neurons display a so-called 'pacemaker activity' that is intrinsically generated by a concerted interplay of a range of voltage-gated ion channels, however, with still unclear individual contributions (Khaliq and Bean, 2010; Liss and Roeper, 2010; Surmeier *et al.*, 2012). Pacemaker activity of SN DA neurons is inhibited by dopamine itself, via therapeutically relevant dopamine D2-autoreceptors, through activation of inwardly rectifying potassium channels (GIRK2) (Luscher and Slesinger, 2010; Anzalone *et al.*, 2012; Gantz *et al.*, 2013). Mouse models with mutations in *PARK* genes, which lead to familial forms of Parkinson's disease (Gasser *et al.*, 2011), show altered substantia nigra dopamine D2-autoreceptor responses (Goldberg *et al.*, 2005;

Tong *et al.*, 2009), further linking D2-autoreceptor function to Parkinson's disease pathology.

The contribution of, in particular, the $\text{Ca}_v1.3$ L-type- Ca^{2+} channels to pacemaker activity is more controversial than that of D2-autoreceptors. More precisely, a suggested postnatal age-dependent switch from a HCN (hyperpolarization-activated cyclic nucleotide gated) channel driven pacemaker in juvenile postnatal mouse SN DA neurons to a metabolically more challenging and $\text{Ca}_v1.3$ L-type- Ca^{2+} channel driven pacemaker in adult SN DA neurons, which would render them more vulnerable to Parkinson's disease triggers, is highly disputed (Chan *et al.*, 2007; Puopolo *et al.*, 2007; Guzman *et al.*, 2009; Drion *et al.*, 2011; Surmeier *et al.*, 2012). No other calcium mediated physiological functions of L-type- Ca^{2+} channels in SN DA neurons are known. However, blood-brain barrier permeable L-type- Ca^{2+} channel blockers (e.g. isradipine) do protect highly vulnerable SN DA neurons from degeneration in Parkinson's disease and its chronic rodent models (Ritz *et al.*, 2010; Ilijic *et al.*, 2011; Surmeier *et al.*, 2012), presumably by reducing substantia nigra dopamine calcium load and metabolic mitochondrial stress during pacemaking (Guzman *et al.*, 2010; Surmeier and Schumacker, 2013); and they are already in clinical trials for pharmacological neuroprotective Parkinson's disease therapy (Parkinson Study Group, 2013; ClinicalTrials.gov Identifier: NCT00909545). In contrast to wild-type, SN DA neurons from adult *PARK7* (DJ-1) knockout mice display not only increased L-type- Ca^{2+} channel-generated metabolic stress levels, but also rapidly desensitizing somatodendritic D2-autoreceptor responses (Goldberg *et al.*, 2005; Guzman *et al.*, 2010), suggesting a possible link between L-type- Ca^{2+} channels and D2-autoreceptor function. D2 receptor signalling in general is complex, depending on additional factors such as Ca^{2+} and the neuronal calcium sensor 1 (NCS-1) (Kabbani *et al.*, 2002), and the signalling mechanisms of somatodendritic D2-autoreceptors remain poorly understood (Beaulieu and Gainetdinov, 2011; Ford, 2014).

Here, we detected dramatically higher messenger RNA levels of D2-autoreceptors and GIRK2 in SN DA neurons from *post-mortem* Parkinson's disease brains compared to those of controls. To address a possible homeostatic plasticity of D2-autoreceptor responses, in age-dependent interplay with L-type- Ca^{2+} channels, we analysed isolated somatodendritic D2-autoreceptor

responses as well as related L-type-Ca²⁺ channel function in SN DA neurons from juvenile (postnatal Day 13) and adult (postnatal Day 90) mice *in vitro* – under control conditions and 3–4 days after inducing a transient elevation of extracellular dopamine levels by a single *in vivo* injection of either L-DOPA or cocaine. We detected prominent desensitization of D2-autoreceptor responses in juvenile SN DA neurons that were absent in adult mice. Non-desensitizing, adult-like D2-autoreceptor responses could be induced in juvenile substantia nigra (but not ventral tegmental area) dopaminergic neurons by transient high-dopamine states *in vivo*. Ca_v1.3 L-type-Ca²⁺ channel function and Ca²⁺-dependent NCS-1/D2-autoreceptor interactions were crucial for this effect. *Ncs1* messenger RNA levels were significantly higher in mouse juvenile SN DA neurons after *in vivo* cocaine, as well as in surviving human SN DA neurons in patients with Parkinson's disease. Thereby, our findings provide a novel functional link in SN DA neurons between Ca_v1.3 L-type-Ca²⁺ channels and D2-autoreceptor activity, controlled by NCS-1 levels. They indicate that this signalling complex (Ca_v1.3/NCS-1/D2-autoreceptors) is tuning substantia nigra dopamine activity (and likely long-term vulnerability), in response to changes in extracellular dopamine levels in mice (cocaine, L-DOPA) as well as in humans (Parkinson's disease). Thus, pharmacological modulation of this signalling network could emerge as a promising novel approach for modulating substantia nigra dopamine activity, as well as their vulnerability.

Materials and methods

Animals

Juvenile (postnatal Days 12–14) and adult (approximately postnatal Day 90) male C57BL/6, Ca_v1.2 DHP^{-/-} mice and Ca_v1.3 knockout mice were bred and genotyped as described previously (Platzer *et al.*, 2000; Sinnegger-Brauns *et al.*, 2009). NCS-1^{-/-} mice were kindly provided by Prof. Olaf Pongs, Hamburg. Mice were injected: subcutaneously with a mix of 100 mg/kg L-DOPA methyl-ester hydrochloride (Sigma) and 200 mg/kg of the peripheral decarboxylase-inhibitor benserazide hydrochloride (Sigma) or intraperitoneally with 15 mg/kg cocaine hydrochloride (Sigma) in saline (0.9% NaCl, Braun), or with saline (subcutaneously or intraperitoneally) as control. In contrast to cocaine, the L-DOPA administration for mice is less well-established. In divergence to a 4:1 ratio used in humans, we used L-DOPA and benserazide in 1:2 ratio, as in most studies for rodents (due to presumably faster metabolism). L-DOPA injections have been administered subcutaneously as this results in more consistent responses; response failures after intraperitoneal, but not subcutaneous, injections have been described (Lundblad *et al.*, 2004, 2005; Lindgren *et al.*, 2007; Prinz *et al.*, 2013). Because L-DOPA and cocaine injections both led to the same effects on mouse substantia nigra dopamine D2-autoreceptor response, we continued our study with more robust and, for rodents, well-established cocaine paradigm (Saal *et al.*, 2003). All animal procedures were approved by the German Regierungspräsidentium Tübingen (Aktenzeichen: 35/9185.81-3; TV-No. 921 and 1043) or Darmstadt (V54-19c20/15-F40/28).

Ultraviolet laser microdissection and reverse transcription quantitative polymerase chain reaction analysis

UV laser microdissection of dopamine neurons from mouse and human brains using an LMD6000 system (Leica Microsystems), or cytoplasm harvest were performed essentially as described (Grundemann *et al.*, 2011). *Post-mortem* human midbrains were obtained by the German Brain Bank (BrainNet). All available information on the respective Parkinson's disease and control patient's disease state, comorbidity, age, and tissue quality is summarized in Supplementary Table 1. We were not able to obtain information about the therapy of our analysed patients with Parkinson's disease. However, it is likely that they were treated with L-DOPA (and/or D2-agonist).

All experiments on human material were approved by the ethic commission of Ulm University (247/10-UBB/bal).

Quantification of messenger RNA levels from UV laser microdissected samples was carried out by real time quantitative PCR after reverse transcription with a GeneAmp 7900HT (Applied Biosystems). For real-time quantitative PCR assays, TaqMan[®] probes were labelled with FAM as reporter and a non-fluorescent quencher. Quantitative PCR assay details, and assay-specific standard curve parameters are given in Supplementary Table 2. Standard curves for quantitative PCR quantification were generated using serial dilutions of complementary DNA (3, 0.3 and 0.03 ng), derived from human substantia nigra tissue RNA (Ambion, Applied Biosystems), or mouse midbrain messenger RNA (RNeasy[®] MINI kit, Qiagen), as template in duplicates in at least $n = 3$ independent quantitative PCR runs. The complementary DNA amount per cell in relation to the standard was calculated according to:

$$cDNA \text{ amount per cell} = \frac{S[(C_t - Y_{intercept})/slope]}{No_{cells} \bullet cDNA \text{ fraction}}$$

Where: S = serial dilution factor of the standard curve (i.e. 10), No_{cells} = number of harvested neurons per sample (i.e. 10 for mouse and 15 for human SN DA neurons), $cDNA \text{ fraction}$ = fraction of the UV laser microdissection complementary DNA reaction sample used as template in the individual quantitative PCR reactions (i.e. 5/17 for mouse, 1/11 for human substantia nigra dopaminergic neurons). The unit-magnitude corresponds to the respective standard used, which defines the unit at the y-intercept (i.e. pg equivalents of standard complementary DNA, derived from substantia nigra tissue messenger RNA). As primers for D1 and D5 dopamine receptors were not intron-spanning, a DNase digestion step (DNA-free[™]kit, Ambion[®]) of UV laser microdissected samples was included before complementary DNA synthesis (Liss and Roeper, 2004).

In vitro electrophysiological recordings from juvenile and adult mouse brain slices

Mouse brain slice preparations and *in vitro* electrophysiological patch clamp recordings from SN DA neurons in 250 μm coronal midbrain slices of juvenile and adult mice were carried out as described (Lammel *et al.*, 2008). To block fast synaptic transmission, 10 μM DNQX (6,7 dinitroquinoxaline-2,3-dione, Tocris) and 10 μM gabazine (SR95531 hydrobromide, Tocris) were added to artificial CSF. To analyse dopamine auto-inhibition of SN DA neurons *in vitro*, a minimum of 5 min stable baseline activity either in gramicidine 100 μg/ml (Sigma) perforated patch or in cell-attached configuration were recorded. This was followed by application of dopamine hydrochloride

(Sigma) in artificial CSF via bath-perfusion for 15 min and a 20 min washout. Whole-cell recordings were performed with patch pipettes containing 0.1 mM EGTA (Sigma) in internal solution (135 mM K-glucuronate, 5 mM KCl, 10 mM HEPES, 2 mM MgCl₂, 0.1 mM NaGTP, 5 mM MgATP, all from Sigma). To test the effects of dopamine antagonists on SN DA neurons, we performed cell-attached recordings of baseline activity for 10 min, followed by application of 100 μM sulpiride (Tocris) in artificial CSF for another 10 min via bath perfusion. For L-type-Ca²⁺ channel and NCS-1 pharmacology, slices were pre-incubated for at least 40 min in artificial CSF containing 300 nM isradipine (Tocris) or 10 μM D2R/NCS-1 interfering peptide (DNIP)/scrambled DNIP peptide (Genscript; Saab *et al.*, 2009) before and during cell-attached recordings. GIRK currents in SN DA neurons were evoked by local application (10 s) of 100 μM dopamine hydrochloride (Sigma) and recorded in voltage-clamp whole-cell configuration (membrane potential clamped at −50 mV, analysed with digital filter 500 mHz) in the presence of 100 μM tolbutamide (Sigma) and 20 μM ZD 7288 (Tocris) to block ATP-sensitive potassium channels (K-ATP) and hyperpolarization-activated cyclic nucleotide-gated channels, respectively. Excitatory postsynaptic currents were recorded at −70 mV and +40 mV from SN DA neurons in whole-cell patch-clamp, and AMPAR/NMDAR ratios were calculated as described previously (Lammel *et al.*, 2011). Half maximal effective concentrations (EC₅₀ values) of dopamine for activity inhibition were calculated from individual cells by applying increasing dopamine concentrations (0.001–100 μM, Sigma), each for 3 min. Mean frequency at each concentration was normalized to the respective baseline frequency for each cell. Individual and mean concentration–response curves were fitted to the Hill equation (curves fitted with constraints; max = 1, min = 0) to obtain EC₅₀ values. For functional SN DA cell-type identification, cells were re-patched, and SN DA neuron-specific basal electrophysiological properties (frequency, action potentials, and sag component) were recorded in whole-cell configuration (Lammel *et al.*, 2008; Ungless and Grace, 2012).

Immunoelectron microscopy

Double-labelling immunoelectron microscopy was performed and relative abundance of GIRK2 and D2-autoreceptor immunoreactivity in TH-positive SN DA neurons in control and cocaine injected juvenile wild-type mice was determined by quantification of immunolabelling in 60 μm coronal slices, as described by Koyrakh *et al.* (2005). Primary monoclonal antibody anti-TH (1–2 μg/ml, Calbiochem) was visualized by immunoperoxidase reaction and primary guinea pig polyclonal anti-GIRK2 and rabbit polyclonal anti-D2R antibody (not splice variant specific, 1–2 μg/ml; Koyrakh *et al.*, 2005; Narushima *et al.*, 2006; Aguado *et al.*, 2008) by the silver-intensified immunogold reaction. Secondary antibody mixtures included goat anti-rabbit (Fab fragment; diluted 1:100) coupled to 1.4 nm gold (Nanoprobes), goat anti-guinea pig (Fab fragment; diluted 1:100) coupled to 1.4 nm gold (Nanoprobes), and biotinylated goat anti-mouse (diluted 1:100; Vector Laboratories). Electron photomicrographs for ultrastructural analyses were captured with a CCD camera (Mega View III; Soft Imaging System) using a Jeol-1010 electron microscope (Jeol).

Data analysis

Data analysis and graphical representations were performed using FitMaster software (HEKA Electronics), GraphPad Prism 5 (GraphPad Software, Inc.), Igor Pro 6 (Wavemetrics Inc.) and Neuroexplorer (Nex Technologies) for electrophysiological experiments and SDS 2.3 (Applied Biosystems) for quantitative PCR. Statistical analysis was performed in Igor Pro 6, the R project for statistical computing and

GraphPad Prism 5 (GraphPad Software, Inc.). Statistical significance was set at $P < 0.05$. Mann-Whitney tests were performed in GraphPad Prism 5 software and used to determine statistical differences ($*P < 0.05$, $**P < 0.01$ and $***P < 0.001$). The coefficient of variation of the interspike interval was calculated using a Gaussian fit over the histogram with Neuroexplorer and Igor Pro 6. Coefficient of variation as [%] = (SD fit) / (mean fit) × 100. If not otherwise stated all data are given as mean ± SEM.

Results

Substantia nigra dopaminergic neurons of patients with Parkinson's disease display elevated levels of D2-autoreceptors and GIRK2 messenger RNA compared with controls

Given the important therapeutic role of D2 receptor agonists in Parkinson's disease, and of D2-autoreceptors and their main downstream target GIRK2 for D2-autoreceptor responses that control substantia nigra dopamine function in health and disease, we quantified messenger RNA levels of D2-autoreceptors and GIRK2 in laser-microdissected human SN DA neurons from *post-mortem* Parkinson's disease brains and controls using reverse transcription real-time quantitative PCR (Fig. 1A and B, and Supplementary Table 1). We detected significantly higher messenger RNA levels for D2-autoreceptors and GIRK2 messenger RNA in remaining SN DA neurons from Parkinson's disease brains compared to those of controls (Fig. 1B). D2-autoreceptor/GIRK2 ratios were not changed (Fig. 1C). These findings point to an altered, sensitized D2-autoreceptor/GIRK2 signalling control of activity of SN DA neurons in Parkinson's disease. To address and define a possible dopamine-dependent homeostatic plasticity of D2-autoreceptor/GIRK2 pacemaker control in SN DA neurons, we analysed isolated somatodendritic dopamine D2-autoreceptor responses in mouse SN DA neurons. As age-dependent, postnatal differences in mouse substantia nigra dopamine pacemaker control have been described (Chan *et al.*, 2007), we analysed juvenile (postnatal Day 13) as well as adult (postnatal Day 90) mice.

Dopamine D2-autoreceptor-mediated pacemaker inhibition of SN DA neurons is pronounced after postnatal maturation

We compared the unperturbed kinetics and dopamine sensitivities of D2-autoreceptor-mediated pacemaker inhibition in SN DA neurons after postnatal maturation in *in vitro* brain slices of juvenile (postnatal Day 13) and adult (postnatal Day 90) mice (Fig. 2A and B). Stimulation of somatodendritic D2-autoreceptors through dopamine bath application reversibly inhibited spontaneous pacemaker activity after initial membrane potential hyperpolarization (Fig. 2A). To ensure full D2-autoreceptor activation, we applied 100 μM dopamine for 15 min. The kinetics of the dopamine-evoked D2-autoreceptor-




Figure 1 Elevated D2-autoreceptor (AR) and GIRK2 messenger RNA levels in remaining human SN DA neurons from patients with Parkinson's disease (PD). (A) Neuromelanin-positive human SN DA neurons (SN DA) (before and after UV laser microdissection) from *post mortem* Parkinson's disease and control midbrain sections (for details, see Supplementary Table 1). (B) Real-time quantitative PCR results identified significantly higher relative messenger RNA levels for D2-autoreceptors and *GIRK2* in human SN DA neurons from Parkinson's disease brains compared to controls (ctr) (D2-autoreceptors: controls, $n = 26$, 1 ± 0.19 ; Parkinson's disease, $n = 39$, 4.25 ± 0.73 ; $****P < 0.0001$; Mann-Whitney test; *GIRK2*: controls, $n = 25$, 1 ± 0.25 ; Parkinson's disease, $n = 34$, 6.31 ± 1.12 ; $****P < 0.0001$; Mann-Whitney test). (C) D2/GIRK2 ratios were not different in SN DA neurons from controls and Parkinson's disease patients. All data are shown as the mean \pm SEM.

mediated pacemaker inhibition differed significantly between adult and juvenile SN DA neurons (Fig. 2A and B). Adult SN DA neurons did not recover pacemaker activity in the presence of saturating dopamine concentrations. However, SN DA neurons from juvenile mice displayed a prominent desensitization of the dopamine-induced pacemaker inhibition, regaining 40% of their baseline firing frequency at the last minute of dopamine application (Fig. 2E and Supplementary Table 3). The sensitivity of SN DA neurons to the dopamine-induced inhibition was ~ 2 -fold higher in adult neurons compared to those of juvenile mice (dopamine EC_{50} values: juvenile, $n = 10$: $6.57 \pm 1.09 \mu\text{M}$; adult, $n = 11$: $3.29 \pm 0.48 \mu\text{M}$; $P = 0.03$). The maximal dopamine-induced membrane hyperpolarization was not significantly different in juvenile and adult SN DA neurons (Supplementary Table 4). Pacemaker frequency and its precision (expressed as coefficient of variation of the interspike interval, in %) were similar between adult and juvenile SN DA neurons under basal conditions in the absence of dopamine (Fig. 2C and D, and Supplementary Table 5). Inhibition of D2-autoreceptor function by the D2 antagonist sulpiride ($100 \mu\text{M}$) had no effect on pacemaker frequency or precision of juvenile or adult SN DA neurons, arguing against an intrinsic dopamine tonus, and thus D2-autoreceptor activity *in vitro* under our recording conditions (Fig. 2C and D, and Supplementary Table 6).

A single dose of L-DOPA or cocaine *in vivo* induces adult-like, stable D2-autoreceptor responses in SN DA neurons *in vitro*

We next studied possible effects of pathophysiologically and Parkinson's disease-therapeutically relevant transient high-

dopamine states on mouse substantia nigra dopamine D2-autoreceptor responses by injecting them *in vivo* either with L-DOPA (the precursor of dopamine, and still the gold standard in Parkinson's disease therapy), cocaine (enhancing extracellular dopamine levels by inhibiting dopamine re-uptake through the plasmalemma dopamine transporter; Ritz *et al.*, 1987), or saline (for controls), and analysed D2-autoreceptor responses 4 or 3 days later, respectively. L-DOPA and cocaine pretreatment strongly affected D2-autoreceptor responses in juvenile SN DA neurons (Fig. 2A, B and E): it converted their desensitizing D2-autoreceptor responses into non-desensitizing D2-autoreceptor responses that resembled the adult phenotype (Fig. 2A, B and E, and Supplementary Table 3). This reduction of desensitization of D2-autoreceptor responses after *in vivo* induction of transient high-dopamine states was a hallmark of juvenile SN DA neurons and was not observed in juvenile (mesolimbic) ventral tegmental area dopaminergic neurons (Fig. 2E and Supplementary Table 3). Neither *in vivo* L-DOPA nor cocaine treatment significantly altered the basal spike-frequency or spike-timing precision (coefficient of variation of the interspike interval) of juvenile or adult substantia nigra dopamine pacemaker (Fig. 2C and D, and Supplementary Table 5). As L-DOPA and cocaine injections both induce transient high-dopamine states by a distinct molecular mechanisms, but had the same effects on juvenile substantia nigra dopamine D2-autoreceptor responses, our data strongly suggest that D2-autoreceptor responses show a dopamine-dependent plasticity. As the cocaine injection protocol, in contrast to the L-DOPA protocol, is well-established for mice (see 'Materials and methods' section for details), we have continued our study with the cocaine paradigm to induce transient *in vivo* high-dopamine states (Saal *et al.*, 2003).




Figure 2 *In vivo* transient high-dopamine states (induced by L-DOPA or cocaine) cause adult-like non-desensitizing dopamine D2-autoreceptor responses in juvenile SN DA neurons (SN DA). (A) Representative SN DA neuron pacemaker activity recordings from *in vitro* brain slices of adult and juvenile wild-type mice, pretreated (4 or 3 days, respectively) *in vivo* with saline, L-DOPA, or cocaine (perforated/cell-attached patch-clamp). Dopamine bath application is indicated by red bars. (B) Normalized frequencies plotted against time for all analysed SN DA neurons revealed pronounced desensitization of D2-autoreceptor responses in juveniles ($n = 14$) compared to adults ($n = 10$), as well as adult-like, non-desensitizing D2-autoreceptor responses in juvenile SN DA neurons after *in vivo* L-DOPA ($n = 15$) or *in vivo* cocaine (coc; $n = 12$), compared to controls (ctr; saline, $n = 14$). (C) Bar graphs show no change in mean frequencies before and after sulpiride application in SN DA neurons from controls, as well as after saline, L-DOPA, or cocaine *in vivo* injections, before dopamine application. (D) Pacemaker precision [given as coefficient of variation of interspike interval values, (CV ISI)] of juvenile and adult SN DA

(continued)

Functional expression of D2-auto-receptors and GIRK2 is not altered in juvenile SN DA neurons after *in vivo* induction of a transient high-dopamine state

We next addressed the underlying molecular mechanism of altered, adult-like D2 autoreceptor responses in juvenile SN DA neurons after *in vivo* induction of a transient high-dopamine state by cocaine. We confirmed that mouse SN DA neurons predominantly express GIRK2 messenger RNA alone, together with D2 dopamine receptor messenger RNA (Fig. 3A, D2 receptor and GIRK2 were co-expressed in 100% of analysed SN DA neurons, $n = 85$, without other GIRKs in $\sim 80\%$). Other dopamine receptor subunits were not detected or only in a small fraction of SN DA neurons (Fig. 3A; D1: $n = 1$, D5: $n = 2$, D3: $n = 3$, D4: $n = 0$ out of 39 TH- and D2-positive SN DA neurons). Although most SN DA neurons co-expressed both the short D2s and the long D2I receptor-splice variants ($\sim 65\%$, Fig. 3A), 35% expressed only one splice variant (only D2I: $n = 23$ of 85; only D2s: $n = 6$ of 85). As all analysed SN DA neurons showed typical functional D2-autoreceptors responses, we conclude that both D2s and D2I alone can act as functional D2-autoreceptors in mouse SN DA neurons. To probe whether the relative expression of the D2-autoreceptor splice variants in SN DA neurons was altered after cocaine injection, we quantified D2 short and long variants, and calculated D2I/s ratios in SN DA neurons from juvenile mice with and without cocaine treatment. We found no change in D2s or D2I messenger RNA levels or D2I/s ratios in juvenile SN DA neurons after cocaine (Fig. 3A). We also quantified D3 and D4 messenger RNA in SN DA neurons from juvenile mice without and after cocaine. However, we detected no expression of D3 or D4 using real-time quantitative PCR, neither in controls or after cocaine ($n = 6$ probes each, data not shown). These data rule out that altered expression of D2 splice variants or other D2-type receptors (D3, D4) are causing the observed functional change in substantia nigra dopamine D2-autoreceptor desensitization after *in vivo* cocaine. The reduced desensitization of D2-autoreceptors in juvenile SN DA neurons after *in vivo* induced transient high-dopamine states was also not associated with an enhanced expression of either D2-autoreceptors or GIRK2 (Fig. 3). D2-autoreceptor messenger RNA and immunoreactivity (Fig. 3A and B) was unaffected after cocaine. GIRK2 messenger RNA (Fig. 3A) and its functional channel activity (Fig. 3C) were downregulated rather than upregulated. We also detected no change in the dopamine transporter and vesicular monoamine transporter (*Vmat2*, also known as *Slc18a2*) messenger RNA ratio in juvenile SN DA neurons after

cocaine, the transporters that define dopamine uptake and release capacity, respectively (Fig. 3D).

Ca_v1.3 L-type calcium channel function and internal Ca²⁺ is crucial for non-desensitizing D2-autoreceptor responses induced by high-dopamine states *in vivo*, in juvenile SN DA neurons

As calcium signalling is not only critical for the control of D2-receptor desensitization in general, but also triggers selective substantia nigra dopamine degeneration in Parkinson's disease, and as L-type-Ca²⁺ channel-blockers protect for Parkinson's disease in humans and mice, we evaluated a potential role of internal calcium ([Ca²⁺]_i) and L-type-Ca²⁺ channels for the expression of the high dopamine-dependent, non-desensitizing D2-autoreceptors in juvenile SN DA neurons. We first tested if the cocaine effect on D2-autoreceptors depended on unperturbed [Ca²⁺]_i by buffering [Ca]_i with 0.1 mM EGTA using whole cell dialysis (Fig. 4A and Supplementary Table 3). EGTA buffering did not affect the kinetics of dopamine autoinhibition in juvenile substantia nigra dopamine neurons in saline treated animals compared to those recorded in perforated patch. However, after cocaine treatment, D2-autoreceptor responses still displayed prominent desensitization, in contrast with the non-desensitizing responses after cocaine recorded in perforated patch (Fig. 4A). These data indicate a complex dependence of cocaine-induced D2-autoreceptor potentiation on intracellular Ca²⁺. They also emphasize the importance of analysing D2-autoreceptors signalling in non-perturbed recording configurations, as the cocaine effect is absent in whole cell recordings.

We next tested if L-type-Ca²⁺ channels could provide the Ca²⁺ source for Ca²⁺-dependent control of D2-autoreceptor responses, by continuous treatment of slices with the selective L-type Ca²⁺ channel blocker isradipine (300 nM), which is already in clinical trials as neuroprotective therapy for Parkinson's disease (Striessnig *et al.*, 2006; Simuni *et al.*, 2010; Surmeier *et al.*, 2012), before, during and after dopamine application. Isradipine completely abolished the effect of *in vivo* cocaine treatment on D2-autoreceptor desensitization in juvenile SN DA neurons (Fig. 4B and F, and Supplementary Table 3), revealing a crucial role of these channels for the expression of the here described cocaine effect.

Another important observation was that although 300 nM isradipine modified the cocaine response, it did not significantly reduce basal pacemaker frequency or its precision in juvenile or adult SN DA neurons (Fig. 4D and E, and Supplementary Table 5).

Figure 2 Continued

neurons was also not altered after sulpiride application, as well as after saline, L-DOPA, or cocaine *in vivo* injections. (E) D2-autoreceptor responses of mouse substantia nigra and ventral tegmental area dopaminergic neurons represented as mean activity of respective dopaminergic neurons at the last minute of dopamine application (15 min). Data are shown as the mean \pm SEM. Data values detailed in Supplementary Tables 3, 5 and 6.




Figure 3 Functional expression of D2-autoreceptors (AR) [D2 short/long variants (s/l), D3, D4] and GIRK2 is not altered in juvenile SN DA neurons after *in vivo* induction of a transient high-dopamine state. (A, *top*) Agarose gel electrophoresis of multiplex nested reverse transcription PCR products identified D2 receptor (both splice variants) and GIRK2 as the molecular components of the D2-autoreceptors in SN DA neurons. Note the absence of D3 and D4 receptor, as well as GIRK1. *Bottom*: Quantitative analysis (reverse transcription quantitative PCR) of laser-microdissected (LMD) juvenile SN DA neurons showed no change in the ratio of D2 long and D2 short splice variant expression levels, after cocaine in single SN DA neurons (saline, $n = 15$, 1 ± 0.12 ; cocaine, $n = 15$, 0.85 ± 0.07 ; $P = 0.29$; Mann-Whitney test). Further, no changes in D2-autoreceptors (relative messenger RNA levels normalized to saline controls: saline, $n = 18$, 1 ± 0.14 ; cocaine, $n = 19$, 0.90 ± 0.08 ; $P = 0.73$; Mann-Whitney test), D3 or D4 receptor messenger RNA (data not shown, as no signal was detected. D3 receptor: saline, $n = 6$; cocaine: $n = 6$; D4 receptor: saline, $n = 6$; cocaine, $n = 6$) were detected. *GIRK2* messenger RNA levels were also decreased rather than increased after *in vivo* cocaine (saline, $n = 19$, 1 ± 0.15 ; cocaine, $n = 20$, 0.62 ± 0.07 ; $*P = 0.04$; Mann-Whitney test; note the significantly altered D2-autoreceptor/GIRK2 ratios after cocaine due to decreased GIRK2-levels; saline, $n = 18$, 1 ± 0.08 ; cocaine, $n = 19$, 1.57 ± 0.12 ; $**P = 0.002$; Mann-Whitney test). All data given as mean \pm SEM normalized to controls. (B, *left*) Pre-embedding double-immunolabelling at electron microscopy level for D2-autoreceptors and GIRK2 in TH-positive SN DA neurons 3 days after saline/cocaine *in vivo* injection of juvenile wild-type mice ($n = 3$ animals each; arrows point to immunogold labelling

(continued)

In SN DA neurons, the pore-forming α 1-subunits of both brain L-type-Ca²⁺ channels, Ca_v1.2 and Ca_v1.3, are expressed (Olson *et al.*, 2005; Sinnegger-Brauns *et al.*, 2009). Since isradipine inhibits both isoforms, we cannot distinguish which of these channels mediate its pharmacological effect on SN DA neurons. To address this question, we took advantage of homozygous Ca_v1.2 DHP^{-/-} mutant mice. These mice express Ca_v1.2 L-type-Ca²⁺ channels with normal function, but their high sensitivity to isradipine is eliminated by a single mutation within the DHP-binding pocket (Sinnegger-Brauns *et al.*, 2004; Striessnig and Koschak, 2008). Thus, Ca_v1.2-mediated isradipine effects are absent in these mice, whereas drug effects through Ca_v1.3 are unaffected. In SN DA neurons from Ca_v1.2 DHP^{-/-} mutant mice the effect of isradipine was indistinguishable from wild-type (Fig. 4C and F, and Supplementary Table 3). As the isradipine concentration used was very low (300 nM), and no other targets are known to be affected at this concentration, Ca_v1.3 remains the only plausible molecular transducer for the cocaine-induced, calcium-dependent, and isradipine-sensitive change in substantia nigra dopamine D2-autoreceptor responses.

We also assessed the role of Ca_v1.3 more directly in Ca_v1.3-deficient mice (Ca_v1.3 knockout mice) (Platzer *et al.*, 2000). In this model, although basal substantia nigra dopamine pacemaker activity remained unaffected, the constitutive absence of Ca_v1.3 activity throughout development prevented the postnatal wild-type substantia nigra dopamine D2-autoreceptor phenotypes, as the non-desensitizing 'adult' D2-autoreceptor responses were present in SN DA neurons from both juvenile as well as adult Ca_v1.3 knockout mice (Supplementary Fig. 1 and Supplementary Table 7). Because of absence of the juvenile desensitizing D2-autoreceptor phenotype, we did not further study *in vivo* cocaine effects in Ca_v1.3 knockout mice.

Taken together the findings from both mouse models reveal a novel functional role of Ca_v1.3 L-type-Ca²⁺ channels for both the development of age-dependent D2-autoreceptor differences in SN DA neurons, and for the switch of juvenile desensitizing D2-autoreceptor response to those of adult substantia nigra dopamine neurons in response to a transient *in vivo* high-dopamine state.

At the messenger RNA level, we found no change in expression of Ca_v1.2 and Ca_v1.3 L-type-Ca²⁺ channel pore-forming α 1-subunits in mouse SN DA neurons after *in vivo* cocaine (Fig. 4G). Also, we found no change in expression levels of Ca_v1.2 and

Ca_v1.3 in remaining human SN DA neurons of patients with Parkinson's disease compared to controls (Fig. 4G).

As Ca_v1.2 and Ca_v1.3 expression were not altered, we probed for altered AMPA/NMDAR ratios in juvenile cocaine-treated SN DA neurons after *in vivo* cocaine as an alternative calcium source, causing synaptic plasticity after cocaine in adult ventral tegmental area dopamine neurons (Ungless *et al.*, 2001; Lammel *et al.*, 2011). However, we detected no change in AMPAR/NMDAR-mediated excitatory postsynaptic currents after *in vivo* cocaine in juvenile SN DA neurons (Supplementary Fig. 2).

Cocaine induced, adult-like, D2-autoreceptor responses in juvenile substantia nigra dopaminergic neurons are mediated by NCS-1

We next explored a possible contribution of NCS-1 for Ca_v1.3 dependent control of D2-autoreceptor responses in juvenile SN DA neurons, in response to *in vivo* transient high-dopamine, induced by cocaine. NCS-1 can directly interact with postsynaptic D2 receptors in a Ca²⁺-dependent manner and inhibit receptor desensitization by preventing its phosphorylation through G protein-coupled receptor kinase 2 (GRK2, also known as ADRBK1), and thus beta-arrestin-mediated receptor internalization (Kabbani *et al.*, 2002). We analysed mouse substantia nigra dopamine D2-autoreceptor responses in the presence of a membrane-permeable peptide, DNIP (D2R/NCS-1 interfering peptide), which physically blocks NCS-1 binding to D2 receptor (Saab *et al.*, 2009). As control, a membrane-permeable, scrambled version of the peptide (scrambled DNIP) was used, which did not influence basal pacemaker activity or the kinetics of the D2-autoreceptor response of SN DA neurons after *in vivo* cocaine (Figs 5A, 5B and Supplementary Tables 3 and 5). Similar to 300 nM of the substantia nigra dopamine- and Parkinson's disease-protective L-type-Ca²⁺ channel-blocker isradipine, 10 μ M DNIP completely abolished the cocaine effect on D2-autoreceptor responses in juvenile SN DA neurons. With NCS-1/D2-autoreceptor interaction blocked, D2-autoreceptors displayed rapid desensitization, similar to that seen without cocaine pretreatment (Fig. 5A and B, and Supplementary Table 3). To prove that NCS-1 function is necessary and sufficient for modulating D2-autoreceptor desensitization in SN DA neurons, we analysed D2-autoreceptor responses in a

Figure 3 Continued

at extrasynaptic plasma membrane and crossed arrows at intracellular sites). *Right*: No significant differences in immunoreactivity or subcellular localization of D2-autoreceptor protein ($n = 3$ animals each; saline, 1107.3 ± 8.7 ; cocaine, 1034.3 ± 33.5 ; $P = 0.2$; Mann-Whitney test), and GIRK2 protein ($n = 3$ animals each; saline, 1408.3 ± 9.1 ; cocaine, 1396.3 ± 7.1 ; $P = 0.4$; Mann-Whitney test). (C, *left*) Representative traces of dopamine evoked GIRK currents in juvenile SN DA neurons recorded in whole-cell voltage-clamp *in vitro* in the presence of 100 μ M tolbutamide and 20 μ M ZD7288 to block ATP-sensitive potassium channels (K-ATP) and hyperpolarization-activated cyclic nucleotide-gated (HCN) channels. *Right*: Mean dopamine-evoked GIRK current amplitudes were not increased, but significantly reduced after *in vivo* cocaine in juvenile SN DA neurons (saline, $n = 12$, 101.48 ± 7.69 pA; cocaine, $n = 10$, 75.88 ± 12.74 pA; $*P = 0.03$; Mann-Whitney test). (D) DAT/VMAT2 messenger RNA ratios in juvenile SN DA neurons, defining their individual dopamine-release capacity, were not altered after *in vivo* cocaine (normalized ratios: saline, $n = 8$, 1 ± 0.14 ; cocaine, $n = 8$, 1.23 ± 0.13 ; $P = 0.29$; Mann-Whitney test). Data are given as mean \pm SEM.




Figure 4 Ca^{2+} and $\text{Ca}_v1.3$ L-type- Ca^{2+} channels are crucial for adult-like, non-desensitizing D2-autoreceptor responses, induced by a transient high-dopamine state *in vivo*, in juvenile SN DA neurons. (A) Mean activity of substantia nigra dopaminergic neurons at the last minute of dopamine application [15 min, recorded in perforated patch (pp) juvenile: saline, $n = 14$; cocaine, $n = 12$] compared to those recorded in whole cell (wc) configuration with 0.1 mM EGTA (whole cell: juvenile: saline, $n = 6$; cocaine, $n = 7$) to buffer free $[\text{Ca}^{2+}]_i$ in juvenile SN DA neurons. Note the absence of cocaine-induced stabilization of D2-autoreceptor responses in whole cell compared to unperturbed perforated patch recordings. (B) Experiments as in Fig. 1 but in the presence of the L-type- Ca^{2+} channel blocker isradipine (300 nM, represented with blue bar). Isradipine completely blocked changes in kinetics of D2-autoreceptor desensitization of SN DA neurons, after transient high-dopamine state induced by cocaine, in wild-type mice (saline, $n = 7$; cocaine, $n = 7$). (C) Similar experiments as in B, but using $\text{Ca}_v1.2$ DHP^{-/-} mice (with $\text{Ca}_v1.2$ L-type- Ca^{2+} channels, insensitive to isradipine; saline, $n = 6$; cocaine, $n = 9$). Application of 300 nM isradipine still abolished cocaine D2-autoreceptor response potentiation, pointing to the crucial role of $\text{Ca}_v1.3$

(continued)

general NCS-1^{-/-} mouse, lacking NCS-1 in all cells. Substantia nigra dopaminergic neurons of NCS-1^{-/-} mice displayed rapidly desensitizing D2-autoreceptors responses already under control conditions that were not converted to a sensitized response after *in vivo* cocaine, confirming that NCS-1 is necessary and sufficient for preventing D2-autoreceptor desensitization in mouse SN DA neurons (Fig. 5A and B, and Supplementary Tables 3 and 5).

To address why Ca²⁺-dependent NCS-1/D2-autoreceptors interactions seem to be more prominent after cocaine (and thus D2-autoreceptors desensitization was reduced), we quantified NCS-1 messenger RNA. We found that NCS-1 messenger RNA levels were ~2-fold higher in juvenile SN DA neurons after *in vivo* cocaine ($P = 0.0001$; Fig. 5C). This could explain enhanced L-type-Ca²⁺ channel-dependent Ca²⁺-stimulated NCS-1/D2-autoreceptor signalling, without increased expression of L-type-Ca²⁺ channels, but due to increased NCS-1 availability. This would allow SN DA neurons to adapt the degree of their dopamine D2-autoreceptor activity to homeostatic dopamine challenges, by tuning NCS-1 expression. To address if a similar increased NCS-1 expression is also present in human SN DA neurons in Parkinson's disease, hinting to a corresponding homeostatic adaptive change in D2-autoreceptors signalling, we quantified messenger RNA levels of NCS-1 in human SN DA neurons from controls and endstage Parkinson's disease brains (Fig. 5C, right). Indeed, similar to mice after cocaine, we detected ~2-fold higher messenger RNA levels of NCS-1 in remaining SN DA neurons from Parkinson's disease brains (Fig. 5C). Thus, increased NCS-1/D2-autoreceptor signalling, downstream of Ca_v1.3, may also fine-tune human substantia nigra dopamine neuron activity in Parkinson's disease. It might (compensatory) allow remaining dopamine to reduce firing, activity-associated rhythmic cellular calcium load, and thus vulnerability of SN DA neurons.

Discussion

In summary, our data indicate a flexible D2-autoreceptor signalling in SN DA neurons, controlling their pacemaker activity, during maturation, and in response to pathophysiological *in vivo* dopamine challenges in mice, which could also be translated to human Parkinson's disease pathology. Our findings describe the neuronal calcium sensor NCS-1 as a novel link between two major molecular players, modulating the activity of SN DA neurons as well as their vulnerability to degeneration in Parkinson's disease: Ca_v1.3

L-type-Ca²⁺ channels and D2-autoreceptors (Beaulieu and Gainetdinov, 2011; Surmeier *et al.*, 2012; Ford, 2014). We show that the degree of somatodendritic D2-autoreceptor desensitization in juvenile mouse SN DA neurons, in response to *in vivo* transient high-dopamine states, depends on Ca_v1.3 channels, as well as on the calcium dependent NCS-1 interaction with D2-autoreceptors. Our findings link flexible D2-autoreceptor desensitization with Ca_v1.3 L-type-Ca²⁺ channel function, through functional expression of NCS-1, and point to a role of this signalling network for adaptive changes in substantia nigra dopamine neuron activity due to altered extracellular dopamine levels. Our data also provide a fresh look at the molecular mechanisms involved in substantia nigra dopamine neuron activity control, as well as their pathophysiology in Parkinson's disease and its dopamine-mimetic therapy.

Dopamine D2 receptors are important for proper function of dopamine midbrain neurons, and are important pharmacological targets for therapy of schizophrenia (antagonist; Ginovart and Kapur, 2012), and in particular Parkinson's disease (agonist; Yamamoto and Schapira, 2008). Somatodendritic D2-autoreceptors are coupled to GIRK2 potassium channels, and reduce activity and excitability of SN DA neurons in response to locally released dopamine in the midbrain (Beckstead and Williams, 2007; Gantz *et al.*, 2013; Ford, 2014). Dysfunction of these somatodendritic D2-autoreceptors is linked to Parkinson's disease pathology (Ford, 2014); however their exact functions in health and disease are by far not as well examined as the function of pre- and particularly postsynaptic D2 receptors (Beckstead *et al.*, 2007; Ford *et al.*, 2010; Bello *et al.*, 2011; Anzalone *et al.*, 2012; Ford, 2014).

To date, only the short D2-autoreceptor splice variant, D2s has been identified as the D2-autoreceptor (Usiello *et al.*, 2000; Beaulieu and Gainetdinov, 2011; Ford, 2014). Our combined functional and molecular single cell analysis strongly suggests that the long D2l splice variant alone can build functional D2-autoreceptors in SN DA neurons. Recent reports confirm this claim as expression of either D2s or D2l alone in SN DA neurons of D2 receptor knockout mice restored D2 agonist-induced activation of GIRKs (Neve *et al.*, 2013). In accordance with the literature, reviewed in Ford (2014), we also show that the other D2-type receptors (D3, D4) are not important for D2-autoreceptor function, or its homeostatic plasticity in mouse SN DA neurons. We did not quantify excitatory acting D1-type dopamine receptors (D1 and D5), as their messenger RNAs were almost never detected in SN DA neurons (Fig. 3A), they are not coupled to

Figure 4 Continued

L-type-Ca²⁺ channels in this process. (D and E) Isradipine (300 nM) did not reduce basal pacemaker frequencies or activity pattern (coefficient of variation of interspike interval) of juvenile SN DA neurons. (F) Activity of SN DA neurons at the last minute of dopamine application. Data values detailed in Supplementary Tables 3 and 5. (G) Quantitative analysis (reverse transcription quantitative PCR) of laser-microdissected (LMD) mouse and human SN DA neurons detected no change in messenger RNA levels of Ca_v1.2 or Ca_v1.3, the pore-forming alpha subunit of L-type-Ca²⁺ channels in the brain, after *in vivo* cocaine (mice, *left*) or in Parkinson's disease compared to controls (humans, *right*). Normalized expression levels: mouse: Ca_v1.2: saline, $n = 18$, 1 ± 0.01 ; cocaine, $n = 18$, 1.01 ± 0.02 ; $P = 0.92$; Ca_v1.3: saline, $n = 17$, 1 ± 0.17 ; cocaine, $n = 18$, 0.81 ± 0.16 ; $P = 0.37$; Mann-Whitney test. Human: Ca_v1.2: control, $n = 4$, 1 ± 0.01 ; Parkinson's disease, $n = 5$, 0.90 ± 0.11 ; $P = 0.43$; Ca_v1.3: control, $n = 16$, 1 ± 0.16 ; Parkinson's disease, $n = 33$, 1.14 ± 0.14 ; $P = 0.64$; Mann-Whitney test). Data are shown as mean \pm SEM.




Figure 5 Adult-like, non-desensitizing D2-autoreceptor responses in SN DA neurons are mediated by the neuronal calcium sensor 1 (NCS-1). (A, left) Normalized frequency plots show that non-desensitizing D2-autoreceptor responses of juvenile SN DA neurons after *in vivo* cocaine were completely blocked by 10 μ M DNIP (cell-permeant D2R/NCS-1 interfering peptide, $n = 8$), whereas the scrambled version had no effect (10 μ M srDNIP, $n = 6$). (Right) Respective analysis (as in A) of a general NCS-1 knockout mouse (NCS-1^{-/-}) demonstrated that NCS-1 is crucial for non-desensitizing D2-autoreceptor responses of juvenile SN DA neurons. (B) Activity of SN DA neurons from (A) at the last minute (15 min) of dopamine application (note the significantly pronounced D2-autoreceptor desensitization in NCS-1^{-/-} already under control conditions). Data values detailed in Supplementary Table 3. (C) Reverse transcription quantitative PCR analysis of mouse and human SN DA neurons identified 2-fold higher *Ncs-1* messenger RNA levels after cocaine (saline, $n = 36$, 1 ± 0.08 ; cocaine, $n = 32$, 1.94 ± 0.15 ; $***P = 0.0001$; Mann-Whitney test), as well as in Parkinson's disease, compared with controls (control, $n = 27$, 1 ± 0.20 ; Parkinson's disease, $n = 33$, 2.60 ± 0.27 ; $****P < 0.0001$; Mann-Whitney test). (D) Cartoon depicting molecular components of a novel Ca_v1.3/NCS-1/D2-autoreceptor/GIRK2 signalling network in SN DA neurons, which allows homeostatic adaptation of their inhibitory dopamine D2 responses, and thus their pacemaker activity.

inhibitory GIRK2 receptors, and have only been functionally described postsynaptically (Vallone *et al.*, 2000; Beaulieu and Gainetdinov, 2011).

We have previously shown that D2-autoreceptor responses are not homogeneous within distinct types of substantia nigra and

ventral tegmental area dopaminergic neurons from adult mice (Lammel *et al.*, 2008). Now we show significantly elevated messenger RNA levels of D2-autoreceptors, as well as GIRK2 in remaining human SN DA neurons in Parkinson's disease, pointing to altered D2-autoreceptor function in the course of the disease. We

addressed a general possibility of a non-static, but plastic, D2-autoreceptor response in SN DA neurons in mice, and found that somatodendritic D2-autoreceptor responses of SN DA neurons are not uniform during postnatal maturation, but display prominent desensitization in juvenile mice (postnatal Day 13), which is absent in adults (postnatal Day 90). Furthermore we show that adult-like, non-desensitizing D2-autoreceptor responses could be induced in juvenile SN DA neurons by a transient high-dopamine state *in vivo* (via L-DOPA or cocaine). Most importantly, we identified that Ca_v1.3 L-type-Ca²⁺ channels are crucial for controlling dopamine-dependent D2-autoreceptor desensitization in mouse SN DA neurons, as they provide the calcium source for a Ca²⁺ dependent NCS-1/D2-autoreceptor interaction, which prevents D2-receptor desensitization (Beaulieu and Gainetdinov, 2011). That defines, to our knowledge, a first physiological function of L-type-Ca²⁺ channels in SN DA neurons. Our Ca_v1.3 knockout and Ca_v1.2 DHP^{-/-} analysis ruled out Ca_v1.2 and identified Ca_v1.3 as the crucial L-type-Ca²⁺ channel for this function.

The here identified role of Cav1.3 L-type-Ca²⁺ channels, for NCS-1 mediated modulation of D2-autoreceptors and activity of SN DA neurons, has not yet been described. Postsynaptically, in striatal medium spiny neurons, D2 receptors upstream of L-type-Ca²⁺ channels can reduce L-type-Ca²⁺ channel activity through a calcineurin signalling cascade (Hernandez-Lopez *et al.*, 2000; Olson *et al.*, 2005). Also postsynaptically, in hippocampal neurons, reduced desensitization of D2-receptors, due to NCS-1/D2-autoreceptor interaction and thereby prevention of phosphorylation through GRK2, and thus receptor internalization, has been reported (Jo *et al.*, 2008; Saab *et al.*, 2009). Accordingly, we hypothesize that after induction of transient elevated dopamine levels, by *in vivo* L-DOPA or cocaine, D2-autoreceptor responses desensitize less in juvenile SN DA neurons due to increased NCS-1 expression and its higher availability for binding to D2-autoreceptors, leading to reduced GRK2 phosphorylation and D2-autoreceptor internalization in response to prolonged dopamine exposure. The *in vitro* Ca²⁺ buffering experiments (EGTA) as well as the DNIP experiments and the analysis of the *Ncs1*^{-/-} mice further support this hypothesis. We hypothesize accordingly that in remaining SN DA neurons in Parkinson's disease, due to increased expression of NCS-1, D2 and GIRK2, inhibitory D2-autoreceptor responses are also sensitized.

At the messenger RNA level, Ca_v1.3 (or Ca_v1.2) expression was not increased, neither in mouse SN DA neurons after cocaine nor in human SN DA neurons in Parkinson's disease. However, our data do not exclude altered Ca_v1.3 activity. Evidence for enhanced Ca_v1.3 protein expression in SN DA neurons in Parkinson's disease compared to controls has been described in immunohistochemical experiments (Hurley *et al.*, 2013). However, although L-type-Ca²⁺ channel activity is necessary, increased calcium entry through L-type-Ca²⁺ channels is not necessarily required to account for the functional D2-autoreceptor plasticity observed in mice, as the detected higher NCS-1 levels would provide enhanced calcium-dependent NCS-1/D2 signalling, preventing desensitization. Indeed, NCS-1 expression is stimulated by calcineurin (Hamasaki-Katagiri and Ames, 2010), a phosphatase that itself is stimulated by D2 receptor activity (Greengard *et al.*, 1999). Thus, a transient increase in dopamine, that leads to increased

D2-autoreceptors stimulation, would be sufficient to increase calcium-dependent NCS-1 expression, e.g. via calcineurin-stimulation, and thus to increased calcium-dependent NCS-1/D2 interaction, in a negative feedback loop.

Our finding that pharmacological L-type-Ca²⁺ channel-block did not significantly reduce basal pacemaker frequency or its precision in juvenile or adult mouse SN DA neurons clearly argues—at least under our recording conditions—against a crucial role of L-type-Ca²⁺ channels in SN DA neurons for pacemaker activity. This disputes a previously reported switch from HCN channel to a metabolically more challenging L-type-Ca²⁺ channel-driven pacemaker in SN DA neurons, that would render them particularly vulnerable to degeneration during postnatal maturation (Bean, 2007; Chan *et al.*, 2007; Puopolo *et al.*, 2007; Drion *et al.*, 2011; Surmeier *et al.*, 2012)

We also detected no cocaine-induced changes in AMPA/NMDA ratios in juvenile SN DA neurons, as described for adult ventral tegmental area dopamine neurons (Ungless *et al.*, 2001; Lammel *et al.*, 2011). In contrast, we observed a cocaine-induced, Ca_v1.3, calcium and NCS-1-dependent substantia nigra-specific 'enhancement' of D2-autoreceptor function *in vitro*, which could act as an excitability brake and thus counteract or prevent excitotoxicity and degeneration of substantia nigra dopamine neurons (Dong *et al.*, 2009). The finding that SN DA neurons from PARK mice (with particularly high vulnerable SN DA neurons) display D2-autoreceptor responses with significantly enhanced desensitization (Goldberg *et al.*, 2005; Tong *et al.*, 2009) would support this interpretation.

Although L-type-Ca²⁺ channels seem not to be crucial for substantia nigra dopamine pacemaking, there is accumulating evidence that the pacemaker activity of SN DA neurons drives somatodendritic Ca²⁺ entry through L-type-Ca²⁺ channels, thus generating a neurotoxic Ca²⁺ load that triggers their degeneration (Chan *et al.*, 2007; Puopolo *et al.*, 2007; Khaliq and Bean, 2010; Surmeier and Schumacker, 2013). Reduction of substantia nigra dopamine activity through a sensitized D2-autoreceptor response could therefore counteract this neurotoxic Ca²⁺ load. This would especially occur in response to transient high dopamine states, induced e.g. by dopamine-mimetic therapy. Thus, Ca_v1.3/Ca²⁺ dependent NCS-1 modulation of D2-autoreceptor desensitization may emerge as a homeostatic response in human SN DA neurons in Parkinson's disease, reducing electrical activity and protecting surviving SN DA neurons from acute overexcitability (to compensate the progressive dopamine loss) and excitotoxicity (Dong *et al.*, 2009; Liss and Roeper, 2010; Callaghan *et al.*, 2012).

In this view, our data also raise an important issue regarding the Parkinson's disease protective effects of L-type-Ca²⁺ channel blockers in humans (Ritz *et al.*, 2010; Marras *et al.*, 2012; Pasternak *et al.*, 2012). We have no information about the therapy of our patients with Parkinson's disease; however, it is likely that they were treated with L-DOPA (and/or D2 agonist). If in Parkinson's disease, pathophysiology and/or dopamine-mimetic treatment induces—potentially protective—sensitized D2-autoreceptor responses in aged SN DA neurons, then this may be inhibited by treatment with L-type-Ca²⁺ channel blockers. The resulting enhancement of substantia nigra activity would counteract their reduction of overall Ca²⁺ load and related metabolic stress, believed to be the neuroprotective mechanism of

L-type-Ca²⁺ channel blockers in Parkinson's disease (Surmeier et al., 2012; Surmeier and Schumacker, 2013). However, if in adult SN DA neurons, another calcium source, distinct from L-type-Ca²⁺ channels, supports the NCS-1-dependent D2-autoreceptor responses (as we have first evidence for Ca_v1.3 knockout mice), this would not be the case, and it could offer another potential drug target for Parkinson's disease therapy.

As blood–brain barrier permissive L-type-Ca²⁺ channel blockers are already in clinical trials for neuroprotective Parkinson's disease therapy (ClinicalTrials.gov Identifier: NCT00909545; Simuni et al., 2010; Parkinson Study, 2013), but as their general neuroprotective mechanism, as well as the functional roles of L-type-Ca²⁺ channels in SN DA neurons are still unclear, our data provide an important first molecular step towards a better understanding of L-type-Ca²⁺ channel function in SN DA neurons, and thus for tailored pharmacological neuroprotective therapy. All molecular components of the Ca_v1.3/NCS-1/D2-autoreceptor signalling network are already accessible in humans to pharmacological modulation. However, they are all expressed, in addition to in SN DA neurons, in a variety of other neuronal and non-neuronal cells. Thus, future studies need to address the specific acute and chronic pathophysiological consequences, as well as side-effects of pharmacological modulation of Ca_v1.3/NCS-1/D2-autoreceptor-signalling in SN DA neurons.

Acknowledgements

We thank Desiree Spaich and Stefanie Schulz for excellent technical support, Olaf Pongs for providing the NCS-1 KO mice, and Leica Microsystems for providing the LMD6000. We thank Ema Ilijic and D James Surmeier for neurotoxic pilot experiments and discussions. We thank Frances M. Ashcroft, Dirk Isbrandt, and Jochen Roeper for discussion and/or critical reading the manuscript. ED, CP, BL contributed *in vitro* electrophysiology. SL, RM AMPA/NMDA ratios. JuS provided pilot *in vivo* electrophysiological experiments. FS, ED, MF, JD, AH, BL RT-PCR and analysis. MW provided D2R and GIRK2 antibodies, RL provided immunogold histochemistry. JS provided Ca_v1.2 DHP^{-/-} and Ca_v1.3 KO mice. ED, JS, and BL designed the study and wrote the manuscript. The authors declare no competing financial, personal, or professional interests that could be construed to have influenced our paper.

Funding

The study was supported by the DFG, SFB497 and LI 1745/1 (BL), the Austrian Science Fund (FWF SFB F4412, F4402; BL, JS), the BMBF (NGFN, 01GS08134, BL), the Spanish Ministry of Education and Science (BFU-2009-08404), and the CONSOLIDER program (CSD2008-00005) to RL, and the Alfred Krupp prize (BL).

Supplementary material

Supplementary material is available at *Brain* online.

References

- Aguado C, Colon J, Ciruela F, Schlaudraff F, Cabanero MJ, Perry C, et al. Cell type-specific subunit composition of G protein-gated potassium channels in the cerebellum. *J Neurochem* 2008; 105: 497–511.
- Anzalone A, Lizardi-Ortiz JE, Ramos M, De Mei C, Hopf FW, laccarino C, et al. Dual control of dopamine synthesis and release by presynaptic and postsynaptic dopamine D2 receptors. *J Neurosci* 2012; 32: 9023–34.
- Bean BP. Neurophysiology: stressful pacemaking. *Nature* 2007; 447: 1059–60.
- Beaulieu JM, Gainetdinov RR. The physiology, signaling, and pharmacology of dopamine receptors. *Pharmacol Rev* 2011; 63: 182–217.
- Beckstead MJ, Ford CP, Phillips PE, Williams JT. Presynaptic regulation of dendrodendritic dopamine transmission. *Eur J Neurosci* 2007; 26: 1479–88.
- Beckstead MJ, Williams JT. Long-term depression of a dopamine IPSC. *J Neurosci* 2007; 27: 2074–80.
- Bello EP, Mateo Y, Gelman DM, Noain D, Shin JH, Low MJ, et al. Cocaine supersensitivity and enhanced motivation for reward in mice lacking dopamine D2-autoreceptors. *Nat Neurosci* 2011; 14: 1033–8.
- Bjorklund A, Dunnett SB. Dopamine neuron systems in the brain: an update. *Trends Neurosci* 2007; 30: 194–202.
- Callaghan RC, Cunningham JK, Sykes J, Kish SJ. Increased risk of Parkinson's disease in individuals hospitalized with conditions related to the use of methamphetamine or other amphetamine-type drugs. *Drug and alcohol dependence* 2012; 120: 35–40.
- Chan CS, Guzman JN, Ilijic E, Mercer JN, Rick C, Tkatch T, et al. 'Rejuvenation' protects neurons in mouse models of Parkinson's disease. *Nature* 2007; 447: 1081–6.
- Collier TJ, Kanaan NM, Kordower JH. Ageing as a primary risk factor for Parkinson's disease: evidence from studies of non-human primates. *Nat Rev Neurosci* 2011; 12: 359–66.
- Dong XX, Wang Y, Qin ZH. Molecular mechanisms of excitotoxicity and their relevance to pathogenesis of neurodegenerative diseases. *Acta Pharmacol Sin* 2009; 30: 379–87.
- Drion G, Massotte L, Sepulchre R, Seutin V. How modeling can reconcile apparently discrepant experimental results: the case of pacemaking in dopaminergic neurons. *PLoS Comput Biol* 2011; 7: e1002050.
- Ford CP. The role of D2-autoreceptors in regulating dopamine neuron activity and transmission. *Neuroscience* 2014; 23: pii: S0306-4522(14)00037-2.
- Ford CP, Gantz SC, Phillips PE, Williams JT. Control of extracellular dopamine at dendrite and axon terminals. *J Neurosci* 2010; 30: 6975–83.
- Gantz SC, Bunzow JR, Williams JT. Spontaneous inhibitory synaptic currents mediated by a G protein-coupled receptor. *Neuron* 2013; 78: 807–12.
- Gasser T, Hardy J, Mizuno Y. Milestones in PD genetics. *Mov Disord* 2011; 26: 1042–8.
- Gazewood JD, Richards DR, Clebak K. Parkinson disease: an update. *Am Fam Physician* 2013; 87: 267–73.
- Ginovart N, Kapur S. Role of dopamine D(2) receptors for antipsychotic activity. *Handb Exp Pharmacol* 2012; 212: 27–52.
- Goldberg MS, Pisani A, Haburcak M, Vortherms TA, Kitada T, Costa C, et al. Nigrostriatal dopaminergic deficits and hypokinesia caused by inactivation of the familial Parkinsonism-linked gene DJ-1. *Neuron* 2005; 45: 489–96.
- Greengard P, Allen PB, Nairn AC. Beyond the dopamine receptor: the DARPP-32/protein phosphatase-1 cascade. *Neuron* 1999; 23: 435–47.
- Grundemann J, Schlaudraff F, Liss B. UV-laser microdissection and mRNA expression analysis of individual neurons from postmortem Parkinson's disease brains. *Methods Mol Biol* 2011; 755: 363–74.
- Guzman JN, Sanchez-Padilla J, Chan CS, Surmeier DJ. Robust pacemaking in SN DA neurons. *J Neurosci* 2009; 29: 11011–9.

- Guzman JN, Sanchez-Padilla J, Wokosin D, Kondapalli J, Ilijic E, Schumacker PT, et al. Oxidant stress evoked by pacemaking in dopaminergic neurons is attenuated by DJ-1. *Nature* 2010; 468: 696–700.
- Hamasaki-Katagiri N, Ames JB. Neuronal calcium sensor-1 (Ncs1p) is up-regulated by calcineurin to promote Ca²⁺ tolerance in fission yeast. *J Biol Chem* 2010; 285: 4405–14.
- Hernandez-Lopez S, Tkatch T, Perez-Garci E, Galarraga E, Bargas J, Hamm H, et al. D2 dopamine receptors in striatal medium spiny neurons reduce L-type Ca²⁺ currents and excitability via a novel PLC[β]1-IP3-calcineurin-signaling cascade. *J Neurosci* 2000; 20: 8987–95.
- Hurley MJ, Brandon B, Gentleman SM, Dexter DT. Parkinson's disease is associated with altered expression of Ca_v1 channels and calcium-binding proteins. *Brain* 2013; 136 (Pt 7): 2077–97.
- Ilijic E, Guzman JN, Surmeier DJ. The L-type channel antagonist isradipine is neuroprotective in a mouse model of Parkinson's disease. *Neurobiol Dis* 2011; 43: 364–71.
- Jo J, Heon S, Kim MJ, Son GH, Park Y, Henley JM, et al. Metabotropic glutamate receptor-mediated LTD involves two interacting Ca(2+) sensors, NCS-1 and PICK1. *Neuron* 2008; 60: 1095–111.
- Kabbani N, Negyessy L, Lin R, Goldman-Rakic P, Levenson R. Interaction with neuronal calcium sensor NCS-1 mediates desensitization of the D2 dopamine receptor. *J Neurosci* 2002; 22: 8476–86.
- Khalik ZM, Bean BP. Pacemaking in dopaminergic ventral tegmental area neurons: depolarizing drive from background and voltage-dependent sodium conductances. *J Neurosci* 2010; 30: 7401–13.
- Koyrakh L, Lujan R, Colon J, Karschin C, Kurachi Y, Karschin A, et al. Molecular and cellular diversity of neuronal G-protein-gated potassium channels. *J Neurosci* 2005; 25: 11468–78.
- Lammel S, Hetzel A, Hackel O, Jones I, Liss B, Roeper J. Unique properties of mesoprefrontal neurons within a dual mesocorticolimbic dopamine system. *Neuron* 2008; 57: 760–73.
- Lammel S, Ion DI, Roeper J, Malenka RC. Projection-specific modulation of dopamine neuron synapses by aversive and rewarding stimuli. *Neuron* 2011; 70: 855–62.
- Lindgren HS, Rylander D, Ohlin KE, Lundblad M, Cenci MA. The “motor complication syndrome” in rats with 6-OHDA lesions treated chronically with L-DOPA: relation to dose and route of administration. *Behav Brain Res* 2007; 177: 150–9.
- Liss B, Roeper J. Correlating function and gene expression of individual basal ganglia neurons. *Trends Neurosci* 2004; 27: 475–81.
- Liss B, Roeper J. Ion channels and regulation of dopamine neuron activity. In: Iversen LL, Iversen SDI, Dunnett SB, Björklund A, editors. *Dopamine handbook*. Oxford, UK: Oxford University press; 2010. p. 118–38.
- Lundblad M, Picconi B, Lindgren H, Cenci MA. A model of L-DOPA-induced dyskinesia in 6-hydroxydopamine lesioned mice: relation to motor and cellular parameters of nigrostriatal function. *Neurobiol Dis* 2004; 16: 110–23.
- Lundblad M, Usiello A, Carta M, Hakansson K, Fisone G, Cenci MA. Pharmacological validation of a mouse model of L-DOPA-induced dyskinesia. *Exp Neurol* 2005; 194: 66–75.
- Luscher C, Slesinger PA. Emerging roles for G protein-gated inwardly rectifying potassium (GIRK) channels in health and disease. *Nat Rev Neurosci* 2010; 11: 301–15.
- Marras C, Gruneir A, Rochon P, Wang X, Anderson G, Brotchie J, et al. Dihydropyridine calcium channel blockers and the progression of parkinsonism. *Ann Neurol* 2012; 71: 362–9.
- Narushima M, Uchigashima M, Hashimoto K, Watanabe M, Kano M. Depolarization-induced suppression of inhibition mediated by endocannabinoids at synapses from fast-spiking interneurons to medium spiny neurons in the striatum. *Eur J Neurosci* 2006; 24: 2246–52.
- Neve KA, Ford CP, Buck DC, Grandy DK, Neve RL, Phillips TJ. Normalizing dopamine D2 receptor-mediated responses in D2 null mutant mice by virus-mediated receptor restoration: comparing D2 and D2. *Neuroscience* 2013; 248C: 479–87.
- Olanow CW, Schapira AH. Therapeutic prospects for Parkinson disease. *Ann Neurol* 2013; 74: 337–47.
- Olson PA, Tkatch T, Hernandez-Lopez S, Ulrich S, Ilijic E, Mugnaini E, et al. G-protein-coupled receptor modulation of striatal Ca_v1.3 L-type Ca²⁺ channels is dependent on a Shank-binding domain. *J Neurosci* 2005; 25: 1050–62.
- Parkinson Study GroupPhase II safety, tolerability, and dose selection study of isradipine as a potential disease-modifying intervention in early Parkinson's disease (STEADY-PD). *Mov Disord* 2013; 28: 1823–31.
- Pasternak B, Svanstrom H, Nielsen NM, Fugger L, Melbye M, Hviid A. Use of calcium channel blockers and Parkinson's disease. *Am J Epidemiol* 2012; 175: 627–35.
- Platzer J, Engel J, Schrott-Fischer A, Stephan K, Bova S, Chen H, et al. Congenital deafness and sinoatrial node dysfunction in mice lacking class D L-type Ca²⁺ channels. *Cell* 2000; 102: 89–97.
- Prinz A, Selesnew LM, Liss B, Roeper J, Carlsson T. Increased excitability in serotonin neurons in the dorsal raphe nucleus in the 6-OHDA mouse model of Parkinson's disease. *Exp Neurol* 2013; 248: 236–45.
- Puopolo M, Raviola E, Bean BP. Roles of subthreshold calcium current and sodium current in spontaneous firing of mouse midbrain dopamine neurons. *J Neurosci* 2007; 27: 645–56.
- Rice ME, Patel JC, Cragg SJ. Dopamine release in the basal ganglia. *Neuroscience* 2011; 198: 112–37.
- Ritz B, Rhodes SL, Qian L, Schernhammer E, Olsen JH, Friis S. L-type calcium channel blockers and Parkinson disease in Denmark. *Ann Neurol* 2010; 67: 600–6.
- Ritz MC, Lamb RJ, Goldberg SR, Kuhar MJ. Cocaine receptors on dopamine transporters are related to self-administration of cocaine. *Science* 1987; 237: 1219–23.
- Saab BJ, Georgiou J, Nath A, Lee FJ, Wang M, Michalon A, et al. NCS-1 in the dentate gyrus promotes exploration, synaptic plasticity, and rapid acquisition of spatial memory. *Neuron* 2009; 63: 643–56.
- Saal D, Dong Y, Bonci A, Malenka RC. Drugs of abuse and stress trigger a common synaptic adaptation in dopamine neurons. *Neuron* 2003; 37: 577–82.
- Schultz W. Multiple functions of dopamine neurons. *F1000 Biol Rep* 2010; 2: 2.
- Shulman JM, De Jager PL, Feany MB. Parkinson's disease: genetics and pathogenesis. *Annu Rev Pathol* 2011; 6: 193–222.
- Simuni T, Borushko E, Avram MJ, Miskevics S, Martel A, Zadikoff C, et al. Tolerability of isradipine in early Parkinson's disease: a pilot dose escalation study. *Mov Disord* 2010; 25: 2863–6.
- Sinnesger-Brauns MJ, Hetzenauer A, Huber IG, Renstrom E, Wietzorek G, Berjukov S, et al. Isoform-specific regulation of mood behavior and pancreatic beta cell and cardiovascular function by L-type Ca²⁺ channels. *J Clin Invest* 2004; 113: 1430–9.
- Sinnesger-Brauns MJ, Huber IG, Koschak A, Wild C, Obermair GJ, Einzinger U, et al. Expression and 1,4-dihydropyridine-binding properties of brain L-type calcium channel isoforms. *Mol Pharmacol* 2009; 75: 407–14.
- Striessnig J, Koschak A. Exploring the function and pharmacotherapeutic potential of voltage-gated Ca²⁺ channels with gene knockout models. *Channels (Austin)* 2008; 2: 233–51.
- Striessnig J, Koschak A, Sinnesger-Brauns MJ, Hetzenauer A, Nguyen NK, Busquet P, et al. Role of voltage-gated L-type Ca²⁺ channel isoforms for brain function. *Biochem Soc Trans* 2006; 34 (Pt 5): 903–9.
- Sulzer D, Surmeier DJ. Neuronal vulnerability, pathogenesis, and Parkinson's disease. *Mov Disord* 2013; 28: 715–24.
- Surmeier DJ, Guzman JN, Sanchez J, Schumacker PT. Physiological phenotype and vulnerability in Parkinson's disease. *Cold Spring Harb Perspect Med* 2012; 2: a009290.
- Surmeier DJ, Schumacker PT. Calcium, bioenergetics, and neuronal vulnerability in Parkinson's disease. *J Biol Chem* 2013; 288: 10736–41.
- Tong Y, Pisani A, Martella G, Karouani M, Yamaguchi H, Pothos EN, et al. R1441C mutation in LRRK2 impairs dopaminergic

- neurotransmission in mice. *Proc Natl Acad Sci USA* 2009; 106: 14622–7.
- Ungless MA, Grace AA. Are you or aren't you? Challenges associated with physiologically identifying dopamine neurons. *Trends Neurosci* 2012; 35: 422–30.
- Ungless MA, Whistler JL, Malenka RC, Bonci A. Single cocaine exposure *in vivo* induces long-term potentiation in dopamine neurons. *Nature* 2001; 411: 583–7.
- Usiello A, Baik JH, Rouge-Pont F, Picetti R, Dierich A, LeMeur M, et al. Distinct functions of the two isoforms of dopamine D2 receptors. *Nature* 2000; 408: 199–203.
- Vallone D, Picetti R, Borrelli E. Structure and function of dopamine receptors. *Neurosci Biobehav Rev* 2000; 24: 125–32.
- Yamamoto M, Schapira AH. Dopamine agonists in Parkinson's disease. *Expert Rev Neurother* 2008; 8: 671–7.