

UC Berkeley

Other Recent Work

Title

Phylogenetic signal in plant pathogen-host range

Permalink

<https://escholarship.org/uc/item/8bd0b511>

Publication Date

2007-06-07

Peer reviewed

Phylogenetic signal in plant pathogen-host range

Gregory S. Gilbert, and Campbell O. Webb

PNAS published online Mar 14, 2007;

doi:10.1073/pnas.0607968104

This information is current as of March 2007.

Supplementary Material

Supplementary material can be found at:
www.pnas.org/cgi/content/full/0607968104/DC1

This article has been cited by other articles:
www.pnas.org#otherarticles

E-mail Alerts

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).

Rights & Permissions

To reproduce this article in part (figures, tables) or in entirety, see:
www.pnas.org/misc/rightperm.shtml

Reprints

To order reprints, see:
www.pnas.org/misc/reprints.shtml

Notes:

Phylogenetic signal in plant pathogen–host range

Gregory S. Gilbert^{*†‡} and Campbell O. Webb[§]

^{*}Environmental Studies Department, University of California, Santa Cruz, CA 95064; [†]Smithsonian Tropical Research Institute, Balboa, Ancón, Republic of Panamá; and [§]Arnold Arboretum of Harvard University, Harvard University Herbaria, 22 Divinity Avenue, Cambridge, MA 02138

Edited by Rodolfo Dirzo, Stanford University, Stanford, CA, and approved January 24, 2007 (received for review September 11, 2006)

What determines which plant species are susceptible to a given plant pathogen is poorly understood. Experimental inoculations with fungal pathogens of plant leaves in a tropical rain forest show that most fungal pathogens are polyphagous but that most plant species in a local community are resistant to any given pathogen. The likelihood that a pathogen can infect two plant species decreases continuously with phylogenetic distance between the plants, even to ancient evolutionary distances. This phylogenetic signal in host range allows us to predict the likely host range of plant pathogens in a local community, providing an important tool for plant ecology, design of agronomic systems, quarantine regulations in international trade, and risk analysis of biological control agents. In particular, the results suggest that the rate of spread and ecological impacts of a disease through a natural plant community will depend strongly on the phylogenetic structure of the community itself and that current regulatory approaches strongly underestimate the local risks of global movement of plant pathogens or their hosts.

fungal pathogen | plant disease ecology | tropical forest |
plant quarantine | host specificity

Most species of plant pathogen can attack a broad diversity of plant species (1), but the number of plant species with which a pathogen interacts in a local community is generally much lower (2). However, we have limited abilities to predict which species within a plant assemblage are most likely to be susceptible to a particular pathogen. Existing databases of pathogen–host range (e.g., ref. 1) have limited value for quantitative assessments because they are based primarily on haphazard records of pathogens on economically important plants. More importantly, only plant species that are susceptible to particular pathogens were recorded but not which plants are resistant. Nevertheless, several economically and scientifically important issues require predicting the likely host range of plant pathogens. The idea of selectivity among local plant species underlies the theory for the role of natural enemies in the maintenance of plant diversity (3–5) and biological invasions (6, 7). Host selectivity is used in studying plant disease epidemics (8, 9), estimating fungal biodiversity (10), managing agriculture and forestry systems (11, 12), and in risk analysis for global movement of plants and pathogens (13, 14).

Conventional wisdom is that two closely related plant species should be more likely to be susceptible to the same plant pathogens than would plants that are evolutionarily distant, because the morphological and chemical traits of plants that regulate interactions with pathogens are often phylogenetically conserved (15). Indeed, recent work has shown a strong phylogenetic signal in host range of herbivorous insects in tropical rain forest (16, 17), and there is qualitative support for such a signal for fungal pathogens (7). The presumption of phylogenetic signal in host range often underlies important agronomic and economic decisions. For instance, risk assessment for the release of exotic natural enemies for biological control of weedy plants is predicated on host specificity. Empirical host-range testing follows the “centrifugal phylogenetic method” (14, 18–20), with testing being most intense on local species in the same genus as the target host, less intense in the same family, and with still fewer tests on hosts at greater phylogenetic distances. In a second

example, the U.S. regulatory Animal and Plant Health Inspection Service (APHIS) has developed a draft policy whereby exotic plant species from the same genus as a known host of a quarantine pathogen would not be authorized for planting in the U.S. pending further risk analysis (13). The APHIS policy uses a “step–function” phylogenetic model of pathogen host range, where plants outside the genus are not considered to pose a risk for pathogen spread. However, neither of these applications of phylogenetic signal is based on robust empirical data. Evaluating and improving these models requires quantitative estimates of the strength and shape of phylogenetic signal in the host range in plant pathogens (21, 22), based on empirical evaluation of both susceptibility and resistance of phylogenetically diverse plant species to a diversity of plant pathogens.

We used experimental, *in situ* inoculations to evaluate the host range of 53 necrotrophic (tissue-killing) plant pathogenic fungi on a diversity of plant species in two experiments in Panama: first, in an artificial assemblage of tree species in a reforestation nursery (Nursery), and second in a natural assemblage of species in a semideciduous lowland moist tropical forest (Forest). The Forest inoculations permit assessment under conditions where historical plant–pathogen interactions may have led to local natural selection on either hosts or pathogens (or both) at a particular site, whereas the Nursery inoculations permit phylogenetic analysis separate from such local rapid evolutionary responses. We analyzed the likelihood that a pathogen would cause disease on a target host as a function of the phylogenetic distance between the target plant and the plant species from which the pathogen was isolated. This is a systematic test of the phylogenetic signal in the host range of foliar plant pathogens.

Results

In both Nursery and Forest studies, the proportion of plant species that developed disease declined continuously with phylogenetic distance (estimated time of independent evolution) between the source and target host species from 28 to 287 My (Fig. 1), with the steepest decline occurring in the most closely related pairs. The responses were similar in the two studies (slopes not different, $P = 0.6$), whether the test was performed with a haphazard collection of plant species (Nursery), or with a natural plant community (Forest), where reciprocal selection for pathogenicity and susceptibility would likely have occurred between pathogens and plants over many generations. In our taxa, congeneric pairs ($n = 9$) were separated by a median distance of 53.6 My (range: 28–54 My), with 66.7% of the cross-inoculations causing disease, and confamilials ($n = 94$) were separated by a median distance of 85 My (range: 34–135),

Author contributions: G.S.G. and C.O.W. designed research; G.S.G. performed research; G.S.G. and C.O.W. analyzed data; and G.S.G. and C.O.W. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS direct submission.

Abbreviations: ha, hectare; masl, meters above sea level; MEA, malt extract agar.

[†]To whom correspondence should be addressed at: Environmental Studies, 1156 High Street, 405 ISB, University of California, Santa Cruz, CA 95064. E-mail: ggilbert@ucsc.edu.

This article contains supporting information online at www.pnas.org/cgi/content/full/0607968104/DC1.

© 2007 by The National Academy of Sciences of the USA




Fig. 1. Proportion of target plant species that developed disease symptoms after inoculation with fungal pathogens from source plant species. Logistic regressions were performed on results of individual pathogen–host combinations, but data were grouped into 5-My source-target phylogenetic distance classes for illustration. Circles (Forest study) and squares (Nursery study) indicate the proportion of inoculations within a 5-My class that produced disease symptoms. Lines indicate the predicted proportion symptomatic based on logistic regression for the Forest study (solid line), the Nursery study (dashed line), or all data combined (dotted line). Fine dotted lines show extrapolation of predictions to shorter phylogenetic distances than represented by experimental data. Size of symbol indicates the number of inoculation pairs included in the distance class (grouped as <10, 10–30, or >30 inoculations). Logistic regressions were performed on raw (not binned) data. The logistic fit still shows slope when pairs of <135 My are excluded (dash-dot line). The logistic regression formulas are: forest: $\text{logit}(S) = 2.2327 - 1.3428 \times (\log_{10}(\text{distance} + 1))$, $\chi^2 = 6.96$, $P = 0.0084$, $n = 578$; Nursery: $\text{logit}(S) = 3.4096 - 1.7562 \times (\log_{10}(\text{distance} + 1))$, $\chi^2 = 7.66$, $P = 0.0056$, $n = 384$; Combined: $\text{logit}(S) = 2.9113 - 1.5944 \times (\log_{10}(\text{distance} + 1))$, $\chi^2 = 16.71$, $P = 0.0001$, $n = 962$; and combined, >135 My: $\text{logit}(S) = 5.93699 - 2.85376 \times \log_{10}(\text{distance} + 1)$, $\chi^2 = 8.09$, $P = 0.004$, $n = 821$; where $\text{Prob}(\text{symptomatic}) = \frac{\exp(\text{logit}(S))}{1 + \exp(\text{logit}(S))}$.

with disease in 43.6% of cross-inoculations, whereas the remaining, more distant pairings ($n = 859$) were separated by a median distance of 233 My (range: 50–288), with 29.9% susceptible. Even when all plant pairs with phylogenetic distances <135 My were removed from analysis (including all congeneric and con-familial pairs), there was still a highly significant effect of phylogenetic distance on the likelihood of symptom development (Fig. 1). Thus, the phylogenetic signal in pathogen–host range is not lost at phylogenetic distance beyond taxonomic levels like genus and family but extends to ancient phylogenetic distances between unrelated angiosperms.

We currently lack good general estimates for breadth of host range for necrotrophic plant pathogenic fungi. Our results suggest that most pathogens in a tropical forest are pathogenic on a large number of locally available hosts but that most local plant species will be resistant. Only 2 of the 53 tested pathogens (3.8%) were restricted to a single host species (Fig. 2). Most species had a moderate number of potential host species (median 27.7% of host species tested), with a few pathogens appearing to be very broad generalists (Fig. 2).

Discussion

The phylogenetic signal we measured in pathogen–host range has significant applications in numerous areas; here, we outline




Fig. 2. Frequency distribution of observed host breadth. Shown is the proportion of target hosts that produced symptoms after inoculation, excluding the original source host species, for 40 fungal pathogens in the forest (gray bars), and 13 in the Nursery (white bars). Each strain was tested on 11–36 host species (median 16). Only two strains (of 53) appeared to be specific to one plant species ($n = 11$ and 15 hosts tested). For all strains combined, mean host range breadth = 0.311 ± 0.241 , median = 0.2727.

the implications for epidemiology, ecology, biodiversity, agronomy, and quarantine risk analysis.

The rate of spread of a fungal plant pathogen increases with the density of suitable hosts (23); for pathogens able to infect multiple species, this density is, in turn, a function of the number of susceptible species in a community. Each plant species in the forest will be linked to others through polyphagous pathogens, and the number of “cohosts” in a local forest will be a function of the phylogenetic structure of plant species in the community. We estimated how many host species in a particular forest would be expected to be susceptible to a hypothetical pathogen with a particular host, using the phylogenetic structure of the 315 species in the well studied, nearby 50-hectare (ha) Forest Dynamics Plot on Barro Colorado Island and the phylogenetic signal estimated in our Forest study (Fig. 1). For a host species with a large number of close relatives on the plot (e.g., *Inga marginata*, Fabaceae, 25% quartile phylogenetic distance to all other heterospecifics = 188.5 My), there were 14 other species (4.5% of pool) with which it had a >50% chance of having a pathogen in common. In contrast, for a species with few close relatives (e.g., *Attalea butyraceae*, Arecaceae, 25% quartile distance = 287.7 My), there were no other species on the plot with such a high likelihood. At the 30% probability level for having a pathogen in common, there were 115 likely cohosts for pathogens of *Inga* compared with only 10 for *Attalea*. The variation in likely number of cohosts across plant species demonstrates the potential for ecologically significant differences in disease epidemiology depending on the phylogenetic structure of the plant community.

Plant pathogens with limited local host ranges are thought to help maintain plant diversity in forest communities (4, 24). In our study, most local plant species were resistant to any given pathogen, but species-specialists or even genus-specialists were rare. Analyzing tropical herbivore communities, Novotny *et al.*

(17) found that a lack of strict host specificity and gradual decrease in insect species overlap with phylogenetic distance between plant pairs. Most herbivores fed on several closely related congeneric plant species, but many fed on plants from multiple genera and families. Our data suggest that, because close relatives are more likely to be susceptible to the same pathogens, pathogens may be most effective at maintaining diversity of higher taxonomic levels ["phylodiversity" (22)] rather than species diversity *per se*. An important next step is to evaluate how the severity of damage varies among pathogens of different host ranges.

Efforts to estimate biodiversity have often used extrapolations based on estimates of host specificity of insects or fungi (10, 25). Incorporation of the phylogenetic signal in host sharing in insects (simplified to 100% host sharing in herbivorous insects for all plants of the same genus), led to significantly reduced estimates of global insect diversity compared with earlier efforts (17). Our more sensitive measure of host breadth provides a continuous model for estimating fungal diversity. Such calculations would require strong (and not yet available) estimates of pathogen diversity within individual species but should significantly reduce current estimates of global fungal diversity.

In designing mixed-species forestry plantations or intercropped agronomic systems (11, 26), the mixing of host species that are phylogenetically distant should decrease the probability that hosts would share common pathogens, effectively reducing apparent host density for density-dependent pathogens. In our system, species pairs with phylogenetic distances of >250 My would reduce by half the probability of hosts sharing pathogens compared with mixtures of species from the same family. In an agricultural example, when maize (*Zea mays*) was intercropped with the distantly related (288 My) common bean (*Phaseolus vulgaris*), the incidence of common rust (*Puccinia sorghi*) on maize was greatly reduced, whereas maize suffered high rust incidence when intercropping with the congeneric sorghum (*Sorghum bicolor*) (55 My) (11). The causes for differences in disease levels in the two systems were not determined in the study, but *Sorghum* has been reported as a host for the pathogen (1).

Finally, the observed data provide a quantitative assessment of the assumption of a phylogenetic signal in host range used in risk assessment for biological control agents, biological invaders, and quarantine decisions. Our data suggest that arbitrary cutoffs at the genus or family level will underestimate host ranges of plant pathogens and their associated risks. Analysis of likely hosts based on a continuous logistic function of estimated phylogenetic distance may provide a more realistic evaluation of risks from introduced pathogens.

These results represent a quantitative assessment of phylogenetic signal in the host range of plant pathogens. They provide a benchmark for evaluating the robustness of existing tools for understanding the spread, impacts, and evolutionary biology of plant pathogens as well as the basis for development of novel predictive tools in plant ecology and risk analysis.

Methods

Study Sites. We evaluated the phylogenetic signal of host range of a 53 diverse plant pathogenic fungi in two studies in central Panama. In the first study (Nursery), we used seedlings of 45 species of tropical forest trees grown for reforestation projects at the PROENA nursery in Gamboa, Colón Province [09°07.225N, 079°42.242W, 73 meters above sea level (masl)]. Although all of the species in the nursery are native to forests in Panama, they are not a natural assemblage, permitting a test of host range in the absence of local selection pressure for particular plant–pathogen pairs. No more than half of the nursery species have been reported in any one of the three large forest plots located in central Panama (midisthmus Barro Colorado Island, 50 ha; Caribbean side Fort Sherman, 6 ha; Pacific side

Cocoli, 4 ha) (<http://ctfs.si.edu/>), and 31% of the species were not reported in any of the three plots. Plants were grown under partial shade with overhead misting.

In the second study (Forest), three plots (50-m radius, 0.785 ha) were located in the forest near the town of Gamboa and along Pipeline Road, in the Parque Nacional Soberanía, Colón Province, Republic of Panama. The forest in this area is semi-deciduous, lowland tropical moist forest. Here, host range testing was conducted on species from within the immediate neighborhood of pathogen isolation where past selection may have shaped the host ranges of local pathogens. The three Forest plots were all separated from each other by at least 1 km (Plot 1 09°07.258N, 079°41.840W, 80 masl; Plot 2 09°08.213N, 079°43.448W, 75 masl; Plot 3 09°09.026N, 079°43.987W, 89 masl). Annual precipitation in Gamboa is 2,133 mm, with 87% of the rain falling between May and November (Panama Canal Authority data, 1897–2005, http://striweb.si.edu/esp/physical_monitoring/index_phy_mon.htm). All of the forest sites have histories of some anthropogenic disturbance dating to canal construction, but all show a mature, multilayered canopy structure.

Plots were systematically surveyed for plant composition in the understory and overstory. Plant identifications were based on comparison with specimens in the herbaria at the Smithsonian Tropical Research Institute and the University of Panama. Plant taxonomy follows that of Correa *et al.* (27), except where in conflict with the Angiosperm Phylogeny Group (28). Vouchers of unusual species have been deposited with the herbarium of the University of Panama.

Pathogen Isolation. We evaluated the phylogenetic signal of host range of a diversity of plant pathogenic fungi in two studies in central Panama. We focused on culturable, necrotrophic fungal pathogens to facilitate cross-inoculations, and because biotrophic pathogens like rusts and smuts are very rare in tropical forests (29) and were not observed at our sites. In both the Nursery and Forest sites, representative leaves of each disease type were collected, placed in plastic bags, and returned to the laboratory for processing within 1 h. Two triangular pieces of tissue were collected from the edge of the symptomatic area by using a hole punch for mounting small insects (area, 7 mm²). These leaf pieces were placed together in a mesh tea strainer spoon and immersed for 1 min in 70% ethanol and then for 1 min in 10% commercial bleach (0.525% sodium hypochlorite) to surface sterilize. They were then transferred to malt extract agar with chloramphenicol (MEA-chlor: 2% malt extract, 1.5% agar, 0.02% chloramphenicol), and incubated at ambient air-conditioned (~23°C) laboratory conditions. After 3–4 days, fungal growth from the leaf pieces was transferred to a new plate of MEA. If multiple fungal morphotypes were apparent, each was separately transferred. Pathogens were isolated during the rainy season: in August 2005 at the Nursery, and in November and December 2005 for the Forest pathogens.

In the first study (Nursery), we isolated into pure culture 13 strains of fungi from diseased leaves of 12 species of plants within the Nursery. In the second study (Forest), we isolated 40 foliar fungal pathogens from a diversity of plants in each of three plots (12 pathogens from 11 plant species in Plot 1, 16 from 16 species in Plot 2, and 12 from 12 species in Plot 3). These fungi include at least 19 species from 11 genera in 6 orders of Ascomycetes. A list of the fungi used in these experiments and the hosts from which they were isolated is available electronically in [supporting information \(SI\) Data Set 1](#). Each fungal strain was preserved by placing mycelial plugs in 2-ml cryovials and covering with sterile water and by growing in slant-culture in glass vials on MEA. Dried fungal cultures as well as pressed, symptomatic leaves of the original source plants were deposited as vouchers in the herbarium of the University of Panama.

