UC Irvine

Western Journal of Emergency Medicine: Integrating Emergency Care with Population Health

Title

Left Upper Quadrant Abdominal Pain

Permalink

https://escholarship.org/uc/item/8cd761fd

Journal

Western Journal of Emergency Medicine: Integrating Emergency Care with Population Health, 13(6)

ISSN

1936-900X

Authors

Nordt, Sean Patrick Bowns, Christopher Moran, Julieta et al.

Publication Date

2012

DOI

10.5811/westjem.2012.1.11737

Copyright Information

Copyright 2012 by the author(s). This work is made available under the terms of a Creative Commons Attribution-NonCommercial License, available at https://creativecommons.org/licenses/by-nc/4.0/

Peer reviewed

Left Upper Quadrant Abdominal Pain

Sean Patrick Nordt, MD, PharmD*
Christopher Bowns, MD†
Julieta Moran, MD†
H. Brendan Kelleher, MD†
Stuart Swadron, MD*

- **Sean Patrick Nordt, MD, PharmD*** * University of Southern California, Department of Emergency Medicine, Los Angeles, Christopher Bowns. MD[†] California
 - [†] Los Angeles County + University of Southern California Medical Center, Department of Emergency Medicine, Los Angeles, California

Supervising Section Editor: Sean O. Henderson, MD

Submission history: Submitted January 10, 2012; Accepted March 8, 2012

Full text available through open access at http://escholarship.org/uc/uciem_westjem

DOI: 10.5811/westjem.2012.1.11737

We present a case of acute appendicitis from mobile cecum presenting with left upper quadrant abdominal pain. [West J Emerg Med. 2012;13(6):495-496]

A 58-year-old male with history of hypertension presented with constant left upper quadrant (LUQ) pain for 3 days. He denied fevers, chills, had no anorexia or association with eating, and no changes in bowel or bladder function. He was nontoxic appearing with normal vital signs. Abdomen was soft with tenderness to palpation in LUQ with normal bowel sounds. There were no masses, guarding, or other areas of tenderness. Remainder of the examination was normal. White blood count was 10,600/mm³ with 74.9% neutrophils with otherwise normal indices. Serum chemistry, urinalysis, and liver function tests, including lipase, were all in normal range. C-reactive protein (CRP) was 217 mg/L (reference range: 0-7 mg/L). A computerized tomography (CT) with intravenous contrast was obtained.




Figure 1. Axial image identifying left-sided enlarged appendix (arrow).

Diagnosis: Left-sided Appendicitis

The CT demonstrated a mobilized mesocecum in the left upper quadrant with a large, inflamed thick-walled appendix consistent with acute appendicitis. He underwent a laparotomy. The colon and cecum were found in the left upper quadrant with an inflamed, enlarged appendix. No malrotation or other congenital abnormalities were seen. Histology demonstrated a gangrenous appendix. He tolerated surgery well and was discharged home two days later.

Left-sided acute appendicitis is predominantly seen with



Figure 2. Coronal image identifying left upper quadrant acute appendicitis (arrow).

congenital anomalies, e.g., situs inversus totalis and midgut malrotation. The majority of these present with left lower quadrant pain. The patient presented had LUQ pain due to a hypermobile colon. Though not specific, an elevated CRP has been suggested as a marker of acute appendicitis. The elevated CRP and clinical concern led to obtaining the CT.

Address for Correspondence: Sean Patrick Nordt, MD, PharmD, Department of Emergency Medicine, University of Southern California, 1200 North State Street, Unit 1, Room 1011, Los Angeles, CA 90033. Email: spnordt@hotmail.com.

Conflicts of Interest: By the WestJEM article submission agreement, all authors are required to disclose all affiliations, funding sources and financial or management relationships that could be perceived as potential sources of bias. The authors disclosed none.

REFERENCES:

- Akbulut S, Ulku A, Senol A, et al. Left-sided appendicitis: review of 95 published cases and a case report. World J Gastroeterol 2010;16:5598-5602.
- Andersson REB. Meta-analysis of the clinical and laboratory diagnosis of appendicitis. Br J Surg 2004;2004:28-37.