

UCSF

UC San Francisco Previously Published Works

Title

Draft Genome Sequence of Mycobacterium heckeshornense Strain RLE

Permalink

<https://escholarship.org/uc/item/8cs6q1qk>

Journal

Microbiology Resource Announcements, 3(4)

ISSN

2576-098X

Authors

Greninger, Alexander L

Cunningham, Gail

Chiu, Charles Y

et al.

Publication Date

2015-08-27

DOI

10.1128/genomea.00930-15

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Draft Genome Sequence of *Mycobacterium heckeshornense* Strain RLE

Alexander L. Greninger, Gail Cunningham, Charles Y. Chiu, Steve Miller

Department of Laboratory Medicine, UCSF, San Francisco, California, USA

We report here the draft genome sequence of *Mycobacterium heckeshornense* strain RLE isolated from a sputum sample from a patient with shortness of breath. This is the first draft genome sequence of *M. heckeshornense*.

Received 8 July 2015 Accepted 21 July 2015 Published 27 August 2015

Citation Greninger AL, Cunningham G, Chiu CY, Miller S. 2015. Draft genome sequence of *Mycobacterium heckeshornense* strain RLE. *Genome Announc* 3(4):e00930-15. doi:10.1128/genomeA.00930-15.

Copyright © 2015 Greninger et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Steve Miller, steve.miller@ucsf.edu.

Mycobacterium heckeshornense is a slowly growing scotochromogenic mycobacterium that was originally isolated in 1994 from a cavitary lung lesion and is named after the Heckeshorn Lung Clinic in Berlin (1). *M. heckeshornense* has since been isolated from a broad range of clinical cases, such as septicemia, osteomyelitis, diskitis, peritonitis, tenosynovitis, and lymphadenitis, as well as from porcine and feline lymph nodes (2–8). It is phylogenetically and phenotypically close to *Mycobacterium xenopi*, which is a common source of confusion and misidentification (1, 5, 9). We sequenced the first draft genome of *M. heckeshornense* isolated from a 2008 sputum sample from a 50-year-old male with HIV and CD4 count of <60 who presented with shortness of breath and cough and was concurrently diagnosed with *Pneumocystis jirovecii* pneumonia.

DNA from *M. heckeshornense* strain RLE was extracted using the Qiagen EZ1 kit, and paired-end libraries were prepared using the Nextera XT DNA library kit, followed by 2 × 80-bp sequencing on Illumina MiSeq. The sequences were adapter and quality (Q20) trimmed using cutadapt, *de novo* assembled using SPAdes version 3.5, metagenomically screened for contaminating sequence with SURPI, and annotated via Prokka version 1.1 (10–13). A total of 8,091,536 paired-end reads were recovered after trimming. *De novo* assembly yielded 191 contigs of >500 bp for a total assembly size of 5,010,173 bp, with an N_{50} of 95,235 bp, an average coverage of 126×, and a total of 4,882 coding sequences. Contiguity was interrupted due to the short read length, high G+C content (66%), and presence of many high-copy-number integrases, transposases, and tyrosine recombinases that aligned with between 70 and 85% identity at the nucleotide level to a variety of actinobacteria, such as *Gordonia bronchialis* DSM 43247, *Rhodococcus aetherivorans* strain lcdP1, *Mycobacterium kansasii* 824, *Mycobacterium marinum* strain M, and *Rhodococcus jostii* RHA1.

BLASTn analysis of the complete 16S sequence from strain RLE showed 100% identity to *M. heckeshornense* type strain S369 (NCBI reference sequence NR_028759) and *Mycobacterium cf. xenopi* (i.e., a *Mycobacterium* sp. very similar to *Mycobacterium xenopi*) strain Hymi_Wue Tb_939/99 (GenBank accession no. AJ243481) and *Mycobacterium sydneyensis* (GenBank accession no. AF101243), as has been noted for *M. heckeshornense* (14). As no publications are available for these two strains nor have any

sequences been deposited, they likely should be renamed as strains of *M. heckeshornense*. BLASTn of the full-length *rpoB* gene revealed 91% nucleotide identity to *rpoB* genes from *Mycobacterium branderi* ATCC 51789, *Mycobacterium celatum* ATCC 51130, *Mycobacterium kyorinense* KUM 060204, and a variety of *Mycobacterium avium* strains, including 104, 2285, and DJO-44271. No high-confidence antibiotic resistance genes were identified by the Comprehensive Antibiotic Resistance Database analysis (15).

Nucleotide sequence accession numbers. This whole-genome shotgun project has been deposited at DDBJ/ENA/GenBank under the accession no. LFOF00000000. The assembly described in this paper is the first version, LFOF01000000.

ACKNOWLEDGMENT

This work was supported by the University of California, San Francisco Department of Laboratory Medicine.

REFERENCES

- Roth A, Reischl U, Schönfeld N, Naumann L, Emler S, Fischer M, Mauch H, Loddenkemper R, Kroppenstedt RM. 2000. *Mycobacterium heckeshornense* sp. nov., a new pathogenic slowly growing *Mycobacterium* sp. causing cavitary lung disease in an immunocompetent patient. *J Clin Microbiol* 38:4102–4107.
- Ahmed RA, Miedzinski LJ, Shandro C. 2010. *Mycobacterium heckeshornense* infection in HIV-infected patient. *Emerg Infect Dis* 16:1801–1803. <http://dx.doi.org/10.3201/eid1611.091226>.
- Van Ingen J, Wisselink HJ, van Solt-Smits CB, Boeree MJ, van Soolingen D. 2010. Isolation of mycobacteria other than *Mycobacterium avium* from porcine lymph nodes. *Vet Microbiol* 144:250–253. <http://dx.doi.org/10.1016/j.vetmic.2009.12.047>.
- Elze J, Grammel L, Richter E, Aupperle H. 2013. First description of *Mycobacterium heckeshornense* infection in a feline immunodeficiency virus-positive cat. *J Feline Med Surg* 15:1141–1144. <http://dx.doi.org/10.1177/1098612X13490711>.
- Carpenter RJ, Graf PC. 2015. Pott's disease? AIDS-associated *Mycobacterium heckeshornense* spinal osteomyelitis and diskitis. *J Clin Microbiol* 53:716–718. <http://dx.doi.org/10.1128/JCM.02686-14>.
- Chan WW, Murray MC, Tang P, Romney MG. 2011. *Mycobacterium heckeshornense* peritonitis in a peritoneal dialysis patient: a case report and review of the literature. *Clin Microbiol Infect* 17:1262–1264. <http://dx.doi.org/10.1111/j.1469-0691.2010.03449.x>.
- Godreuil S, Marchandin H, Terru D, Le Moing V, Chammas M, Vincent V, Jumas-Bilak E, Van De Perre P, Carriere C. 2006. *Mycobacterium heckeshornense* tenosynovitis. *Scand J Infect Dis* 38:1098–1101. <http://dx.doi.org/10.1080/00365540600606606>.

8. McBride SJ, Taylor SL, Pandey SK, Holland DJ. 2009. First case of *Mycobacterium heckeshornense* lymphadenitis. *J Clin Microbiol* 47: 268–270. <http://dx.doi.org/10.1128/JCM.00890-08>.
9. Morimoto K, Kazumi Y, Maeda S, Yoshimori K, Yoshiyama T, Ogata H, Kurashima A, Kudoh S. 2011. *Mycobacterium heckeshornense* lung infection that was diagnosed as *Mycobacterium xenopi* disease by DNA-DNA hybridization (DDH). *Intern Med* 50:1251–1253. <http://dx.doi.org/10.2169/internalmedicine.50.4628>.
10. Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068–2069. <http://dx.doi.org/10.1093/bioinformatics/btu153>.
11. Naccache SN, Federman S, Veeraraghavan N, Zaharia M, Lee D, Samayoa E, Bouquet J, Greninger AL, Luk K-C, Enge B, Wadford DA, Messenger SL, Genrich GL, Pellegrino K, Grard G, Leroy E, Schneider BS, Fair JN, Martínez MA, Isa P, Crump JA, DeRisi JL, Sittler T, Hackett J, Miller S, Chiu CY. 2014. A cloud-compatible bioinformatics pipeline for ultrarapid pathogen identification from next-generation sequencing of clinical samples. *Genome Res* 24:1180–1192. <http://dx.doi.org/10.1101/gr.171934.113>.
12. Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.journal* 17:10–12. <http://dx.doi.org/10.14806/ej.17.1.200>.
13. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <http://dx.doi.org/10.1089/cmb.2012.0021>.
14. Richter E, Niemann S, Ruesch-Gerdes S, Harmsen D. 2001. Description of *Mycobacterium heckeshornense* sp. nov. *J Clin Microbiol* 39:3023–3024. <http://dx.doi.org/10.1128/JCM.39.8.3023-3024.2001>.
15. McArthur AG, Waglechner N, Nizam F, Yan A, Azad MA, Baylay AJ, Bhullar K, Canova MJ, Pascale G, Ejim L, Kalan L, King AM, Koteva K, Morar M, Mulvey MR, O'Brien JS, Pawlowski AC, Piddock LJ, Spanogiannopoulos P, Sutherland AD, Tang I, Taylor PL, Thaker M, Wang W, Yan M, Yu T, Wright GD. 2013. The Comprehensive Antibiotic Resistance Database. *Antimicrob Agents Chemother* 57:3348–3357. <http://dx.doi.org/10.1128/AAC.00419-13>.