

# UC San Diego

## UC San Diego Previously Published Works

### Title

Associations between Weight-Adjusted Waist Index and Abdominal Fat and Muscle Mass: Multi-Ethnic Study of Atherosclerosis

### Permalink

<https://escholarship.org/uc/item/8f11s58j>

### Journal

Diabetes & Metabolism Journal, 46(5)

### ISSN

2233-6079

### Authors

Kim, Ji Yoon  
Choi, Jimi  
Vella, Chantal A  
[et al.](#)

### Publication Date

2022-09-30

### DOI

10.4093/dmj.2021.0294

Peer reviewed



# Associations between Weight-Adjusted Waist Index and Abdominal Fat and Muscle Mass: Multi-Ethnic Study of Atherosclerosis

Ji Yoon Kim<sup>1</sup>, Jimi Choi<sup>1</sup>, Chantal A. Vella<sup>2</sup>, Michael H. Criqui<sup>3</sup>, Matthew A. Allison<sup>3</sup>, Nam Hoon Kim<sup>1</sup>

<sup>1</sup>Division of Endocrinology and Metabolism, Department of Internal Medicine, Korea University College of Medicine, Seoul, Korea,

<sup>2</sup>Department of Movement Sciences, College of Education, Health and Human Sciences, University of Idaho, Moscow, ID,

<sup>3</sup>Division of Preventive Medicine, Department of Family and Preventive Medicine, University of California San Diego, La Jolla, CA, USA

**Background:** The weight-adjusted waist index (WWI) reflected body compositional changes with aging. This study was to investigate the association of WWI with abdominal fat and muscle mass in a diverse race/ethnic population.

**Methods:** Computed tomography (CT) data from 1,946 participants for abdominal fat and muscle areas from the Multi-Ethnic Study of Atherosclerosis (785 Whites, 252 Asians, 406 African American, and 503 Hispanics) were used. Among them, 595 participants underwent repeated CT. The WWI was calculated as waist circumference (cm) divided by the square root of body weight (kg). The associations of WWI with abdominal fat and muscle measures were examined, and longitudinal changes in abdominal composition measures were compared.

**Results:** In all race/ethnic groups, WWI was positively correlated with total abdominal fat area (TFA), subcutaneous fat area, and visceral fat area, but negatively correlated with total abdominal muscle area (TMA) and abdominal muscle radiodensity ( $P < 0.001$  for all). WWI showed a linear increase with aging regardless of race and there were no significant differences in the WWI distribution between Whites, Asians, and African Americans. In longitudinal analyses, over 38.6 months of follow-up, all abdominal fat measures increased but muscle measures decreased, along with increase in WWI. The more the WWI increased, the more the TFA increased and the more the TMA decreased.

**Conclusion:** WWI showed positive associations with abdominal fat mass and negative associations with abdominal muscle mass, which likely reflects the abdominal compositional changes with aging in a multi-ethnic population.

**Keywords:** Anthropometry; Body composition; Obesity; Sarcopenia

## INTRODUCTION

Excess body fat, assessed regionally or generally, is recognized as a cause and predictor of various metabolic and cardiovascular diseases [1,2]. Therefore, measurement or assessment of body fat has been a critical issue in medical science [3,4]. On the other hand, cumulative evidence has emphasized the role of muscle mass in the development of several diseases [5]. Specifically, low muscle mass and/or strength has been shown to

be an independent indicator of fragility, age-related disorders, and even mortality [6]. However, most anthropometric indices do not distinguish between muscle mass and fat mass and thus tend to be proportional to both [7,8]. This is one of the reasons for the seemingly paradoxical relationship between obesity and some health outcomes [9,10].

A newly proposed anthropometric index, weight-adjusted waist index (WWI), was notable for its ability to assess fat and muscle mass reciprocally. Specifically, in elderly individuals

Corresponding author: Nam Hoon Kim  <https://orcid.org/0000-0002-9926-1344>  
Division of Endocrinology and Metabolism, Department of Internal Medicine, Korea University College of Medicine, 73 Incheon-ro, Seongbuk-gu, Seoul 02841, Korea  
E-mail: [pourlife@korea.ac.kr](mailto:pourlife@korea.ac.kr)

Received: Oct. 22, 2021; Accepted: Jan. 14, 2022

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

over 65 years, WWI was positively associated with total and abdominal fat measures, but negatively associated with appendicular skeletal muscle mass [11]. Furthermore, a linear positive association of WWI with mortality, as well as cardiometabolic morbidity was observed [12], which was not seen with body mass index (BMI) or waist circumference (WC). However, the associations were only validated in Korean elderly, an East Asian race. Given that there exists substantial difference of body composition by races or ethnic groups, we do not have answers whether the association of WWI with fat and muscle mass is also applicable to the other ethnic groups than East Asian, yet.

Given the aforementioned, the current analysis evaluated the associations between WWI and both abdominal fat and muscle mass measures in different racial and ethnic groups, as well as with the longitudinal changes in abdominal fat and muscle mass in the multi-ethnic population cohort.

## METHODS

### Study population

The Multi-Ethnic Study of Atherosclerosis (MESA) is a community-based longitudinal cohort study with 6,814 men and women aged 45 to 84 years at baseline, recruited from six United States communities. At this visit, it was comprised of 38% Caucasians, 28% African Americans, 22% Hispanics, and 12% Asians (primarily Chinese). The first examination began in July 2000, and the subsequent examinations were conducted regularly. The latest (6th) examination was from September 2016 to June 2018. Details of the study protocol and design have been published previously [13].

The components of the first examination included personal history, demographic data, socioeconomic status, medications and medical history, occupational and leisure-time physical activity, anthropometry, blood pressure measurement, phlebotomy, spot urine collection for microalbuminuria, and cardiac computed tomography (CT) for coronary calcification and magnetic resonance imaging scans for cardiac structure and function. Informed consent was obtained from all participants at each study clinic.

The abdominal body composition ancillary study consisted of a random subset of 1,975 MESA participants who underwent abdominal CT for measurement of aortic calcium during visit 2 and visit 3 examinations (2002 to 2005). These scans were then interrogated for different body composition mea-

asures to include visceral, subcutaneous, and intermuscular fat, as well as lean muscle mass in the abdomen (as described below). This data was used for cross-sectional analyses in the current study. Among the participants who underwent CT during second visit (visit 2), 595 underwent a follow-up CT scan during visit 4 (2005 to 2007), which was used for the longitudinal analyses in this study.

The MESA studies were approved by the Institutional Review Board of each study site and all participants provided written informed consent. The Institutional Review Board of Korea University Anam Hospital also approved this study (IRB number 2021AN0501).

### Anthropometric index and variables from abdominal CT scans

Anthropometric measurements included height, body weight, WC, and hip circumference. BMI was calculated as weight (kg) divided by the square of the height (m<sup>2</sup>). WWI was calculated as WC (cm) divided by the square root of weight in kilograms ( $\sqrt{\text{kg}}$ ).

In the MESA body composition ancillary study, abdominal slices at vertebral level L4/L5 from the abdominal CT were processed using Medical Image Processing, Analysis, and Visualization version 4.1.2 software (MIPAV, National Institutes of Health, Bethesda, MD, USA) that measured the areas of fat, muscle mass, and total tissue by a semiautomated method. The densities of all anatomical structures were measured and expressed in Hounsfield units (HU) for each tissue type. Fat tissue and skeletal muscle tissue were identified as being between -190 and -30 HU and between 0 and 100 HU, respectively [14]. Any densities other than these two HU ranges were labeled as undefined tissue type. Total abdominal fat area (TFA) was subdivided into subcutaneous fat area (SFA) and visceral fat area (VFA) according to the corresponding anatomical location. Muscle area was defined within their unique fascial planes as psoas, rectus abdominis, paraspinal muscle group, and oblique muscle group. Total abdominal muscle area (TMA) was subdivided into muscles of locomotion (psoas muscle) and muscles of stability (rectus abdominis, paraspinal muscle group, and oblique muscle group). Muscle density was the average HU within each muscle's distinct fascial plane. Inter-rater and intra-rater reliability for total abdominal, subcutaneous, and visceral cavity areas were 0.99 for all measurements. Inter- and intra-rater reliability for all muscle groups ranged from 0.93 to 0.98.

### Other variables

Sociodemographic and lifestyle information including age, sex, race, current occupation, family income, insurance status, marital status, detailed history of cigarette smoking and alcohol consumption was obtained from standardized questionnaires. Current and past medical history including medications, treatments, and medical conditions was also obtained. Diabetes mellitus was defined as fasting glucose  $\geq 126$  mg/dL or use of glucose-lowering agents. Hypertension was defined as systolic blood pressure  $\geq 140$  mm Hg or diastolic blood pressure  $\geq 90$  mm Hg or use of anti-hypertensive medications. Using the Typical Week Physical Activity Survey, participants self-reported their time spent in sedentary behavior and in physical activities.

### Statistical analysis

Demographics, laboratory measures, and body composition measures using abdominal CT scans at baseline were presented as mean (standard deviation) or number (percentage). A Pearson correlation analysis was performed to investigate the linear relationship between WWI and both abdominal fat and muscle mass measures. The results are presented as correlation coefficients and corresponding 95% confidence intervals based on Fisher's *r*-to-*z* transformation. To compare the distribution of WWI and BMI according to race, age and sex adjusted *z*-score for WWI and BMI were calculated by regressing WWI or BMI on age and sex, respectively and standardized residuals of the regression models. Density plot of age-sex or sex adjusted WWI *z*-score drawn by smooth curves that goes through the tops of the histogram bars were displayed according to race (Caucasian, Asian, African American, and Hispanic) and age (45–54, 55–64, 65–74, and 75–84 years). The difference in mean WWI by race and age was evaluated after adjusting for age and sex (separately) using analysis of covariance. Means of fat and muscle measures according to quartiles of WWI were compared by one-way analysis of variance and used Tukey's method for *post hoc* analysis.

Five hundred and ninety-five participants underwent repeated abdominal CT during the 4th MESA examination. The mean interval between the 2nd and 4th examinations was  $38.6 \pm 3.3$  months. The longitudinal changes in the anthropometric index and abdominal fat and muscle area measures were assessed using the paired *t*-test. To evaluate the association between the change in anthropometric index and change in body composition measures, means of percentage changes

in TFA or TMA were compared by quartiles of change in WWI or BMI using analysis of variance with adjustment for age.

Statistical significance was set at  $P < 0.05$ . All statistical analyses were performed using the SAS software version 9.4 (SAS Institute Inc., Cary, NC, USA) and R version 4.0.2 (R Foundation for Statistical Computing, Vienna, Austria).

**Table 1.** Baseline characteristics of participants ( $n = 1,946$ )

Variable	Value
Age, yr	64.7 $\pm$ 9.7
Sex	
Female	962 (49.4)
Male	984 (50.6)
Race	
White	785 (40.3)
Asian	252 (12.9)
African American	406 (20.9)
Hispanic	503 (25.8)
Body mass index, kg/m <sup>2</sup>	28.1 $\pm$ 5.2
Waist circumference, cm	98.3 $\pm$ 14.1
Fasting glucose, mg/dL	98.3 $\pm$ 27.7
Cholesterol, mg/dL	
Total	189.7 $\pm$ 35.5
High-density lipoprotein	51.5 $\pm$ 15.2
Low-density lipoprotein	112.0 $\pm$ 31.4
Triglyceride, mg/dL	133.6 $\pm$ 95.0
Creatinine, mg/dL	0.98 $\pm$ 0.22
Systolic blood pressure, mm Hg	124.1 $\pm$ 20.8
Hypertension	914 (47.4)
Diabetes mellitus	277 (14.2)
Current smoker	225 (11.6)
Current alcohol consumption	1,004 (51.7)
MVPA (MET), min/wk	4,939.9 $\pm$ 4,778.8
Weight-adjusted waist index, cm/ $\sqrt{\text{kg}}$	11.17 $\pm$ 0.91
Abdominal computed tomography measures, cm <sup>2</sup>	
Total abdominal fat area	419.5 $\pm$ 160.2
Subcutaneous fat area	253.7 $\pm$ 117.7
Visceral fat area	147.9 $\pm$ 69.1
Total abdominal muscle area	98.3 $\pm$ 27.6
Locomotor abdominal muscle area	23.7 $\pm$ 7.4
Stability abdominal muscle area	74.6 $\pm$ 21.8

Values are presented as mean  $\pm$  standard deviation or number (%). MVPA, moderate and vigorous physical activity; MET, metabolic equivalent.

## RESULTS

The baseline characteristics of all participants are shown in Table 1. The study participants comprised 785 Caucasians (40.3%), 252 Asians (12.9%), 406 African Americans (20.9%), and 503 Hispanics (25.8%), with mean age, BMI, and WC of 64.7 years, 28.1 kg/m<sup>2</sup>, and 98.3 cm, respectively. WWI values were approximately normally distributed with a mean of 11.17, standard deviation of 0.91 and range of 8.34 to 14.44 cm/√kg (Supplementary Fig. 1). WWI increased with age in both men and women, and this trend was consistent in all race/ethnic groups (Supplementary Table 1).

### WWI is positively associated with abdominal fat area measures but negatively associated with abdominal muscle area measures in all race/ethnic groups

Correlation analyses showed that abdominal fat area measures,

including TFA, SFA, and VFA, positively correlated with WWI, whereas all abdominal muscle area measures negatively correlated with WWI, regardless of race/ethnicity ( $P < 0.001$  for all analyses) (Supplementary Table 2). Total abdominal muscle radiodensity negatively correlated with WWI, which indicated that the fat component within abdominal muscles was positively associated with WWI.

The association of WWI with abdominal fat and muscle area measures was examined by comparing the mean values and z-scores according to WWI quartiles (Table 2, Supplementary Table 3). The mean values of abdominal fat area z-scores showed increasing trends from the lowest quartile to the highest quartile of WWI, whereas those of muscle area z-scores showed decreasing trends with increasing quartiles of WWI. These results were consistent regardless of ethnicity ( $P$  for trends  $< 0.001$  for all ethnic groups), and age (Fig. 1). Fig. 2 is a representative figure showing the contrasting relationship be-


**Table 2.** Mean values of muscle and fat measure z-scores by quartiles of WWI, BMI, and WC

All ( $n = 1,946$ )	z-TFA	z-SFA	z-VFA	z-TMA	z-TMR
Quartiles of WWI					
1st	$-0.64 \pm 0.79^a$	$-0.55 \pm 0.83^a$	$-0.65 \pm 0.74^a$	$0.04 \pm 0.94^a$	$0.39 \pm 0.91^a$
2nd	$-0.17 \pm 0.69^b$	$-0.16 \pm 0.70^b$	$-0.16 \pm 0.85^b$	$0.13 \pm 1.03^a$	$0.12 \pm 0.93^b$
3rd	$0.20 \pm 0.82^c$	$0.17 \pm 0.84^c$	$0.17 \pm 0.92^c$	$-0.04 \pm 0.95^b$	$-0.09 \pm 0.96^c$
4th	$0.84 \pm 1.09^d$	$0.74 \pm 1.17^d$	$0.66 \pm 0.99^d$	$-0.14 \pm 1.06^b$	$-0.44 \pm 1.03^d$
<i>P</i> value <sup>e</sup>	$< 0.001$	$< 0.001$	$< 0.001$	$< 0.001$	$< 0.001$
<i>P</i> value <sup>f</sup>	$< 0.001$	$< 0.001$	$< 0.001$	$< 0.001$	$< 0.001$
Quartiles of BMI					
1st	$-0.93 \pm 0.54^a$	$-0.87 \pm 0.55^a$	$-0.83 \pm 0.63^a$	$-0.32 \pm 0.88^a$	$0.35 \pm 0.90^a$
2nd	$-0.23 \pm 0.46^b$	$-0.20 \pm 0.47^b$	$-0.28 \pm 0.69^b$	$-0.05 \pm 0.92^b$	$0.15 \pm 0.93^b$
3rd	$0.31 \pm 0.49^c$	$0.25 \pm 0.6^c$	$0.28 \pm 0.78^c$	$0.22 \pm 1.01^c$	$-0.01 \pm 0.97^c$
4th	$1.40 \pm 0.88^d$	$1.33 \pm 1.01^d$	$0.86 \pm 0.98^d$	$0.17 \pm 1.10^c$	$-0.52 \pm 0.99^d$
<i>P</i> value <sup>e</sup>	$< 0.001$	$< 0.001$	$< 0.001$	$< 0.001$	$< 0.001$
<i>P</i> value <sup>f</sup>	$< 0.001$	$< 0.001$	$< 0.001$	$< 0.001$	$< 0.001$
Quartiles of WC					
1st	$-0.94 \pm 0.54^a$	$-0.84 \pm 0.58^a$	$-0.88 \pm 0.59^a$	$-0.22 \pm 0.92^a$	$0.46 \pm 0.85^a$
2nd	$-0.24 \pm 0.44^b$	$-0.22 \pm 0.49^b$	$-0.27 \pm 0.66^b$	$-0.01 \pm 0.92^b$	$0.17 \pm 0.94^b$
3rd	$0.34 \pm 0.50^c$	$0.28 \pm 0.62^c$	$0.27 \pm 0.79^c$	$0.12 \pm 1.02^b$	$-0.12 \pm 0.95^c$
4th	$1.41 \pm 0.85^d$	$1.31 \pm 1.00^d$	$0.92 \pm 0.94^d$	$0.11 \pm 1.10^b$	$-0.54 \pm 0.98^d$
<i>P</i> value <sup>e</sup>	$< 0.001$	$< 0.001$	$< 0.001$	$< 0.001$	$< 0.001$
<i>P</i> value <sup>f</sup>	$< 0.001$	$< 0.001$	$< 0.001$	$< 0.001$	$< 0.001$


Values are presented as mean  $\pm$  standard deviation.

WWI, weight-adjusted waist index; BMI, body mass index; WC, waist circumference; TFA, total abdominal fat area; SFA, subcutaneous fat area; VFA, visceral fat area; TMA, total abdominal muscle area; TMR, total abdominal muscle radiodensity.

<sup>a,b,c,d</sup>Different letters indicate significant differences between groups ( $P < 0.05$ ) by Tukey's method, <sup>e</sup>*P* value by ANOVA, <sup>f</sup>*P* value for linear trend test.



**Fig. 1.** (A) Total abdominal fat area (TFA) and (B) total abdominal muscle area (TMA) by quartiles of weight-adjusted waist index (WWI) in adults older and younger than 65 years. <sup>a</sup>Statistically different from the 1st quartile group, <sup>b</sup>Statistically different from the 1st and 2nd quartile groups, <sup>c</sup>Statistically different from the 1st, 2nd, and 3rd quartile groups.



**Fig. 2.** Association between weight-adjusted waist index (WWI) and (A) total abdominal fat area (TFA) and (B) total abdominal muscle area (TMA).

tween abdominal fat and muscle mass measures with WWI. In contrast, BMI or WC did not reflect the fat and muscle area measures differently. Not only fat area but also muscle area showed positive correlation with WC and BMI (Table 2).

**Age, not race/ethnicity, determines WWI**

Given that the trend for increasing WWI with increasing age, we examined the distribution of WWI according to age in all participants in the MESA cohort. Fig. 3A shows the distribu-

tion curves by age groups and reveals that WWI is positively associated with age. Compared to the 45 to 54 years age group, mean WWI was significantly higher with increasing age groups ( $P=0.001$  for 55–64 years group and  $P<0.001$  for 65–74 and 75–84 years groups). This relationship was consistent in each ethnic group (Supplementary Fig. 2); however, there were no significant associations between BMI and age.

On the other hand, distribution of WWI was similar between ethnic groups. For example, WWI of Asians did not sig-




Fig. 3. Distribution of body mass index (BMI) and weight-adjusted waist index (WWI) by (A) age and (B) race/ethnicity.

nificantly deviate from that of Caucasians and African Americans ( $P=0.512$ , and  $P=0.720$ , respectively) (Fig. 3B). However, BMI of Asians was significantly lower than that of other ethnic groups ( $P<0.001$  for all).

#### WWI reflects body compositional changes with aging


We examined the longitudinal changes in CT-derived fat and muscle area measures and their associations with anthropometric index. Over 38.6 months, the mean TFA and VFA increased, but the TMA decreased in all participants (Supplementary Table 4). Mean SFA increased significantly only in men, not in women. During the same period, BMI did not change, but the WC and WWI significantly increased. Increase in WWI was commonly observed in all age groups, but the direction of changes in BMI and WC differed with age (Supplementary Table 5).

To examine whether longitudinal changes in WWI were associated with increasing fat mass and decreasing muscle area, we examined the percentage changes of TFA and TMA according to quartiles of changes in WWI (delta WWI) (Fig. 4). We found that as WWI increased, TFA increased ( $P<0.001$ ) and TMA decreased, although changes in TMA were not statistically significant ( $P=0.963$ ). In contrast, as BMI increased, both TFA and TMA increased ( $P<0.001$  and  $P=0.056$ , respectively).

#### DISCUSSION

In this study, we found that WWI was positively associated with abdominal fat measures, but negatively with abdominal muscle mass measures, regardless of race/ethnicity. Also, we showed that WWI increased with age in all races, but there was





**Fig. 4.** The association between change of (A) weight-adjusted waist index (WWI), (B) body mass index (BMI) and age-adjusted change of body composition measures. TFA, total abdominal fat area; TMA, total abdominal muscle area.

a small difference in the mean values and distribution between races. Finally, we determined that WWI also reflects abdominal composition changes with aging, i.e., increase in abdominal fat mass and decrease in muscle mass. Collectively, WWI could be an indicator of fat and muscle composition changes related to aging commonly applicable to multiethnic groups.

As the role of muscle or lean mass in the development of age-related disorders has been elucidated, measurement of muscle mass has become increasingly important as a measure of body composition. However, estimation of muscle mass without a special device is difficult, and no anthropometric index so far has been sufficiently verified as an indicator of muscle mass [15,16]. Similar to a previous paper on WWI in Korean elderly people, the results presented above indicate that WWI was differentially associated with fat and muscle mass in a multi-ethnic population. However, this study is limited because the analyses were based solely on abdominal CT scan-derived measures, and not on the whole-body composition data. Although we found weaker associations of WWI with muscle mass measures compared to that of a previous Korean study, we showed the similar patterns with the previous study. These findings suggest that the WWI could serve as a body composition index, which is not limited to East Asians, and can be commonly applied in diverse populations.

Furthermore, we found that WWI was negatively associated with abdominal muscle radiodensity (represented by HU), indicating a positive association between WWI and fat infiltra-

tion in the muscles. Interestingly, the correlation of abdominal muscle radiodensity with WWI ( $r=-0.516$ ) was higher than that with BMI ( $r=-0.242$ ) or WC ( $r=-0.300$ ) (data not shown in the results). Because abdominal muscle radiodensity has been found to be an indicator of myosteatosis, metabolic derangement such as insulin resistance and alterations in adipokines, as well as mortality in cancer patients [17-20], the high correlation of WWI with abdominal muscle mass radiodensity suggests WWI could be an indicator of diverse health outcomes. Although we were unable to assess the associations of WWI with other organ radiodensities, such as liver radiodensity, these associations should be tested in future studies for the broader applications of WWI as a health indicator.

Another significant finding of this study was that the racial difference of WWI distribution was small. BMI, a widely-used anthropometric index, revealed fundamental differences between races with different body shape or different fat distribution; hence, different standards are applied according to race [21,22]. For example, there are different BMI and WC cut-points for defining obesity in Asians compared to Caucasians [23,24]. In this multiethnic cohort, we also found certain differences by races (e.g., mean BMI was significantly lower in Asians than in Whites). However, in the case of WWI, there was no significant difference in the mean and distribution between Caucasians and Asians. This suggests that WWI could be a universal health index applicable to various races/ethnic groups. This finding can be explained partially by the fact that



WWI reflects the ratio of fat and muscle mass, but not the absolute fat amount, and shows a direct proportional relationship with aging.

Regardless of race, WWI was observed to increase with age. In addition, the longitudinal increase in WWI reflected the age-related changes in abdominal fat and muscle composition. Since this is a short-term change spanning over an average of 38 months, a long-term study will be needed to establish an association that is more reliable.

The previous study on WWI only included elderly individuals over 65 years of age, and it was unclear whether the bidirectional association of WWI with fat and muscle mass measures would be reproducible in younger people. This study included 977 participants younger than 65 years (50.2% of total participants) and found that the associations of WWI with fat and muscle measures did not differ with age. Therefore, our data indicated that WWI could be generally applicable to adults of various races and ethnicities, although we still need more evidence with respect to younger adults and adolescents.

This study has several limitations. First, as mentioned earlier, we did not have relevant data regarding the overall body fat and appendicular muscle mass of participants. However, we assumed that the relationship between WWI and various muscle measures, including appendicular muscle mass, would be similar based on previous study results. Second, the number of Asians was relatively small in the current cohort data. Third, the association of WWI with health outcomes remains to be determined. Further studies will be needed to provide the relevant information regarding compatible cut-offs for WWI.

In conclusion, this study showed that WWI had a differential association with fat and muscle mass measures in a multi-ethnic population. Moreover, WWI was closely associated with age and age-related body compositional changes. Further investigation is needed for the validation of this index as a predictor of various health outcomes.

## SUPPLEMENTARY MATERIALS

Supplementary materials related to this article can be found online at <https://doi.org/10.4093/dmj.2021.0294>.

## CONFLICTS OF INTEREST

No potential conflict of interest relevant to this article was reported.

## AUTHOR CONTRIBUTIONS

Conception or design: N.H.K.

Acquisition, analysis, or interpretation of data: J.C., C.A.V., M.H.C., M.A.A., N.H.K.

Drafting the work or revising: J.Y.K., N.H.K.

Final approval of the manuscript: N.H.K.

## ORCID

Ji Yoon Kim <https://orcid.org/0000-0001-6626-2124>

Nam Hoon Kim <https://orcid.org/0000-0002-9926-1344>

## FUNDING

None

## ACKNOWLEDGMENTS

The authors thank the all investigators, staffs, and participants of the MESA study.

## REFERENCES

1. Amato MC, Guarnotta V, Giordano C. Body composition assessment for the definition of cardiometabolic risk. *J Endocrinol Invest* 2013;36:537-43.
2. Britton KA, Massaro JM, Murabito JM, Kreger BE, Hoffmann U, Fox CS. Body fat distribution, incident cardiovascular disease, cancer, and all-cause mortality. *J Am Coll Cardiol* 2013; 62:921-5.
3. Snijder MB, van Dam RM, Visser M, Seidell JC. What aspects of body fat are particularly hazardous and how do we measure them? *Int J Epidemiol* 2006;35:83-92.
4. Romero-Corral A, Lopez-Jimenez F, Sierra-Johnson J, Somers VK. Differentiating between body fat and lean mass-how should we measure obesity? *Nat Clin Pract Endocrinol Metab* 2008;4:322-3.
5. Choi KM. Sarcopenia and sarcopenic obesity. *Endocrinol Metab (Seoul)* 2013;28:86-9.
6. Fielding RA, Vellas B, Evans WJ, Bhasin S, Morley JE, Newman AB, et al. Sarcopenia: an undiagnosed condition in older adults: current consensus definition: prevalence, etiology, and consequences. International working group on sarcopenia. *J Am Med Dir Assoc* 2011;12:249-56.

7. Romero-Corral A, Somers VK, Sierra-Johnson J, Jensen MD, Thomas RJ, Squires RW, et al. Diagnostic performance of body mass index to detect obesity in patients with coronary artery disease. *Eur Heart J* 2007;28:2087-93.
8. Romero-Corral A, Somers VK, Sierra-Johnson J, Thomas RJ, Collazo-Clavell ML, Korinek J, et al. Accuracy of body mass index in diagnosing obesity in the adult general population. *Int J Obes (Lond)* 2008;32:959-66.
9. Kim NH, Lee J, Kim TJ, Kim NH, Choi KM, Baik SH, et al. Body mass index and mortality in the general population and in subjects with chronic disease in Korea: a nationwide cohort study (2002-2010). *PLoS One* 2015;10:e0139924.
10. Antonopoulos AS, Oikonomou EK, Antoniadou C, Tousoulis D. From the BMI paradox to the obesity paradox: the obesity-mortality association in coronary heart disease. *Obes Rev* 2016;17:989-1000.
11. Kim NH, Park Y, Kim NH, Kim SG. Weight-adjusted waist index reflects fat and muscle mass in the opposite direction in older adults. *Age Ageing* 2021;50:780-6.
12. Park Y, Kim NH, Kwon TY, Kim SG. A novel adiposity index as an integrated predictor of cardiometabolic disease morbidity and mortality. *Sci Rep* 2018;8:16753.
13. Bild DE, Bluemke DA, Burke GL, Detrano R, Diez Roux AV, Folsom AR, et al. Multi-Ethnic Study of Atherosclerosis: objectives and design. *Am J Epidemiol* 2002;156:871-81.
14. Goodpaster BH, Thaete FL, Kelley DE. Composition of skeletal muscle evaluated with computed tomography. *Ann N Y Acad Sci* 2000;904:18-24.
15. Cruz-Jentoft AJ, Baeyens JP, Bauer JM, Boirie Y, Cederholm T, Landi F, et al. Sarcopenia: European consensus on definition and diagnosis: report of the European Working Group on Sarcopenia in Older People. *Age Ageing* 2010;39:412-23.
16. Rolland Y, Czerwinski S, Abellan Van Kan G, Morley JE, Cesari M, Onder G, et al. Sarcopenia: its assessment, etiology, pathogenesis, consequences and future perspectives. *J Nutr Health Aging* 2008;12:433-50.
17. Amini B, Boyle SP, Boutin RD, Lenchik L. Approaches to assessment of muscle mass and myosteatosis on computed tomography: a systematic review. *J Gerontol A Biol Sci Med Sci* 2019;74:1671-8.
18. Tilves C, Zmuda JM, Kuipers AL, Carr JJ, Terry JG, Wheeler V, et al. Associations of thigh and abdominal adipose tissue radiodensity with glucose and insulin in nondiabetic African-ancestry men. *Obesity (Silver Spring)* 2020;28:404-11.
19. Vella CA, Cushman M, Van Hollebeke RB, Allison MA. Associations of abdominal muscle area and radiodensity with adiponectin and leptin: the multiethnic study of atherosclerosis. *Obesity (Silver Spring)* 2018;26:1234-41.
20. Sjoblom B, Gronberg BH, Wentzel-Larsen T, Baracos VE, Hjerstad MJ, Aass N, et al. Skeletal muscle radiodensity is prognostic for survival in patients with advanced non-small cell lung cancer. *Clin Nutr* 2016;35:1386-93.
21. Rahman M, Temple JR, Breitkopf CR, Berenson AB. Racial differences in body fat distribution among reproductive-aged women. *Metabolism* 2009;58:1329-37.
22. Zhu S, Heymsfield SB, Toyoshima H, Wang Z, Pietrobelli A, Heshka S. Race-ethnicity-specific waist circumference cutoffs for identifying cardiovascular disease risk factors. *Am J Clin Nutr* 2005;81:409-15.
23. Obesity: preventing and managing the global epidemic: report of a WHO consultation. *World Health Organ Tech Rep Ser* 2000;894:i-xii, 1-253.
24. Seo MH, Lee WY, Kim SS, Kang JH, Kang JH, Kim KK, et al. 2018 Korean Society for the Study of Obesity guideline for the management of obesity in Korea. *J Obes Metab Syndr* 2019;28:40-5.