

UC Davis

UC Davis Previously Published Works

Title

Draft Genome Sequences of Salmonella Lysozyme Gene Knockout Mutants

Permalink

<https://escholarship.org/uc/item/8f44g7rw>

Journal

Microbiology Resource Announcements, 5(23)

ISSN

2576-098X

Authors

Arabyan, Narine
Huang, Bihua C
Weimer, Bart C

Publication Date

2017-06-08

DOI

10.1128/genomea.00519-17

Peer reviewed



Draft Genome Sequences of *Salmonella* Lysozyme Gene Knockout Mutants

Narine Arabyan,^{a,b} Bihua C. Huang,^{a,b}  Bart C. Weimer^{a,b}

Department of Population Health and Reproduction, School of Veterinary Medicine, University of California, Davis, Davis, California, USA^a; 100K Pathogen Genome Project, UC Davis, Davis, California, USA^b

ABSTRACT Lysozyme enzymes hydrolyze the β -1,4-glycosidic bond in oligosaccharides. These enzymes are part of a broad group of glucoside hydrolases that are poorly characterized; however, they are important for growth and are being recognized as emerging virulence factors. This is the release of four lysozyme-encoding-gene-deletion mutants in *Salmonella enterica* serovar Typhimurium LT2.

Lysozyme enzymes belong to the glucoside hydrolase 24 (GH24) family (1). GHs play an important role during infection by altering the host glycan structure to gain access to the host epithelial cells by binding to terminal monosaccharides to initiate glycan degradation (2). Lysozyme enzymes recognize host GlcNAc containing glycans in the form of N-glycans, O-glycans, glycolipids, glycoproteins, and glucosaminoglycans during infection (3) for digestion, and hence may be new virulence factors due to cleavage of the β -1,4-glycosidase bond. These GlcNAc molecules are linked to monosaccharides in the glycan via a β -1,4-glycosidic bond (4) that can be cleaved by enzymes from *Salmonella* with lysozyme activity during host association.

Lysozymes with β -1,4-glycosidase activity are also involved during the secretion of proteins, which is central for the virulence of all pathogenic bacteria (1). Gram-negative organisms translocate proteins across the peptidoglycan that is composed of linear chains of N-acetylglucosamine (GlcNAc) and N-acetylmuramic acid (MurNAc), and the alternating sugars are connected by β -1,4-glycosidic bonds (5–7). The peptidoglycan structure is a physical barrier for the assembly of macromolecular complexes and for the transport of proteins. For this reason, all bacterial lysozymes degrade the peptidoglycan to allow the assembly of type III or type IV secretion systems essential for virulence, flagella, or conjugation (8, 9). This remodeling creates gaps in the peptidoglycan necessary for the assembly of these macromolecular systems. Intracellular pathogenic bacteria, such as *Brucella abortus*, use lysozyme during the early stages of intracellular replication (8).

Four *Salmonella enterica* serovar Typhimurium LT2 lysozyme mutants (Δ STM1028, Δ STM2612, Δ STM2715.S, and Δ STM3605 mutants) were constructed in the Weimer laboratory (UC Davis, Davis, CA) (2), as described by Datsenko and Wanner (10). Cultures were grown on 1.5% Luria-Bertani (LB) agar (Difco, Franklin Lakes, NJ) with 10 μ g/ml chloramphenicol at 37°C and lysed (11); genomic DNA (gDNA) was extracted (12) and checked for quality (13); and sequencing libraries were constructed using the Kapa HyperPlus kit, with enzymatic-based fragmentation (13), and indexed with Weimer 384 TS-LT DNA barcodes (Integrated DNA Technologies, Inc., Coralville, IA, USA) at 192 genomes/lane. The final libraries had average sizes of 350 to 450 bp (14, 15). All genomes were sequenced on an Illumina HiSeq 4000 using PE150 (13, 16, 17) at the UC Davis DNA Technologies Core (Davis, CA). Genome sequences were *de novo* assembled using CLC Workbench version 6.5.1 (Qiagen), with default parameters.

This work was done as part of the 100K Pathogen Genome Project (<http://www.100kgenomes.org>), which is a large-scale sequencing consortium that uses next-generation sequencing methods to create genome databases for use in public health,

Received 24 April 2017 Accepted 28 April 2017 Published 8 June 2017

Citation Arabyan N, Huang BC, Weimer BC. 2017. Draft genome sequences of *Salmonella* lysozyme gene knockout mutants. Genome Announc 5:e00519-17. <https://doi.org/10.1128/genomeA.00519-17>.

Copyright © 2017 Arabyan et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Bart C. Weimer, bcweimer@ucdavis.edu.

TABLE 1 *Salmonella enterica* serovar Typhimurium LT2 deletion mutants with lysozyme activity

GenBank accession no.	SRA accession no.	Isolate name	Gene deleted	Enzyme activity	No. of contigs	Coverage (×)	Total genome size (bp)	No. of CDSs ^a
MZNN00000000	SRR5288766	BCW8410	$\Delta STM1028$	Lysozyme	68	156	4,894,775	4,816
MZNO00000000	SRR5288765	BCW8422	$\Delta STM2612$	Lysozyme	66	138	4,894,815	4,814
MZNP00000000	SRR5288764	BCW8423	$\Delta STM2715.S$	Prophage lysozyme	67	138	4,894,604	4,807
MZYU00000000	SRR5288741	BCW8430	$\Delta STM3605$	Phage endolysin	59	79	4,893,277	4,803

^aCDSs, coding sequences.

food safety, and environmental science, where it is critical to capture genome diversity. This project is focused on sequencing genomes of bacteria from the environment, plants, animals, and humans worldwide, providing new insights into the genetic diversity of pathogens and the microbiome.

Accession number(s). All sequences are publicly available and can be found at the 100K Project bioproject (NCBI PRJNA186441) in the Sequence Read Archive (<http://www.ncbi.nlm.nih.gov/sra>), and genome assemblies can be found in NCBI GenBank (see accession numbers in Table 1).

ACKNOWLEDGMENTS

B.C.W. is grateful for the funding that was contributed from Mars, Inc., NIH (grants 1R01HD065122-01A1 and NIH-U24-DK097154) and an Agilent Technologies Thought Leader Award.

REFERENCES

- Mushegian AR, Fullner KJ, Koonin EV, Nester EW. 1996. A family of lysozyme-like virulence factors in bacterial pathogens of plants and animals. *Proc Natl Acad Sci U S A* 93:7321–7326. <https://doi.org/10.1073/pnas.93.14.7321>.
- Arabyan N, Park D, Foutouhi S, Weis AM, Huang BC, Williams CC, Desai P, Shah J, Jeannotte R, Kong N, Lebrilla CB, Weimer BC. 2016. *Salmonella* degrades the host glycoalkaloid leading to altered infection and glycan remodeling. *Sci Rep* 6:29525. <https://doi.org/10.1038/srep29525>.
- Jacobs H, Mink SN, Duke K, Bose D, Cheng ZQ, Howlett S, Ferrier GR, Light RB. 2005. Characterization of membrane *N*-glycan binding sites of lysozyme for cardiac depression in sepsis. *Intensive Care Med* 31: 129–137. <https://doi.org/10.1007/s00134-004-2487-y>.
- Stanley P, Schachter H, Taniguchi N. 2009. *N*-Glycans. In Varki A, Cummings RD, Esko JD, Freeze HH, Stanley P, Bertozzi CR, Hart GW, Etzler ME (ed), *Essentials of glycobiology*, 2nd ed. Cold Spring Harbor Press, Cold Spring Harbor, NY.
- Allaoui A, Sansonetti PJ, Parsot C. 1993. MxiD, an outer membrane protein necessary for the secretion of the *Shigella flexneri* *Ipa* invasins. *Mol Microbiol* 7:59–68. <https://doi.org/10.1111/j.1365-2958.1993.tb01097.x>.
- Kirby AJ. 1987. Mechanism and stereoelectronic effects in the lysozyme reaction. *CRC Crit Rev Biochem* 22:283–315. <https://doi.org/10.3109/10409238709086959>.
- Pei J, Grishin NV. 2005. COG3926 and COG5526: a tale of two new lysozyme-like protein families. *Protein Sci* 14:2574–2581. <https://doi.org/10.1110/ps.051656805>.
- Del Giudice MG, Ugalde JE, Czubener C. 2013. A lysozyme-like protein in *Brucella abortus* is involved in the early stages of intracellular replication. *Infect Immun* 81:956–964. <https://doi.org/10.1128/IAI.01158-12>.
- Koraimann G. 2003. Lytic transglycosylases in macromolecular transport systems of Gram-negative bacteria. *Cell Mol Life Sci* 60:2371–2388. <https://doi.org/10.1007/s00018-003-3056-1>.
- Datsenko KA, Wanner BL. 2000. One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products. *Proc Natl Acad Sci U S A* 97:6640–6645. <https://doi.org/10.1073/pnas.120163297>.
- Jeannotte R, Lee E, Kong N, Ng W, Kelly L, Weimer BC. 2014. High-throughput analysis of foodborne bacterial genomic DNA using Agilent 2200 TapeStation and genomic DNA ScreenTape system. Agilent Technologies application note. Agilent Technologies, Santa Clara, CA. <https://www.agilent.com/cs/library/applications/5991-4003EN.pdf>.
- Kong N, Ng W, Lee V, Kelly L, Weimer BC. 2013. Production and analysis of high molecular weight genomic DNA for NGS pipelines using Agilent DNA extraction kit (p/n 200600). Agilent Technologies application note. Agilent Technologies, Santa Clara, CA. <https://www.agilent.com/cs/library/applications/5991-3722EN.pdf>.
- Weis AM, Huang BC, Storey DB, Kong N, Chen P, Arabyan N, Gilpin B, Mason C, Townsend AK, Smith WA, Byrne BA, Taff CC, Weimer BC. 2017. Large-scale release of *Campylobacter* draft genomes: resources for food safety and public health from the 100K pathogen genome project. *Genome Announc* 5(1):e00925-16. <https://doi.org/10.1128/genomeA.00925-16>.
- Kong N, Ng W, Foutouhi A, Huang BH, Kelly L, Weimer BC. 2014. Quality control of high-throughput library construction pipeline for KAPA HTP library using an Agilent 2200 TapeStation. In Agilent Technologies application note. Agilent Technologies, Santa Clara, CA. <http://www.agilent.com/cs/library/applications/5991-5141EN.pdf>.
- Kong N, Thao K, Huang C, Appel M, Lappin S, Knapp L, Kelly L, Weimer BC. 2014. Automated library construction using KAPA library preparation kits on the Agilent NGS workstation yields high-quality libraries for whole-genome sequencing on the Illumina platform. Agilent Technologies application note. Agilent Technologies, Santa Clara, CA. <http://www.agilent.com/cs/library/applications/5991-4296EN.pdf>.
- Lüdeke CH, Kong N, Weimer BC, Fischer M, Jones JL. 2015. Complete genome sequences of a clinical isolate and an environmental isolate of *Vibrio parahaemolyticus*. *Genome Announc* 3(2):e00216-15. <https://doi.org/10.1128/genomeA.00216-15>.
- Weis AM, Clothier KA, Huang BC, Kong N, Weimer BC. 2016. Draft genome sequences of *Campylobacter jejuni* strains that cause abortion in livestock. *Genome Announc* 4(6):e01324-16. <https://doi.org/10.1128/genomeA.01324-16>.