

UC Davis

UC Davis Previously Published Works

Title

Conserved Transcriptional Unit Organization of the Cag Pathogenicity Island among Helicobacter pylori Strains

Permalink

<https://escholarship.org/uc/item/8q82656x>

Authors

Ta, Linda H
Hansen, Lori M
Sause, William E
et al.

Publication Date

2012

DOI

10.3389/fcimb.2012.00046

Peer reviewed



Conserved transcriptional unit organization of the *cag* pathogenicity island among *Helicobacter pylori* strains

Linda H. Ta¹, Lori M. Hansen¹, William E. Sause², Olga Shiva³, Aram Millstein¹, Karen M. Ottemann², Andrea R. Castillo^{2*} and Jay V. Solnick^{1*}

¹ Departments of Medicine and Microbiology & Immunology, Center for Comparative Medicine, University of California, Davis, Davis, CA, USA

² Department of Microbiology and Environmental Toxicology, University of California Santa Cruz, Santa Cruz, CA, USA

³ Department of Biology, Eastern Washington University, Cheney, WA, USA

Edited by:

D. Scott Merrell, Uniformed Services University, USA

Reviewed by:

Jeong-Heon Cha, Yonsei University, South Korea

Hilde De Reuse, Institut Pasteur, France

*Correspondence:

Andrea R. Castillo, Department of Biology, Eastern Washington University, Cheney, WA 99004, USA.
e-mail: acastillo@ewu.edu;

Jay V. Solnick, Center for Comparative Medicine, University of California Davis, Davis, CA 95616, USA.
e-mail: jvsolnick@ucdavis.edu

The *Helicobacter pylori* *cag* pathogenicity island (*cag* PAI) encodes a type IV secretion system that is more commonly found in strains isolated from patients with gastroduodenal disease than from those with asymptomatic gastritis. Genome-wide organization of the transcriptional units in *H. pylori* strain 26695 was recently established using RNA sequence analysis (Sharma et al., 2010). Here we used quantitative reverse-transcription polymerase chain reaction of open reading frames and intergenic regions to identify putative *cag* PAI operons in *H. pylori*; these operons were analyzed further by transcript profiling after deletion of selected promoter regions. Additionally, we used a promoter-trap system to identify functional *cag* PAI promoters. The results demonstrated that expression of genes on the *H. pylori* *cag* PAI varies by nearly five orders of magnitude and that the organization of *cag* PAI genes into transcriptional units is conserved among several *H. pylori* strains, including, 26695, J99, G27, and J166. We found evidence for 20 transcripts within the *cag* PAI, many of which likely overlap. Our data suggests that there are at least 11 operons: *cag1-4*, *cag3-4*, *cag10-9*, *cag8-7*, *cag6-5*, *cag11-12*, *cag16-17*, *cag19-18*, *cag21-20*, *cag23-22*, and *cag25-24*, as well as five monocistronic genes (*cag4*, *cag13*, *cag14*, *cag15*, and *cag26*). Additionally, the location of four of our functionally identified promoters suggests they are directing expression of, in one case, a truncated version of *cag26* and in the other three, transcripts that are antisense to *cag7*, *cag17*, and *cag23*. We verified expression of two of these antisense transcripts, those antisense to *cag17* and *cag23*, by reverse-transcription polymerase chain reaction. Taken together, our results suggest that the *cag* PAI transcriptional profile is generally conserved among *H. pylori* strains, 26695, J99, G27, and J166, and is likely complex.

Keywords: *cag* PAI, operon structure, expression

INTRODUCTION

Helicobacter pylori is a Gram-negative bacterium that infects the stomachs of approximately half the human population. Although infection is typically asymptomatic throughout the lifetime of the host, it causes peptic ulcer disease in about 10% of those infected and gastric adenocarcinoma in about 1–3% (Kusters et al., 2006). The best-studied bacterial factor associated with clinical sequelae of *H. pylori* infection is the cytotoxin associated gene pathogenicity island (*cag* PAI). Patients infected with *H. pylori* strains that contain the *cag* PAI are at increased risk for both peptic ulcer and gastric cancer (Kusters et al., 2006). Experimental studies in gerbils (Rieder et al., 2005), mice (Arnold et al., 2011), and rhesus macaques (Hornsby et al., 2008) have also demonstrated the pro-inflammatory effects of the *cag* PAI.

The 40-kb *cag* PAI contains on average 27 genes, several of which encode a type IV secretion apparatus that is required for translocation of the effector molecules CagA (*cag26*) and peptidoglycan into host epithelial cells (Segal et al., 1997; Odenbreit et al., 2000; Rohde et al., 2003; Viala et al., 2004). Of the 27 genes

on the *cag* PAI, 18 are required for the translocation of CagA into host cells and 15 are required to induce transcription of the pro-inflammatory cytokine IL-8 (Fischer et al., 2001; Shaffer et al., 2011). CagA is reliant on the secretion chaperone protein CagF (*cag22*) for recruitment to the type IV translocation channel (Pattis et al., 2007). Upon translocation into the cell, CagA is phosphorylated at C-terminal tyrosine residues by c-Src and other kinases, which results in the activation of receptor tyrosine kinase (RTK)-like signaling pathways (Segal et al., 1997; Selbach et al., 2002). Both phosphorylated and unphosphorylated CagA contribute to *H. pylori* pathogenesis via multiple mechanisms, including the disruption of the cytoskeleton, interruption of cellular signaling, and interference with adhesion between adjacent cells (Backert and Selbach, 2008).

Several studies have provided a glimpse of the *cag* PAI transcriptional unit organization. One initial study employed a urease transcription fusion to check for promoters in nine *cag* PAI DNA regions that were upstream of groups of co-directional genes (Joyce et al., 2001). This analysis determined that there were at least

five promoters on the *cag* PAI. Another early study identified the promoters responsible for regulating *cagA* and *cagB* (Spohn et al., 1997). A more recent genome-wide transcriptional unit analysis that used RNA sequencing identified 14 transcriptional units within the *cag* PAI. Additionally, they found many potential small regulatory RNAs (Sharma et al., 2010). Other studies have suggested that several *cag* PAI genes are differentially regulated *in vivo* compared to *in vitro* (Joyce et al., 2001; Boonjakuakul et al., 2005; Castillo et al., 2008b). In one such study, an *in vivo* induced promoter called *Pivi66*, was internal to the *cag7* gene (Castillo et al., 2008b), which suggested that promoters may not always be within intergenic regions.

Here we sought to determine the conservation of operon structure in the *cag* PAI among *H. pylori* strains, and to identify promoters responsible for the transcription of *cag* PAI genes in strains 26695, J99, and G27, whose genomes are sequenced (Tomb et al., 1997; Alm et al., 1999), and in strain J166 that we and others have used to infect rhesus macaques (Hornsby et al., 2008). Operon structure was first predicted by a gene expression analysis that used quantitative reverse-transcriptase polymerase chain reaction (qRT-PCR) for both open reading frames (ORFs) and intergenic regions. The predicted putative operons were further defined by qRT-PCR after deletion of selected promoter regions. Since our transcription analyses suggested a potentially complex operon structure, we augmented these studies with a non-biased promoter-trap study that identified *cag* PAI promoters as DNA regions capable of directing expression of a heterologous reporter. Our results demonstrate that there is remarkable consistency across strains in the expression of genes in the *cag* PAI, which is organized into at least 20 transcriptional units.

MATERIALS AND METHODS

BACTERIAL STRAINS AND CULTURE

Helicobacter pylori strains 26695 (Tomb et al., 1997), J99 (Alm et al., 1999), J166 (Hornsby et al., 2008), and ACHP17 (mG27 *HP0294/295::res1-aphA3-res1*; Castillo et al., 2008a) were used for these studies. DNA and RNA for qRT-PCR were prepared from strains cultured on *Brucella* agar (Difco Laboratories, Detroit, MI, USA) containing 5% bovine calf serum (Invitrogen Life Technologies, Carlsbad, CA, USA) supplemented with 5 μ g/mL trimethoprim, 10 μ g/mL vancomycin, 2.5 IU/mL polymyxin B, 2.5 μ g/mL amphotericin B (TVPA, all from Sigma, St. Louis, MO, USA) and incubated at 37°C with an atmosphere that contained 5% CO₂. Plate grown bacteria were then transferred to *Brucella* broth containing bovine calf serum with TVPA and incubated at 37°C in 5% CO₂ with gentle rotation at 60 rpm. The OD₆₀₀ was determined for each culture 18–24 h after inoculation. The promoter reporter *H. pylori* strain ACHP17 and strain G27 from which RNA was isolated for RT-PCR were grown under microaerobic conditions (10% CO₂, 5% O₂, and 85% N₂) at 37°C on columbia blood agar plates with 4% (w/v) columbia agar base, 5% (w/v) defibrinated horse blood (Hemostat labs), 0.2% (w/v) β -cyclodextrin, 10 μ g/mL vancomycin, 50 μ g/mL cycloheximide, 5 μ g/mL cefsulodin, 8 μ g/mL amphotericin B, 2.5 IU/mL polymyxin and 5 μ g/mL trimethoprim. *H. pylori* strains were stored at –80°C in brain heart infusion media supplemented with 10% fetal bovine serum, 1% (w/v) β -cyclodextrin, 25% glycerol, and 5% dimethyl sulfoxide.

Escherichia coli strain DH10B (Grant et al., 1990) was grown at 37°C in Luria–Bertani (LB) broth (1% w/v tryptone, 0.5% w/v yeast extract, 0.5% w/v NaCl), with 100 μ g/mL ampicillin. *E. coli* was also grown on solid LB media consisting of LB broth with 1.5% (w/v) agar. All antibiotics were purchased from Sigma-Aldrich, Fisher, or ISC BioExpress. All culture media were purchased from Remel, Fisher, or Difco unless otherwise indicated.

RNA AND DNA EXTRACTION

At OD₆₀₀ 0.4–0.5 (early exponential growth phase) 2 mL aliquots were taken from *H. pylori* liquid cultures and centrifuged at 16,000 \times g for 30 s at room temperature. Supernatants were removed and 1 mL of TriZol (Invitrogen) was immediately added. Samples were vortexed and RNA was extracted according to the manufacturer's directions. RNA was treated with DNase I (Roche Applied Sciences, Mannheim, Germany), purified using an RNeasy clean up kit (QIAGEN, Inc., Valencia, CA, USA), and suspended in ultra pure water (Invitrogen) at a concentration of 20 ng/ μ L.

DNA was extracted from plate grown bacteria using a DNeasy Tissue Kit (QIAGEN). DNA samples were diluted in ultra pure water to a concentration of 5 ng/ μ L and stored at –20°C.

RT-PCR TO DETECT PROMOTERS, PIII, PIX, AND PXII

Reverse-transcriptase polymerase chain reactions were carried out using the Super Script One-step RT-PCR kit with Platinum Taq (Invitrogen). One hundred or 250 ng of RNA was used as a template for each RT-PCR reaction. For the reverse-transcription step (55°C for 30 min), only the oligonucleotide that was complementary to the putative transcript was included in the reaction, PIIIR (5'-cctagcgaccaaagcgatgaa-3'), PIXR (5'-gaaactgctaagaatcagtg-3'), and PXIIR (5'-cgctcattaatcaaatagaacaaagc-3'). The reverse-transcriptase in these reactions was then inactivated by incubation at 94°C for 5 min. Prior to starting the PCR program (35 cycles, 94°C/30 s, 55°C/30 s, 72°C/30 s) the reactions were briefly incubated on ice (~1 min) while the second oligonucleotides, PIIIF (5'-cattgtggtcttcccgaagc-3'), PIXF (5'-cactcttgccataaagccc-3'), and PXIIF (5'-ctgagacgacaagctatgatttc-3') were added. Oligonucleotides for our positive control were HP188F (5'-ccactataaaagagatcttcaagcggaagc-3') and HP187R (5'-gcttgccctcggtgtctgcac-3'); HP187R was present in the RT reaction and both HP187R and HP188F were present in the PCR reaction. As a control for amplification, each set of oligonucleotides was used in a PCR reaction with DNA as the template. Additionally, each set of oligonucleotides was used in an RT-PCR reaction with the RNA template and Platinum Taq only. This control was done to verify our RNA samples were DNA free.

qRT-PCR AND AGAROSE GEL ELECTROPHORESIS

Quantitative real time RT-PCR was performed with primer pairs specific for each *cag* gene (Table A1 in Appendix) and for each intergenic region (Table A2 in Appendix), using methods essentially as described (Boonjakuakul et al., 2004, 2005). In brief, RT and PCR were performed in a single 20 μ L reaction mixture using the thermostable recombinant *Tth* (*rTth*) DNA polymerase (Applied Biosystems) with 100 ng RNA extracted as described above. In the presence of Mn(OAc)₂, *rTth* has reverse-transcriptase activity and DNA polymerase activity. Two-step amplification was

performed with 45 cycles at 95°C for 20 s followed by 59.5°C for 1 min. Accumulation of PCR product was detected during each cycle by excitation of SYBR green at 490 nm. Relative fluorescence was characterized by a cycle threshold (Ct) value, which was defined as the crossover point of the kinetic curve with an arbitrary fluorescence level set at 150 relative fluorescence units. The absence of contaminating DNA was examined by performing the RT-PCR with MgCl₂, in which *rTth* has DNA polymerase but no RT activity. All qRT-PCR products were electrophoresed on a 2% agarose (Invitrogen) gel to verify correct product size. Transcript abundance was calculated only if the observed Ct with RNA template was less than that of the no-template control, and there was a band of the appropriate size on an agarose gel. Otherwise, transcript was considered absent. All transcript copy numbers were normalized to 16S RNA and the data presented represents the average of duplicate wells.

CONSTRUCTION OF *cag* PAI PROMOTER DELETION MUTANTS

The chloramphenicol resistance conferring *cat* gene from plasmid pNR9589 (Wang and Taylor, 1990) and 1–2 kb DNA fragments of the genes directly flanking the region targeted for deletion were PCR amplified (oligonucleotides in **Table A3** in Appendix) with compatible restriction sites. All three fragments were digested with the appropriate enzymes and ligated with compatibly digested pBluescript SK– (Stratagene, La Jolla, CA, USA) to generate a shuttle plasmid with fragments of the *cag* PAI flanking the *cat* gene. The shuttle plasmid was amplified in *E. coli* Top10 (Invitrogen, Carlsbad, CA, USA), sequence verified, and then used to transform *H. pylori* strain J166 by a standard natural transformation procedure (Salama et al., 2001). *H. pylori* transformants were selected on *Brucella* agar plates with TVPA and 4 µg/mL chloramphenicol. Correct replacement of *cag* PAI DNA regions with the *cat* gene was verified using PCR and DNA sequence analyses.

GENERATING THE *H. PYLORI cag* PAI LIBRARY OF PUTATIVE PROMOTERS

Genomic DNA was isolated from *H. pylori* J166 and mG27 (Wizard genomic prep kit, Promega). The DNA region representing the *cag* PAI was amplified from each strain as a set of 13 PCR products of ~2.5 kb in length with 600 bp of overlap between adjacent PCR products (oligonucleotides in **Table A4** in Appendix). For each strain, the PCR products were pooled, partially digested with *Sau3A*, and ligated to *Bgl*III digested *pcat-T-tnpR* (Castillo et al., 2008a) to generate recombinant plasmids, *pcat-T-caglibmG27-tnpR* and *pcat-T-caglibJ166-tnpR*. After ligation, the recombinant plasmids were transformed into *E. coli* DH10B and the *E. coli* were plated on LB agar with ampicillin. For these strains, ~2193 (*pcat-T-caglibmG27-tnpR*) or 5000 (*pcat-T-caglibmJ166-tnpR*) individual ampicillin resistant (Amp^R) colonies were pooled, grown overnight, and treated (Qiagen miniprep extraction kit, Qiagen) to extract the recombinant plasmids. For a subset of colonies from each library, individual recombinant plasmids were analyzed for the presence and size of a *H. pylori cag* PAI insert. All recombinant plasmids analyzed contained inserts and had an average insert size of 469 bp for *pcat-T-caglibmG27-tnpR* and 96 bp for *pcat-T-caglibmJ166-tnpR*.

To isolate putative promoters, *H. pylori* strain ACHP17 was transformed using natural transformation (Salama et al., 2001) with either *pcat-T-caglibmG27-tnpR* or *pcat-T-caglibJ166-tnpR*, and transformants were selected based on their resistance to chloramphenicol (Cm) on CBA plus 13 µg/mL Cm. Cm resistant (Cm^R) transformants were passed twice on Cm prior to being analyzed for kanamycin sensitivity (Km^S) on CBA plus 15 µg/mL kanamycin.

To examine the diversity of the *cag* PAI library clones in *H. pylori*, 10–30 Cm^R clones were selected from each library and the region upstream of *tnpR* was sequenced using primers *rrnB1* and *tnpRbk75* (Castillo et al., 2008a). The average insert size was 232 bp for *pcat-T-caglibmG27-tnpR* and 100 bp for *pcat-T-caglibJ166-tnpR*. PCR amplicons were sequenced and compared to the 26695 and G27 genomes to assess randomness of the cloned regions. The number of transformants needed to obtain 100% coverage of the *cag* PAI for each library was determined using the formula $N = \ln[1 - P/\ln(1 - I/G)]$ (N = number of independent clones, I = size of averaged cloned fragment, G = size of target genome, and P = probability). These calculations suggested that 791 *pcat-T-caglibmG27-tnpR* and 1840 *pcat-T-caglibJ166-tnpR* transformants would be required for complete coverage of the *cag* PAI.


RESULTS

We used both transcription profiling and a functional genetic approach to define *cag* PAI operons and the putative promoters that regulate them. First, we performed qRT-PCR to determine the mRNA copy number within each ORF and each intergenic region on the *cag* PAI. Our assumption was that contiguous genes transcribed in the same direction, with the presence of intergenic message and similar mRNA copy number, would likely form an operon. Selected putative operons were then further analyzed by deletion of the promoter region and reanalysis of mRNA copy number of downstream genes. We then augmented these analyses by using a non-biased promoter-trap system to find active promoters within the *cag* PAI.

CO-EXPRESSION OF *cag* PAI GENES BASED ON GENE AND INTERGENIC TRANSCRIPT COPY NUMBER

We first calculated the transcript copy number for each gene and intergenic region within the *cag* PAI of *H. pylori* J99, 26695, and J166, using methods described previously (Boonjakuakul et al., 2004, 2005). Briefly, three factors were used to calculate copies per cell: (a) a 10-fold change in starting template concentration corresponds to a 3.3-cycles change in Ct ($2^{3.3} = 10$); (b) 100 ng of RNA equals 10⁶ *H. pylori* cells, and (c) the empirically derived observation that a Ct of 19 corresponds to 1 × 10⁵ copies of starting DNA template (assuming 1 copy per bacterial chromosome). We have previously shown that calculation of mRNA copies/cell using Ct corrected for primer efficiency yields values that are essentially identical to those obtained by the more conventional method using standard curves (Boonjakuakul et al., 2004).

Transcript levels for all genes on the *cag* PAI for each *H. pylori* strain are shown in **Figure 1**. For clarity, intergenic transcript is shown only as present (adjacent bars representing gene transcript levels are shaded identically) or absent (adjacent bars are shaded differently). For example, intergenic transcript was



detected between *cag1* and *cag2* (both black bars) and between *cag10* and *cag9* (both gray bars), but not between *cag15* (gray bar) and *cag14* (black bar). Transcript levels varied within each strain by as much as five orders of magnitude, ranging from about 10 copies/cell to as low as 1 copy per 10,000 cells. These estimates are consistent with our previous studies (Boonjakuakul et al., 2004, 2005) and with estimates of gene expression levels in *Saccharomyces cerevisiae* (Kang et al., 2000) and *E. coli* (Young and Bremer, 1975). The highest transcript abundance was found for *cag26* and for *cag25*. Since *cag26* encodes an effector protein, CagA, secreted via the type IV secretion system, and *cag25* encodes a virB2 ortholog that is thought to encode a pilin protein that forms a multimeric structure (Andrzejewska et al., 2006), it is not surprising that these genes are highly expressed. Although in general, the expression level of genes on the *cag* PAI was similar across the three strains analyzed, there is some variation that appears to occur within the operons predicted by these experiments (Figure 2).

We reasoned that adjacent genes transcribed with ORFs in the same direction, with the presence of intergenic transcript, might represent a single transcriptional unit, particularly if the transcript abundance was similar across genes. Therefore, we initially considered the possibility that the following may represent *cag* PAI operons (numbered in the direction of transcription): *cag1-4*, *cag10-5*, *cag11-12*, *cag16-17*, *cag21-18*, and *cag25-22* (Figure 1). However, there were sometimes marked differences in transcript abundance of genes within these putative operons (e.g., *cag25-22*, Figures 1 and 2). This might occur due to differential decay of the transcript or possibly because the gene is part of more than one transcriptional unit. To address these possibilities, we deleted the genomic region immediately upstream of the translational start of the first gene in each of six putative operons in *H. pylori* strain J166, a region likely to contain the promoter, and then measured *cag* PAI gene transcript abundance. We reasoned that deletion of this region should decrease the expression level of all genes in the transcriptional unit, and leave others unchanged.

Deletion of the putative promoter regions upstream of *cag1*, *cag10*, *cag11*, *cag16*, *cag21*, and *cag25* had differential effects on the expression of downstream genes when compared to the isogenic wild type *H. pylori* J166 strain (Figure 3). Deletion of the region upstream of *cag1* reduced expression of *cag1-3* by three orders of magnitude and *cag4* by only 1.5 orders of magnitude. By contrast, expression of *cag5*, a gene transcribed in the opposite direction of this putative operon, remained essentially unchanged. Deletion of the region upstream of *cag10* reduced expression of both *cag10* and *cag9* by similar levels and had no effect on expression of *cag8-7*. Deletion of the putative promoters upstream of *cag11*, *cag16*, and *cag21* reduced expression of the downstream genes, *cag11-12*, *cag16-17*, and *cag21-18*, but in each case to different levels, ranging from 1 to 3 orders of magnitude (Figure 3). Finally, deletion of the region upstream of *cag25* reduced expression of the downstream genes *cag25-23* to different levels and had no effect on the expression of *cag22*. In some cases, these results make clear predictions about operon structure. For example, our original prediction of *cag10-5* and *cag25-22* as operons was incorrect, since in each case one or more downstream genes did not




FIGURE 2 | Composite gene expression (mRNA copies/cell, normalized to 16S rRNA) for each gene on the *cag* PAI of *H. pylori* strains J166 (open circles), J99 (closed circles), and 26695 (open triangles). Our data represents the average of duplicate reactions.

change appreciably in the promoter knockouts. Thus, *cag10-5* consists of at least two operons, *cag10-9* and *cag8-7*, which also may be organized into one or more transcriptional units. Similarly, *cag25-22* appears to have only *cag25-24* on one transcriptional unit, with *cag22* and perhaps *cag23* on separate transcripts. The variable change we observed in *cag* PAI gene expression after deletion of the predicted upstream promoter again suggests that either the transcripts are being degraded or that there are additional promoters controlling expression of these genes. To identify additional promoters that may contribute to the more complex expression pattern we observed here, we undertook a non-biased promoter-trap approach.

NON-BIASED PROMOTER-TRAP IDENTIFIES ADDITIONAL *cag* PAI PROMOTERS

We next employed a functional identification of *cag* PAI promoters strategy based on the ability of short cloned regions of the *cag* PAI to direct expression of a heterologous promoter. We used a *tnpR* transcriptional reporter developed for *Vibrio cholerae* (Camilli et al., 1994) that had been previously modified to function in *H. pylori* (Camilli et al., 1994; Castillo et al., 2008b). We constructed libraries of putative *cag* PAI promoters using both *H. pylori* strains J166 and G27 as template for PCR; we cloned the *Sau*3A-digested fragments upstream of the promoterless *tnpR* gene in pCT-*tnpR*. If the cloned *cag* PAI region contained a promoter, we predicted it would direct *tnpR* expression and the creation of the TnpR protein. TnpR in turn would catalyze the removal of an unlinked kanamycin resistance (Km^R) cassette and convert the *H. pylori* reporter strain ACHP17 from Km^R to Km^S .


For promoter identification, *H. pylori* strain ACHP17 bearing the *res1-kan-res1* marker was transformed with *pcat-T-caglibmG27-tnpR* or *pcat-T-caglibJ166-tnpR* to Cm^R , followed by screening for retention or loss of the *res1-kan-res1* cassette.

We screened 1060 *H. pylori* *pcat-T-caglibmG27-tnpR* transformants and 1274 *H. pylori* *pcat-T-caglibmJ166-tnpR* transformants, representing 100 and 71% coverage, respectively. This analysis identified 34 and 27 transformants, respectively, that were sensitive to kanamycin and thus had expressed *tnpR*. After removing redundant clones, we determined that the DNA sequences upstream of *tnpR* in these Km^S transformants correspond to 14 unique loci (Table 1). Eleven and four promoters were identified through the screening of our *pcat-T-caglibmG27-tnpR* and *pcat-T-caglibmJ166-tnpR* libraries, respectively; one promoter, PIII (Table 1; Figure 4), was isolated from both libraries.

PROMOTER-TRAP-IDENTIFIED PROMOTERS

We next mapped our promoter-trap-identified promoters onto the *cag* PAI map and compared these promoters to those found in by our initial qRT-PCR analysis and also to the work of others. Several of the promoter-trap-identified promoters were located in *cag* PAI regions that were either predicted by the qRT-PCR or promoter deletion analyses (Table 1). These include the promoters upstream of *cag10* (PVIII), *cag11* (PII), *cag21* (PI), and *cag25* (PV; Figures 3 and 4; Table 1). The promoter-trap approach also identified several possible promoters that were located within operons that might account for the variable gene expression observed after deleting the main promoter (Figures 3 and 4). These include PIV, PX, and PXIII that are located in genomic positions to suggest they contribute to the expression of *cag4*, *cag8-7*, and *cag23-22* (Figures 3 and 4).

The other putative promoters identified in our promoter-trap study are either consistent with *cag* PAI transcripts predicted by other groups, or as of yet, unique. We identified a promoter that is upstream of *cag26* (PXI) and one that is within, and in the same direction as, *cag26* (PXIV). The promoter upstream of *cag26* was identified in work done by Spohn et al. (1997) and more recently



by Sharma et al. (2010) as a promoter that drives expression of *cag26* (Figure 4; Spohn et al., 1997; Sharma et al., 2010). We also identified a unique putative promoter (PVI) that overlaps *cag3* and the adjacent upstream region and is in the correct direction to promote expression of a polycistronic mRNA including *cag3* and *cag4* (Figure 4). Two of our putative promoters were located within *cag7*, one in the same direction (PVII) and one antisense (PIX) to *cag7* (Figure 4). We hypothesize the promoter located within *cag7* contributes to expression of *cag6-5* and the promoter that is antisense to *cag7* may direct expression of a regulatory sRNA. Neither of these promoters has been identified by other studies. Finally, the last two putative promoters we identified, PIII and PXII, were within and antisense to *cag23* and in the 3' end of *cag18* and may direct expression of sRNAs that are antisense to *cag23* and *cag17*, respectively. These promoters are also unique to this study.

Although our transcription, promoter deletion and promoter-trap analyses do not completely overlap, they show reasonable agreement in predicting transcripts and operon structure and are generally consistent with operon structure predicted by others (Table 1, discussion). Taken together our data suggests the existence of at least 20 *cag* PAI transcripts (Figure 4).

PIII AND PXII DIRECT EXPRESSION OF ANTISENSE TRANSCRIPTS

To determine if the promoters PIII, PIX, and PXII direct expression of transcripts that are antisense to *cag23*, *cag7*, and *cag17*, respectively, we carried out additional RT-PCR reactions on RNA isolated from *H. pylori* strain G27. The oligonucleotides (PIIIR, PIXR, and PXIIR) used in the reverse-transcription reactions were located ~100–150 nt downstream of PIII, PIX, and PXII and were antisense to the putative transcripts. For the subsequent PCR reactions in which reverse-transcriptase had been inactivated, the sense oligonucleotides, PIIIF, PIXF, and PXII were added. Amplicons were detected downstream of PIII and PXII in the RT-PCR reactions and were absent in the corresponding polymerase only controls, suggesting that these promoters do in fact direct expression of transcripts (Figure 5). We did not detect a transcript downstream of PIX in our experiments; while it is possible that PIX is not a promoter, it is more likely that the transcript is regulated or is in very low abundance. The promoter-trap system by which PIX was identified was designed to capture low abundant and transient expression events.

DISCUSSION

In this study we used transcript profiling coupled with putative promoter deletion and a non-biased promoter-trap system to analyze expression of *cag* PAI genes and their organization into transcriptional units across several *H. pylori* strains. We found that *cag* PAI gene expression varies by nearly five orders of magnitude across the *cag* PAI, and that expression of *cag* PAI genes is similar across strains 26695, J99, and J166. Based on transcript profiling of *cag* PAI ORFs and intergenic regions, we initially placed *cag* PAI genes into six polycistrons and four monocistrons. However, subsequent promoter deletions coupled with transcript profiling and promoter-trap promoter identification studies suggested *cag* PAI operon structure was much more complex. Our data suggests that there are at least 11 operons: *cag1-4*, *cag3-4*, *cag10-9*, *cag8-7*, *cag6-5*, *cag11-12*, *cag16-17*, *cag19-18*, *cag21-20*, *cag23-22*, and *cag25-24*, as well as five monocistronic genes (*cag4*, *cag13*, *cag14*, *cag15*, *cag26*). Additionally, the location of four of our promoter-trap-identified promoters suggests they direct expression of, in one case, a truncated version of *cag26* and in the other three, transcripts that are antisense to *cag7*, *cag17*, and *cag23*. Using RT-PCR we verified the presence of transcripts that are antisense to *cag17* and *cag23*.


CONSERVATION OF *cag* PAI GENE EXPRESSION AMONG *H. PYLORI* STRAINS

Our transcript profiling of *cag* PAI ORFs and intergenic regions of three *H. pylori* strains, 26695, J99, and J166, suggested that *cag* PAI expression is generally conserved among strains. There were some genes, however, whose expression showed appreciable differences across strains. Potential reasons for these differences may be attributed to one or a combination of the following: (1) difficulty in accurate quantitation of low abundance transcripts, (2) differential stability of the transcripts, and (3) differential strength of the promoters. We suspect that the differences in *cag15* expression between strains may be due to its very low expression *in vitro* (Joyce et al., 2001). The reduced expression of *cag12*, *cag13*, and *cag19* in *H. pylori* strain 26695 compared to that of

Table 1 | Chromosomal location of putative cag PAI promoters.

Putative promoter (length; nt) ± strand	Cag PAI library, genome source	G27 genome position	26695 genome position	Sharma et al. (2010) ^a , transcription start site (TSS)	Joyce et al. (2001) ^b	This study ^c
I (244)–	G27	542756-542513	574379-574136	TSS575200–	Y	Y (P)
II (38)+	G27	530249-530286	563751-563788	TSS564140+ TSS564347+	Y	Y (P)
III (91)+	G27 and J166	544137-544227	575760-575850	None	N	N
IV (104)+	G27	515270-515371	548689-548790	TSS549427+	N	Y (T)
V (71)–	G27	547735-547665	579357-579287	TSS579114–		Y (P)
VI (314)+	G27	514575-514888	547996-548307	None	N	N
VII (234)–	G27	525984-525661	558996-558763	None	N	N
VIII (635)–	G27	530667-530918	564409-563785	TSS564078– TSS564329–	Y	Y (P)
IX (132)+	G27	521375-521506	554800-554931	None	N	N
X (269)–	G27	528142-527874	561644-561372	TSS561595–	N	Y (T)
XI (419)+	G27	547665-548083	579287-579744	TSS579817+	N	N
XII (54)–	J166	539314-539259	570937-570882	None	N	N
XIII (68)–	J166	546797-546736	578420-578359	None	N	Y (T)
XIV (70)+	J166	548866-548927	580539-580604	None	ND	N

^aTranscription start sites (TSS) predicted by Sharma et al. (2010) that are downstream of, or within, the promoter-trap-identified promoter. The 26695 genomic position and DNA strand (+ or –) of the TSS are indicated. None, no TSS near the functionally identified promoter. ^bY, promoter was previously identified by Joyce et al. (2001). N, promoter was not previously identified by Joyce et al. (2001). ^cP, the promoter-trap-identified promoter was within the genomic region deleted in our promoter deletion analysis; T, it is in a location that is consistent with the changes we observe in cag gene expression; N, it was not within a genomic region deleted in our promoter deletion analysis or indicated by changes in cag PAI gene expression.



J99 and J166 is more likely attributed to transcript instability and differences in promoter strength. Our expression findings should allow researchers to more confidently apply our and other cag PAI expression data to unique clinically isolated *H. pylori* strains.

DIFFERENT STUDIES PREDICT SIMILAR cag PAI OPERON STRUCTURE

Our findings are generally consistent with previous predictions of cag PAI promoters, expression and operon structure. First, our promoter-trap and promoter deletion studies identified four of the five cag PAI promoters, upstream of cag1, cag10, cag21, and cag25 (not cag15), that were predicted by Joyce et al. (2001) in

the *H. pylori* Alston strain. However our transcript profiling of cag PAI ORFs and intergenic regions did predict the promoter upstream of the cag15 (Figure 1). The failure of our promoter-trap to identify the promoter upstream of cag15 was not surprising, as Joyce et al. (2001) found that this promoter was only induced in co-culture with epithelial cells or in mice. A similar profile of promoters between the clinically isolated Alston strain and 26695, J99, J166, and G27 again supports conservation of cag PAI operon structure and expression between *H. pylori* strains.

Our promoter analyses also identified promoters upstream of cag25 (cagB) and cag26 (cagA) that were in positions similar to




FIGURE 5 | Reverse-transcriptase polymerase chain reaction (RT-PCR) identifies transcripts downstream of promoter-trap-identified promoters PIII and PXII.

RT-PCR was carried out for the three promoters that potentially directed expression of antisense transcripts, PIII, PIX, and PXII. Three reactions were included for each promoter, a DNA template + DNA polymerase (DNA), an RNA template + DNA polymerase (T) and an RNA template + reverse-transcriptase and DNA polymerase (RT). Amplicons of the correct size were detected for PIII and PXII in the RT reactions but not in the T reactions. An amplicon was not detected for PIX in the RT or T reactions. This supports expression of transcripts downstream of PIII and PXII, but not PIX. The * indicates the 100-bp marker of the 100-bp ladder.

those previously reported by Spohn et al. (1997) for *H. pylori* strain G27, and by Sharma et al. (2010) for *H. pylori* strain 26695. Spohn et al. (1997) identified two transcription start points upstream of *cag25* that are ~200 bp upstream of what we and Sharma et al. (2010) found for *cag25*. All three studies predicted the same start point that is upstream of *cag26*, but we found an additional promoter that is located within *cag26*. The significance of multiple start sites upstream of *cag25* and within *cag26* are, as of yet, unclear. However, a recent study suggests discreet roles for the amino- and carboxy-terminus of Cag26 (CagA) and it is interesting to speculate this promoter could separate Cag26 function by creating a truncated protein (Pelz et al., 2011).

Our transcript profiles obtained from our work were also consistent with many of the 14 *cag* PAI operons identified in the *H. pylori* genome-wide transcript analysis conducted by Sharma et al. (2010). In common, we predicted five polycistrons: *cag1-4*, *cag6-5*, *cag8-7*, *cag11-12*, and *cag16-17*, and the three monocistrons: *cag4*, *cag13*, and *cag26*. Our promoter locations are consistent with their transcripts that start at *cag10*, *cag14*, and *cag25*, but our data did not predict that the transcripts extended to *cag7*, *cag13*, and *cag18*, respectively. We also did not find functional promoters upstream of *cag17* and *cag18* that would suggest they were also expressed as monocistrons. However, in addition to the truncated *cag26* transcript mentioned above, we also identified the following set of transcripts that were not identified by Sharma et al. (2010), including the polycistrons *cag3-4*, *cag21-20*, *cag19-18* and three transcripts that were antisense to *cag7*, *cag17*, and *cag23*. A transcript for *cag15* was also not identified by Sharma et al. (2010), likely due to its very low abundance *in vitro* (Joyce et al., 2001). We speculate that these discrepancies are due to potential issues with transcript abundance and stability here and in Sharma et al. (2010) and incomplete screening of our *cag* PAI promoter libraries.

INCOMPLETE SCREENING OF *cag* PAI PROMOTER LIBRARIES

Outstanding observations in our screening of the *pcat-T-caglibJ166-tnpR* and *pcat-T-caglibG27-tnpR* libraries in *H. pylori* ACHP17 were that we only identified four promoters grouped at the 3'-end of the *cag* PAI from *pcat-T-caglibJ166-tnpR* and that we did not identify promoters from the central region of the *cag* PAI from *pcat-T-caglibG27-tnpR*. We hypothesize that this was due to a combination of two things: (1) incomplete representation of the *cag* PAI region in both of our libraries and then (2) restriction modification system differences that were apparent in transforming our G27 based reporter strain ACHP17 with J166 *cag* PAI DNA. Although our library screening calculations (see Materials and Methods) suggested that we had screened 100% of the *H. pylori* G27 *cag* PAI and 71% of the *H. pylori* J166 *cag* PAI, our control experiments with 10 or 30 randomly selected *H. pylori* transformants, respectively, suggested that our libraries were biased; the *H. pylori caglibG27-tnpR* library was biased toward the left and right ends of the *cag* PAI and the *H. pylori caglibJ166* library was biased toward the right side of the *cag* PAI. Nonetheless, this methodology was very effective at identifying promoters in positions where we observed slight differences in expression of adjacent genes. Specific amplification of *cag* PAI regions (e.g., *cag12-17*) that were underrepresented in our *cag* PAI libraries will ensure better representation of the G27 *cag* PAI region in our library for future *in vivo* analyses.

COMPARING *cag* PAI EXPRESSION *IN VITRO* AND *IN VIVO*

This and previous studies have contributed to building a more complete expression profile of the clinically important *cag* PAI of *H. pylori* grown *in vitro* (Spohn et al., 1997; Joyce et al., 2001; Sharma et al., 2010). The promoters identified by these *in vitro* studies can now be analyzed for their potential regulation during *H. pylori* infection of a host. It is clear in at least two cases that *in vitro* predicted promoters, those upstream of *cag15* and *cag21*, are expressed at higher levels when co-cultured with an epithelial cell monolayer and in mice (Joyce et al., 2001). While we anticipate that a subset of our *in vitro* identified promoters will be regulated *in vivo* and may contribute to virulence, other studies suggest that there is a set of promoters or transcripts uniquely expressed *in vivo* (Scott et al., 2007; Castillo et al., 2008b); analysis of *H. pylori* transcripts isolated from gerbil stomachs predicted that *cag25* is expressed as a monocistron *in vivo* (Scott et al., 2007) and a promoter-trap study identified a unique promoter, *Pivi66*, within *cag7* (Castillo et al., 2008b). Analysis of our *H. pylori cat-T-caglibmG27-tnpR* library in rodents has the potential to identify additional *in vivo* induced *cag* PAI promoters.

ACKNOWLEDGMENTS

This work was supported by Public Health Service Grant (AI42081) from the NIH to Jay V. Solnick, a Research Scholar Grant (RSG-05-249-01-MBC) from the American Cancer Society to Karen M. Ottemann, and an Eastern Washington University Foundation Grant to Andrea R. Castillo. We thank Michael Hornsby, Wendy Axsen, and Thanh Vo for their technical assistance with experiments.

REFERENCES

- Alm, R. A., Ling, L. S., Moir, D. T., King, B. L., Brown, E. D., Doig, P. C., Smith, D. R., Noonan, B., Guild, B. C., DeJonge, B. L., Carmel, G., Tummino, P. J., Caruso, A., Uria-Nickelsen, M., Mills, D. M., Ives, C., Gibson, R., Merberg, D., Mills, S. D., Jiang, Q., Taylor, D. E., Vovis, G. F., and Trust, T. J. (1999). Genomic-sequence comparison of two unrelated isolates of the human gastric pathogen *Helicobacter pylori*. *Nature* 397, 176–180.
- Andrzejewska, J., Lee, S. K., Olbermann, P., Lotzing, N., Katzowitzsch, E., Linz, B., Achtman, M., Kado, C. I., Suerbaum, S., and Josenhans, C. (2006). Characterization of the pilin ortholog of the *Helicobacter pylori* type IV cag pathogenicity apparatus, a surface-associated protein expressed during infection. *J. Bacteriol.* 188, 5865–5877.
- Arnold, I. C., Lee, J. Y., Amieva, M. R., Roers, A., Flavell, R. A., Sparwasser, T., and Muller, A. (2011). Tolerance rather than immunity protects from *Helicobacter pylori*-induced gastric preneoplasia. *Gastroenterology* 140, 199–209.
- Backert, S., and Selbach, M. (2008). Role of type IV secretion in *Helicobacter pylori* pathogenesis. *Cell. Microbiol.* 10, 1573–1581.
- Boonjakuakul, J. K., Canfield, D. R., and Solnick, J. V. (2005). Comparison of *Helicobacter pylori* virulence gene expression in vitro and in the rhesus macaque. *Infect. Immun.* 73, 4895–4904.
- Boonjakuakul, J. K., Syvanen, M., Suryaprasad, A., Bowlus, C. L., and Solnick, J. V. (2004). Transcription profile of *Helicobacter pylori* in the human stomach reflects its physiology in vivo. *J. Infect. Dis.* 190, 946–956.
- Camilli, A., Beattie, D. T., and Mekalanos, J. J. (1994). Use of genetic recombination as a reporter of gene expression. *Proc. Natl. Acad. Sci. U.S.A.* 91, 2634–2638.
- Castillo, A. R., Arevalo, S. S., Woodruff, A. J., and Ottemann, K. M. (2008a). Experimental analysis of *Helicobacter pylori* transcriptional terminators suggests this microbe uses both intrinsic and factor-dependent termination. *Mol. Microbiol.* 67, 155–170.
- Castillo, A. R., Woodruff, A. J., Connolly, L. E., Sause, W. E., and Ottemann, K. M. (2008b). Recombination-based in vivo expression technology identifies *Helicobacter pylori* genes important for host colonization. *Infect. Immun.* 76, 5632–5644.
- Fischer, W., Puls, J., Buhrdorf, R., Gebert, B., Odenbreit, S., and Haas, R. (2001). Systematic mutagenesis of the *Helicobacter pylori* cag pathogenicity island: essential genes for CagA translocation in host cells and induction of interleukin-8. *Mol. Microbiol.* 42, 1337–1348.
- Grant, S. G., Jessee, J., Bloom, F. R., and Hanahan, D. (1990). Differential plasmid rescue from transgenic mouse DNAs into *Escherichia coli* methylation-restriction mutants. *Proc. Natl. Acad. Sci. U.S.A.* 87, 4645–4649.
- Hornsby, M. J., Huff, J. L., Kays, R. J., Canfield, D. R., Bevins, C. L., and Solnick, J. V. (2008). *Helicobacter pylori* induces an antimicrobial response in rhesus macaques in a Cag pathogenicity island-dependent manner. *Gastroenterology* 134, 1049–1057.
- Joyce, E. A., Gilbert, J. V., Eaton, K. A., Plaut, A., and Wright, A. (2001). Differential gene expression from two transcriptional units in the cag pathogenicity island of *Helicobacter pylori*. *Infect. Immun.* 69, 4202–4209.
- Kang, J. J., Watson, R. M., Fisher, M. E., Higuchi, R., Gelfand, D. H., and Holland, M. J. (2000). Transcript quantitation in total yeast cellular RNA using kinetic PCR. *Nucleic Acids Res.* 28, e2.
- Kusters, J. G., Van Vliet, A. H., and Kuipers, E. J. (2006). Pathogenesis of *Helicobacter pylori* infection. *Clin. Microbiol. Rev.* 19, 449–490.
- Odenbreit, S., Puls, J., Sedlmaier, B., Gerland, E., Fischer, W., and Haas, R. (2000). Translocation of *Helicobacter pylori* CagA into gastric epithelial cells by type IV secretion. *Science* 287, 1497–1500.
- Pattis, I., Weiss, E., Laugks, R., Haas, R., and Fischer, W. (2007). The *Helicobacter pylori* CagF protein is a type IV secretion chaperone-like molecule that binds close to the C-terminal secretion signal of the CagA effector protein. *Microbiology* 153, 2896–2909.
- Pelz, C., Steininger, S., Weiss, C., Coscia, F., and Vogelmann, R. (2011). A novel inhibitory domain of *Helicobacter pylori* protein CagA reduces CagA effects on host cell biology. *J. Biol. Chem.* 286, 8999–9008.
- Rieder, G., Merchant, J. L., and Haas, R. (2005). *Helicobacter pylori* cag-type IV secretion system facilitates corpus colonization to induce precancerous conditions in Mongolian gerbils. *Gastroenterology* 128, 1229–1242.
- Rohde, M., Puls, J., Buhrdorf, R., Fischer, W., and Haas, R. (2003). A novel sheathed surface organelle of the *Helicobacter pylori* cag type IV secretion system. *Mol. Microbiol.* 49, 219–234.
- Salama, N. R., Otto, G., Tompkins, L., and Falkow, S. (2001). Vacuolating cytotoxin of *Helicobacter pylori* plays a role during colonization in a mouse model of infection. *Infect. Immun.* 69, 730–736.
- Scott, D. R., Marcus, E. A., Wen, Y., Oh, J., and Sachs, G. (2007). Gene expression in vivo shows that *Helicobacter pylori* colonizes an acidic niche on the gastric surface. *Proc. Natl. Acad. Sci. U.S.A.* 104, 7235–7240.
- Segal, E. D., Lange, C., Covacci, A., Tompkins, L. S., and Falkow, S. (1997). Induction of host signal transduction pathways by *Helicobacter pylori*. *Proc. Natl. Acad. Sci. U.S.A.* 94, 7595–7599.
- Selbach, M., Moese, S., Meyer, T. F., and Backert, S. (2002). Functional analysis of the *Helicobacter pylori* cag pathogenicity island reveals both VirD4-CagA-dependent and VirD4-CagA-independent mechanisms. *Infect. Immun.* 70, 665–671.
- Shaffer, C. L., Gaddy, J. A., Loh, J. T., Johnson, E. M., Hill, S., Hennig, E. E., McClain, M. S., McDonald, W. H., and Cover, T. L. (2011). *Helicobacter pylori* exploits a unique repertoire of type IV secretion system components for pilus assembly at the bacteria-host cell interface. *PLoS Pathog.* 7, e1002237. doi:10.1371/journal.ppat.1002237
- Sharma, C. M., Hoffmann, S., Darfeuille, F., Reignier, J., Findeiss, S., Sittka, A., Chabas, S., Reiche, K., Hackermuller, J., Reinhardt, R., Stadler, P. F., and Vogel, J. (2010). The primary transcriptome of the major human pathogen *Helicobacter pylori*. *Nature* 464, 250–255.
- Spohn, G., Beier, D., Rappuoli, R., and Scarlato, V. (1997). Transcriptional analysis of the divergent CagAB genes encoded by the pathogenicity island of *Helicobacter pylori*. *Mol. Microbiol.* 26, 361–372.
- Tomb, J. F., White, O., Kerlavage, A. R., Clayton, R. A., Sutton, G. G., Fleischmann, R. D., Ketchum, K. A., Klenk, H. P., Gill, S., Dougherty, B. A., Nelson, K., Quackenbush, J., Zhou, L., Kirkness, E. F., Peterson, S., Loftus, B., Richardson, D., Dodson, R., Khalak, H. G., Glodek, A., Mckenney, K., Fitzgerald, L. M., Lee, N., Adams, M. D., Hickey, E. K., Berg, D. E., Gocayne, J. D., Utterback, T. R., Peterson, J. D., Kelley, J. M., Cotton, M. D., Weidman, J. M., Fujii, C., Bowman, C., Watthey, L., Wallin, E., Hayes, W. S., Borodovsky, M., Karp, P. D., Smith, H. O., Fraser, C. M., and Venter, J. C. (1997). The complete genome sequence of the gastric pathogen *Helicobacter pylori*. *Nature* 388, 539–547.
- Viala, J., Chaput, C., Boneca, I. G., Cardona, A., Girardin, S. E., Moran, A. P., Athman, R., Memet, S., Huerre, M. R., Coyle, A. J., Distefano, P. S., Sansonetti, P. J., Labigne, A., Bertin, J., Philpott, D. J., and Ferrero, R. L. (2004). Nod1 responds to peptidoglycan delivered by the *Helicobacter pylori* cag pathogenicity island. *Nat. Immunol.* 5, 1166–1174.
- Wang, Y., and Taylor, D. E. (1990). Chloramphenicol resistance in *Campylobacter coli*: nucleotide sequence, expression, and cloning vector construction. *Gene* 94, 23–28.
- Young, R., and Bremer, H. (1975). Analysis of enzyme induction in bacteria. *Biochem. J.* 152, 243–254.

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 02 November 2011; accepted: 17 March 2012; published online: 20 April 2012.

Citation: Ta LH, Hansen LM, Sause WE, Shiva O, Millstein A, Ottemann KM, Castillo AR and Solnick JV (2012) Conserved transcriptional unit organization of the cag pathogenicity island among *Helicobacter pylori* strains. *Front. Cell. Inf. Microbiol.* 2:46. doi: 10.3389/fcimb.2012.00046

Copyright © 2012 Ta, Hansen, Sause, Shiva, Millstein, Ottemann, Castillo and Solnick. This is an open-access article distributed under the terms of the Creative Commons Attribution Non Commercial License, which permits non-commercial use, distribution, and reproduction in other forums, provided the original authors and source are credited.

APPENDIX

Table A1 | Open reading frame primer pairs selected for real time RT-PCR.

Primer	Strain ^a	Forward primer (5'-3')	Reverse primer (5'-3')
Cag1	U	GCTATGGGGATTGTTGGGATAA	GCTTCAGTTGGTTCGTTGGTAA
Cag2	A	TGTAAGGGCGTTTTACGAGAA	TTTGGGATATTTAGGATTTTGTGAA
Cag2	B	ACTGTAAGGGCGTTTTACGAGAA	GGGCGTGTTTTACCAATGTAA
Cag2	C	GAATTTGTCCAATAGGGGATTTTTA	AACAGAGAGATTGCCTTTTTTTGTG
Cag3	A, C	GACACCTTGAATGTGAATGACAAA	GTTGTAATACCCATTGACTTGCTCTAA
Cag3	B	AAACAAGAGCGATGGGAACCTTA	TAGGGGCGAACACACTTCA
Cag4	A, B	GCGAGAAAAATCCCTTAAAGACA	GTGTTTCATTCCCCATTGTAA
Cag4	C	CCATCACTTTC AAGCAATACGA	GAGCGTTTTAGAAATAGGTAGGGTAGAA
Cag5	A, C	CGGACTAGAGATATAGGAGCGAATAA	GCCACTGCCTGCCTACAA
Cag5	B	TAGTAGGAGCAATCAAGCCAATAAA	TAGGGACATAGGAGCGAATAAAAA
virB11	U	CCTCTAAGGCATGCTACTGAAGAA	TCGCTAAATTGCTGCTCAAAA
Cag6	U	GAAAGCACGTATCAAAAATGAACTAA	CAGATAAGAAGCCACTAGGTCTGAA
Cag7	U	AAGTCAGAAGAAATAACTGACGACTCTAA	TCACGATAAGAACAGCGACTACAA
Cag8	U	AGCAATGAACAGATTATCAACAAGA	GTAGTAGTTGTAGTTTCTAGGCACGG
Cag9	U	TCTCATTGTTCTAATTGGTTGAAA	CTTGTGGCTAATGGTGTGCTAA
Cag10	U	AACGAAGAAGTCTTGATTGATGAAA	CTGTGTATCGATCAATGCCATAA
Cag11	U	CACCTAGCAACTCACAGAGCAA	CCCACCCATACACAATCCTAA
Cag12	A, C	TGTTTTAATCGGTGCGACAA	GAGCGTAATCTTTTTAGAATGGTGAA
Cag12	B	CAAACCTCAAAGACACACCATTCA	TTGTTGTTGGGATTATCATTGTAGTTA
Cag13	U	AATAACATGCGAAAACCTTCTCAA	CTCCATAGTCTCACTCTCAAGCAA
Cag14	U	ACGCATTAGAGATCCGAACAA	CCATTCTTCAACACTTCTGCCATAA
Cag15	U	CAGGGGTGATTTTAGTTTATCCAA	TATGCTGAGAGAAAACGAGTAGCAA
Cag16	U	GAAGAAGTGGCTGCAAAAAGAA	CATAGGCATAAGGGTTAGGAAGAA
Cag17	A, C	TCAAAGACATGACGACGAAGAA	GCTCTTGCCCTATCATTTCGTAA
Cag17	B	TCGCTCTTTATTCTTTGGTTGTTTA	TGCTCTACTCTGCCCTATCA
Cag18	U	CCAACCAACAAGTGTCTCAAAA	TCAATAACGCTAAATCTCCTCTCAA
Cag19	A, C	GACTTTTTGTGGTTTGTCTCTGAA	CGCCAAGCAAGATGTCTGAA
Cag19	B	GAATGGCTTTTTCTTTGGCA	TTTTTGTGGCTTGTCTTTGAA
Cag20	U	GCTGCTAACCAACAATACAACCAA	CTAAGATACCGCTCATCATTTCAA
Cag21	U	GGGGCTTGTTTCTAGAGATCAACTAA	GAAAGGATTGTTTGGACCGTAA
Cag22	A	CTTGCCCATCGTTTATTTCCCTTA	ACCTTACCGCTCTTTATGATTTTTCTA
Cag22	B, C	TTTATGTTTATGCTTACTTTCATGCTAGAA	CGCTCATATCAATCTGAATCCAA
Cag23	U	GCTAGTCATAGAGCAAGAGTTCAAAA	CACAATAACAATCGCTACAATCAAA
Cag24	U	GTATGGGTTAGCAAATGACGATAAA	TTAAGGACTCTATTGACAATCACGAA
Cag25	U	CAAGAATCACTGACAGCTACAAGAA	ATACCGCTGCCACCGCTAA
Cag26	U	ATAAAGCGATCAAAAATCCTACCAA	GGGGGTTGTATGATTTTTCCATAA

^a Strain specificity: A-26695, B-J99, C-J166, U-universal.

Table A2 | Intergenic primer pairs selected for real time RT-PCR.

Primer pair ^a	Strain ^b	Forward primer (5'-3')	Reverse primer (5'-3')
Cag1/2	A	CGTTACCAACGAACCAACTGA	TCATTGAGCCATTTATTTCTCGTAA
Cag2/3	A	GTGTAAATGGACTGCTAGTGTTGGA	AGCGATACAGCGGTTGCTA
Cag2/3	B	ATGTCCCAAATGCAACAGATTAA	CGGCTTCACCTATTTCTTTAGCATAA
Cag2/3	C	TTCACCTCCAATGAGTTTTTTTACA	CGGCTTCACCTATTTCTTTAGCATA
Cag3/4	A	GTGTGGTTATGAGAGCGTTCAA	ATTTTTCTCGCCTGTTGTTCAA
Cag3/4	B	AACTGCTTTCACCACTAAGGGA	GGATTTTTCTCGTGTGTTGTTCA
Cag3/4	C	TGTGTGGTTATGAGAGCGTTCA	GGTGAGATTTTCGTATTGCTTGA
Cag4/5	A	TTCTCAAGTGCGATATAACGAGTAGA	TCTTTAGTGCTGTGGGTTCAA
Cag4/5	B	CTAAGAGCGATGGTTGGCAA	AAAATATGATCTTTGTCTTGCATGAA
Cag4/5	C	GGGGAATGAAACACAACCCTAA	GAAGGCCAAAAGCCTATTCCAA
Cag5/virB1 1	U	ACCCTTTCTTTTCAGCCCATCTATAA	TGTCCATATCAACCACCACAA
virB11/Cag 6	U	AGGATTTAATGCCGCTTCTTTTAA	CAATGAGATGGTCCAAGATATAGGGA
Cag6/7	U	GTTCCATTGCTGTTTCCCTTCA	AATCACCACAAGCCCCAAA
Cag7/8	A, B	GGTGAATCTTGTGGGCTTTTTTA	AGGATTGAGATGGTATAGAGTTAATGAAA
Cag7/8	C	TTCAAGTTTATCGTTTTTCTTATTCA	ATGGTATAGAGTTAATGAAATTGCAGAA
Cag8/9	U	AATAACCAAGACAGAAACAGCCAA	GATGGTAGCAGAAATGGATAGAGAAA
Cag9/10	U	AATAGCTTTCACCAATTAGGAACAA	AACTCTTCTCAAGAAAATCTTATCATCAA
Cag10/11	U	CCCAACCAATTTTCATCAATCA	GTTTGAAGCAATCCGCTACTTACA
Cag11/12	A, B	AAACAAGGCGGTGCAGAA	GGTGTGCTTTTGTGATTTGTCATTTAA
Cag11/12	C	GCTTCATAGGTATGGGCTATTTGA	TGTTTACCACCTTTTTTTCGCATA
Cag12/13	U	AAATCAGAAGTTTGCTCAGTGGTAA	CGCTAATCTAAAACCATTGAACAA
Cag13/14	U	AGCGGTCATAATTCAAAGAGCAA	CAACAAGGCAATAGATTACTAGCT GAA
Cag14/15	U	ACCAATCGCAAACAATCAAA	GATATGGTGGTGGTTTTCCAA
Cag15/16	A, C	AATACCAACAAGCCGCATACAA	CTCCAAACGCAACCAATGA
Cag15/16	B	AATACCAACAAGCCGCATACAA	TGGATCAGATTAGGGATTATTGGAA
Cag16/17	U	CGATCCTATGATGAGCGACAA	GATAGCGTTTAAGCCCCCATAA
Cag17/18	U	GCTCTAAATCTGAACTGCCAATA	GAACAAAGTAAGCGACAATACCTACA
Cag18/19	U	GGCTAGTGGTTGAAAAATCTCATCTA	AAAGAGAAACGACAGCAAGAAACA
Cag19/20	U	ATAACGCCATTAGCCCCTTTTAA	GGGTGCCAACTAAAATAATCGTGAA
Cag20/21	A, B	GATTAGTAAATCCCACAACAATAGGAATAA	TTTTTACCACCGATCTTAGGGTATTAA
Cag20/21	C	GCGGTCATTTGCGGATTAGTA	GGCGATTACGGTCCAAACA
Cag21/22	U	AGCGTTAAACATGCCAATGATAA	GATATGAGCGTTGAAGCTAAAAGAA
Cag22/23	U	TCACCTCCATTTCTTCTTATGAA	GTTTTAATTTGAGGGGCATTCTTA
Cag23/24	U	CGCTTTTGAACCTCTTGCTCTA	TCAGCACGACCAACAACAA
Cag24/25	U	CATACGAACTGAAAACAACGAGACTTA	ATATTGGCGAGAGTGGAGGA

^aOne primer was in the intergenic space and the other was sometimes located in an adjacent gene. ^bStrain specificity: A-26695, B-J99, C-J166, U-universal.

Table A3 | Primer pairs used to construct promoter knockouts.

Primer pair	Forward primer (5'-3')	Reverse primer (5'-3')
Cag1-U	TATGCGGCCGCCGCTTTCACTAACGCTTCCACTA	GCACTGCAGGCGCAAGAATACAGCATTGGGC
Cag1-D	AGTGATCCATGGCTGACACAATCAATACAACG	CTGCTCGAGTTCATCATCCACATTCTTGTGAAG
Cag10-U	ACACTCGAGTCCACTCACATCATAGCCATGCA	ACACTGCAGCCTAGCAACTCACAGAGCAATGA
Cag10-D	ACAGGATCCGCCCTTGATAGATTGGCTAAACTCA	ACAGCGGCCGCCGACAAAAGCAAGCATGGCTGTA
Cag11-U	ACAGCGGCCGCCGACAAAAGCAAGCATGGCTGTA	ACAGGATCCGCCCTTGATAGATTGGCTAAACTCA
Cag11-D	ACACTGCAGCCTAGCAACTCACAGAGCAATGA	ACACTCGAGTCCACTCACATCATAGCCATGCA
Cag16-U	AACCTCGAGGAGTCTTACTTGTGGGACACTC	AACGGATCCATAGGCTGTTCAATACAGCTCTATC
Cag16-D	AACCTGCAGAGCTCATTGGTTGCGTTTGGAG	AACCGGCCGCTTCTCTCAAAGCGTTAGTGGCG
Cag21-U	AACCGGCCGCCAACCTTATCACAGGAGATGAACC	AACGAGCTTAGCATTGAGACTATCTATGAGACC
Cag21-D	AACGGATCCGCTTGGTGTCTTATCATTGGCATG	AACCTCGAGGATGTAATCAAGGTAAGTCAAATGCG
Cag25-U	AGTGCGGCCGCCCTTGCTAAAGCCAAATTCATGCC	AGTCTGCAGCCTTCCAATACAGCTTGATTGTCA
Cag25-D	AGTGATCCCGCACAGAATCACTGACAGCTACAAGA	AGTCTCGAGGAGAATAGTTGTTAGTAAGGATCAC
CAT-1 ^a	AACGGATCCGCGGACAACGAGTAAAAGAG	AACCTGCAGGCGAGGACGCACTACTCTCG
CAT-2 ^b	AACGGATCCGCGGACAACGAGTAAAAGAG	AACGAGCTCGAGGACGCACTACTCTCG
CAT-3 ^c	AAC <u>CTGCAGG</u> CGGACAACGAGTAAAAGAG	AACGGATCCGAGGACGCACTACTCTCG

U, upstream arm; D, downstream arm.

^{a,b,c} Knockouts were constructed with amplification of the CAT gene using primer pairs CAT-1 (cag1, cag10, cag25), CAT-2 (cag21), or CAT-3 (cag11, cag16), which differed only in the 5' restriction sites (underlined).

Table A4 | Primer pairs used to generate amplicons for cag PAI libraries.

Oligo upstream	Sequence	Oligo downstream	Sequence
J166		J166	
MWG268	cgctcaaacctgaaagatcaa	MWG653	tagggcgcaacacactca
MWG607	ctaaagagaccaagaagaggctaaa	MWG18571	cactatggagactgcgaaa
MWG406	ctaagagcagtggttgcaa	MWG463	cctctaagcagctactgaagaa
MWG103	tggacaatcatatcaatcaaatcttta	MWG1198	gtttgagcagtagaagaagc
MWG988	acaagaggagcttttaatacaca	MWG19145	tctcattgtcctaattggttgaaaa
MWG1008	aacgagctccatagaatctttgaaccaatctag aacga	MWG028	tgcggtttgttttgattagaa
MWG410	aatgtaagtagcggattgctcaa	MWG19155	acgcattagagatccgaacaa
MWG416	agcgttcataattcaagagcaaa	MWG657	tgctctactctgcctatca
MWG063	cagcttcaattttgatcccaatc	MWG475	gactttttgtttgtctctgaa
MWG476	cgccaagcaagatgtctgaa	MWG436	gttttaattgagggcattccta
MWG18880	ctagcatgaagtaagcataaacataaactaa	MWG17626	gctaataatccacttaaatccaaa
MWG734	aacctcgagaccttgagatacaagcttttctgtg	MWG807	gacagatttcaagacagcttca
MWG17643	aacctcgagccttactttatggtgagccataac	MWG9165	ttagaataatcaacaacatcacgcat
G27		G27	
A-cag	gcgagcggcagatgtgatctggc	B-cag	ggaacgccaccgttggtataaagac
C-cag	gattggatcgtaatgcttcaaatcc	MWG9165	ttagaataatcaacaacatcacgcat
D-cag	cctgtatctgtccctagctc	MWG807	gacagatttcaagacagcttca
F-cag	ggactccattgttctaatgg	E-cag	gatgatggggtgatccttactaacaac
Q-cag	gacccgttagggaattataatc	R-cag	cctatcaataacaacataagcgag
O-cag	cagagcagtcataattcaagagc	P-cag	tgctctgctctgcctatca
M-cag	gtaaggtagcggattgcttcaacaag	N-cag	cagcattagagatccgaacaagc
L-cag	gctttaagactcttttagcttc	I-cag	gatgggaaattgagcatgactg
J-cag	gaagctaagagagagctttaaagc	K-cag	gtaaagaaccgagtttgtaaac
G-cag	ccttagcaccattcctgccataacc	H-cag	gaaggaagctcaatgagattgtc

Several additional oligonucleotide pairs were required to amplify the entire *H. pylori* G27 strain cag PAI region, due to poor or no amplicon presence in some PCR reactions.

Table A5 | Putative promoter sequences that direct expression of the reporter *tnpR*.

Promoter	Length (nt)	Sequence
I	244	GATCAAAAAAATCAAACAAAAATAACGATTGAGTGGCGTTAATGCGCTAGAATAGTGCTAAAAATAAGAATAAAGGAATCAAAA GTATGAAAACGAATTTTTATAAAATTAATACTATTGCTTGGTGTCTTATCATTGGCATGTTTAACGCTCCGCTTAACGCTGACC AAAACACTGATATAAAGATATTAGTCTGAAGATATGGCACTAAATAGCGTGGGGCTTGTTTCTAGAGATC
II	38	GATCGTTTGACAATTTTAAATCTCCTGTGTATCGATC
III	91	GATCTGTTGCTTTATTGTCAAAAAGCCATTGAAATTCACCATTGGTTGATTGCAAAAA GCGCTAATCGCGCGACAAGCCCAT TAGGATC
IV	104	GATCCCAATGCCACACGCTTGATAAGGGAGCGTCAATTGATGAGAACAAGCTTTTT GAACAACAAAACGCGCGTATTTCAA CTACGCCAACGATGTGATC
V	71	GATCAGCTTGGGTTTGTCTGCTGTTTTAGGTTTCAACCTGAGACGATTAATAA TACATCAAAGATC
VI	314	GATCCTAAATATCCCAATGCAATGGATTGATGAAAAGAAAAAGAATTTCAAAAACAATGAGTTTTTACAGCTGCATTACTT ACCTTAAATGCAATGGAATTTGTCTCTATATCAATTCTGAAAAAAAAGGAACTAATGTTTAGAAAACCTAGCAACCGCTGTATC GCTCATAGGCTTACTAACCTCTAACACTCTTTATGCTAAAGAAATAAGTGAAGCCGATAAGTGCATTAAGGCCACTAAAGAACT AAAGAGACCAAGAAAGAAGTAAACGACTCAAAAAAGAAGCTAAACAGCGCCAACAGATC
VII	234	GATCAAGAGATTATCAAAGGAAGCAAAAAAATACATTATTAGTGGCATTGTAGTCGCTGCTCTTATCGTGATTATTTATTTCT AGAAGCATTTTTACTACTTTTATACCTTTGGAAGATAAAAGCTCTCGTTTTAGCAAAGACAGGAATCTTTATGTCAATGATGAAATC CAAATAAGGCAAGAGTATAACCGATTGCTGAAAGAACGGAATGAAAAAGGCAATATGATC
VIII	635	GATCCATGATGCTCTGTTGTATCGTTCATGAAATTCCTTTCAAGAATTAATTTAGAAATTTGTTTTGATATTATACCATTCTCTCTCT GAGTTGTGATTGTCTTATCTTTGAATTAGGCGCTTCAAAATTTTACTACTGATTACGACTGCTTACTTATTGCTCTTACTTTTTG AGTTGCATCGTGTTCATCTTGTCTTGTGTTGAAGCAATCCGCTACCTTACATTTATTATAAGGAATCTTTGTTCAACGCCTTATCC AAAAAGTTTTTTATTAAGTTTTTTCAAATATATTTTTACAGAAATTTTGTATACTATACTGAAATGTTTTAAGGAGTTTTTG ATGAAAAAATTTCTCAAGTATTCTCAAGTTTTGCAACCAACAAACAATAACTAAACAATAAGCTGTCGCATGTTAGGGAAAA AAAACGAGGAAGTCTTGATTGATGAAAAATTTGGTTGGGGGTGTGATAGCCCTTGATAGATTGGCTAAACTCAATAAGGCCAATA GGACTTTCAAAAGGGCTTTTTATCTCTCTATGGCGCTCAATGTGCGCGCTGTAACGAGTATTGTGATGATGATGCCTTTGAAGAA AACGGATATTTTTGTTTATGGCATTGATC
IX	132	GATCATAGTGCCGTTTCATGTTCCATACATCTTTGGCTACAACCCCACTACTATACCT GTGAGAGTGGCATCTACTTTAGAAGTCA- GAGTGATTTCATAGGGGTGTATTGCGCT AAAACAATGTGGGATC
X	269	GATCTATGTTAAAGGCTAGCCGCTTTATTCTGTGTTACAATTACAAATATTTTTTAAGAGGAATTTGTTGATGGGGCGGGCATTG TTTTAAAAAATTTGTTGGCTGTTCTGTCTTGGTTATTATTTTTATCTAGCGTAATAGAAGCAGCACCTGACATTAATAATTTAATC GTGGTAGGGTGAAGTGGTGAATAAGAAGATTGCTTATTTGGGAGATGAAAAACCTATTACGATTTGGACTTCATTAGACAATGTT ACTGTGATC
XI	419	GATCTTTGATGATTTTTTAATCGTCTCAGGTTGAAACCTAAAAACAAGCAGAAACAACCCCAAGCTGATCAGAGTGAGAATAAA GCTCCATTTAAGCAACTCCATAGACCACTAAAGAACTTTTTTTGAAGCTGTCTTTGAAAATCTGTCTATTGATTTGTTTCCATG TGAATCACAACGCTTAATTGCAAAATATACTTTATGGTAAGCATGACACACAACCAACCATTTTAGAACGCTTCATGCACTC ACCTTGATTCCAACATATTTAAGCATTGCATTTGATTTATTCTTGAAGGTTCAATTTCTATTCTTTCTTTGTTAAATTCGTTTCATT TTAGCAAATTTTTGTTAATTGTGGTAAAAATGTGAATCGTTCCTAGCCTTTAGACGCCTGCAACGATC
XII	54	GATCCCTAGAACAAGTAAGCGCAATACCTACAAGAAAGGCAATCAAGTAAGATC
XIII	68	GATCCAATCATTGAAAAATCTTTGATGAAAAGGGTAAAGAAATGGGATTGAATGTAG AATTACGATC
XIV	70	GATCCTACTGGTGGGGATTGGTTGGATTTTTCTCTCATTATATTTGACAAAAACA ATCTCCGATC

Table A6 | Key to cag PAI gene names.

26695 ORF	Gene number	Gene name	T4SS homolog
HP0520	<i>cag1</i>	<i>cagC</i>	
HP0521	<i>cag2</i> /hypothetical	Hypothetical	
HP0522	<i>cag3</i>	<i>cagA</i>	
HP0523	<i>cag4</i>	<i>cagy</i>	VirB1
HP0524	<i>cag5</i>	<i>cagβ</i>	VirD4
HP0525		<i>cagα</i>	VirB11
HP0526	<i>cag6</i>	<i>cagZ</i>	
HP0527	<i>cag7</i>	<i>cagY</i>	VirB10
HP0528	<i>cag8</i>	<i>cagX</i>	VirB9
HP0529	<i>cag9</i>	<i>cagV</i>	VirB6
HP0530	<i>cag10</i>	<i>cagW</i>	VirB8
HP0531	<i>cag11</i>	<i>cagU</i>	
HP0532	<i>cag12</i>	<i>cagT</i>	VirB7
HP0533	Hypothetical	Hypothetical	Hypothetical
HP0534	<i>cag13</i>	<i>cagS</i>	
HP0535	<i>cag14</i>	<i>cagQ</i>	
HP0536	<i>cag15</i>	<i>cagP</i>	
HP0537	<i>cag16</i>	<i>cagM</i>	
HP0538	<i>cag17</i>	<i>cagN</i>	
HP0539	<i>cag18</i>	<i>cagL</i>	VirB5
HP0540	<i>cag19</i>	<i>cagI</i>	
HP0541	<i>cag20</i>	<i>cagH</i>	
HP0542	<i>cag21</i>	<i>cagG</i>	
HP0543	<i>cag22</i>	<i>cagF</i>	
HP0544	<i>cag23</i>	<i>cagE</i>	VirB3/VirB4
HP0545	<i>cag24</i>	<i>cagD</i>	
HP0546	<i>cag25</i>	<i>cagC</i>	VirB2
HP0547	<i>cag26</i>	<i>cagA</i>	