

UC San Diego

UC San Diego Previously Published Works

Title

Chrononutrition and Cardiometabolic Health: An Overview of Epidemiological Evidence and Key Future Research Directions.

Permalink

<https://escholarship.org/uc/item/8hp0x01f>

Journal

Nutrients, 16(14)

Authors

Raji, Oluwatimilehin

Kyeremah, Esther

Sears, Dorothy

et al.

Publication Date

2024-07-19

DOI

10.3390/nu16142332


Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Review

Chrononutrition and Cardiometabolic Health: An Overview of Epidemiological Evidence and Key Future Research Directions

Oluwatimilehin E. Raji¹, Esther B. Kyeremah¹, Dorothy D. Sears^{2,3,4,5} , Marie-Pierre St-Onge^{6,7} 
and Nour Makarem^{1,*} 

¹ Department of Epidemiology, Mailman School of Public Health, Columbia University Irving Medical Center, New York, NY 10032, USA; oer2105@cumc.columbia.edu (O.E.R.); ebk2150@cumc.columbia.edu (E.B.K.)

² College of Health Solutions, Arizona State University, Tempe, AZ 85287, USA; dorothy.sears@asu.edu

³ Department of Medicine, University of California San Diego School of Medicine, La Jolla, CA 92093, USA

⁴ Department of Family Medicine, University of California San Diego School of Medicine, La Jolla, CA 92093, USA

⁵ Center for Circadian Biology, University of California San Diego, La Jolla, CA 92093, USA

⁶ Department of Medicine, Columbia University Irving Medical Center, New York, NY 10032, USA; ms2554@cumc.columbia.edu

⁷ Sleep Center of Excellence, Columbia University Irving Medical Center, New York, NY 10032, USA

* Correspondence: nm2968@cumc.columbia.edu

Abstract: Chrononutrition is a rapidly evolving field of nutritional epidemiology that addresses the complex relationship between temporal eating patterns, circadian rhythms, and metabolic health, but most prior research has focused on the cardiometabolic consequences of time-restricted feeding and intermittent fasting. The purpose of this topical review is to summarize epidemiological evidence from observational and intervention studies regarding the role of chrononutrition metrics related to eating timing and regularity in cardiometabolic health preservation and cardiovascular disease prevention. Observational studies are limited due to the lack of time-stamped diet data in most population-based studies. Findings from cohort studies generally indicate that breakfast skipping or the later timing of the first eating occasion, a later lunch and dinner, and a greater proportion of caloric intake consumed in the evening are associated with adverse cardiometabolic outcomes, including higher risk for coronary heart disease, hypertension, type 2 diabetes, obesity, dyslipidemia, and systemic inflammation. Randomized controlled trials are also limited, as most in the field of chrononutrition focus on the cardiometabolic consequences of time-restricted feeding. Overall, interventions that shift eating timing patterns to earlier in the day and that restrict evening caloric intake tend to have protective effects on cardiometabolic health, but small sample sizes and short follow-up are notable limitations. Innovation in dietary assessment approaches, to develop low-cost validated tools with acceptable participant burden that reliably capture chrononutrition metrics, is needed for advancing observational evidence. Culturally responsive pragmatic intervention studies with sufficiently large and representative samples are needed to understand the impact of fixed and earlier eating timing schedules on cardiometabolic health. Additional research is warranted to understand the modifiable determinants of temporal eating patterns, to investigate the role of chrononutrition in the context of other dimensions of diet (quantity, quality, and food and nutrition security) in achieving cardiometabolic health equity, and to elucidate underlying physiological mechanisms.

Keywords: chrononutrition; temporal eating patterns; circadian rhythms; eating timing; eating regularity; cardiometabolic health; cardiovascular health; cardiovascular disease prevention



Citation: Raji, O.E.; Kyeremah, E.B.; Sears, D.D.; St-Onge, M.-P.; Makarem, N. Chrononutrition and Cardiometabolic Health: An Overview of Epidemiological Evidence and Key Future Research Directions. *Nutrients* **2024**, *16*, 2332. <https://doi.org/10.3390/nu16142332>

Academic Editor: Javier Gómez-Ambrosi

Received: 6 June 2024

Revised: 10 July 2024

Accepted: 11 July 2024

Published: 19 July 2024



Copyright: © 2024 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Chrononutrition is a rapidly evolving field of nutritional epidemiology that addresses the complex relationship between temporal eating patterns, circadian rhythms,

and metabolic health to elucidate how eating timing metrics influence health preservation and chronic disease risk [1–3]. The timing of food intake relative to circadian rhythms, representing the innate 24 h cycles in human behavior, physiology, and metabolism, may be particularly important for preserving cardiometabolic health (CMH) and preventing cardiovascular disease (CVD). While light exposure is the predominant zeitgeber, food intake also entrains the circadian system, and research shows that metabolic health is influenced not just by the quantity and quality of dietary intake, but also by the timing, regularity, and energy distribution of eating occasions in the 24 h day and across days [3,4]. Indeed, aspects of chrononutrition, stemming from environmental, cultural, and physiological factors as well as personal preference and lifestyle choices that alter eating behaviors, have been linked to the etiology of CVD, type 2 diabetes, metabolic syndrome, hypertension, and obesity [5].

Enhancing cardiometabolic disease prevention approaches with strategies grounded in chrononutrition could play a pivotal role in reducing the public health burden of these conditions [1]. Through targeted health education and promotion measures, individuals can be informed about the implications of eating timing on health and guided towards establishing and preserving consistent eating routines synchronous with their circadian rhythms. Such measures could potentially delay the emergence of metabolic risk factors linked to cardiometabolic and cardiovascular diseases or aid in their management. For example, “Life’s Essential 8”, a cardiovascular health construct developed by the American Heart Association (AHA) to guide primordial prevention efforts, includes diet as a metric [6]. However, this metric mainly addresses dietary quality and excludes dietary dimensions related to the timing of meals and other eating occasions. Screening for and addressing eating timing and regularity could potentially provide a more comprehensive assessment of nutritional health and unlock the true potential of dietary approaches in primordial prevention efforts aimed at CMH preservation.

Despite the immense promise of this field for addressing the chronic disease burden and persistent CMH inequities, many knowledge gaps remain. This topical review summarizes the epidemiologic evidence, encompassing both observational and intervention studies, on temporal eating patterns in relation to CMH. Given that multiple prior reviews have focused on time-restricted feeding and nightly fasting duration in relation to CMH [3,7,8], we focused this review on associations of eating timing, meal skipping, and eating timing regularity with cardiometabolic outcomes and CVD. We also highlight knowledge gaps and important research directions necessary to advance the field and translate chrononutrition research into public health policy and action.

2. Eating Timing and Regularity Patterns in the US and Other Populations

Despite the widely held view that breakfast holds paramount importance as the day’s initial meal, research indicates that 20% to 30% of US adults regularly omit this eating occasion [9]. Further, there has been a decline in the reported consumption of breakfast and lunch over the past 40 years [10]. Similar trends are observed in Asian populations; for instance, a multicenter Japanese study showed that ~23% of adults skipped breakfast [11], and ~45% of Korean adults do not report caloric intake during the morning hours [12]; this proportion is even higher among Korean office workers (~70%) who report skipping breakfast due to time constraints [13]. In the Mediterranean population, breakfast, lunch, and dinner are typically consumed later than in the US and other populations [14–16]. For example, lunch is consumed, on average, between 2:00 and 3:00 pm, compared to 1:00 and 2:00 pm in the US population [14,15]. Notably, within the framework of the Mediterranean diet, lunch holds particular importance, accounting for approximately 40% of the daily caloric consumption compared to about 24% in the US [17,18].

In terms of the distribution of calories across the 24 h day, data from a smartphone application show that US adults eat around the clock and that calorie consumption extends beyond typical meal hours [19]. There is also a tendency for later eating timing, as >75% of caloric intake occurs after noon with ~38% of calories consumed after 6:00 pm. A tendency

for substantial nighttime eating was also observed in the National Health and Nutrition Examination (NHANES) survey, whereby over a third of individuals reported consuming $\geq 25\%$ of their daily caloric intake after 7:00 pm and $\sim 10\%$ reported consuming $\geq 50\%$ of caloric intake after 7:00 pm or any calorie intake after 11:00 pm [20].

In addition to eating timing within the 24 h day, eating timing across days has emerged as a novel potential contributor to CMH [19–22]. The term ‘eating jet lag’ describes the difference in mealtimes between weekdays and weekends [21,22]. In a 2021 study examining racially and ethnically diverse US women, the timing of the first eating occasion was ~ 1.5 h later on weekends than on weekdays [22]. Similarly, breakfast timing was ~ 2 h later on weekends relative to weekdays in a study of young Spanish and Mexican adults [21]. In NHANES and the Continuing Survey of Food Intakes by Individuals, night eating (i.e., $\geq 25\%$ kcal after 7:00 pm, $\geq 50\%$ kcal after 7:00 pm, and any eating after 11:00 pm) was more common during the weekend than weekday [20]. Consistent with these findings, data from a smartphone app revealed that US adults have erratic eating timing patterns with substantial day-to-day variation, particularly for timing of the first eating occasion with later eating start times being observed on weekends compared to weekdays (10:26 vs. 09:21 am) [19].

Sociocultural, economic, and environmental factors can substantially influence when someone eats and could help to explain differences in temporal eating patterns observed across populations [23]. In Mediterranean cultures, lunch accounts for a substantial proportion of daily caloric intake, despite the later overall timing of lunches and dinners; whereas in the USA and in central and northern European countries, dinner and after dinner snacks tend to account for the largest percentage of daily caloric intake [10,16,23]. Further, the emphasis on breakfast consumption could vary across cultures with breakfast skipping predisposing to later eating timing and higher caloric intake in the evening and generally being linked to having a lower socioeconomic status [24,25]. Relatedly, being an essential worker and/or a shift worker is also associated with later and irregular eating timing patterns [25,26]. This can help to explain the higher prevalence of adverse cardiometabolic outcomes among essential workers and shift workers and among minoritized populations that tend to be overrepresented in these occupations.

Individual behavioral factors can also influence the timing of food intake, the most notable of which is sleep. Having a shorter sleep duration and going to bed later provides more opportunities for later eating timing; in addition, irregular sleep schedules are likely correlated with irregular eating schedules [23]. This is exacerbated by light exposure at night, which can delay sleep and extend eating periods later into the night. Other dietary dimensions can also influence meal timing; for instance, food insecurity can lead to less structured mealtimes and irregular eating timing patterns [27]. In addition, there is likely a bidirectional association between chrononutrition metrics and diet quality, with later eating times being linked to unhealthy food choices and exacerbating the adverse health effects of poor diet quality [28,29]. However, there is a dearth of evidence on the behavioral underpinnings of temporal eating patterns, particularly the influence of sleep health, physical activity, and other diet dimensions on when someone eats. Other individual-level characteristics such as age, sex, chronotype, genetic predisposition and health status could also shape eating timing patterns, with later and less structured eating times observed among those with later chronotypes (representing the behavioral manifestation of endogenous circadian rhythms), in younger age groups, and among individuals with obesity [23,30]. The multi-level predictors of chrononutrition metrics influencing cardiometabolic outcomes and the attainment of health equity and warranting further investigation are summarized in Figure 1.




Figure 1. Determinants of eating timing and regularity.

3. Overview of Mechanisms Linking Temporal Eating Patterns to Cardiometabolic Health

The influence of eating timing on CMH operates through various mechanisms related to circadian rhythm disruption, metabolic and inflammatory processes, hormonal imbalances, and other behavioral factors (Figure 2) [3,31,32]. Although the rhythm of the central circadian clock (located in the suprachiasmatic nucleus of the hypothalamus) aligns predominantly with photic cues, the rhythms of peripheral tissues are primarily governed by external cues such as food intake, a powerful zeitgeber [33,34]. Adverse temporal eating patterns, characterized by delayed and irregular eating timing, can disrupt the synchrony of these peripheral clocks with the central clock, resulting in circadian misalignment [34]. Circadian misalignment disrupts metabolic regulation or homeostasis, resulting in impaired glucose control, increased insulin levels, insulin resistance, and a glucose response that mimics a prediabetic state [35]. Notably, late eating has been shown to decrease glucose tolerance and impair pancreatic beta cell function in individuals carrying the MTNR1B type 2 diabetes risk variant [36]. Eating timing also entrains blood pressure (BP) levels and circadian patterns [37]. Short-term circadian misalignment has been shown to elevate BP by modulating the autonomic nervous system, reducing sympathetic activity while increasing parasympathetic activity. Circadian misalignment also diminishes the secretion of melatonin, which is known to lower BP [38]. Additionally, later and irregular eating timing patterns are associated with higher C-reactive protein (CRP), a marker of systemic inflammation linked to high BP and other cardiovascular complications [32,39].

Later eating timing and circadian misalignment from mismatched eating patterns and endogenous rhythms can also alter appetite-regulating hormones, increasing the risk of obesity and its metabolic sequelae [35]. For example, inverting behavioral cycles by ~12 h can suppress leptin and lead to increased appetite and decreased energy expenditure. Overall, later eating timing alters appetite-regulating hormones by reducing 24 h serum leptin and increasing the 24 h ghrelin–leptin ratio, promoting increased hunger [40]. It is also associated with decreased energy expenditure during the day and changes in gene expression that favor increased lipid storage and promote obesity [40]. Lastly, the timing of high-energy meals can impact fat accumulation and metabolism through the adipose tissue clock, potentially contributing to dyslipidemia and obesity [41].




Figure 2. Mechanisms linking temporal eating patterns to cardiometabolic disease. Created with BioRender.com (accessed on 19 December 2023).

Other potential mechanisms requiring further investigation include the impact of eating timing on the gut microbiome and gene expression. Alterations in the composition, function, and rhythms of the human gastrointestinal microbiota, as well as changes in clock and metabolic gene expression, have been reported in response to eating timing cues in emerging studies [42,43]. For example, later eating timing has been shown to induce changes in the microbiota similar to those observed in women with obesity or in older age [44]. In terms of the interplay of eating timing with genetic factors, the *PERIOD2* (*PER2*) genes have been associated with Night Eating Syndrome in prior research [45]. In addition, the minor allele (G) of the *CLOCK* rs4580704 SNP, associated with obesity, has been shown to be more prevalent among late eaters (after 3:00 pm) compared to early eaters (before 3:00 pm) and has been linked to consumption of a later lunch [17]. Indeed, it appears that the heritability of food timing may vary by meal; a Spanish study of twins showed that it ranged from 56% for breakfast, 38% for lunch, to being undetectable for dinner [46]. Another study of twin pairs showed ~24% heritability for the timing of breakfast, and lower heritability estimates for lunch and dinner timing (18–22%) [47]. There is also evidence of a potential shared genetic architecture between food timing and chronotype that warrants further investigation [46,47].

There is likely a bidirectional association between eating timing and mental health, which can influence eating habits and the quality of food intake [48–50]. Shift workers, who often experience misalignment between their internal circadian rhythms and daily eating schedules, are at increased risk of depression and anxiety [49,51]. Consuming meals during daytime hours versus during both the daytime and nighttime has been shown to reduce levels of depression-like and anxiety-like moods [52]. Furthermore, the timing and regularity of eating patterns may influence other health behaviors that impact CMH [53–56]. For example, both skipping breakfast and irregular eating patterns have been linked to lower sleep quality and reduced physical activity [55,57]. Lastly, eating timing may influence diet quality; individuals who consume food during the nighttime tend to have a higher energy intake from snacks, a greater percentage of their daily energy intake coming from fats, and lower dietary diversity scores [58].

4. Summary of Observational Evidence

4.1. Breakfast Skipping and Timing of First Eating Occasion

In the prospective Health Professionals Follow-up Study (N = 26,902), men who omitted breakfast exhibited a 27% elevated coronary heart disease (CHD) risk, and the association was mediated by BMI, hypertension, diabetes, and hypercholesterolemia [59]. Indeed, breakfast skipping has been linked to greater type 2 diabetes risk, higher total and LDL cholesterol levels, body weight, fat mass, and abdominal adiposity and lower HDL cholesterol, though most studies focus on obesity risk [60–62]. For instance, a landmark study of >50,000 Seventh-day Adventists in the US and Canada revealed that individuals who consumed their largest meal at breakfast vs. dinner had a significant decrease in their body mass index (BMI) over a 7 year follow-up period [63].

However, there are some contradictory findings to the aforementioned studies. Among Japanese adults without obesity, breakfast skipping was associated with 28% and 57% higher odds of developing metabolic syndrome and obesity, respectively, but only when in combination with eating late-night dinners (within 2 h of bedtime) [11]. A cross-sectional analysis of healthy Korean adults revealed that those who infrequently ate breakfast had lower odds of having elevated serum triglycerides compared to those who regularly or frequently consumed breakfast [64]. This could be explained by the larger proportion of energy intake from fats and smaller proportion from carbohydrates observed among the infrequent breakfast eaters.

In addition to breakfast consumption, the timing of the first eating occasion has also been linked to CMH [65,66]. Morning eating, regardless of whether or not the meal is identified as breakfast and independent of fasting duration and number of eating episodes, has been linked to 27% and 31% lower odds of metabolic syndrome in Asian men and women, respectively [12]. Among US women encompassing different life stages, later timing of the first eating occasion was associated with poorer cardiovascular health, defined using the AHA's Life's Simple 7 score, as well as with higher waist circumference, diastolic BP, and fasting glucose levels [65]. Conversely, a prospective study conducted with older adults revealed that a timing of first eating occasion after 09:00 am might be linked to a reduced long-term risk of developing type 2 diabetes, particularly among older individuals with impaired fasting glucose, suggesting that associations may vary by life stage and chronic disease status [66].

Only one prospective cohort investigated eating timing patterns in relation to incident CVD [67]. In a 2023 analysis that encompassed >100,000 French adults who participated in the NutriNet-Santé cohort, each 1 h delay in timing of the first meal was associated with 6% higher risk of overall CVD [67]. Notably, every 1 h increase in nightly fasting duration in that study predicted a 7% lower risk of cerebrovascular diseases, suggesting that there may be an interplay between eating timing and the span of the daily eating period. In addition, associations were stronger among women, particularly for the timing of the first meal whereby an eating start time later than 9 am was associated with up to a 35% higher risk of CVD and cerebrovascular diseases. The authors attribute these sex differences to the robust associations between temporal eating patterns and cardiometabolic health among women as reported in the aforementioned studies, and also to potential sexual dimorphisms in the anatomy and physiology of the circadian system [67–70].

4.2. Early Lunch vs. Late Lunch

In a post hoc analysis of 420 Mediterranean adults with overweight and obesity who were followed-up for 20 weeks while undergoing a weight loss intervention, consuming lunch earlier in the day (before vs. after 3:00 pm) was associated with pronounced weight loss and lower HOMA-IR, even when caloric consumption and energy expenditure were held constant across both groups [17]. Weight outcomes were not significantly affected by breakfast or dinner times, highlighting the potential health benefits of the Mediterranean diet's emphasis on lunch as a principal daily meal [17].

4.3. Nighttime Eating and Evening Caloric Intake

In a 16-year prospective study of 26,902 middle-aged male health professionals, a 55% higher coronary heart disease (CHD) risk was observed among those who ate late at night, but this was defined as eating after going to bed and does not capture the association of late dinner or snacking with cardiovascular risk [59]. The association was mediated by cardiometabolic risk factors, and several studies have since linked evening caloric intake or late-night eating with cardiometabolic outcomes [59].

In the Korean NHANES dataset, night eating, defined as eating after 9:00 pm, was associated with 48% higher odds of metabolic syndrome in men only, suggesting that there may be sex differences in these associations [12]. However, higher nighttime eating levels have been linked to CMH in women as well [39,65]. For instance, each 10% increase in calorie intake between 5:00 pm and midnight was associated with 3% higher CRP levels among 2650 women from NHANES [39]. Consistent with these findings, a greater proportion of daily calories consumed at the self-identified largest evening meal (“dinner” and/or “supper” in an Automated Self-Administered 24 h (ASA24) dietary record) was associated with higher BP in a community-based cohort of 116 New York women [65]. In contrast, an analysis using the NHANES-III dataset (N = 18,407) found an inverse relationship between nighttime eating, defined as consuming $\geq 25\%$ of daily calories after 7:00 pm, and BMI, even after adjusting for total caloric intake, with nighttime eaters having a predicted BMI 0.44 units lower than those who were not significant nighttime eaters [20]. However, because this analysis relied on a 24 h recall, it may not capture associations of habitual nighttime eating with BMI.

In ~900 middle-aged to older adults, a higher % of daily energy consumed at night (within 2 h before bed time) was associated with 82% higher odds of having overweight and obesity, and ~5 fold higher odds were observed in those with later chronotypes [71]. In contrast, a higher % of daily energy intake consumed during the morning window (within 2 h after wake time) was associated with 47% lower odds of having overweight or obesity, particularly among those with an earlier chronotype [71]. These data suggest that lower energy intake after waking up and higher intakes before bedtime are associated with higher BMI and highlights that associations between eating timing and adiposity likely vary by chronotype.

Finally, in the NutriNet-Santé study, later timing of the last meal was also associated with an elevated CVD risk, particularly among women [67]. In that study, every 1 h increment in timing of the last meal was associated with an 8% higher CVD risk. A last meal timing between 8:00 p.m. and 9:00 p.m. and after 9:00 p.m. was associated with a 19% and 28% higher CVD risk, respectively. Similar to the results for the timing of the first meal in that study, the associations of last meal timing with incident CVD and cerebrovascular disease were stronger among women. Further, the time interval between the last meal and bedtime, as a proxy for the circadian timing of food intake, was inversely associated with the risk of overall CVD [67].

4.4. Eating Timing Variability and Eating Jetlag

An aspect of chrononutrition that is in the nascent stages of characterization is the role of variability in daily eating timing patterns in CMH. In a longitudinal study of racially and ethnically diverse US women (N = 115) with 7-day food records collected using the NIH’s ASA24, significant associations were demonstrated between day-to-day variability of multiple eating timing metrics, captured from the standard deviation of these variables, and eating jetlag, captured from weekday–weekend differences in these metrics, with CMH indicators [22]. Specifically, greater day-to-day variability and eating jetlag in eating start time, span of the daily eating period, and evening caloric intake (% calories consumed after 5:00 p.m. and 8:00 p.m.) were associated with clinically meaningful increases in BMI, waist circumference, HbA1c, and BP [22]. In a subsequent analysis, every 30 min difference in weekday–weekend eating end time was related to a 13% higher CRP level, and every 1 h weekday–weekend difference in eating duration was associated with a 45% elevation

in CRP [32]. Consistent with these findings, a positive association between eating jetlag and BMI, independent of chronotype and social jet lag, was observed in a Spanish study of >1000 students; there was a threshold effect, whereby >3.5 h of eating jetlag was associated with a 1.34 kg/m² higher BMI [21]. Notably, eating jetlag in this study was captured by the weekday–weekend difference in eating midpoint and was primarily driven by the later timing of breakfast on weekends compared to weekdays.

5. Summary of Intervention Studies

5.1. Meal Skipping

Evidence from randomized controlled trials (RCT) addressing the impact of breakfast skipping or not eating before noon on body weight and adiposity metrics is mixed, but it is suggestive of a possible adverse effect on glycemic regulation and blood lipids [55,72–78]. In a multisite, 16-week, 3-parallel-arm RCT of otherwise healthy adults with overweight and obesity (N = 309), breakfast skipping had no impact on weight loss [72]. Despite this, compliance to the intervention was >90%, suggesting that recommendations to adjust eating timing may be effective at changing eating habits in free-living settings. Comparable outcomes were observed in smaller trials [55,73]. In lean adults (N = 33), no impact was observed on weight loss, but those who ate breakfast had a 10% increase in adipose tissue insulin sensitivity [73]. Similarly, in adults with obesity (N = 23), there was no significant difference in body weight changes between a breakfast consuming group (≥ 700 kcal before 11:00 a.m. daily, with ~50% consumed within 2 h of waking up) and a breakfast skipping group (abstained from any caloric intake until noon), but the regular omission of breakfast was shown to reduce insulin sensitivity [55]. In a crossover trial (N = 17) that utilized three isocaloric 24 h interventions (comprising a breakfast-skipping day, a dinner-skipping day, and a conventional three-meal-structure day), HOMA-IR and glucose concentrations after lunch were significantly higher after breakfast skipping than dinner skipping [74]. Skipping breakfast also resulted in increased fat oxidation and an intensified inflammatory response in peripheral blood cells after lunch, suggesting that foregoing breakfast can lead to chronic low-grade inflammation and glycemic dysregulation [74].

Despite the null findings for weight loss in the aforementioned studies, in a crossover trial of young adult Japanese males (N = 10), skipping breakfast actually led to a modest yet significant weight gain, and beginning on the sixth day of breakfast omission, an elevated mean 24 h blood glucose was observed [75]. Similarly, a parallel arm RCT (N = 36) found that the breakfast skipping group (control) exhibited a more substantial decrease in weight loss compared to the two breakfast intervention groups but also showed a significant elevation in total cholesterol concentrations [76]. This is consistent with a prior study showing that breakfast omission is linked to increased fasting total and LDL cholesterol levels in healthy lean women [77]. In contrast, a crossover study of 22 patients with diabetes, which compared a meal schedule comprising breakfast, lunch, and dinner to one that skipped breakfast, revealed that individuals who skipped breakfast had significant weight loss but experienced a greater glycemic response after lunch and dinner and had impaired insulin responses to meals [78]. Taken together, these data suggest that the impact of breakfast skipping on weight may vary by chronic disease status, but that breakfast skipping is consistently related to poorer glycemic control.

5.2. Eating Timing

In a meta-analysis of twelve RCTs on time-restricted eating, encompassing 730 adults with obesity or overweight, although both early and late time-restricted eating resulted in moderate reductions in body weight and insulin resistance, early time-restricted eating was more effective in improving insulin resistance [8]. However, there were no significant differences in the effects of early vs. later time-restricted eating on fasting blood glucose, BP, and lipid profiles. On the other hand, a systematic review and meta-analysis of nine RCTs that examined the impact of eating timing, in the context of an energy reduced diet, showed that an energy intake distribution that favors the early vs. late timing of food

intake is associated with greater weight loss (~1.23 kg) and with lower fasting glucose, insulin resistance, and LDL cholesterol [79].

Earlier timing of the largest meal may be particularly relevant for improving CMH; an RCT of 93 women with overweight and obesity that investigated the effect of a 12-week calorie restricted diet (~1400 kcal/day) with high caloric intake (~50% of total kcal) consumed at breakfast versus dinner showed that the breakfast group had greater weight loss outcomes, including reduced waist measurements, more favorable glucose and lipid profiles, and experienced longer-lasting satiety [80]. In a similar RCT of 60 women with polycystic ovary syndrome, the breakfast group exhibited a 7% and 54% reduction in glucose and insulin levels, respectively, while no association was observed in the dinner group [81]. A randomized crossover trial of 32 women comparing late lunch (at 4:30 pm) to early lunch (at 1:00 pm) reported a 46% increase in glucose levels above the baseline suggesting that a later lunch may reduce glucose tolerance [82].

Lastly, an experimental study of female students (N = 12), found that consuming supper at a regular time (6:00 p.m.) versus later (11:00 p.m.) was associated with less efficient digestion and the absorption of carbohydrates at breakfast [83]. This increased efficiency in the late supper group could be attributed to the extended orocecal transit time of the chyme following a late supper, and resulted in elevated blood glucose levels following breakfast the subsequent day [83].

6. Research Gaps, Challenges, and Limitations

A major obstacle to accurately interpreting the evidence on eating timing and regularity in relation to CMH is the heterogeneity in how eating timing, regularity, and the extent of evening eating are defined and measured, making it challenging to compare findings across both observational and intervention studies. For instance, some studies evaluate eating timing (e.g., timing of last eating occasion) while others evaluate meal timing (e.g., timing of self-defined supper or dinner). Similarly, studies evaluating the distribution of caloric intake across the 24 h day may examine the distribution of calories across time bins or meals, at the largest self-identified evening meal (e.g., dinner) or past certain clock times (e.g., % kcal consumed after 5:00 pm or 8:00 pm). In studies of eating regularity, the standard deviation of multiple eating timing metrics has been used to assess day-to-day variability [32], while weekday–weekend differences have been used to capture eating jetlag [21]. Collectively, these variables measure distinct chrononutrition concepts, limiting the ability to draw definitive conclusions and make public health recommendations.

Another important aspect to consider is the methodologies used to assess temporal eating patterns. First, most population-based cohort studies use food frequency questionnaires that lack time-stamped diet data, making the chrononutrition dimension of diet difficult to integrate into observational studies of diet and CMH. Second, most studies rely on self-report, which limits the number of days of diet assessment due to participant burden considerations and introduces measurement error. Studies typically rely on a limited number of 24 h recalls, often collected on non-consecutive days or at a single time point, which are not necessarily representative of habitual intakes. Low-cost, validated diet assessment methods with acceptable participant burden are needed to implement chrononutrition metrics measurement in population studies, enable cross-study comparisons, and support the meaningful translation of research on temporal eating patterns into interventions and public health guidelines aimed at promoting optimal CMH.

Although there is promising emerging evidence from intervention studies, which allow the establishment of causality, regarding the CMH benefits of early and regular eating schedules, modest sample sizes and short follow-up durations are notable limitations. Further, the recruitment of racially and ethnically diverse samples would enhance the generalizability of findings. These methodological limitations, coupled with the inconsistent findings across the heterogeneous intervention protocols and the limited data on underlying physiological mechanisms, are barriers to making evidence-based public health guidelines related to eating timing and regularity.

7. Research Priorities and Future Research Directions

To advance this field, longitudinal population-based cohort studies of racially and ethnically diverse participants encompassing different life stages are needed to evaluate chrononutrition metrics in relation to healthy longevity. The first step to achieve this is the development of innovative chrononutrition assessment tools that are easily administered in research and public health settings. Ideally, the assessment of temporal eating patterns would span 7–14 consecutive days, but at minimum should capture two weekdays and one weekend day to estimate habitual eating timing and regularity metrics. In the meantime, a promising existing tool is the NIH’s ASA-24, which enables the collection of multiple, automatically coded, self-administered electronic food records and 24 h recalls, as well as sleep data using the novel sleep module [84]. Another convenient tool is the novel “Chrononutrition Questionnaire for General and Shift Work Populations”, a 10–15 min questionnaire for obtaining information on shift work schedules, eating timing, regularity, and frequency, wake and sleep times, and chronotype [85]. These tools enable the investigation of the interplay between eating timing and regularity, multidimensional sleep health, and chronotype, as well as the circadian timing of food intake (i.e., eating timing relative to biological time vs. societal clock time) in relation to CMH, which are all key knowledge gaps in this field. Alternatively, novel tools such as picture-based smartphone applications or wearable technologies that detect eating timing could be developed and validated. In addition, a scientific consensus framework for chrononutrition metrics, akin to the multidimensional sleep health framework [86], should be developed to adequately and holistically capture the impact of dietary rhythms on health. Such a framework could encompass eating duration, eating timing, and eating regularity metrics.

Notably, most observational and intervention studies focus on obesity risk and weight loss as outcomes [60]; examining other aspects of CMH as outcomes and elucidating underlying physiological mechanisms and biological pathways is another important research direction. In addition, evaluating temporal eating patterns in relation to diurnal patterns of cardiometabolic markers (e.g., incorporating ambulatory BP monitoring and continuous glucose monitoring) may elucidate novel targets for improving CMH using chrononutrition approaches. Importantly, siloed approaches to research on sleep, circadian rhythms, and diet should be avoided. Future observational studies should evaluate the interplay of chrononutrition metrics, sleep and circadian health, and chronotype in CMH. On the other hand, future intervention studies should consider tailoring eating schedules to chronotype and addressing chrononutrition metrics as part of comprehensive multidimensional dietary or circadian health interventions and alongside sleep interventions for optimizing health impact. Insights from behavioral, dissemination, and implementation sciences should be leveraged to design feasible interventions that work in real-world settings and promote adherence. This includes research to understand how wearables, smartphone apps, daily reminders or nudges, educational content, social support, and working with a health professional or peer health educator could enhance participation, engagement, and retention in such studies.

Given that behavioral and endogenous circadian rhythms vary across life stages and the robustness of circadian rhythms tends to dampen with aging [87,88], adopting a life course approach in future research is crucial to understanding how eating timing and regularity impact CMH preservation, cardiovascular resilience, and healthy longevity. Finally, little is known about the social determinants of health in relation to chrononutrition metrics and how temporal eating patterns contribute to CMH inequities [89]. Chrononutrition metrics are intertwined with sleep and circadian health, which are known to be significant contributors to racial and ethnic disparities in CMH [90–92], suggesting that chrononutrition approaches may represent a novel and timely opportunity to promote CMH equity. The research priorities necessary to advance this field and facilitate the translation of chrononutrition research to clinical practice and public health guidelines are summarized in Figure 3.




Figure 3. Summary of research priorities on eating timing and regularity and cardiometabolic health.

8. Implications for Clinical Practice and Public Health Guidelines and Policy

Upon addressing the aforementioned research gaps, the field of chrononutrition holds great promise for advancing health equity and improving CMH at the individual and population level. At the individual level, personalized chrononutrition interventions addressing the circadian timing of food intake, i.e., eating timing relative to an individual's biological clock, chronotype, and circadian rhythm, could improve CMH. In clinical and research settings, this approach would involve prescribing a fixed eating schedule customized based on dim light melatonin onset or relative to sleep timing, as a less expensive, more feasible proxy. Importantly, such interventions should be contextual, and multiple levels of influence should be addressed to maximize health benefits. For example, a chrononutrition intervention that also addresses food insecurity would be necessary to ensure consistent mealtimes. The integration of chrononutrition interventions with other lifestyle approaches, such as interventions aimed at improving diet quality or promoting sleep health, could also maximize the health benefit of such interventions.

At the population level, the integration of the concept of "timing" into public health guidelines would enhance the effectiveness of lifestyle approaches for improving health. Diet guidelines could encourage keeping a consistent eating start time and eating end time each day and recommend that the timing of the last meal occurs before a certain clock time or within a specific interval prior to bedtime. However, additional questions need to be addressed before such quantitative public health recommendations can be made. For example, by how many hours or minutes can eating timing vary day to day before we start observing adverse health effects? Is there a clock time after which eating has adverse health effects for the majority of the population? When should the last eating occasion occur relative to bedtime for optimal health? Similar to individual-level interventions, community-level and population-level chrononutrition interventions should be contextual, addressing social determinants of health and other domains of influence (Figure 1). Implementation science frameworks should be leveraged and advanced to develop culturally responsive, sustainable, and scalable chrononutrition interventions that promote health equity [93].

9. Conclusions

Observational and intervention studies suggest that earlier and more regular eating timing patterns may promote CMH, but many knowledge gaps remain to be addressed before these aspects of chrononutrition can be formally integrated into public health guidelines and approaches for chronic disease prevention (Table 1). Currently, it is known that an unhealthy diet is the leading cause of death in the US, particularly from cardiovascular and

cardiometabolic disease, but these estimates are based on solely on what (not when) people eat [94]. Elucidating the impact of chrononutrition dimensions on health will be key for quantifying the true impact of the multidimensional diet on the chronic disease burden and unlocking the full potential of dietary approaches for addressing persistent CMH inequities by designing and implementing flexible and feasible interventions in real-world settings.

Table 1. Summary of key research findings and future research directions.

Chrononutrition Metrics	Key Findings	Research Limitations and Future Directions
Breakfast Consumption	Breakfast skipping generally associated with higher risk of coronary heart disease, type 2 diabetes (higher glucose, insulin, and insulin resistance), hypertension, obesity (higher BMI and central adiposity), and hypercholesterolemia (higher total and LDL cholesterol levels and decreased HDL-cholesterol levels) [59–63]. Mixed evidence on weight, ranging from no impact of breakfast skipping on weight [55,72,73] to breakfast skipping leading to weight gain [75,76] or loss [78]. Breakfast skipping may have an adverse effect on glycemic markers and insulin resistance [64,74,78].	<p>Observational data limitations:</p> <ul style="list-style-type: none"> • Cross-sectional design of most studies. • Limited number of studies per chrononutrition metric. • Reliance on few non-consecutive or single 24 h recalls to assess habitual eating timing patterns. • Heterogenous definitions of eating timing and regularity metrics. • Residual confounding. • Lack of racially and ethnically diverse cohort studies encompassing participants from different life stages. <p>RCT data limitations:</p> <ul style="list-style-type: none"> • Small sample sizes and homogenous study populations. • Limited generalizability. • Short follow-up duration. • Focus on weight loss as an outcome. <p>Future Research Directions:</p> <ul style="list-style-type: none"> • Development of innovative dietary assessment tools that are validated, low cost, and readily administered. • Evaluating eating timing and regularity in relation to novel outcomes (e.g., glycemic variability and BP diurnal pattern). • Identifying modifiable multi-level determinants of temporal eating patterns (e.g., environmental, sociocultural, and psychosocial factors) and designing multi-level contextual chrononutrition interventions for healthy longevity. • Investigating the interplay between eating timing and regularity, multidimensional sleep health, and chronotype in cardiometabolic health preservation. • Understanding the association of circadian timing of food intake, i.e., eating timing relative to biological timing (captured from dim light melatonin onset or using sleep timing as a practical proxy), in relation to cardiometabolic outcomes. • Investigating the role of chrononutrition in achieving cardiometabolic health equity. • Elucidating novel underlying physiological mechanisms.
Timing of Breakfast or First Eating Occasion	Morning eating is associated with lower odds of metabolic syndrome, while later timing of first eating occasion is linked to poorer overall cardiovascular health, central adiposity, higher BP, and poorer glycemic control [8,12,65,79]. Earlier timing of meals is associated with lower CVD risk, particularly among women [65,67,80,81].	
Timing of Lunch	Earlier timing of lunch linked to greater weight loss, better glycemic control, and lower insulin resistance [17,82].	
Timing of Dinner or Last Eating Occasion and Extent of Evening Caloric Intake	Later eating timing and greater caloric intake in the evening linked to higher cardiovascular disease risk, metabolic syndrome, systemic inflammation, poorer glycemic control, and higher BP [59,67,71,83].	
Eating Regularity and Eating Jetlag	Increased day-to-day variability and weekday–weekend differences in timing of the first and last eating occasion, span of the daily eating period, and evening caloric intake is associated with higher BMI, waist circumference, BP, HbA1c, and CRP [21,22,32].	

Author Contributions: Conceptualization, N.M.; methodology, O.E.R., E.B.K., D.D.S., M.-P.S.-O. and N.M.; investigation, O.E.R., E.B.K., D.D.S., M.-P.S.-O. and N.M.; resources, N.M.; data curation, O.E.R.; writing—original draft preparation, O.E.R., E.B.K. and N.M.; writing—review and editing, N.M., D.D.S. and M.-P.S.-O.; visualization, O.E.R. and E.B.K.; supervision, N.M.; project administration, N.M.; funding acquisition, N.M. All authors have read and agreed to the published version of the manuscript.

Funding: This research and the APC was funded by NHLBI Grant # R00-HL148511, American Heart Association Grant #855050, New York Regional CDTR Pilot and Feasibility Grant (DK111022), and NIMHD Grant # P50MD017341 (sub-project ID: 8126). The funders had no role in the design and conduct of the study, in the collection, analysis, and interpretation of the data, and in the preparation, review, or approval of the manuscript.

Acknowledgments: Figure 2 was created with the use of Biorender.com.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Kessler, K.; Pivovarov-Ramich, O. Meal Timing, Aging, and Metabolic Health. *Int. J. Mol. Sci.* **2019**, *20*, 1911. [[CrossRef](#)] [[PubMed](#)]
2. Flanagan, A.; Bechtold, D.A.; Pot, G.K.; Johnston, J.D. Chrono-nutrition: From Molecular and Neuronal Mechanisms to Human Epidemiology and Timed Feeding Patterns. *J. Neurochem.* **2021**, *157*, 53–72. [[CrossRef](#)] [[PubMed](#)]
3. Katsi, V.; Papakonstantinou, I.P.; Soulaïdopoulos, S.; Katsiki, N.; Tsioufīs, K. Chrononutrition in Cardiometabolic Health. *J. Clin. Med.* **2022**, *11*, 296. [[CrossRef](#)] [[PubMed](#)]
4. Almoosawi, S.; Vingeliene, S.; Gachon, F.; Voortman, T.; Palla, L.; Johnston, J.D.; Van Dam, R.M.; Darimont, C.; Karagounis, L.G. Chronotype: Implications for Epidemiologic Studies on Chrono-Nutrition and Cardiometabolic Health. *Adv. Nutr.* **2019**, *10*, 30–42. [[CrossRef](#)] [[PubMed](#)]
5. Kahleova, H.; Levin, S.; Barnard, N.D. Vegetarian Dietary Patterns and Cardiovascular Disease. *Prog. Cardiovasc. Dis.* **2018**, *61*, 54–61. [[CrossRef](#)] [[PubMed](#)]
6. Lloyd-Jones, D.M.; Allen, N.B.; Anderson, C.A.M.; Black, T.; Brewer, L.C.; Foraker, R.E.; Grandner, M.A.; Lavretsky, H.; Perak, A.M.; Sharma, G.; et al. Life’s Essential 8: Updating and Enhancing the American Heart Association’s Construct of Cardiovascular Health: A Presidential Advisory From the American Heart Association. *Circulation* **2022**, *146*, 5. [[CrossRef](#)] [[PubMed](#)]
7. Schuppeli, B.; Peters, B.; Ottawa, A.; Pivovarov-Ramich, O. Time Restricted Eating: A Dietary Strategy to Prevent and Treat Metabolic Disturbances. *Front. Endocrinol.* **2021**, *12*, 683140. [[CrossRef](#)] [[PubMed](#)]
8. Liu, J.; Yi, P.; Liu, F. The Effect of Early Time-Restricted Eating vs Later Time-Restricted Eating on Weight Loss and Metabolic Health. *J. Clin. Endocrinol. Metab.* **2023**, *108*, 1824–1834. [[CrossRef](#)] [[PubMed](#)]
9. Cho, S.; Dietrich, M.; Brown, C.J.P.; Clark, C.A.; Block, G. The Effect of Breakfast Type on Total Daily Energy Intake and Body Mass Index: Results from the Third National Health and Nutrition Examination Survey (NHANES III). *J. Am. Coll. Nutr.* **2003**, *22*, 296–302. [[CrossRef](#)]
10. Kant, A.K.; Graubard, B.I. 40-Year Trends in Meal and Snack Eating Behaviors of American Adults. *J. Acad. Nutr. Diet.* **2015**, *115*, 50–63. [[CrossRef](#)]
11. Kutsuma, A.; Nakajima, K.; Suwa, K. Potential Association between Breakfast Skipping and Concomitant Late-Night-Dinner Eating with Metabolic Syndrome and Proteinuria in the Japanese Population. *Scientifica* **2014**, *2014*, 253581. [[CrossRef](#)] [[PubMed](#)]
12. Ha, K.; Song, Y. Associations of Meal Timing and Frequency with Obesity and Metabolic Syndrome among Korean Adults. *Nutrients* **2019**, *11*, 2437. [[CrossRef](#)] [[PubMed](#)]
13. Jung, J.; Kim, A.-S.; Ko, H.-J.; Choi, H.-I.; Hong, H.-E. Association between Breakfast Skipping and the Metabolic Syndrome: The Korea National Health and Nutrition Examination Survey, 2017. *Medicina* **2020**, *56*, 396. [[CrossRef](#)] [[PubMed](#)]
14. Barragán, R.; Fernández-Carrión, R.; Asensio-Márquez, E.M.; Ortega-Azorín, C.; Álvarez-Sala, A.; Pérez-Fidalgo, A.; Sorlí, J.V.; Portolés, O.; González-Monje, I.; St-Onge, M.P.; et al. Timing of Meals and Sleep in the Mediterranean Population: The Effect of Taste, Genetics, Environmental Determinants, and Interactions on Obesity Phenotypes. *Nutrients* **2023**, *15*, 708. [[CrossRef](#)] [[PubMed](#)]
15. Kant, A.K. Eating Patterns of US Adults: Meals, Snacks, and Time of Eating. *Physiol. Behav.* **2018**, *193*, 270–278. [[CrossRef](#)] [[PubMed](#)]
16. Huseinovic, E.; Winkvist, A.; Slimani, N.; Park, M.; Freisling, H.; Boeing, H.; Buckland, G.; Schwingshackl, L.; Weiderpass, E.; Rostgaard-Hansen, A.; et al. Meal Patterns across Ten European Countries—Results from the European Prospective Investigation into Cancer and Nutrition (EPIC) Calibration Study. *Public Health Nutr.* **2016**, *19*, 2769–2780. [[CrossRef](#)] [[PubMed](#)]
17. Garaulet, M.; Gómez-Abellán, P.; Albuquerque-Béjar, J.J.; Lee, Y.-C.; Ordovás, J.M.; Scheer, F.A.J.L. Timing of Food Intake Predicts Weight Loss Effectiveness. *Int. J. Obes.* **2013**, *37*, 604–611. [[CrossRef](#)] [[PubMed](#)]
18. Reid, K.J.; Baron, K.G.; Zee, P.C. Meal Timing Influences Daily Caloric Intake in Healthy Adults. *Nutr. Res.* **2014**, *34*, 930–935. [[CrossRef](#)]
19. Gill, S.; Panda, S. A Smartphone App Reveals Erratic Diurnal Eating Patterns in Humans That Can Be Modulated for Health Benefits. *Cell Metab.* **2015**, *22*, 789–798. [[CrossRef](#)]
20. Striegel-Moore, R.H.; Franko, D.L.; Thompson, D.; Affenito, S.; Kraemer, H.C. Night Eating: Prevalence and Demographic Correlates. *Obesity* **2006**, *14*, 139–147. [[CrossRef](#)]
21. Zerón-Rugiero, M.; Hernández, Á.; Porras-Loaiza, A.; Cambras, T.; Izquierdo-Pulido, M. Eating Jet Lag: A Marker of the Variability in Meal Timing and Its Association with Body Mass Index. *Nutrients* **2019**, *11*, 2980. [[CrossRef](#)]

22. Makarem, N.; Sears, D.D.; St-Onge, M.; Zuraikat, F.M.; Gallo, L.C.; Talavera, G.A.; Castaneda, S.F.; Lai, Y.; Aggarwal, B. Variability in Daily Eating Patterns and Eating Jetlag Are Associated with Worsened Cardiometabolic Risk Profiles in the American Heart Association Go Red for Women Strategically Focused Research Network. *J. Am. Heart Assoc.* **2021**, *10*, e022024. [[CrossRef](#)]
23. Dashti, H.S.; Scheer, F.A.J.L.; Saxena, R.; Garaulet, M. Timing of Food Intake: Identifying Contributing Factors to Design Effective Interventions. *Adv. Nutr.* **2019**, *10*, 606–620. [[CrossRef](#)] [[PubMed](#)]
24. Drewnowski, A.; Rehm, C.D.; Vieux, F. Breakfast in the United States: Food and Nutrient Intakes in Relation to Diet Quality in National Health and Examination Survey 2011–2014. A Study from the International Breakfast Research Initiative. *Nutrients* **2018**, *10*, 1200. [[CrossRef](#)]
25. Johnson, D.A.; Cheng, P.; FarrHenderson, M.; Knutson, K. Understanding the Determinants of Circadian Health Disparities and Cardiovascular Disease. *Chronobiol. Int.* **2023**, *40*, 83–90. [[CrossRef](#)]
26. Ansu Baidoo, V.Y.; Zee, P.C.; Knutson, K.L. Racial and Ethnic Differences in Eating Duration and Meal Timing: Findings from NHANES 2011–2018. *Nutrients* **2022**, *14*, 2428. [[CrossRef](#)]
27. Schuler, B.R.; Bauer, K.W.; Lumeng, J.C.; Rosenblum, K.; Clark, M.; Miller, A.L. Poverty and Food Insecurity Predict Mealtime Structure: Mediating Pathways of Parent Disciplinary Practices and Depressive Symptoms. *J. Child Fam. Stud.* **2020**, *29*, 3169–3183. [[CrossRef](#)] [[PubMed](#)]
28. Wang, P.; Tan, Q.; Zhao, Y.; Zhao, J.; Zhang, Y.; Shi, D. Night Eating in Timing, Frequency, and Food Quality and Risks of All-Cause, Cancer, and Diabetes Mortality: Findings from National Health and Nutrition Examination Survey. *Nutr. Diabetes* **2024**, *14*, 5. [[CrossRef](#)]
29. Lesani, A.; Barkhidarian, B.; Jafarzadeh, M.; Akbarzade, Z.; Djafarian, K.; Shab-Bidar, S. Time-Related Meal Patterns and Breakfast Quality in a Sample of Iranian Adults. *BMC Nutr.* **2023**, *9*, 13. [[CrossRef](#)] [[PubMed](#)]
30. Vera, B.; Dashti, H.S.; Gómez-Abellán, P.; Hernández-Martínez, A.M.; Esteban, A.; Scheer, F.A.J.L.; Saxena, R.; Garaulet, M. Modifiable Lifestyle Behaviors, but Not a Genetic Risk Score, Associate with Metabolic Syndrome in Evening Chronotypes. *Sci. Rep.* **2018**, *8*, 945. [[CrossRef](#)]
31. Garaulet, M.; Madrid, J.A. Chronobiological Aspects of Nutrition, Metabolic Syndrome and Obesity. *Adv. Drug Deliv. Rev.* **2010**, *62*, 967–978. [[CrossRef](#)] [[PubMed](#)]
32. Makarem, N.; Zuraikat, F.M.; Caceres, B.; Sears, D.D.; St-Onge, M.-P.; Lai, Y.; Aggarwal, B. Variable Eating Patterns: A Potential Novel Risk Factor for Systemic Inflammation in Women. *Ann. Behav. Med.* **2023**, *57*, 93–97. [[CrossRef](#)] [[PubMed](#)]
33. Poggiogalle, E.; Jamshed, H.; Peterson, C.M. Circadian Regulation of Glucose, Lipid, and Energy Metabolism in Humans. *Metabolism* **2018**, *84*, 11–27. [[CrossRef](#)] [[PubMed](#)]
34. Ruddick-Collins, L.C.; Morgan, P.J.; Johnstone, A.M. Mealtime: A Circadian Disruptor and Determinant of Energy Balance? *J. Neuroendocrinol.* **2020**, *32*, e12886. [[CrossRef](#)] [[PubMed](#)]
35. Scheer, F.A.J.L.; Hilton, M.F.; Mantzoros, C.S.; Shea, S.A. Adverse Metabolic and Cardiovascular Consequences of Circadian Misalignment. *Proc. Natl. Acad. Sci. USA* **2009**, *106*, 4453–4458. [[CrossRef](#)] [[PubMed](#)]
36. Garaulet, M.; Lopez-Minguez, J.; Dashti, H.S.; Vetter, C.; Hernández-Martínez, A.M.; Pérez-Ayala, M.; Baraza, J.C.; Wang, W.; Florez, J.C.; Scheer, F.A.J.L.; et al. Interplay of Dinner Timing and *MTNR1B* Type 2 Diabetes Risk Variant on Glucose Tolerance and Insulin Secretion: A Randomized Crossover Trial. *Diabetes Care* **2022**, *45*, 512–519. [[CrossRef](#)] [[PubMed](#)]
37. Zhang, D.; Colson, J.C.; Jin, C.; Becker, B.K.; Rhoads, M.K.; Pati, P.; Neder, T.H.; King, M.A.; Valcin, J.A.; Tao, B.; et al. Timing of Food Intake Drives the Circadian Rhythm of Blood Pressure. *Function* **2020**, *2*, zqaa034. [[CrossRef](#)] [[PubMed](#)]
38. Morris, C.J.; Purvis, T.E.; Hu, K.; Scheer, F.A.J.L. Circadian Misalignment Increases Cardiovascular Disease Risk Factors in Humans. *Proc. Natl. Acad. Sci. USA* **2016**, *113*, E1402–E1411. [[CrossRef](#)] [[PubMed](#)]
39. Marinac, C.R.; Sears, D.D.; Natarajan, L.; Gallo, L.C.; Breen, C.I.; Patterson, R.E. Frequency and Circadian Timing of Eating May Influence Biomarkers of Inflammation and Insulin Resistance Associated with Breast Cancer Risk. *PLoS ONE* **2015**, *10*, e0136240. [[CrossRef](#)]
40. Vujović, N.; Piron, M.J.; Qian, J.; Chellappa, S.L.; Nedeltcheva, A.; Barr, D.; Heng, S.W.; Kerlin, K.; Srivastav, S.; Wang, W.; et al. Late Isocaloric Eating Increases Hunger, Decreases Energy Expenditure, and Modifies Metabolic Pathways in Adults with Overweight and Obesity. *Cell Metab.* **2022**, *34*, 1486–1498.e7. [[CrossRef](#)]
41. Garaulet, M.; Gómez-Abellán, P. Timing of Food Intake and Obesity: A Novel Association. *Physiol. Behav.* **2014**, *134*, 44–50. [[CrossRef](#)] [[PubMed](#)]
42. Kaczmarek, J.L.; MUSAAD, S.M.; Holscher, H.D. Time of Day and Eating Behaviors Are Associated with the Composition and Function of the Human Gastrointestinal Microbiota. *Am. J. Clin. Nutr.* **2017**, *106*, 1220–1231. [[CrossRef](#)] [[PubMed](#)]
43. Jakubowicz, D.; Wainstein, J.; Landau, Z.; Raz, I.; Ahren, B.; Chapnik, N.; Ganz, T.; Menaged, M.; Barnea, M.; Bar-Dayan, Y.; et al. Influences of Breakfast on Clock Gene Expression and Postprandial Glycemia in Healthy Individuals and Individuals with Diabetes: A Randomized Clinical Trial. *Diabetes Care* **2017**, *40*, 1573–1579. [[CrossRef](#)] [[PubMed](#)]
44. Collado, M.C.; Engen, P.A.; Bandín, C.; Cabrera-Rubio, R.; Voigt, R.M.; Green, S.J.; Naqib, A.; Keshavarzian, A.; Scheer, F.A.J.L.; Garaulet, M. Timing of Food Intake Impacts Daily Rhythms of Human Salivary Microbiota: A Randomized, Crossover Study. *FASEB J.* **2018**, *32*, 2060–2072. [[CrossRef](#)] [[PubMed](#)]
45. Garaulet, M.; Corbalán-Tutau, M.D.; Madrid, J.A.; Baraza, J.C.; Parnell, L.D.; Lee, Y.-C.; Ordovas, J.M. PERIOD2 Variants Are Associated with Abdominal Obesity, Psycho-Behavioral Factors, and Attrition in the Dietary Treatment of Obesity. *J. Am. Diet. Assoc.* **2010**, *110*, 917–921. [[CrossRef](#)] [[PubMed](#)]

46. Lopez-Minguez, J.; Dashti, H.S.; Madrid-Valero, J.J.; Madrid, J.A.; Saxena, R.; Scheer, F.A.J.L.; Ordoñana, J.R.; Garaulet, M. Heritability of the Timing of Food Intake. *Clin. Nutr.* **2019**, *38*, 767–773. [[CrossRef](#)] [[PubMed](#)]
47. De Castro, J.M. Heritability of Diurnal Changes in Food Intake in Free-Living Humans. *Nutrition* **2001**, *17*, 713–720. [[CrossRef](#)]
48. Wilson, J.E.; Blizzard, L.; Gall, S.L.; Magnussen, C.G.; Oddy, W.H.; Dwyer, T.; Sanderson, K.; Venn, A.J.; Smith, K.J. An Eating Pattern Characterised by Skipped or Delayed Breakfast Is Associated with Mood Disorders among an Australian Adult Cohort. *Psychol. Med.* **2020**, *50*, 2711–2721. [[CrossRef](#)] [[PubMed](#)]
49. Tahara, Y.; Makino, S.; Suiko, T.; Nagamori, Y.; Iwai, T.; Aono, M.; Shibata, S. Association between Irregular Meal Timing and the Mental Health of Japanese Workers. *Nutrients* **2021**, *13*, 2775. [[CrossRef](#)]
50. Chang, Z.-S.; Boolani, A.; Conroy, D.A.; Dunietz, T.; Jansen, E.C. Skipping Breakfast and Mood: The Role of Sleep. *Nutr. Health* **2021**, *27*, 373–379. [[CrossRef](#)]
51. Walker, W.H.; Walton, J.C.; DeVries, A.C.; Nelson, R.J. Circadian Rhythm Disruption and Mental Health. *Transl. Psychiatry* **2020**, *10*, 28. [[CrossRef](#)]
52. Qian, J.; Vujovic, N.; Nguyen, H.; Rahman, N.; Heng, S.W.; Amira, S.; Scheer, F.A.J.L.; Chellappa, S.L. Daytime Eating Prevents Mood Vulnerability in Night Work. *Proc. Natl. Acad. Sci. USA* **2022**, *119*, e2206348119. [[CrossRef](#)]
53. Pot, G.K. Sleep and Dietary Habits in the Urban Environment: The Role of Chrono-Nutrition. *Proc. Nutr. Soc.* **2018**, *77*, 189–198. [[CrossRef](#)] [[PubMed](#)]
54. Garrido, A.L.F.; Duarte, A.D.S.; Santana, P.T.; Rodrigues, G.H.; Pellegrino, P.; Nogueira, L.F.R.; Cipolla-Neto, J.; Moreno, C.R.D.C.; Marqueze, E.C. Eating Habits, Sleep, and a Proxy for Circadian Disruption Are Correlated with Dyslipidemia in Overweight Night Workers. *Nutrition* **2021**, *83*, 111084. [[CrossRef](#)]
55. Chowdhury, E.A.; Richardson, J.D.; Holman, G.D.; Tsintzas, K.; Thompson, D.; Betts, J.A. The Causal Role of Breakfast in Energy Balance and Health: A Randomized Controlled Trial in Obese Adults. *Am. J. Clin. Nutr.* **2016**, *103*, 747–756. [[CrossRef](#)] [[PubMed](#)]
56. Roshanmehr, F.; Hayashi, K.; Tahara, Y.; Suiko, T.; Nagamori, Y.; Iwai, T.; Shibata, S. Association between Breakfast Meal Categories and Timing of Physical Activity of Japanese Workers. *Foods* **2022**, *11*, 2609. [[CrossRef](#)]
57. Katagiri, R.; Asakura, K.; Kobayashi, S.; Suga, H.; Sasaki, S.; the Three-generation Study of Women on Diets and Health Study Group. Low Intake of Vegetables, High Intake of Confectionary, and Unhealthy Eating Habits Are Associated with Poor Sleep Quality among Middle-aged Female Japanese Workers. *J. Occup. Health* **2014**, *56*, 359–368. [[CrossRef](#)] [[PubMed](#)]
58. Hernandez, E.; Kim, M.; Kim, W.G.; Yoon, J. Nutritional Aspects of Night Eating and Its Association with Weight Status among Korean Adolescents. *Nutr. Res. Pract.* **2016**, *10*, 448. [[CrossRef](#)] [[PubMed](#)]
59. Cahill, L.E.; Chiuve, S.E.; Mekary, R.A.; Jensen, M.K.; Flint, A.J.; Hu, F.B.; Rimm, E.B. Prospective Study of Breakfast Eating and Incident Coronary Heart Disease in a Cohort of Male US Health Professionals. *Circulation* **2013**, *128*, 337–343. [[CrossRef](#)]
60. Paoli, A.; Tinsley, G.; Bianco, A.; Moro, T. The Influence of Meal Frequency and Timing on Health in Humans: The Role of Fasting. *Nutrients* **2019**, *11*, 719. [[CrossRef](#)]
61. Deshmukh-Taskar, P.; Nicklas, T.A.; Radcliffe, J.D.; O’Neil, C.E.; Liu, Y. The Relationship of Breakfast Skipping and Type of Breakfast Consumed with Overweight/Obesity, Abdominal Obesity, Other Cardiometabolic Risk Factors and the Metabolic Syndrome in Young Adults. The National Health and Nutrition Examination Survey (NHANES): 1999–2006. *Public Health Nutr.* **2013**, *16*, 2073–2082. [[CrossRef](#)] [[PubMed](#)]
62. Witbracht, M.; Keim, N.L.; Forester, S.; Widaman, A.; Laugero, K. Female Breakfast Skippers Display a Disrupted Cortisol Rhythm and Elevated Blood Pressure. *Physiol. Behav.* **2015**, *140*, 215–221. [[CrossRef](#)] [[PubMed](#)]
63. Kahleova, H.; Lloren, J.I.; Mashchak, A.; Hill, M.; Fraser, G.E. Meal Frequency and Timing Are Associated with Changes in Body Mass Index in Adventist Health Study 2. *J. Nutr.* **2017**, *147*, 1722–1728. [[CrossRef](#)] [[PubMed](#)]
64. Min, C.; Noh, H.; Kang, Y.-S.; Sim, H.J.; Baik, H.W.; Song, W.O.; Yoon, J.; Park, Y.-H.; Joung, H. Skipping Breakfast Is Associated with Diet Quality and Metabolic Syndrome Risk Factors of Adults. *Nutr. Res. Pract.* **2011**, *5*, 455. [[CrossRef](#)] [[PubMed](#)]
65. Makarem, N.; Sears, D.D.; St-Onge, M.-P.; Zuraikat, F.M.; Gallo, L.C.; Talavera, G.A.; Castaneda, S.F.; Lai, Y.; Mi, J.; Aggarwal, B. Habitual Nightly Fasting Duration, Eating Timing, and Eating Frequency Are Associated with Cardiometabolic Risk in Women. *Nutrients* **2020**, *12*, 3043. [[CrossRef](#)] [[PubMed](#)]
66. Carew, A.S.; Mekary, R.A.; Kirkland, S.; Theou, O.; Siddiqi, F.; Urquhart, R.; George, M.; Blanchard, C.; Biggs, M.L.; Djoussé, L.; et al. Prospective Study of Breakfast Frequency and Timing and the Risk of Incident Type 2 Diabetes in Community-Dwelling Older Adults: The Cardiovascular Health Study. *Am. J. Clin. Nutr.* **2022**, *116*, 325–334. [[CrossRef](#)] [[PubMed](#)]
67. Palomar-Cros, A.; Andreeva, V.A.; Fezeu, L.K.; Julia, C.; Bellicha, A.; Kesse-Guyot, E.; Hercberg, S.; Romaguera, D.; Kogevinas, M.; Touvier, M.; et al. Dietary Circadian Rhythms and Cardiovascular Disease Risk in the Prospective NutriNet-Santé Cohort. *Nat. Commun.* **2023**, *14*, 7899. [[CrossRef](#)] [[PubMed](#)]
68. Van Cauter, E.; Leproult, R.; Kupfer, D.J. Effects of Gender and Age on the Levels and Circadian Rhythmicity of Plasma Cortisol. *J. Clin. Endocrinol. Metab.* **1996**, *81*, 2468–2473. [[CrossRef](#)] [[PubMed](#)]
69. Bailey, M.; Silver, R. Sex Differences in Circadian Timing Systems: Implications for Disease. *Front. Neuroendocrinol.* **2014**, *35*, 111–139. [[CrossRef](#)] [[PubMed](#)]
70. Nicolaides, N.C.; Chrousos, G.P. Sex Differences in Circadian Endocrine Rhythms: Clinical Implications. *Eur. J. Neurosci.* **2020**, *52*, 2575–2585. [[CrossRef](#)]
71. Xiao, Q.; Garaulet, M.; Scheer, F.A.J.L. Meal Timing and Obesity: Interactions with Macronutrient Intake and Chronotype. *Int. J. Obes.* **2019**, *43*, 1701–1711. [[CrossRef](#)]

72. Dhurandhar, E.J.; Dawson, J.; Alcorn, A.; Larsen, L.H.; Thomas, E.A.; Cardel, M.; Bourland, A.C.; Astrup, A.; St-Onge, M.-P.; Hill, J.O.; et al. The Effectiveness of Breakfast Recommendations on Weight Loss: A Randomized Controlled Trial. *Am. J. Clin. Nutr.* **2014**, *100*, 507–513. [[CrossRef](#)]
73. Betts, J.A.; Richardson, J.D.; Chowdhury, E.A.; Holman, G.D.; Tsintzas, K.; Thompson, D. The Causal Role of Breakfast in Energy Balance and Health: A Randomized Controlled Trial in Lean Adults. *Am. J. Clin. Nutr.* **2014**, *100*, 539–547. [[CrossRef](#)]
74. Nas, A.; Mirza, N.; Hägele, F.; Kahlhöfer, J.; Keller, J.; Rising, R.; Kufer, T.A.; Bosy-Westphal, A. Impact of Breakfast Skipping Compared with Dinner Skipping on Regulation of Energy Balance and Metabolic Risk. *Am. J. Clin. Nutr.* **2017**, *105*, 1351–1361. [[CrossRef](#)]
75. Ogata, H.; Kayaba, M.; Tanaka, Y.; Yajima, K.; Iwayama, K.; Ando, A.; Park, I.; Kiyono, K.; Omi, N.; Satoh, M.; et al. Effect of Skipping Breakfast for 6 Days on Energy Metabolism and Diurnal Rhythm of Blood Glucose in Young Healthy Japanese Males. *Am. J. Clin. Nutr.* **2019**, *110*, 41–52. [[CrossRef](#)]
76. Geliebter, A.; Astbury, N.M.; Aviram-Friedman, R.; Yahav, E.; Hashim, S. Skipping Breakfast Leads to Weight Loss but Also Elevated Cholesterol Compared with Consuming Daily Breakfasts of Oat Porridge or Frosted Cornflakes in Overweight Individuals: A Randomised Controlled Trial. *J. Nutr. Sci.* **2014**, *3*, e56. [[CrossRef](#)] [[PubMed](#)]
77. Farshchi, H.R.; Taylor, M.A.; Macdonald, I.A. Deleterious Effects of Omitting Breakfast on Insulin Sensitivity and Fasting Lipid Profiles in Healthy Lean Women 1–3. *Am. J. Clin. Nutr.* **2005**, *81*, 388–396. [[CrossRef](#)] [[PubMed](#)]
78. Jakubowicz, D.; Wainstein, J.; Ahren, B.; Landau, Z.; Bar-Dayana, Y.; Froy, O. Fasting Until Noon Triggers Increased Postprandial Hyperglycemia and Impaired Insulin Response After Lunch and Dinner in Individuals with Type 2 Diabetes: A Randomized Clinical Trial. *Diabetes Care* **2015**, *38*, 1820–1826. [[CrossRef](#)] [[PubMed](#)]
79. Young, I.E.; Poobalan, A.; Steinbeck, K.; O'Connor, H.T.; Parker, H.M. Distribution of Energy Intake across the Day and Weight Loss: A Systematic Review and Meta-analysis. *Obes. Rev.* **2023**, *24*, e13537. [[CrossRef](#)]
80. Jakubowicz, D.; Barnea, M.; Wainstein, J.; Froy, O. High Caloric Intake at Breakfast vs. Dinner Differentially Influences Weight Loss of Overweight and Obese Women: Effect of High-Calorie Breakfast vs. Dinner. *Obesity* **2013**, *21*, 2504–2512. [[CrossRef](#)]
81. Jakubowicz, D.; Barnea, M.; Wainstein, J.; Froy, O. Effects of Caloric Intake Timing on Insulin Resistance and Hyperandrogenism in Lean Women with Polycystic Ovary Syndrome. *Clin. Sci.* **2013**, *125*, 423–432. [[CrossRef](#)] [[PubMed](#)]
82. Bandín, C.; Scheer, F.A.J.L.; Luque, A.J.; Ávila-Gandía, V.; Zamora, S.; Madrid, J.A.; Gómez-Abellán, P.; Garaulet, M. Meal Timing Affects Glucose Tolerance, Substrate Oxidation and Circadian-Related Variables: A Randomized, Crossover Trial. *Int. J. Obes.* **2015**, *39*, 828–833. [[CrossRef](#)] [[PubMed](#)]
83. Tsuchida, Y.; Hata, S.; Sone, Y. Effects of a Late Supper on Digestion and the Absorption of Dietary Carbohydrates in the Following Morning. *J. Physiol. Anthropol.* **2013**, *32*, 9. [[CrossRef](#)] [[PubMed](#)]
84. Shams-White, M.M.; O'Connor, L.E.; O'Connor, S.G.; Herrick, K.A. Development of the Sleep Module for the Automated Self-Administered 24-Hour (ASA24) Dietary Assessment Tool: New Research Opportunities. *J. Acad. Nutr. Diet.* **2022**, *122*, 2017–2022. [[CrossRef](#)]
85. Phoi, Y.Y.; Bonham, M.P.; Rogers, M.; Dorrian, J.; Coates, A.M. Content Validation of a Chrononutrition Questionnaire for the General and Shift Work Populations: A Delphi Study. *Nutrients* **2021**, *13*, 4087. [[CrossRef](#)] [[PubMed](#)]
86. Buysse, D.J. Sleep Health: Can We Define It? Does It Matter? *Sleep* **2014**, *37*, 9–17. [[CrossRef](#)] [[PubMed](#)]
87. Mattis, J.; Sehgal, A. Circadian Rhythms, Sleep, and Disorders of Aging. *Trends Endocrinol. Metab.* **2016**, *27*, 192–203. [[CrossRef](#)] [[PubMed](#)]
88. Cornelissen, G.; Otsuka, K. Chronobiology of Aging: A Mini-Review. *Gerontology* **2017**, *63*, 118–128. [[CrossRef](#)] [[PubMed](#)]
89. Dashti, H.S.; Scheer, F.A.; Jacques, P.F.; Lamon-Fava, S.; Ordovás, J.M. Short Sleep Duration and Dietary Intake: Epidemiologic Evidence, Mechanisms, and Health Implications. *Adv. Nutr.* **2015**, *6*, 648–659. [[CrossRef](#)]
90. Kingsbury, J.H.; Buxton, O.M.; Emmons, K.M.; Redline, S. Sleep and Its Relationship to Racial and Ethnic Disparities in Cardiovascular Disease. *Curr. Cardiovasc. Risk Rep.* **2013**, *7*, 387–394. [[CrossRef](#)]
91. Makarem, N.; St-Onge, M.-P.; Liao, M.; Lloyd-Jones, D.M.; Aggarwal, B. Association of Sleep Characteristics with Cardiovascular Health among Women and Differences by Race/Ethnicity and Menopausal Status: Findings from the American Heart Association Go Red for Women Strategically Focused Research Network. *Sleep Health* **2019**, *5*, 501–508. [[CrossRef](#)] [[PubMed](#)]
92. Hale, L.; Troxel, W.; Buysse, D.J. Sleep Health: An Opportunity for Public Health to Address Health Equity. *Annu. Rev. Public Health* **2020**, *41*, 81–99. [[CrossRef](#)] [[PubMed](#)]
93. Tumilowicz, A.; Ruel, M.T.; Peltó, G.; Pelletier, D.; Monterrosa, E.C.; Lapping, K.; Kraemer, K.; De Regil, L.M.; Bergeron, G.; Arabi, M.; et al. Implementation Science in Nutrition: Concepts and Frameworks for an Emerging Field of Science and Practice. *Curr. Dev. Nutr.* **2019**, *3*, nzy080. [[CrossRef](#)] [[PubMed](#)]
94. The US Burden of Disease Collaborators; Mokdad, A.H.; Ballestrós, K.; Echko, M.; Glenn, S.; Olsen, H.E.; Mullany, E.; Lee, A.; Khan, A.R.; Ahmadi, A.; et al. The State of US Health, 1990–2016: Burden of Diseases, Injuries, and Risk Factors Among US States. *JAMA* **2018**, *319*, 1444. [[CrossRef](#)]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.